HISTORY OF MATHEMATICS

VOLUME I

GENERAL SURVEY OF THE HISTORY OF ELEMENTARY MATHEMATICS

BY

DAVID EUGENE SMITH

DOVER PUBLICATIONS, INC.
NEW YORK NEW YORK

© COPYRIGHT, 1951, BY EVA MAY LUSE SMITH
COPYRIGHT, 1923, BY DAVID EUGENE SMITH
ALL RIGHTS RESERVED

This new Dover edition, first published in 1958, is an unaltered and unabridged republication of the last edition. It is published through special arrangement with GINN AND COMPANY.

PRINTED IN THE UNITED STATES OF AMERICA

357.3

PREFACE

This work has been written for the purpose of supplying teachers and students with a usable textbook on the history of elementary mathematics, that is, of mathematics through the first steps in the calculus. The subject has come to be recognized as an important one in the preparation of teachers of mathematics and in the liberal education of students in colleges and high schools. Although several works upon the history of mathematics are already available, the author feels that a book written from somewhat different standpoints will be found helpful to those who are beginning the study of the subject in our universities, colleges, and normal schools, and because of this belief the present work has been prepared.

A history of mathematics may be constructed on several general plans, each of which may be justified by the purpose in mind. For example, it may be arranged solely with a view to the chronological sequence of events, or as a series of biographies, or according to the branches of mathematics, or as a source book of material [illegible], or with respect to national or racial achievements, or in [illegible] other ways, each of which may have certain advantages.

The general plan adopted in the preparation of this work is that of presenting the subject from two distinct standpoints, the first, as in Volume I, leading to a survey of the growth of mathematics by chronological periods, with due consideration to racial achievements; and the second, as in Volume II, leading to a discussion of the evolution of certain important topics. To attempt to fuse these two features and thus to carry them along together has often been attempted. [illegible]haracterizes, for example, the monumental treatise of Montucla and, to a large extent, that of Cantor. For the teacher, however, this plan is not satisfactory, and the excellent work of Tropfke is an example of the tendency to break away from the mere chronological recital of facts. Long experience in teaching the subject in colleges has convinced the author that a general historical presentation is

desirable for the purpose of relating the development of mathematics to the development of the race, of revealing the science as a great stream rather than a static mass, and of emphasizing the human element, but that this ought to lead to a topical presentation by which the student may understand something of the life history of the special subject which he may be studying, whether it be the elementary theory of numbers, the methods of calculation, the solution of equations, the functions of trigonometry, the common symbolism in use, the various types of elementary geometry, the early steps in the calculus, or one of the various other important topics of elementary mathematics. The general plan can best be understood by a glance at the table of contents in each volume.

Perhaps the chief objection to the general arrangement set forth in Volume I is that the reader may occasionally feel that a mere statement of the subjects in which some particular mathematician was interested is not very illuminating, and that a more extended statement of his achievements would have greater significance. In most cases, however, a further elaboration of the record would destroy the possibility of successfully carrying out the plan of showing the growth of the several leading branches of elementary mathematics by themselves, as in Volume II, at least without a large amount of wearisome repetition. Of the two evils the lesser has been chosen.

In Volume I, which forms the general survey by periods, attention has been given to geographical and racial considerations as well as to chronological sequence. While it is evident that no race or country has any monopoly of genius, and while the limits of successive centuries are only artificial boundaries with no significance in the creation of the masterpieces of any science, nevertheless linguistic and racial influences tend to develop tastes in mathematics as they do in art and in letters, and certain centuries stand out with interesting prominence.

The student will therefore find it to his advantage to give some attention to the geographical distribution of scholars as well as to the general periods in which they lived. While it is impossible to grade countries according to any definite scale of excellence, and while the world has always seen more or less of the migration of scholars from one country to another, it is possible in a general way

to give prominent positions to those national groups which have contributed most to the advancement of the science in each period under discussion.

In this treatment of the subject an attempt has been made to seek out the causes of the advance or the retardation of mathematics in different centuries and with different races, but always with the consciousness that the world has no certain prescription for the creation of the genius and that the causes of any series of historical events are usually very intangible. The effort has also been made to introduce enough of the anecdote to relieve the monotony of mere historical statement and to reveal the mathematician as a human being like others of his race.

While the footnote is often condemned as merely an apology for obscurity or as an exhibition of pedantry, it would be difficult, in a work of this kind, to dispense with its aid. There are two principal justifications for such a device: first, it enables an author to place the responsibility for a statement that may be open to question; and second, it encourages many students to undertake further study, either from secondary sources or, what is more important, from the original writings of the men who rank among the creators of mathematics. With these two points in mind, footnotes have been introduced in such a way as to be used by readers who wish for further aid, and to be neglected by those who wish merely a summary of historical facts. For the student who seeks an opportunity to study original sources, a slight introduction has been made to this field. The text of the book contains almost no quotations in foreign languages, the result being that the reader will not meet with linguistic difficulties in the general narrative. In the notes, however, it is frequently desirable to quote the precise words of an author, and this has been done with reference to such European languages as are more or less familiar. It is not necessary to translate literally all these extracts, since the text itself sets forth the general meaning. Students who have some general knowledge of Latin, French, or German will have little difficulty, and in many cases will have much interest, in seeing various statements in their original form. For special reasons a few notes have been given in Greek, but in every case the meaning is evident from the text.

The footnotes have also permitted of the insertion of various biographical items which would merely burden the text, but which have considerable value to the student. In a general way it may be said that it is a matter of no moment where a man was born or on what day; but a work of this nature must be more than a book to be read,—it must be a work for future reference, and for this reason there may properly be made available, to be used if thought necessary, certain material which will aid the student in his later research. It would be possible to place all such supplementary material at the end of the book, but this would be merely an invitation to ignore it entirely.

No selection of names is ever satisfactory, even to the writer who makes it. In this work there are often included in one period names which would not be considered a century later; while others are omitted, particularly in the last three centuries, that would have been given prominence had their possessors lived at an earlier date. The criterion of selection has been the contribution of the individual to the development of elementary mathematics, his reputation as a scholar, and in particular his work in the creating of tendencies to further the study of particular branches of the subject. For the latter reason certain names have been included which would not otherwise have been considered. In Volume II a few minor names of arithmeticians have been mentioned in connection with the peculiar use of certain terms and the like, with no biographical notes, the latter being of little or no consequence.

In connection with dates before the Christian era the letters B. C. are used; in connection with dates after the beginning of this era no distinguishing letters are added except in a few cases near the beginning of the period, in which the conventional letters A. D. have occasionally been inserted to avoid ambiguity. With some hesitation, but for a purpose which seems valid, dates are frequently given in parentheses after proper names. It is well recognized that a precise date, like 1202 after the name Fibonacci, is of no particular value in itself. It makes no difference, in ordinary cases, whether Fibonacci wrote his *Liber Abaci* in 1202, in 1180, or in 1220, or whether *abacus* is spelled *abbacus*, as in some manuscripts, or in the more correct Latin form. On the other hand, two things are

accomplished by a free use of such dates. In the first place, a reader is furnished with a convenient measuring instrument; he does not have to look in the index or a chronological table in order to see approximately where the particular writer belongs in the world's progress. The casual reader may well be pardoned if he does not recall where Bede, Alcuin, Gerbert, Jordanus, Fibonacci, and Roger Bacon stood chronologically with respect to one another, and in reading a technical history of this kind there is no reason why he should not be relieved of the trouble of consulting an index when he meets with one of these names in the text. In the second place, it needs no psychologist to confirm the familiar principle that the mind comes, without conscious effort, to associate in memory those things which the eye has frequently associated in reading. At the risk, therefore, of disturbing the minds of those who are chiefly interested in the literary aspect of a general statement of the progress of mathematics, many important dates have been repeated, especially where they have not appeared in the pages immediately preceding.

In quoting from other writers the rule has been followed of making the quotation exact in spelling, punctuation, and phraseology. In carrying out this rule it is inevitable that errors should occasionally enter into the transcription, particularly in the case of old dialects; but the effort has been made to give the language precisely as it appears in the original. This accounts for the fact that certain French words in a quotation will sometimes appear without the modern accent, and that a word like Lilāvati may appear with any one of three spellings, depending upon the translator to whose work reference is made, or upon the author using the word.

Use has been made of such international symbols as *s.a.* (*sine anno*, without date of publication), *s.l.* (*sine loco*, without place of publication), *s.l.a.* (without place or date), *c.* (*circa*, about), and *seq.* (*sequens*, following), and of the abbreviations *ed.* (edition, edited by, or edition of), *vol.* (volume), and *p.* (page).

At the close of each chapter there has been given a page of topics for discussion, so arranged as to command more attention than they would have received had they been given in scattered form. These topics are not limited to questions to be answered from the text, but have purposely been made general, suggesting somewhat

more extended fields for study. The student will find it to his advantage if he is thus led to consult encyclopedias, general histories, and such works as are suggested in the bibliographical notes and as are available in libraries to which he may have access. It is by no means expected that an elementary work like this should contain the material for an extended study of any of these topics.

In the selection of illustrations the general plan has been to include only such as will be helpful to the reader or likely to stimulate his interest. It would be undesirable to attempt to give, even if this were possible, illustrations from all the important sources, for this would tend to weary the reader. On the other hand, where the student has no access to a classic that is being described or even to a work which is mentioned as having contributed to the world's progress in some humbler manner, a page in *facsimile* is often of value.

In general the illustrations have been made from the original books or manuscripts in the well-known and extensive library of George A. Plimpton, Esq., who has generously allowed this material to be used for the purpose; from the author's collection of books, manuscripts, mathematical portraits and medals, and early mathematical instruments; from manuscripts in various other libraries; and from such works as those by Professor Breasted.

Long experience in the use of books of reference has led the author to believe that a single index is more convenient than a series of indexes by names, subjects, and titles. Furthermore, readers who have used works like those of Cantor and Tropfke, for example, know the annoyance of a long list of page references after a given name, many of them of no particular significance. In this work, therefore, only a single index is given in each volume, and in each entry the page references are only such as the reader will find of particular value. In each case the first reference after a proper name relates to the biography of the individual, if one is given; the others relate to his leading contributions and are arranged approximately in order of importance.

DAVID EUGENE SMITH

CONTENTS

CHAPTER PAGE

BIBLIOGRAPHY xiii

PRONUNCIATION, TRANSLITERATION, AND SPELLING OF PROPER NAMES xvii

I. PREHISTORIC MATHEMATICS 1

1. IN THE BEGINNING 1
2. PRIMITIVE COUNTING 6
3. GEOMETRIC ORNAMENT 15
4. MYSTICISM 16

TOPICS FOR DISCUSSION 19

II. THE HISTORIC PERIOD DOWN TO 1000 B.C. 20

1. GENERAL VIEW 20
2. CHINA 22
3. INDIA 33
4. BABYLON 35
5. EGYPT 41

TOPICS FOR DISCUSSION 53

III. THE PERIOD FROM 1000 B.C. TO 300 B.C. 54

1. THE OCCIDENT IN GENERAL 54
2. THE GREEKS 55
3. ORIGINS OF GREEK MATHEMATICS 63
4. FROM PYTHAGORAS TO PLATO 69
5. INFLUENCE OF PLATO AND ARISTOTLE 87
6. THE ORIENT 95

TOPICS FOR DISCUSSION 101

CHAPTER PAGE

IV. THE PERIOD FROM 300 B.C. TO 500 A.D. 102

1. The School of Alexandria 102
2. Euclid 103
3. Eratosthenes and Archimedes 108
4. Apollonius and his Successors 116
5. Period of Menelaus 125
6. Ptolemy and his Successors 130
7. Diophantus and his Successors 133
8. The Orient 138

Topics for Discussion 147

V. THE PERIOD FROM 500 TO 1000 148

1. China 148
2. Japan 151
3. India 152
4. Persia and Arabia 164
5. The Christian West 177
6. The Christian East 190
7. Spain 192

Topics for Discussion 193

VI. THE OCCIDENT FROM 1000 TO 1500 194

1. Christian Europe from 1000 to 1200 194
2. Oriental Civilization in the West 205
3. Christian Europe from 1200 to 1300 211
4. Christian Europe from 1300 to 1400 230
5. Christian Europe from 1400 to 1500 242

Topics for Discussion 265

CHAPTER PAGE

VII. THE ORIENT FROM 1000 TO 1500 266

1. China 266
2. Japan 273
3. India 274
4. Persia and Arabia 283

Topics for Discussion 291

VIII. THE SIXTEENTH CENTURY 292

1. General Conditions 292
2. Italy 294
3. France 306
4. England 314
5. Germany 324
6. The Netherlands 338
7. Spain 343
8. Other European Countries 346
9. The Orient 350
10. The New World 353

Topics for Discussion 357

IX. THE SEVENTEENTH CENTURY 358

1. General Conditions 358
2. Italy 361
3. France 370
4. Great Britain 387
5. Germany 416
6. The Netherlands 422
7. Other European Countries 426
8. The Orient 435

Topics for Discussion 443

CHAPTER PAGE

X. THE EIGHTEENTH CENTURY AND AFTER 444

1. General Conditions 444
2. Great Britain 446
3. France 470
4. Germany 501
5. Italy 511
6. Switzerland 519
7. Other European Countries 526
8. United States 531
9. The Orient 533
10. The Historians of Mathematics 539

Topics for Discussion 547

CHRONOLOGICAL TABLE 549

INDEX 571

BIBLIOGRAPHY

The extent of a bibliography in a work of this kind is a matter of judgment. It can easily run to great length if the writer is a bibliophile, or it may have but little attention. The purpose of giving lists of books for further study is that the student may have access to information which the author has himself used and which he believes will be of service to the reader. For this reason the secondary sources mentioned in this work are such as may be available, and in many cases are sure to be so, in the libraries connected with our universities, while the original sources are those which are of importance in the development of elementary mathematics or which may be of assistance in showing certain tendencies.

The first time a book is mentioned the title, date, and place of publication are given, together, whenever it seems necessary, with the abbreviated title which will thereafter be used. To find the complete title at any time, the reader has only to turn to the index, where he will find given the first reference to the book. The abbreviation *loc. cit.* (for *loco citato*, in the place cited) is used only where the work has been cited a little distance back, since any more general use of the term would be confusing. The symbolism "I, 7" has been used for "Vol. I, p. 7" in order to conserve space, although exceptions have been made in certain ambiguous cases, as in the references to Heath's *Euclid*, references to Euclid being commonly by book and proposition, as in the case of Euclid, I, 47.

Although the number of works and articles on the history of mathematics is very great, the student will be able, in the initial stages of his investigation, to consult relatively few. For his convenience the books that he may most frequently use are here listed, special reference being made to those in English, French, and German which are likely to be found in college, university, and city libraries. The student will also find it advantageous to consult the leading encyclopedias.

Allman, G. J., *Greek Geometry from Thales to Euclid*, Dublin, 1889. Referred to as Allman, *Greek Geom.*

Ball, W. W. Rouse, *A Short Account of the History of Mathematics*, 6th ed., London, 1915. A readable survey of the general field. Referred to as Ball *History.*

Bretschneider, C. A., *Die Geometrie und die Geometer vor Eukleides*, Leipzig, 1870. Referred to as Bretschneider, *Die Geometrie.*

Cajori, F., *A History of Elementary Mathematics*, rev. ed., New York, 1917 Referred to as Cajori, *Elem. Math.*

—— *A History of Mathematics*, 2d ed., New York, 1919. Referred to as Cajori, *History.*

Cantor, M., *Mathematische Beiträge zum Kulturleben der Völker*, Halle, 1863. Referred to as Cantor, *Beiträge.*

—— *Vorlesungen über Geschichte der Mathematik*, 4 vols., Leipzig, 1880–1908, with various revisions. The standard general history of mathematics. Referred to as Cantor, *Geschichte.*

Encyklopädie der Mathematischen Wissenschaften, Leipzig, 1898–, with a French translation. Referred to as *Encyklopädie.*

Gow, James, *A Short History of Greek Mathematics*, Cambridge, 1884. Referred to as Gow, *Greek Math.*

Günther, S., and Wieleitner, H., *Geschichte der Mathematik*, 2 vols., Leipzig, 1908–1921. Referred to as Günther-Wieleitner, *Geschichte.* The second volume is the work of Dr. Wieleitner.

Hankel, H., *Zur Geschichte der Mathematik in Alterthum und Mittelalter*, Leipzig, 1874. Referred to as Hankel, *Geschichte.*

Heath, Sir Thomas Little, *A History of Greek Mathematics*, 2 vols., Cambridge, 1921. Referred to as Heath, *History.* Although the following special works by the same author are referred to in the footnotes, they are so important that it seems advisable to include them in this general bibliography.

—— *Apollonius of Perga*, Cambridge, 1896. Referred to as Heath, *Apollonius.*

—— *Archimedes*, Cambridge, 1897. Referred to as Heath, *Archimedes.*

—— *Aristarchus of Samos*, Oxford, 1913. Referred to as Heath, *Aristarchus.*

—— *Aristarchus of Samos. The Copernicus of Antiquity*, London, 1920. Referred to as Heath, *Aristarchus* (abridged).

—— *Diophantus of Alexandria*, 2d ed., Cambridge, 1910. Referred to as Heath, *Diophantus.*

—— *Euclid in Greek, Book I*, Cambridge, 1920. Referred to as Heath, *Euclid in Greek.*

—— *The Thirteen Books of Euclid's Elements*, 3 vols., Cambridge, 1908. Referred to as Heath, *Euclid.*

—— *Greek Mathematics and Science*, pamphlet, Cambridge, 1921. Referred to as Heath, *Address.*

Hilprecht, H. V., *Mathematical, Metrological, and Chronological Tablets from the Temple Library of Nippur*. Philadelphia, 1906. Referred to as Hilprecht, *Tablets*.

Libri, G., *Histoire des Sciences Mathématiques en Italie*, 4 vols., Paris, 1838–1841. Valuable on account of its style and its extensive notes. Referred to as Libri, *Histoire*.

Loria, G., *Guida allo Studio della Storia delle Matematiche*, Milan, 1916. Very valuable for its bibliography of the history of mathematics.

Marie, M., *Histoire des Sciences Mathématiques et Physiques*, 12 vols., Paris, 1883–1888. Biographical, convenient for reference, but inaccurate. Referred to as Marie, *Histoire*.

Mikami, Y., *The Development of Mathematics in China and Japan*, Leipzig, 1913. Referred to as Mikami, *China*. See also Smith-Mikami.

Miller, G. A., *Historical Introduction to Mathematical Literature*, New York, 1916. Serves a purpose in English similar to that of Loria's *Guida* in Italian. Referred to as Miller, *Introduction*.

Montucla, J. E., *Histoire des Mathématiques*, 2d ed., 4 vols., Paris, 1799–1802. Although written in the 18th century, it is a classic that is well worth consulting, particularly for its style. Referred to as Montucla, *Histoire*.

Pauly (A.)-Wissowa (G.), *Real-Encyclopädie der Classischen Altertumswissenschaft*, Stuttgart, 1894–. Best reference work for classical biography and antiquities. Referred to as Pauly-Wissowa.

Poggendorff, J. C., *Handwörterbuch zur Geschichte der exacten Wissenschaften*, 4 vols., Leipzig, 1863–1904. Referred to as Poggendorff.

Smith, David Eugene, *Our Debt to Greece and Rome. Mathematics*, Boston, 1922. Referred to as Smith, *Greece and Rome*.

—— *Rara Arithmetica*, Boston, 1908. being a bibliography of early arithmetics. Referred to as *Rara Arithmetica*.

Smith, D. E., and Karpinski, L. C., *The Hindu-Arabic Numerals*, Boston, 1911. Referred to as Smith-Karpinski.

Smith, D. E., and Mikami, Y., *History of Japanese Mathematics*, Chicago, 1914. Referred to as Smith-Mikami.

Tannery, P., *La Géométrie Grecque*, Paris, 1887. Referred to as Tannery, *Géom. Grecque*.

—— *Mémoires Scientifiques*, edited by J. L. Heiberg and H. G. Zeuthen, 2 vols., Paris, 1912.

—— *Pour l'Histoire de la Science Hellène de Thalès à Empédocle*, Paris, 1887. Referred to as Tannery, *Histoire*.

Tropfke, J., *Geschichte der Elementar-Mathematik in systematischer Darstellung*, 2 vols., Leipzig, 1902, 1903; 2d ed., 1921–. The best history of elementary mathematics. Referred to as Tropfke, *Geschichte*.

Zeuthen, H. G., *Histoire des Mathématiques dans l'Antiquité et le Moyen Age* translated by J. Mascart. Paris, 1902. Referred to as Zeuthen, *Histoire*

In matters of biography the student will find in W. Smith, *Dictionary of Greek and Roman Biography*, London, 3 vols., 1862–1864 (referred to as Smith's *Dict. of Greek and Roman Biog.*), a work of exceptional value, particularly with reference to the Greek mathematicians; and in the *Dictionary of National Biography* he will find the British mathematicians treated in a scholarly manner. For French biographies, Michaud's *Biographie Universelle* (1854–1865), Hoefer's *Nouvelle Biographie Générale* (1857–1866), *La Grande Encyclopédie*, and Larousse's *Grand Dictionnaire Universel du XIX siècle français* are very satisfactory. For German biographies the *Allgemeine Deutsche Biographie*, the Brockhaus *Conversations-Lexikon*, and the Meyer *Grosses Konversations-Lexikon* are helpful. Various earlier and less frequently used works are referred to in the footnotes.

Such special works as those of Matthiessen, Braunmühl, and Dickson are mentioned in the notes from time to time, as well as various other sources of information that may be found in the larger libraries.

Of the journals devoted to the history of mathematics those to which the student will most frequently refer are Boncompagni's *Bullettino di Bibliografia e Storia delle Scienze Matematiche e Fisiche*, Rome, 1868–1887 (referred to as Boncompagni's *Bullettino*), and Eneström's *Bibliotheca Mathematica*, Leipzig, three series, 1885–1915 (referred to as *Bibl. Math.*).

PRONUNCIATION, TRANSLITERATION, AND SPELLING OF PROPER NAMES

General Question. The question of the spelling and transliteration of proper names is always an annoying one for a writer of history. There is no precise rule that can be followed to the satisfaction of all readers. In general it may be said that in this work a man's name has been given as he ordinarily spelled it, if this spelling can be ascertained. To this rule there is the exception that where a name has been definitely anglicized, the English form has been adopted. For example, it would be mere pedantry to use, in a work in English, such forms as Platon and Strabon, although it is proper to speak of Antiphon and Bryson instead of Antipho and Bryso. When in doubt, as in the case of Heron, the preference has been given to the transliteration which most clearly represents the spelling that the man himself used.

In many cases this rule becomes a matter of compromise, and then the custom of a writer's modern compatriots is followed. An example is seen in the case of Leibniz. This spelling seems to be gaining ground in our language, and it has therefore been adopted instead of Leibnitz, even though the latter shows the English pronunciation better than the former. Leibniz himself wrote in Latin, and the family spelled the name variously in the vernacular. There seems, therefore, to be no better plan than to conform to the spelling of those recent German writers who appear to be setting the standard that is likely to be followed.

There is also the difficulty of finding a satisfactory solution in the case of men who were themselves polyglots, who lived in polyglot towns, or who made their homes in more than a single country. This is seen, for example, in the case of the Bernoullis. Jacques Bernoulli lived in Basel, a Swiss city where German was chiefly spoken and where the common spelling of the name of the place is the one

here given. He was of Belgian descent, but he usually wrote either in Latin, in which his first name was spelled Jacobus, or in French, in which he would naturally use the name of Jacques. To call him James, as various English writers have done, would merely confuse an American reader, while to adopt the German Jakob would be to use a form which Bernoulli himself did not adopt in writing. The fact that he preferred to use French as his means of correspondence, when not writing in Latin, makes it desirable to speak of him as Jacques and to follow a similar usage with respect to his brother, Jean Bernoulli.

Another difficulty arises when we consider the Graeco-Latin forms of names in the Renaissance period. In general, if a man commonly used such a form, as was the case with Grammateus, Regiomontanus, and Dasypodius, this form has been used in the text, with the family name given in a footnote. In a case on the border line, like that of Schoner, however, the vernacular form, spelled as the man seems himself to have preferred, has been adopted. It must also be understood that early writers were often not uniform in spelling their own names. Thus, we have Recorde and Record, Widman and Widmann, and Scheubel and Scheybel, and in these cases all that the historian can do is to endeavor to choose that spelling which the writer seems himself to have most commonly used.

A further difficulty is encountered with certain names in regard to which the possessor was himself undecided as to his preference. A typical case is that of Leonardo Pisano Fibonacci,—Leonardo the Pisan, son of Fibonacci, or of Bonacci, or of Bonacius. It would be proper to write his name "l. pisano," since this form appears in one of the early manuscripts; or Bigollo, since he used this nickname; or Leonardo of Pisa, although this combines Italian and English; but the form Fibonacci has been chosen for general use, chiefly because Fibonacci's Series is so frequently mentioned in mathematics. It would not be difficult to show a lack of consistency in many cases, as when the common form of Gemma Frisius (Gemma the Frieslander) is preferred to the family name of Renier, with various spellings. In the case of a name like that of Pacioli, where different forms are used in the various works of the individual, the one seemingly preferred by the majority of historians has been chosen. In the case of a name like Joannes or Johannes the effort

has been made to use the form which the possessor used, or at least the one which was the more commonly employed by his contemporaries when referring to him.

The greatest difficulty in transliteration arises with respect to oriental names. In the first place, we have no international system of transliteration that is generally accepted; and in the second place, it is difficult to know the name which the writer himself preferred. An Arab scholar may have as many as a dozen parts to his name; a Japanese or Hindu writer may have an intimate name and also an official name; a Chinese mathematician may be known only by an ideogram, the pronunciation of the name being lost or varying in different parts of his own country; and certain of these names may have found their way into medieval Latin and have been distorted almost beyond recognition.

If a name is fairly familiar in English, like Omar Khayyam, it has been retained, even if the form is open to criticism. If it has taken an English form but is not so familiar, as in the case of Savasorda, the attempt has been made to use the distorted name and also to adopt the best modern transliteration of the real name from which this is derived. In the cases of such less familiar names as seem to deserve mention, these will neither be read aloud nor be kept in mind by most readers, and hence an abridged form has been given in the text, in as good transliteration as seems possible, the full form being placed in a footnote. The Arabic *al-* has been used instead of *el-* or *ul-*, simply because it is the most common form in English. As a matter of fact, the Arabic pronunciation, like that of the Chinese, is by no means standardized.

The pronunciation of proper names has been given in cases where it is likely to be helpful to the student, and in many cases the accent has been indicated when the name first appears in the text. In such cases the English pronunciation has been taken whenever the name has become thoroughly anglicized, but otherwise the pronunciation has been given as nearly as possible as it stands in the vernacular. In the case of Greek names the original form has usually been given in the notes, partly because of the differences in accent and partly because the Greek alphabet is well enough known to allow the original and frequently interesting form to be understood.

Arabic Names. The standard authority on the transliteration and pronunciation of Arabic names is Suter, a Swiss writer, whose "Die Mathematiker und Astronomen der Araber und ihre Werke" appeared in Volume X of the *Abhandlungen zur Geschichte der Mathematischen Wissenschaften*, and in "Das Mathematiker-Verzeichniss im Fihrist," in Volume VI of the same work. The rules given by this writer have, in general, been followed, except that *j* has been used for *ǧ*, *y* for *j*, *v* for *w*, *kh* for *ch*, and *al-* for *el-*, to conform to English pronunciations and custom. Although the reader will seldom need to pronounce the names, it will be helpful to be able to do so if necessary. The following is a summary of the scheme of transliteration and pronunciation employed:

b, **d**, **f**, **g**, **h**, **j**, **l**, **m**, **n**, **p**, **s**, **sh**, **t**, **th**, **w**, **x**, **z** as in English.

a as in *ask*; **â** as in *father*, the form *â* being used instead of *ā* in Arabic words, partly to conform to the Suter list.

e as in *bed*.

i as in *pin*; **î** as in *pique*.

o as in *obey*.

u as in *put*; **û** as in *rule*.

ḍ, **ṣ**, **ṭ**, **ẓ** as in English but made with the tongue spread so that the sounds are produced largely against the side teeth.

ṇ is generally pronounced by Europeans as simple *n*.

d̲ like *th* in *that*; **t̲** like *th* in *thin*.

ġ is a voiced consonant formed below the vocal chords; it is sometimes compared to a guttural *g* and sometimes to a guttural *r*.

h retains its consonant sound at the end of a word.

ḥ may be compared to the German hard *ch*, as in *nach*.

k as in English; **kh** is the hard German *ch*, as in *nach*.

q like *c* or *k* in *cook*.

r stronger than in English.

v like the English *w*; **y** as in *you*.

ʿ represents the *spiritus lenis* and may be taken simply as separating distinctly two vowels, like the break between the *e*'s in *reentrant*.

A final vowel is shortened before *al* (which then becomes *'l*) or *ibn* (whose *i* is then silent).

In *al* the final *l* often takes the sound of a following consonant, as in *al-Rashid* (*ar-Rashid*).

The accent is on the last syllable containing a long vowel or a vowel followed by two consonants, except that a final long vowel is not usually accented. Otherwise the accent falls on the first syllable.

Hindu Names. The transliteration of Hindu names has changed greatly within a century, and even yet is not internationally standardized. In general, in quoting from earlier English writers, the forms which they used have been followed. Thus, there will be found in the notes various references to Taylor's *Lilawati*, this being the name of the book as the translator used it; or, when the actual title is mentioned, to Colebrooke's translation of the *Lílávatí*, this being the form which this author used; but the modern form *Lilāvati* appears in the text. The effort has been made to follow the best current practice of English orientalists, and in determining the form and pronunciation of Sanskrit words the following equivalents have been used:

b, **d**, **f**, **g**, **h**, **j**, **l**, **m**, **n**, **p**, **v**, **w**, **x**, **z** as in English.

a like *u* in *but*; thus, *pandit*, pronounced *pundit*; **ā** as in *father*, the form *ā* being used instead of *â* in Hindu words.

e as in *they*.

i as in *pin*; **ī** as in *pique*.

o as in *so*.

u as in *put*; **ū** as in *rule*.

c like *ch* in *church* (Italian *c* in *cento*).

ḍ, **ṇ**, **ṣ**, **ṭ** like *d*, *n*, *sh*, *t* made with the tip of the tongue turned up and back into the dome of the palate.

h preceded by *b*, *c*, *t*, *ṭ* does not form a single sound with these letters but is a more or less distinct sound following them, somewhat as in *abhor*; **h** is final consonant *h*.

k as in *kick*.

ṁ, **ṅ** like the French final *m* or *n*, nasalizing the preceding vowel.

ś, English *sh*.

y as in *you*.

' in some transliterations is used to indicate the *spiritus lenis*, a break between two letters.

The accent is as in Latin: if the penult is long, it is accented; if it is short, the antepenult is accented.

Japanese Names. Modern Japanese scholars have carefully transliterated into the Roman alphabet the names of all their leading mathematicians. The letters are pronounced as in English except that *i* is pronounced like *e* in *feel*; *e* as in *grey*; *ai* as in *aisle*; and *ei* like long *a*; but *i* and *e* also take a short sound as in English. Japanese names have only a slight accent.

Chinese Names. There is no uniform system of transliterating and pronouncing Chinese names and terms. The author's colleague, Professor Hirth, in his *Ancient History of China*, followed in general the plan adopted by the Royal Geographical Society of London and the United States Board on Geographic Names, and the present text follows in the main the rules which he has laid down. Briefly stated, the scheme of pronunciation is as follows:

a as in *father*.
e, **é** as in *men*. The accent simply shows that it does not form part of a diphthong.
i as in *pique*. When followed by *n* or by a vowel it is short as in *pin*.
ï, used when *i* is intonated with the adjoining consonant, as in *ji*, or is but faintly heard, when it follows *e*, as in *leï*.
o as in *mote*.
ö like the French *eu* in *jeu* or like the German *ö*.
u like *oo* in *boot*. When preceding *n*, *a*, or *o* it is short.
ü like the French *u*. When preceding *n*, *a*, or *é* it is short.
ai like *i* in *ice*.
au like *ow* in *how*.
eï somewhat like *ey* in *they*.
ou a diphthong with the two vowels distinctly intonated.
ui like *ooi* contracted into a diphthong.
The initials **k**, **p**, **t**, **ch**, **ts**, and **tz** are not so hard as in English. When pronounced as hard as possible they are followed by (') as in *k'au*.
ch like *ch* in *church*. When followed by *i*, the vowel blends with it.
f, **h**, **l**, **m**, **n**, **sh**, **w**, **ng** as in English.
j like *j* in French.
ss like *ss* in *mess*. When followed by *i*, the vowel disappears.
y like *y* in *you*.

Names from Other Languages. In the case of Russian names there has been chosen the transliteration which represents most effectively the English equivalent sounds. For example, the spelling Lobachevsky has been preferred to the German form, Lobatschewski, or to other forms which are not appropriate to our language. The same may be said with respect to other foreign names where the Roman alphabet is not in use or is supplemented by other letters.

3. Geometric Ornament

Early Art. A further prehistoric stage of mathematical development is seen in the use of such simple geometric forms as were suggested by the plaiting of rushes, the first step in the textile art. From this there developed those forms used in clothing, tent cloths, rugs, and drapery which are usually found among primitive peoples. Since the earliest trace of human art that we have thus far found is seen in representations of animals, these being drawn on bone in the Early Stone Age, one might expect to find such figures in early mural decorations, and this is not only the case but is one means of dating the latter with some degree of approximation. The geometric ornament, however, became in due time a favorite one among nearly all early peoples. This may have been because the plaiting of rushes furnishes an easy medium for the representation of geometric forms, but at any rate such forms as the swastika and the Greek key developed at an early period. Such decorations are not confined to the textiles of the people; they are equally prominent in architecture in all parts of the world. They are found on the early monuments of Mexico, on the architectural remains of Peru; on the huts of the savage, and on the early buildings of the historic period in various parts of the Old World, especially on those devoted to the commemoration of the dead or to the worship of the gods.




EGYPTIAN POTTERY OF THE PREDYNASTIC PERIOD

It shows the earliest stage of geometric ornament on pottery. The Predynastic Period extended from *c.* 4000 to *c.* 3400 B.C. From the Metropolitan Museum, New York

The same instinct that leads to geometric decoration of religious structures shows itself in the decoration of personal ornaments and of articles intended for domestic use. This is seen in the handicraft of the Stone Age, it is found in the

rich gold work of early Egypt, and it is equally in evidence in most of the jewelry of modern times. It is not merely the instinct of symmetry that we find in these petrified thoughts of the race; it is quite as much a desire to fathom the mystery and grasp the meaning of the beauty of geometric form.



CYPRUS JUG OF THE PERIOD 3000-2000 B.C.


Pottery of the Early Bronze Age, showing the second stage in geometric ornament. From the Metropolitan Museum, New York

Early Pottery. The early pottery of Egypt and Cyprus shows very clearly the progressive stages of geometric ornament, from rude figures involving parallels to more carefully drawn figures in which geometric design plays a more important part and in which such mystic symbols as the swastika are found. Art was preparing the way for geometry.

4. Mysticism

Religious Mysticism. The beginning of an appreciation of the wonders of mathematics is closely connected with the beginning of religious mysticism. Man wondered at the heavens above him; he wondered at life and he wondered even more at death; all was a mystery. He likewise wondered at the peculiarities of geometric forms and at the strange properties of such numbers as three and seven, the two primes within his limited number realm that were not connected with his common scales of counting. The mystery of form and the mystery of number he connected with the mystery of the universe about him,—the universe in which he felt himself a mere mote in the sunbeam. His sense of wonder at the potency of the sun led him to the orientation of his religious structures; his recogni-

tion of a pole star led him to consider a fourfold division of his horizon, and to speak of the four corners of the earth; and it is not impossible that the swastika and the various other cruciform figures of the ancient civilization are a recognition of this tendency. The number four was looked upon as peculiarly significant by certain American aborigines as well as by the early peoples of Asia, Australia, and Africa, and we may have a relic of this attitude of mind when we speak of "a square man" or one who acts "squarely."



PAINTED JUG FROM CYPRUS
1000–750 B.C.

Pottery of the Early Iron Age in Cyprus, showing a third stage in the use of geometric ornament, with the swastika. The Geometric Period of decoration closed, for the Mediterranean countries, just before the time of Thales. From the Metropolitan Museum, New York

Architecture. Just as we find an instinctive appreciation of the beauties of geometric forms as applied to personal ornaments, so we find it as applied to architecture, not merely with respect to decoration as already mentioned, but in the general structure of temples, of altars, and of tombs. In early India, for example, there seems to have been no study of geometry as such except in connection with forms used in the temple, and this was probably the case in other parts of the earth. A desire to adapt symmetry to architecture is seen in the terraced pyramids of Mexico as well as in those of Egypt; and while these buildings are not prehistoric, they doubtless are the outgrowth of prehistoric forms.

Observations of the Stars. As already mentioned, the primitive man seems to have felt that the secret of the stars was closely bound up with the secret of his destiny. It was

this that led the Babylonian shepherd and the desert nomad to observe the stars, to speculate upon their meaning, and to take the first steps in what developed into a priest lore in the temples along the Nile and in the land of Mesopotamia. It was this, too, that led the early philosophers and poets to consider the stars as lighted lamps suspended in a vast material vault, or as golden nails fixed in a crystal sphere,—ideas perfectly suited to the childhood of the race. When it was that these observations of the heavens led to angle measure, to the recording of such celestial phenomena as eclipses, and to a naming of the signs of the zodiac and the constellations, we cannot say. One writer of prominence[1] places a recognition of the common constellations as early as 17000 B.C., and while this date seems to be very improbable, even though supported by certain historico-astronomical considerations, it is doubtless true that the period of this recognition and of the observance of certain celestial phenomena is very remote. While there is good reason for thinking that these early steps in astronomy were taken in Mesopotamia, the proof is not sufficiently strong to enable us to say that this was unquestionably the case, nor are we able to fix upon the period within any particular century or even within any particular millennium. Similarly, we are unable to state the time or place in which the early peoples began to recognize the constellations or to give them fanciful names. After attaining a certain degree of success in our research, we are lost in the prehistoric clouds.

Lengthen the story all that we can, it is not possible to extend it back more than an imperceptible distance on the great clock face. For if we represent the period of all life on our planet by one revolution of the minute hand, the period of human life will be covered by only half a minute, and recorded history will be represented by less than two seconds. What we definitely know of the history of mathematics covers a period in world development so short as to seem almost infinitesimal.

[1] G. Schlegel, *Uranographie Chinoise*, 2 vols., II, 796 (Leyden, 1875). For a recent discussion of the whole question, see Léopold de Saussure, "Les origines de l'astronomie chinoise," *T'oung Pao*, Vols. X seq. (Leyden).

TOPICS FOR DISCUSSION

1. Geometric forms that were in existence before the advent of life on the planet.

2. Laws of motion that entered into the formation and perpetuation of our solar system.

3. Geometric forms that appear prominently in the vegetable world and in the bodily structure of certain animals.

4. Geometric forms that appear prominently in the products of the labor of the lower animals, with the question of maximum efficiency in any of these cases.

5. The question of animal counting or pseudo-counting as discussed by psychologists.

6. Evidence of primitive counting without any scale.

7. The world's use of scales below five as shown by a study of our language and of savage tribes.

8. Reasons why the scales of five, ten, and twenty were the leading favorites.

9. Reasons why the scale of twelve would have been a particularly good one.

10. Reasons why three and seven have been particularly notable as mystic numbers, with several illustrations.

11. Circumstances which developed a high degree of skill in counting among certain peoples.

12. Reasons which led primitive peoples to the use of geometric forms in ornament.

13. The effect of religious mysticism upon primitive mathematics.

14. Various stages of geometric ornament in Cyprus, Crete, and the mainland of Greece.

15. Possible influence of geometric decoration upon the study of geometry as a science.

16. Geometric decoration that has persisted in all ages, with a study of the probable causes for this persistence.

17. Causes leading to an interest in astronomy among primitive peoples. Features of the ancient astronomy that are still found either in our present study of the science or in folklore.

18. Evidence of the antiquity of astronomical ideas, particularly in Mesopotamia, Egypt, and China, with probable evidence in the case of India and other parts of the East.

CHAPTER II

THE HISTORIC PERIOD DOWN TO 1000 B.C.

1. General View

Sources of our Knowledge. The period down to the arbitrarily selected date 1000 B.C. overlaps the prehistoric period mentioned in Chapter I, the prehistoric gradually merging into the historic in certain parts of the world but not reaching this stage in certain other parts. Various facts which might properly have found place in Chapter I will therefore be related in this chapter, but in a general way we shall now pass to that period in the evolution of the race in which less use need be made of conjecture in the recital of the story of mathematics, although it cannot be wholly eliminated.

The sources of our knowledge are no longer mere tradition, nor does inference from the study of savage tribes constitute so important a basis for our statements. The sources are now, in general, the relics of human activity, largely in the form of inscriptions or manuscripts which actually date from remote centuries, or of copies of such evidences.

Countries Considered. There are four countries which have left to posterity such an abundance of historical material prior to the beginning of the first millennium of the pre-Christian era as to warrant our special consideration. These countries, considered geographically instead of politically, are Egypt, Mesopotamia, China, and India. Each claims for itself a high degree of antiquity, each claims to have been a pioneer in mathematical development, each is ethnographically somewhat of a unit; and in certain respects the claims of each have reasonable foundations. Each had, at least in considerable areas, a salubrious climate in the warm intervals between the several

descents of the ice from the north,—descents which characterize what is commonly called the Glacial Epoch,—and hence each was able to develop an early civilization. Each flourished along one or more important rivers, which not only furnished water for navigation and for domestic purposes but also afforded opportunities for the application of a rude form of mathematics to the irrigation projects which were already in evidence in the early centuries of the historic period. From each, after the ice retreated, a human stream in due time flowed to the more invigorating climate of the north and carried along with it traces of the early mathematical lore which had already begun to develop in the temples, and of the customs established among the primitive traders of the more favored lands.

Antiquity of Mathematics. Although there is evidence showing that the human race has been on the earth for hundreds of thousands of years, the earliest definite remains that have come down to us are from the Early Stone Age, or Paleolithic Age, which seems to have begun in the warm interval after the third descent of the ice. This was not less than 50,000 years before our era and may have been from 50,000 to 75,000 years earlier still. In the remains of this period we find implements which suggest the existence of barter and the need of numbers for counting, although they date from a period thousands of years before there had dawned upon human intelligence any idea of written numerals. We may simply suppose that the presence of such implements and the important discovery of a means for making fire, which seems to have occurred about 50,000 years ago, are evidence of a degree of intelligence high enough to assure some idea of number.

About 15,000 years before our era there is thought to have begun the Middle Stone Age, the period of the fourth descent of the ice. In this period we find the oldest known works of art. These works show such an intellectual advance as to make it quite certain that the world had reached a period when the abstract notion of number must have been in evidence,—a judgment warranted by our knowledge of all primitive peoples today who have reached this stage in art.

The Late Stone Age is relatively a recent period, dating from *c.* 5000 B.C., and by this time there had developed quite elaborate number systems, and the observation of the stars had become a fairly well organized science. That this was the case we know from various historical facts which will be mentioned later.

Advent of Writing. Metal, as distinguished from natural ore, was discovered *c.* 4000 B.C., possibly in the Sinai peninsula, and with this there came a new need for weighing and measuring and a new impetus to a system of barter which was doubtless very old even at that time. About 500 years later, writing is known to have been in use, and the system of ruling over masses of people had become so advanced as to render possible the control of a population of several millions by one government. The bearing of all this upon the development of a number system and upon systematic taxation is apparent. About 3000 B.C. the earliest stone masonry was laid and sea-going ships began to cross the Mediterranean, and a little later the pyramids of Egypt were erected, so that history now enters a period in which mathematics reached out beyond mere counting and into such fields as that of practical geometry, including a primitive kind of leveling and surveying.

2. China

Early Chinese Mathematics. We have no definite knowledge as to where mathematics first developed into anything like a science. Mesopotamia has several strong claims to priority, and so has Egypt. As to China, we have little positive knowledge of its earliest literature, the possibility of corruption of its texts being such as to cast doubts upon its extreme claims. Until native scholars develop a textual criticism commensurate with that which has been developed in the Occident, this uncertainty will continue to exist. The historical period begins with the 8th century B.C., or, at the earliest, with the reign of Wu Wang, the Martial Prince, in 1122 B.C.[1] In beginning with

[1] A. J. Little, *The Far East*, p. 20. Oxford, 1905.

China, therefore, it must not be thought that we should recognize the validity of all the claims that are often advanced for the antiquity of her science.

Basing his opinion upon later historical descriptions of the primitive astronomy of China, Professor Schlegel of The Hague, as already remarked, asserts that the Chinese recognized the constellations as early as 17000 B.C., which was about the close of the Early Stone Age. There is nothing impossible in such a supposition, although it is improbable. The race had developed considerably by that time, and it may well have extended its poetic fancy to the giving of forms to groups of stars which it had looked upon for thousands of years. Professor Schlegel also fixes upon 14700 B.C. as an approximate date of the duodenary zodiac, other scholars asserting that 13000 B.C. is more probable and still others fixing upon 4000 B.C.,—a discrepancy that may well arouse skepticism as to the validity of any of these hypotheses. Schlegel also believes that there is evidence of the extended study of the celestial sphere in China in or about 14600 B.C.

While such claims are generally doubted by competent Sinologists, it is quite likely that the Chinese developed some acquaintance with descriptive astronomy at an early period, and that this development necessitated such knowledge of mathematics as the measure of time and angles and the use of fairly large numbers. Reasonably well-founded tradition gives the probable dates of Fuh-hi,[1] the reputed first emperor of China, as 2852–2738 B.C.,[2] and in his reign there were extensive astronomical observations. In this general period the Chinese are believed to have changed their zodiac into one of twenty-eight animals.

[1] In general, the transliteration of Chinese names is that of Y. Mikami, *The Development of Mathematics in China and Japan* (Leipzig, 1913) (hereafter referred to as Mikami, *China*), and H. A. Giles, *A Chinese Biographical Dictionary* (London, 1898). The transliteration varies greatly with different Sinologists, a title like *I-king* appearing as *Yih-ching*, *Yi-ching*, *Ye King*, *Y-Ching*, and so on. In many cases I have been greatly assisted by my colleague, Professor Friedrich Hirth, one of the greatest living Sinologists.

[2] F. Hirth, *The Ancient History of China*, p. 7 (New York, 1908). Professor Hirth follows Arendt's tables as being the most carefully considered.

Reign of Huang-ti. In the year 2704 B. C.[1] Huang-ti, the Yellow Emperor, began his reign. Under his patronage it is said that Li Shu wrote on astronomy and that Ta-nao established the *Chia-tsŭ*, or sexagesimal system, both of these statements being supported by copies (possibly altered) of ancient records.[2] Even the emperor himself is said to have taken such an interest in mathematics as to write upon astronomy and arithmetic, and in his reign an eclipse of the sun was observed and recorded. Tradition assigns to this period even the decimal system of counting, although it is more likely that some popular work on the subject was written at this time. It was possibly during the reign of the emperor Yau[3] (*c.* 2357–*c.* 2258 B.C.) that two brothers, Ho and Hi, made astronomical observations. They are said to have suffered the displeasure of the emperor through their failure to predict a solar eclipse,[4] an incident showing a state of mathematical advancement quite equal to that in Greece in the time of Thales, some 1500 years later. The story is told in the *Shu-king* (*Canon of History*), an ancient record sometimes attributed to the pen of the emperor himself and sometimes to that of Confucius nearly two thousand years later.[5]

[1] According to Arendt and Hirth. Giles gives 2698 and others give 2697. Huang-ti is said to have died at the age of 111 years.

[2] As stated in Volume II, it is doubtful if the Chinese used anything like a sexagesimal system at this time, although they may have learned from the Sumerians that 60 is a convenient unit for subdivision.


[3] Reputed to have lived nearly a full century. See A. T. de Lacouperie, *The Languages of China before the Chinese*, p. 9 (London, 1887); Hirth, *loc. cit.*, p. 29.

[4] R. Wolf, *Geschichte der Astronomie*, p. 9 (Munich, 1877); *Handbuch der Mathematik, Physik, Geodäsie und Astronomie*, I, 7 (Zürich, 1872).

[5] Hirth, *loc. cit.*, pp. 29, 33, 251, who believes it to be a late work, or at least, if written in the time of Yau or of his immediate successors, to have been greatly modified by later copyists. Some effort has been made to fix the date of the eclipse as May 7, 2165 B.C. Other proposed dates are October 11, 2154 B.C., October 12, 2127 B.C., October 24, 2006 B.C., October 22, 2155 B.C., and October 21, 2135 B.C. The discrepancy between these later dates and those tentatively assigned to Yau, as given above, has little significance in the present state of knowledge as to Chinese chronology. The whole subject is still in the conjectural stage and awaits extended research on the part of capable Sinologists.

It is this emperor Yau and his successor, the emperor Shun, who, it is said, carried farther to the eastward the dominion established by the Bak tribes which had come from western Asia. These tribes had been under the civilizing influences of the people of Susiana, who in turn had received their civilization from Babylon.[1] If this theory proves to be correct, the similarity between certain early forms of astronomy and mathematics in the East and the West is more easily explained.

I-king. Of the "Five Canons" (*Wu-king*) of the Chinese probably the third in point of antiquity is the *I-king*, or *Book of Permutations*.[2] In this appear the *Liang I*, or "two principles" (the male, *yang*, ——; and the female, *ying*, — —) and from these were formed the *Sz' Siang*, or "four figures,"



and the *Pa-kua* (eight-kua) or eight trigrams, the eight permutations of two forms taken three at a time, repetitions being allowed. These *Pa-kua* had various virtues assigned to them and have been used from a very early period until the present for purposes of divination. It was probably Wön-wang (1182–1135 B.C.) who wrote the *I-king*; at any rate it was he who extended the *Pa-kua* into the sixty-four hexagrams now found in this classic.[3]

[1] A. T. de Lacouperie, *loc. cit.*, pp. 9 seq.

[2] It is often called the oldest of the Chinese classics, as in the edition by J. Mohl, *Y-King; Antiquissimus Sinarum liber* (Stuttgart, 1834–1839). In the extensive literature on the *I-king* the following works may be consulted: H. Cordier, *Bibliotheca Sinica*, II, cols. 1372 seq. (Paris, 1905–1906); A. T. de Lacouperie, "The Oldest Book of the Chinese," *Journal of the Royal Asiatic Society*, XIV (N.S.) (London, 1882), 781, reprinted in 1892, with an extensive bibliography; T. McClatchie, "The Symbols of the Yih-King," *The China Review*, I (Hongkong, 1872), 151; J. Edkins, "The Yi king of the Chinese," *Journal of the Royal Asiatic Society*, XVI (N.S.) (London, 1884). 360; H. J. Allen, *Early Chinese History*, chap. viii (London, 1906). The first European edition of the *I-king* appeared at Frankfort in 1724.

[3] Hirth, *loc. cit.*, p. 59. The *Pa-kua* are attributed to Fuh-hi by Liu Hui, who wrote *c.* 250 A.D. The Leibniz theory, set forth in his *Philosophia Sinensium*, § 4, that these symbols had some connection with binary numerals, has no historical foundation in the *I-king* as originally written.

It is hardly conceivable by the Western mind that such a set of symbols should last for thousands of years, that it should be the subject of such a large number of books and monographs as have appeared in explanation of its meaning, and that it should be known today to everyone among the hundreds of millions who have come under the influence of the Chinese philosophy, not merely in China but all through the East.

☰	☱	☲	☳	☴	☵	☶	☷
k'ién heaven	*tui* steam	*li* fire	*chön* thunder	*sün* wind	*k'an* water	*kön* mountain	*k'un* earth
7	6	5	4	3	2	1	0
Heaven Sky	Collected Water	Fire	Thunder	Wind Wood	Water as in Rain Moon	Hills	Earth
S.	S.E.	E.	N.E.	S.W.	W.	N.W.	N.


THE PA-KUA, OR EIGHT TRIGRAMS

From the *I-king*, or *Book of Permutations*. On the ordinary diviner's compass these directions are reversed

An examination of the above interpretation of the *Pa-kua*, the one commonly given by Oriental writers, suggests the Pythagorean doctrines with respect to numbers, and as we proceed we shall find still more to strengthen the belief that the West obtained much of its mysticism from the East.

Although there is no historical evidence that the Chinese looked upon the *Pa-kua* as numerals, based upon the scale of two, it is true that if we take ⚊ for one and ⚋ for zero, the successive trigrams, beginning at the right, have values which we may represent by our numerals as 000, 001, 010, 011, 100, 101, 110, and 111. If these are considered as numbers written on the scale of two, their respective values are 0, 1, 2, 3, 4, 5, 6, and 7.

The *Pa-kua* are found today on the compasses used by the diviners in every city and village of China. They are also



THIBETAN "WHEEL OF LIFE"

From a sheet of block printing done at Lhassa. This portion represents the signs of the zodiac, the *Pa-kua*, and, in the center, a magic square

found on fans, vases, and many other objects of the home, and on talismans of various kinds in common use in Thibet and other parts of the Far East.


The lo-shu and ho-t'u. The *I-king* also states that the *Pa-kua* were footsteps of a dragon horse which appeared on a

4	9	2
3	5	7
8	1	6

MAGIC SQUARE

The rows, columns, and diagonals in this particular magic square have 15 as their respective sums

river bank in the reign of the Emperor Fuh-hi, and that the *lo-shu*, in reality the magic square here shown, was written




THE LO-SHU FROM THE I-KING

This is the world's oldest specimen of a magic square. The black circles are used in representing feminine (even) numbers, the white ones in representing masculine (odd) numbers

upon the back of a tortoise which appeared to Emperor Yu (*c.* 2200 B.C.) when he was embarking on the Yellow River. The *ho-t'u*, also a highly honored mystic symbol, appears in the same work.

It thus appears that the *I-king* is not a work on mathematics, but that it contains the first evidence of an interest in permutations and magic squares that has come down to us. It is



THE HO-T'U FROM THE I-KING


This was never considered so important as the *lo-shu*, lacking as it does the interest of the magic square

reasonable to believe, however, that both these ideas were already ancient when the book was written.

The Chóu-peï. The oldest Chinese work that can be designated as mathematical is the *Chóu-peï*, or the *Chóu-peï Suan-king*,[1]

[1] *Suan-king*, or *Suan-ching*, means "arithmetic classic." Also transliterated in various other ways, such as *Tcheou-pei-swan-king*. See E. Biot, "Traduction et examen d'un ancien ouvrage chinois intitulé Tcheou pei," *Journal Asiatique* (1841), p. 595, with a discussion of dates.

a work relating chiefly to the calendar but containing information referring to ancient mathematics, including some work on shadow reckoning. The author and the date of the work are both unknown, and there is some reason for believing that it has undergone considerable change since it was first written. The fact that Emperor Shï Huang-ti[1] of the Ch'in Dynasty, in 213 B.C., ordered all books burned and all scholars buried, would seem at first thought to have given an opportunity for radically



CHÓU-PEÏ SUAN-KING

A work written in the second millenium B.C. This illustration is from a very early specimen of block printing. It shows the figure of the Pythagorean Theorem, but gives no proof

altering all ancient treatises; but such a sweeping decree could not possibly have been executed, and even if every book had been lost there would have been many who could have repeated the ancient classics verbatim from memory. The probability is that we have about as near the primitive form of these classics as we have of the writings attributed to Boethius, Bede, or Alcuin, or of certain Greek authors whose works we assume

[1] Also transliterated Tsin Chi Hoang-ti and Tsin schè huâng ty (born 259 B.C.; died 210 or 211 B.C.). The claims of such writers as Weber and J. B. Biot for a high grade of mathematical learning in China before this time are contradicted by L. Am. Sédillot, "De l'astronomie et des mathématiques chez les Chinois," Boncompagni's *Bullettino*, I, 161.

as known. In any case it is probable that we have in the *Chóu-peï* a very good record of the mathematics of about 1105 B.C., the year of the death of Chóu-Kung, a party to one of several dialogues which the book records.[1] One of these dialogues is between the prince Chóu-Kung and his minister Shang Kao, and relates to number mysticism, mensuration, and astronomy. Among the stories told of the energy of Chóu-Kung is one relating to his habit of rushing several times from his bath, holding his long, wet hair in his hand, to consult with his officials. Tradition also states that he had a wrist like a swivel, on which his hand could turn completely round,—an odd fiction for those who are interested in stories of mathematicians. A few extracts from the *Chóu-peï* will give some idea of the nature of the work:

The art of numbers is derived from the circle and the square.

Break the line and make the breadth 3, the length 4; then the distance between the corners is 5.[2]

Ah, mighty is the science of number.

Forms are round or pointed; numbers are odd or even. The heaven moves in a circle whose subordinate numbers are odd; the earth rests on a square whose subordinate numbers are even.

One who knows the earth is intelligent, but one who knows the heavens is a wise man. The knowledge comes from the shadow, and the shadow comes from the gnomon.[3]

The Nine Sections. Next in order of antiquity among the mathematical works of China is the *K'iu-ch'ang Suan-shu*, or *Arithmetic in Nine Sections*.[4] This is the greatest of the Chinese classics in mathematics, and for many centuries has been held in the highest esteem in the Orient. As to its authorship and the period in which it was written we are ignorant.

[1] Y. Mikami, *China*, p. 4; W. A. P. Martin, *The Lore of Cathay*, p. 30 (New York, 1901); A. Wylie, *Chinese Researches*, Part III, p. 159 (Shanghai, 1897).

[2] This evidently refers to the right-angled triangle whose three sides are in the ratio 3:4:5, a special case of the Pythagorean Theorem.

[3] The gnomon was the index which cast the shadow on the sundial.

[4] In some editions, *K'iu-ch'ang Suan-shu-ts'au-t'u-shuo*.

We know that not long after the burning of the books (213 B.C.) there appeared a mathematician by the name of Ch'ang Ts'ang, that he collected the writings of the ancients, and that he seems to have edited the *K'iu-ch'ang Suan-shu*. There is a tradition, unsupported by positive proof, that the work was originally prepared by direction of the Chóu-Kung, who, as already stated, died in 1105 B.C., and it has even been asserted that it dates back to the reign of Huang-ti in the 27th century B.C.[1] The evidence of tradition, therefore, places it very early, and it seems probable that it existed, at least in great part, in the period of which we are writing, that is, before 1000 B.C.

Topics in the Nine Sections. The work consists, as the title says, of nine sections, books, or chapters. The titles and the sequence of chapters vary somewhat in different editions, but the following list is substantially correct as given in the revision of the work in the 2d or 3d century B.C.:

1. *Fang-t'ien* (*Squaring the farm*), relating to surveying, with correct rules for the area of the triangle, trapezium (trapezoid),[2] and circle ($\frac{1}{2}c \cdot \frac{1}{2}d$ and $\frac{1}{4}cd$), and with the circle approximations $\frac{3}{4}d^2$ and $\frac{1}{12}c^2$, where π is taken as 3.
2. *Su-mi* (*Calculating the cereals*), relating to percentage and proportion.
3. *Shuai-fen* (*Calculating the shares*), relating to partnership and the Rule of Three.[3]
4. *Shao-Kuang* (*Finding length*), relating to the finding of the sides of figures, and including square and cube roots.
5. *Shang-kung* (*Finding volumes*), relating to volumes.
6. *Chün-shu*, or *Kin-shu* (*Alligation*), relating to motion problems (couriers, hare and hound) and alligation.

[1] A. Wylie, "Jottings of the Science of Chinese Arithmetic," *North China Herald*, 1852, and the *Shanghai Almanac* for 1853; K. L. Biernatzki, "Die Arithmetik der Chinesen," Crelle's *Journal*, Vol. LII (1856).

[2] The meanings of the words *trapezoid* and *trapezium* were curiously interchanged in England and America about a century ago, and the error still persists in America. In this work the two words will be given as above, but the meaning will always be the etymological one for trapezium, a quadrilateral with two parallel sides.

[3] A kind of proportion, discussed at length in Volume II.

7. *Ying-pu-tsu*, or *Ying-nu* (*Excess and deficiency*), relating to the Rule of False Position,[1] the terms "excess" and "deficiency" relating to two concepts that are used in this rule.

8. *Fang-ch'êng* (*Equation*), relating to simultaneous linear equations, with some idea of determinants.

9. *Kou-ku* (*Right triangle*), relating to the Pythagorean Triangle.

These three works constitute the Chinese classics involving mathematics which were probably written in whole or in part before the year 1000 B.C. They show a degree of advancement quite as high as that found in the other ancient countries, and they prove that China was among the pioneers in the establishing of the early science of mathematics.

3. India

Early Hindu Mathematics. When we pass from a consideration of Chinese mathematics to the mathematics of India, Babylonia, and Egypt, we meet with the mental product of an entirely different type of people, or rather of two different types. There were two great branches of the human race affecting the Western World on the one hand, and India, Mesopotamia, and certain adjoining regions on the other hand. The first of these branches is supposed to have wandered from the Northern Grasslands, and constitutes what is known as the Indo-Europeans. In the West its members appear as Celts, Romans, and Greeks, and in Asia Minor it has several representative groups. In the East this same stock is seen in the Medes, Persians, and Hindus. The eastern branch is properly designated as Aryan, from which we have the name "Iran" for Persia. The people were generally highly imaginative, and their work in mathematics developed along such lines as the theory of numbers, geometry, and astronomy.

The second great branch is thought to have had its first habitat in the Southern Grasslands of Arabia, and is represented

[1] A primitive method of solving equations, considered at length in Volume II.

by what is known as the Semitic peoples. These include the inhabitants of Assyria, Babylonia, Phœnicia, and the Phœnician colonies. They dwelt in the paths of trade from East to West, and their work in mathematics developed chiefly along the line of computation,—a line which led to extensive numerical work in the field of astronomy as well as in that of commerce.

If the early mathematical achievements of the Chinese are uncertain as to date and importance, much more so is the early progress of the Hindus. Not only are we without any satisfactory records of the remote past of these people, but we are not infrequently confronted by claims that are preposterous and that are so recognized by Hindu scholars themselves. The first edition of the *Sūrya Siddhānta* of the Swami Press at Meerut, for example, says that the work was "Compiled about 2,165,000 years ago," representing a period about four times as long as it is thought the human race has been in existence. With even more absurdity the Laws of Manu are placed as far back as $6 \times 71 \times 4{,}320{,}000$ years ago,[1] giving almost an appearance of modesty to the ancient Chaldean claims that their astronomical observations began more than 720,000 years ago. As a matter of fact this well-known work on astronomy, the *Sūrya Siddhānta*, was probably written about the 4th or 5th century of our era. So little sympathy had the early native scholars with those outside their own caste that a general literature is wholly lacking, and it has only been through the labors of those from other lands that an all-round view of scientific progress has been attempted. There is, however, sufficient evidence for the belief that primary schools existed very early in India, and that arithmetic and writing were looked upon as the most important of the seventy-two recognized branches of learning, at least in the elementary stages of education.[2]

[1] On the extravagant ideas in the native Hindu chronology see J. C. Marshman, *Abridgment of the History of India*, p. 2 (London, 1893). See also W. Jones, "On the Chronology of the Hindus," in his *Works*, IV, 1 (London, 1807); M. Elphinstone, *History of India*, p. 136 (London, 1849).

[2] A. Hillebrandt, *Alt-Indien*, p. 111 (Breslau, 1899).

Lack of Authentic Records. As to authentic records, India has none written before the first Mohammedan invasion, *c.* 664 A.D.[1] All that we know of her earlier history is what we can glean from her two great epics, the Mahābhārata and the Rāmayāna, and from coins and a few inscriptions. The Mahābhārata relates the skill in numerals possessed by the ancient heroes, and the inscriptions tell us something of the notation used by the Hindus two thousand years ago, but neither gives us any knowledge of the period closing a thousand years before our era. The Vedas, the sacred writings of India, lead us to understand that in this period some attention was given to astronomy, as was the case in contemporary China, Mesopotamia, and Egypt.

All that we can say, therefore, about this period of Hindu mathematics is that there is some evidence from ancient literature that in very early times India paid attention to astronomy and calculation, just as was the case with other advanced peoples of that period.[2]


4. Babylon

Early Babylonian Mathematics. For our purposes Chaldea and Babylonia are synonymous, each name referring to the land extending from the delta of the Tigris and Euphrates northward to Assyria, the hilly, forest-covered district originally surrounding the ancient capital of Assur (Asshur). Indeed, it is convenient at present to consider as one large group all those Semitic peoples descended from the wanderers from the Southern Grasslands who settled in Assyria, in the region about Nineveh, in Asia Minor, and along the Phœnician coast. We shall also find it convenient to include a non-Semitic tribe, the Sumerians, who dwelt in the land of Sumer at the head of the Persian Gulf, directly in one of the chief paths of world

[1] The so-called Mohammedan period did not begin until 1001 A.D.

[2] G. Oppert, *On the Original Inhabitants of Bharatavarṣa or India*, p. 1 (London, 1893); R. C. Dutt, *A History of Civilization in Ancient India* (London. 1893).

commerce. These people, coming from the mountainous region to the east, early developed a numeral system, and numerals used by them in the 28th century B.C. are known to us through certain inscriptions. Dwelling in a low country formed by alluvial deposits, and thus deprived of stone for monumental purposes, the primitive Sumerians resorted to the use of bricks



NUMERALS OF THE 28TH CENTURY B.C.

Sumerian tablet. The numerals at this time were made with the upper end of the scribe's stylus and appear as curved symbols, and as such can easily be recognized. From Breasted's *Ancient Times*

for the preservation of their records. Upon the surface of clay tablets they pressed with a round and pointed stick, the result being a circular, a semicircular, or a wedge-shaped (cuneiform) character. These inscriptions were a mystery to the modern world until the first half of the 19th century, when Grotefend (1802) suggested and Rawlinson (1847) perfected the key

to the rich literature of ancient Mesopotamia. The clay tablets, after being inscribed, were baked by fire or in the sun, and thousands of them are now available for study in various museums. These records of the Sumerians give us the information that nearly 3000 years before Christ their merchants were familiar with bills, receipts, notes, accounts, and systems of measures. In no part of the world have we as clear evidence of commercial mathematics at this early date as is revealed by these Sumerian tablets. Here also we find evidence of an approach to a scientific calendar, although of a later date than similar evidence found in Egypt, and here is probably to be found the first use of a kind of scale of 60 in counting.

Early Calendars. Some knowledge of mathematics must, however, have long preceded the work recorded on these Sumerian tablets. The old Babylonian year began with the vernal equinox, and the first month was named after the Bull. The calendar must, therefore, have been established at a period in which the sun was in Taurus at this equinox, and such a period began about 4700 B.C. A calendar of any kind presupposes a system of numbers and some form of calculation, so that we may safely say that some kind of arithmetic existed in Babylonia in the 4th or 5th millennium B.C. Indeed, so far as the calendar is concerned, it should be said that the Sumerians celebrated the beginning of the year at the vernal equinox as early as 5700 B.C., and possibly even earlier.[1]

Early Babylonia. What is commonly known as Early Babylonia endured from about 3100 to about 2100 B.C. Sargon, the first great ruler, flourished about 2750 B.C., his remarkable career beginning in Akkad, the district just north of Sumer. It was partly due to this proximity of territory that the people of Akkad in particular and of Babylonia in general adopted the business methods, the astronomy,[2] the calendar, the

[1] H. Radau, "Miscellaneous Sumerian Texts from . . . Nippur," in the *Hilprecht Anniversary Volume*, pp. 408, 410. Chicago, 1909.

[2] E. F. Weidner, *Handbuch der Babylonischen Astronomie*, Bd. I. Leipzig, 1915.

measures, and the numerals of the more highly cultivated Sumerians. In Sargon's reign we find a record of eclipses, so that the numeral system must have been well advanced,[1] and for him there was compiled the first great treatise on astrology of which we possess any original fragments.[2]

Among the tablets of about 2400 B.C. that have been deciphered are various specimens dating from the reigns of kings of the third dynasty of Ur[3] and recording the use of a kind of draft or check, the measurement of land in *shars*, the weighing by talents (*gur*), the measurement of liquids by *ka*, the taking of interest, the use of the fractions $\frac{1}{2}$, $\frac{1}{3}$,[4] and $\frac{5}{6}$, and the measurement of both liquids and solids by the *qa* (not identical with the *ka*).

In order to fix clearly in mind the period of which we are speaking there should be mentioned not only the reign of Sargon (*c.* 2750 B.C.) but the remarkable reign of Hammurabi or Hammurapi (*c.* 2100 B.C.), in which the world's first great code of laws, so far as we know, was written, and in which the calendar was reformed. Among the other interesting relics of the time of Hammurabi is the ruin of the oldest known schoolhouse. This was discovered by French archeologists in 1894.[5] In the building were numerous tablets on which the pupils had written their lessons, and it is from such tablets as these that we have part of our knowledge of the arithmetic of the Babylonians.

The general conclusion of archeologists, as will be elaborated on page 40, is that these early Babylonians (in the thousand years of their activity) developed a fair knowledge of computation, of mensuration, and of commercial practice, in spite of an awkward numeral system by which they were handicapped.

[1] See also F. Thureau-Dangin, in the *Hilprecht Anniversary Volume*, p. 156.

[2] G. Bigourdan, *L'Astronomie*, 1920 ed., p. 27. Paris, 1911.


[3] G. A. Barton, *Haverford Library Collection of Cuneiform Tablets*, Part I (Philadelphia, n.d [1905]); *ibid.*, Part II (1909).

[4] These from Barton, *loc. cit.*, Part I. On the taking of interest, the rates running from 20% to $33\frac{1}{3}$%, see E. Huber, "Die altbabylonischen Darlehnstexte," in the *Hilprecht Anniversary Volume*, pp. 189, 217.

[5] For a plan of the building see J. H. Breasted, *Ancient Times*, p. 136 (Boston, 1916); hereafter referred to as Breasted, *Anc. Times*.

Early Assyria. As early as 3000 B.C. a Semitic tribe of nomads settled at Assur, and in due time it too adopted the Sumerian calendar and such of the mathematics of trade as had been developed by these people of the south.

Much later, and after 1200 B.C., the Arameans, or Syrians, established kingdoms in the region to the west of Assyria. They were great merchants, and the Sumerian mathematics of trade, which had worked slowly northward through Babylonia and Assyria, now found place in the new territory. We have bronze weights of this period, showing that whole numbers, fractions, measures, and elementary forms of computation played a considerable part in the daily life of the people.



ARAMEAN WEIGHT FOUND IN ASSYRIA

The weight is of bronze and the inscription is Aramaic. Fifteen of these lion weights were found in Nineveh and testify to the common presence of Aramean merchants in Assyria. From Breasted's *Ancient Times*

Early Chaldea. The desert tribe called the Kaldi came into prominence long after the period now under discussion. It gained a foothold in ancient Sumer and finally (606 B.C.) conquered the Assyrians and established the Chaldean empire in the region of Babylonia. Although their empire lasted only to 539 B.C., they made great progress in science. In particular, astrology was extensively cultivated, the equator was probably divided into 360°, the twelve signs of the zodiac definitely appeared, and mathematics flourished as the handmaid of commerce and astronomy. Thus Babylonia became Chaldea, and Chaldea became the patron of science and art.

Early Cuneiform Tablets. Our first important knowledge of Babylonian arithmetic was derived from two tablets found in 1854 at Senkereh, the ancient Larsam or Larsa, on the Euphrates, by a British geologist, W. K. Loftus. These tablets

contain the squares of numbers from 1 to 60 and the cubes of numbers from 1 to 32.[1] Their date is uncertain, but the evidence seems to show that they were of about the Hammurabi period (*c.* 2100 B.C.).

Since the discovery of the Senkereh tablets there have been unearthed some 50,000 tablets at Nippur, the modern Nuffar, an ancient city lying to the south of Babylon, and among these are many that relate to mathematics.[2] They are apparently from a large library which seems to have been destroyed by the Elamites about 2150 B.C. or a little earlier, and again about 1990 B.C., and they constitute the most extensive mass of ancient mathematical material ever brought to light. The cylinders include multiplication and division tables, tables of squares and square roots, geometric progressions, a few computations, and some work on mensuration. Neugebauer's studies (1935) of a large number of tablets show that the Sumerians and Babylonians could solve special linear, quadratic, cubic, and biquadratic equations and had some knowledge of negative numbers.

Babylonian Geometry. The tablets found at Nippur and elsewhere also give us some knowledge of the Babylonian geometry. From these it seems that as early as 1500 B.C. the Babylonians could find the area of a rectangle, including that of a square; the area of a right-angled triangle; the area of a trapezium (trapezoid); and possibly the area of a circle, the volume of a parallelepiped, and the volume of a cylinder. There is ground for the belief that they knew the law of expansion of $(a+b)^2$, although we have no knowledge as to whether this was inferred from a geometric figure or from their extensive study of square numbers. There is also some reason to believe that they knew the abacus, since it has been suggested that one of their signs (ŠID) may have been derived from a pictograph of such an instrument.

[1] Apparently from 1 to 60 originally, but part of the tablet is broken off.

[2] H. V. Hilprecht, *Mathematical, Metrological, and Chronological Tablets from the Temple library of Nippur* (hereafter referred to as Hilprecht, *Tablets*). Philadelphia, 1906.

Scale of Sixty. One peculiarity of Babylonian arithmetic is the constant use of the number 60,—a use which finally suggested the development of sexagesimal fractions and which still survives in our division of degrees, hours, and minutes into sixty sub-units. It is generally thought that the Babylonians, interested as they were in watching the stars, early came to believe that the circle of the year consisted of 360 days. It is also thought that they knew that the side of the regular inscribed hexagon is equal to the radius of the circle, this property suggesting the division of 360 into six equal parts, and 60 being thus looked upon as a kind of mystic number. This may, indeed, be the origin of this use of 60, but we find other nations using 40, 20, and even 15 in somewhat the same way, with no apparent reason, so that all such customs may have developed from racial notions which were started by some leader or sect with no particular reason in mind. It is more probable that 60 was chosen because of its integral divisors 2, 3, 4, 5, 6, 10, 12, 15, 20, and 30, thus rendering work with its fractional parts very simple.

Although the subject of fractions with the denominator 60 is discussed in Volume II, a brief mention may be made at this time of an important tablet first described in 1920.[1] It dates from *c.* 2000 B.C. and illustrates the Babylonian custom of using either 360 or 60 for the denominator except in the cases of unit fractions and of fractions in which the numerator is 1 less than the denominator. For example, $\frac{6\,0}{8\,6\,0}$ may appear as $\frac{1\,0}{6\,0}$ or as $\frac{1}{6}$ (ŠUŠŠU), $\frac{2\,4\,0}{3\,6\,0}$ as $\frac{4\,0}{6\,0}$ or $\frac{2}{3}$ (ŠINIPU), and $\frac{3\,0\,0}{3\,6\,0}$ as $\frac{5\,0}{6\,0}$ or $\frac{5}{6}$ (PĀRAB).

5. Egypt

Early Egyptian Mathematics. Whatever claims may properly be made for the antiquity of mathematics in various other countries, claims of even greater validity can justly be made for the science in Egypt. Civilization has generally developed along great rivers; the Nile is one of the world's greatest arteries

[1] H. F. Lutz, "A mathematical cuneiform tablet," *American Journal of Semitic Languages*, XXXVI, 249.

of commerce, and its fertile valley is one of the world's greatest gardens. Egypt was a well-protected country, and civilization had a more favorable opportunity for uninterrupted development there than in such lands as Mesopotamia, Phœnicia, India, and China. Furthermore, her art, as shown by wall sculptures, was much farther advanced in the 4th millennium B.C. than it was, say, with the Sumerians, and so there is every reason to feel that her science was also in the lead of that in other lands.

The earliest dated event in human history is the introduction of the Egyptian calendar of twelve months of thirty days each, plus five feast days, in the year 4241 B.C.[1] Such an achievement as the creation of this calendar, a better one than was used in Europe from the time of the Romans until the reform of Gregory XIII (1582), and in some respects better than the one used at present, shows a high development of computation as well as of astronomy. No authentic record of mathematical progress in any other country dates back as far as this; it reaches back even into the Stone Age, more than a thousand years before the earliest stone masonry and long before any people had the slightest idea of an alphabet as we understand the term. An event like this is a silent but powerful witness to the noteworthy arithmetic attainments of its sponsors and to a long series of scientific observations by the temple astronomers. Furthermore, our own calendar may be said to be merely a poor adaptation of this ancient Egyptian one, although containing the great improvement of having as centennial leap years only those of which the numbers representing the hundreds are divisible by four.

Third Millennium B.C. in Egypt. When we approach the year 3000 B.C., toward the close of the Second Dynasty, we find ourselves in a period of rapid development in practical engineering. We have no manuscripts of this period from which to obtain direct information, but in the achievements of the engineers it is possible to recognize a number of interesting

[1] Breasted, *Anc. Times*, p. 45.

facts. Professor Breasted has characterized the development of civilization in the 30th century B.C. in these words:

Hardly more than a generation before this 30th century the first example of hewn stone masonry was laid, and in the generation after this 30th century the Great Pyramid of Gizeh was built. With amazingly accelerated development the Egyptian passed from the earliest example of stone masonry just before 3000 B.C. to the Great Pyramid just after 2900. The great-grandfathers built the first stone masonry wall a generation or so before 3000 B.C., and the great-grandsons erected the Great Pyramid of Gizeh, within a generation after 2900. . . .

One finds it difficult to imagine the feelings of these earliest architects . . . as they paced off the preliminary plan and found an elevation in the surface of the desert which prevented them from sighting diagonally from corner to corner and applying directly a well-known Egyptian method of erecting an accurate perpendicular by means of measuring off a hypotenuse. . . .

The Egyptian engineers early learned to carry a straight line over elevations of the earth's surface, or a plane around the bends of the Nile. In his endeavor to record the varying Nile levels in all latitudes the Egyptian engineer was confronted by nice problems in surveying, even more exacting than those which he met in the Great Pyramid. A study of the surviving nilometers has disclosed the fact[1] that their zero points, always well below lowest water, are all in one plane. This plane inclines as does the flood slope from south to north. The Pharaohs' engineers succeeded in carrying the line in the same sloping plane, around innumerable bends in the river for some seven hundred miles from the sea to the First Cataract.[2]

Accuracy of Early Engineers. Such was the degree of accuracy secured by these early surveyors that Petrie found the maximum error in fixing the length of the sides of the Great Pyramid to be only 0.63 of an inch, or less than $\frac{1}{14000}$ of the total length, and the angle error at the corners to be 12″, or only $\frac{1}{27000}$ of a right angle.

[1] L. Borchardt, *Nilmesser und Nilstandsmarken.*

[2] J. H. Breasted, "The Origins of Civilization," *The Scientific Monthly*, X, 87.

Speculations on the Great Pyramid. As to the speculations relating to the Great Pyramid it is possible to make only a brief statement in this work. That mathematics, and possibly mathematical mysticism of some kind, played an important part in the design of the structure is admitted by all scholars, but precisely what the dominating principle was we do not know. It has been suggested that four equilateral triangles were put together for the pyramidal surface, but this theory is not borne out by measurements, the base being considerably longer than the sloping edge.

A second theory asserts that the ratio of the side to half the height is the approximate value of π, or that the ratio of the perimeter to the height is $2\,\pi$. It is true that this would give the value of π as about 3.14, an approximation that may have been known to the pyramid builders; but this was possibly a mere matter of chance. If one searches in any building, or indeed in any given object, for lines having this ratio, they are not difficult to find. Nevertheless, there probably is some mysticism of this kind in the proportions of the structure.

A third theory makes the claim that the angle of elevation of the passage leading to the principal chamber determines the latitude of the pyramid, approximately 30° N., or that the passage itself pointed to what was then the pole star; but even after making all reasonable allowances in favor of this hypothesis, the angular difference is too great to make out a very strong case.

It is also claimed that the pyramids have a constant angle of slope, and it is true that the three at Gizeh vary but little, being approximately 51° 51′, 52° 20′, and 51°; but others have slopes running from about 45° to 74° 10′.

Recent measurements have been so accurately made that it is probable that further study will reveal in the near future whatever mathematical principles actuated the architects. For the present we may simply dismiss the speculations of such men as Charles Piazzi Smyth[1] as interesting rather than scientific.

[1] An English astronomer; born at Naples, 1819; died 1900; astronomer royal of Scotland (1845–1888). *Our Inheritance in the Great Pyramid* (1864), *Life and Work at the Great Pyramid*, 3 vols. (1867).

Testimony of the Wall Reliefs. The wall reliefs of this general period of the Pyramid builders testify to the collection of taxes, probably in the form of grain, and the issuing of receipts by the officials of the king. Nothing so tangible, showing the applications of elementary arithmetic at this early period, has been found in any other region except Sumeria.



COLLECTION OF TAXES, *c.* 3000 B.C.

Showing the clerks and scribes at the right, with pen and papyrus, and the officials and taxpayers at the left. From Breasted's *Ancient Times*


Reign of Amenemhat III, or Moeris. About 1850 B.C., in the 12th Dynasty, there came to the throne one of the most energetic of all the kings of Egypt, Amenemhat III.[1] In his reign there was carried out an extensive system of irrigation, necessitating a knowledge of leveling, surveying, and mensuration such as had probably never been developed before this time in any other part of the world, except perhaps in Mesopotamia.[2] There is good reason to believe that in this reign, say about 1825 B.C., there was written the original of the oldest elaborate manuscript on mathematics now extant, the Ahmes treatise mentioned a little later. If the conjecture is correct, the unit fraction was already known in Egypt, as also seems to have been the case in Mesopotamia,[3] and the simple equation with a fairly usable symbolism,

[1] The name also appears as Ne-mat-re and as Amenemha. He is the Moeris of Herodotus (II, 148–150), the Marros of Diodorus Siculus (I, 52), and the Mares of Eratosthenes. He is also referred to by Strabo (XVII), Pliny (*Hist. Nat.*, V, 9, 50, and XXXVII, 12, 76), and Pomponius Mela (I, cap. 9). Recent Egyptologists give the date of his reign as 1849–1801 B.C.; others place his reign as *c.* 1986–*c.* 1942.

[2] On the Egyptian work in this line see A. Wiedemann, *Aegyptische Geschichte*, I, 256 (Gotha, 1884); J. Lieblein, "L'Exode des Hébreux," *Proceedings of the Society of Biblical Archæology*, XXI (London, 1899), 55. On the claims of priority in the general development of mathematics, see E. Weyr, *Ueber die Geometrie der alten Aegypter*, p. 4 (Vienna, 1884).

[3] See the author's review of Hilprecht's work in the *Bulletin of the American Mathematical Society*, XIII (2), 392.

arithmetic and geometric series, and the elements of mensuration were already familiar to the élite among the mathematicians of the Nile Valley.



THE OLDEST ASTRONOMICAL INSTRUMENT KNOWN

The original is in the Berlin Museum. Part *A* was a plumb line. By its aid the observer could hold *B* over a given point and sight along the slot to some object like the North Star, thus establishing a meridian line. From Breasted's *Ancient Times*

Mathematics of the Feudal Age. Amenemhat III lived in the so-called Feudal Age of Egypt,—a period which lasted for several centuries, closing about 1800 B.C. To this period belongs the oldest astronomical instrument extant, a forked stick used in sighting for the purpose of obtaining the meridian. Such a work presupposes some ability in calculation and in constructive geometry, and the existence of this ability is still further proved by the Ahmes treatise. There is another treatise, written much earlier than this,[1] in which we find mention of the civil calendar of twelve months of thirty days each, plus five extra days, as already mentioned.

About the close of the Feudal Age a postal service existed in Asia under Egyptian control, requiring some means of payment on the part of those whose convenience it served. At the same time census lists were prepared for use in the taxation of the people, surveys for irrigation projects were made, and the Nilometer served to foretell the beginning and the end of the rise of the river, all of which involved,

[1] That is, in the 30th century B.C. See E. Mahler, "Der Kalender der Babylonier," in the *Hilprecht Anniversary Volume*, pp. 1, 9.

as already stated, the use of a considerable amount of mensuration and computation, and adds to the evidence of an interest in mathematics in this period of Egyptian history.

In the period of the Middle Kingdom (2160–1788 B.C.) business arithmetic was such as to demand bills, accounts, and tax lists. From this period we have various fragments of papyrus rolls which were found in the remains of the libraries of the feudal lords. These are the oldest libraries of papyrus rolls thus far known. Among these remains were found such evidences of the commercial activity above mentioned as are seen in the fragments of papyri found at Kahun and now in London and Berlin.[1]


Ahmes Papyrus. About 1650 B.C. there lived in Egypt a scribe named A'h-mosè, commonly called by modern writers Ahmes.[2] He wrote a work on mathematics; or rather he copied an older treatise, for he says: "This book was copied in the year 33, in the fourth month of the inundation season, under the majesty of the king of Upper and Lower Egypt, 'A-user-Rê', endowed with life, in likeness to writings of old made in the time of the king of Upper and Lower Egypt, Ne-ma'et-Rê'. It is the scribe A'h-mosè who copies this writing." Another manuscript of the same period, containing a number of lines on fractions, is in the British Museum. It was published in 1927. The actual manuscript[3] of Ahmes has come down to us, having been purchased in Egypt about the

[1] The earliest date in the London fragments is in the reign of Amenemhat III. See W. M. Flinders Petrie, *Kahun, Gurob, and Hawara*, chap. vi by F. L. Griffith (London, 1890). This article places the date *c.* 1986–1942 B.C., which is somewhat later than that given by the earlier writers and a little earlier than that given by some of the latest authorities.

[2] "A'h-mosè" was also the name of certain kings. Ahmes I, often known as Amosis or Amasis, came to the throne at the beginning of the 18th Dynasty, when Egypt entered upon a period of empire, and it was he who expelled the Hyksos and pursued them into Palestine.

[3] Since the first edition there have appeared editions of the Rhind Papyrus by T. E. Peet (London, 1923) and A. B. Chace (Oberlin, O., 2 vols., 1927, 1928). The Chace edition is the more elaborate, containing a facsimile of the papyrus, a transcription into hieroglyphic and Latin characters, a complete translation, numerous notes, and an extensive bibliography by R. C. Archibald. The preferred form of the name is given as A'h-mosè and the date as between 1750 and 1580.

middle of the 19th century by the English Egyptologist, A. Henry Rhind (whence the name "Rhind Papyrus"), and having later been acquired by the British Museum. It is one of the oldest mathematical manuscripts on papyrus extant.



A PAGE FROM THE AHMES PAPYRUS

Written *c.* 1550 B.C. The original is in the British Museum

The Ahmes manuscript is not a textbook, but is rather a practical handbook. It contains material on linear equations of such types as $x + \frac{1}{7}x = 19$; it treats extensively of unit fractions; it has a considerable amount of work on mensuration, and it includes problems in elementary series.[1] The

[1] The British Museum published an inexact facsimile of the papyrus in 1898 under the title *Facsimile of the Rhind Mathematical Papyrus*. The standard works on the subject are those of Chace and Peet, mentioned on page 47, and (less valuable) that of A. Eisenlohr, *Ein mathematisches Handbuch der alten Aegypter*, 2d ed. (Leipzig, 1877). See also F. L. Griffith, *Proceedings of the Society of Biblical Archæology*, 1891, 1894; A. Favaro, *Atti della R. Accad. . . . in Modena*, Vol. XIX.

internal evidence shows the work to be a compendium of the contributions of at least two or three authors.[1]

Evidence of Commercial Mathematics. About 1500 B.C. there was built by Queen Hatshepsut[2] the temple known at present as Der al-Bahri. This is not far from Thebes[3] and in 1904 was uncovered and made known to modern scholars. On the walls of this temple is pictured the receipt of tribute from the land of Punt, probably on the Somali coast of Africa,[4] and mention is made of "reckoning with numbers, summing up in millions, hundreds of thousands, tens of thousands, thousands, and hundreds," showing the extent to which numbers were used in commercial matters even before coins were invented. There are certain inscriptions of the same period in the tomb of Rekhmire, at Thebes, giving the tax list of Upper Egypt, and interesting because of the fact that the highest number is 1000 and that $\frac{1}{2}$ is the only fraction used.[5]

Oldest Sundial. From this period or a little later, but from about 1500 B.C., there dates the oldest sundial extant, an Egyptian piece now in the Berlin Museum, showing that the Egyptians had already developed, as we might have inferred from the other mathematical and astronomical knowledge possessed by them, a good system of timekeeping by means of a primitive sun clock. On this clock the shadow shortened as the forenoon advanced, and lengthened from noon to night.

[1] This is seen particularly in the several rules which are evidently followed in the formation of unit fractions.


[2] Hât-shepset, Hatasu, or Hatshepsu, also known as Ramaka (Ma-ka-ra). See E. A. W. Budge, *The Mummy*, p. 30 (Cambridge, 1893).

[3] The No Amon (City of Ammon) of the Bible, also known to the Greeks as Diospolis.

[4] J. H. Breasted, *Ancient Records, Egypt* (Chicago, 1906, 1907), II, pp. 104, 114, 210, 211; IV, pp. 362 seq.; hereafter referred to as Breasted, *Anc. Records*. See also official accounts of the same period in facsimile in Golénischeff, *Les Papyrus hiératiques nos. 1115, 1116A et 1116B de l'Ermitage impérial à St Pétersbourg*, 1913.

[5] Breasted, *Anc. Records*, II, 283. In later inscriptions, as of *c.* 600 B.C., the fractions $\frac{1}{6}$ and $\frac{2}{3}$ appear (*ibid.*, IV, p. 486). Still later, from 19 A.D. to 250 A.D., the papyri tell us of the periodic census introduced apparently by Augustus, with taxes and the records of imports and exports. See A. S. Hunt, "Papyri and Papyrology," *Journal of Egyptian Archæology*, I (1914).

There were six hours in the forenoon and six in the afternoon, from which division of the day came the system of twelve hours later adopted in Europe. Such clocks, in various forms, were afterwards used by the Greeks and gave rise to the sundials of



OLDEST SUNDIAL EXTANT

Egyptian specimen, restored after Borchardt, now in the Berlin Museum. Dates from *c.* 1500 B.C. In the morning the crosspiece was turned to the east, and in the afternoon to the west. From Breasted's *Ancient Times*

later times. The clock above shown bears the name of Egypt's greatest general, Thutmose III, who has justly been called her Napoleon.

Practical Problems. By the time of Seti I (*c.* 1350 B.C.) business calculation had come to require larger numbers than those needed in the time of Ahmes. This is seen from the problems in the Rollin papyrus manuscript now in the Louvre,[1] one of which, line for line as in the text, is as follows:

1601 392,325
together bread 107,893 makes in *ten* 364,371
bread 6121 loaves 1800 *thes* makes in *ten* 21,600
together 385,871
rest 6354
quantity of maize sacks 1601 makes in bread 112,090
makes in *ten* 392,306
brought to the magazine bread 114,064 makes in *ten* 385,971

The meaning is that here are two accounts of 1601 sacks of wheat each, the produce varying in the two cases. The weights

[1] M. F. Chabas, *Aegyptische Zeitschrift*, 1869, p. 85. The manuscript was published by W. Pleyte in 1868, and a new translation by Eisenlohr appeared in 1897 in the *Proceedings of the Society of Biblical Archæology*, XIX, 91, 115 147, 252.

are calculated in *thes* or *ten*, 12 *ten* making 1 *thes*, a *ten* being about 316 grams. In the first case a loaf of unbaked bread weighed 3.63 *ten*, 1.15 kg., or $2\frac{1}{2}$ lb., and after it was baked it weighed 3.37 *ten*, 1.06 kg., or $2\frac{1}{3}$ lb. But the first case also gives the weight of 6121 loaves as 21,600 *ten*, which is at the rate of 3.52 *ten* per loaf when baked, so that the sizes evidently varied. In the second case the bread weighed 3.55 *ten* per loaf, possibly unbaked, and 3.38 *ten* per loaf when delivered. The first account may be represented as follows:

107,893 loaves weigh	364,371 *ten*
6121 loaves weigh	21,600 *ten*
together they weigh	385,971 *ten*
there is left	6,354 *ten*
the total being	392,325 *ten*

The problem in itself is of little moment except as it shows the practical use of large numbers in these early times.

Rameses II divides the Land. At the close of Seti's relatively short reign his son, Rameses II (*c.* 1347 B.C.), known to the Greeks as Sesostris, came to the throne. In his reign a redivision of land took place among the people, and surveying must have attracted much attention.

Herodotus (*c.* 484–*c.* 425 B.C.), referring to information that he had received from the priests, relates the following:

> Sesostris also, they declared, made a division of the soil of Egypt among the inhabitants, assigning square plots of ground of equal size to all, and obtaining his chief revenue from the rent which the holders were required to pay him every year. If the river carried away any portion of a man's lot, he appeared before the king, and related what had happened; upon which the king sent persons to examine, and determine by measurement the exact extent of the loss; and thenceforth only such a rent was demanded of him as was proportionate to the reduced size of his land. From this practice, I think, geometry first came to be known in Egypt, whence it passed into

Greece. The sundial, however, and the gnomon, with the division of the day into twelve parts, were received by the Greeks from the Babylonians.[1]

Harris Papyrus. Rameses IV came to the throne *c.* 1167 B.C. and immediately prepared a remarkable document setting forth the great works of his father, Rameses III (1198–1167 B.C.), including a list of his extensive gifts to the gods. The list shows the proportion of the wealth of ancient Egypt held by the temples and is of value in giving the numerals of the period. This document, known as the Harris Papyrus, is still extant[2] and affords the best example of practical accounts that has come down to us from the ancient world.

That surveying played a prominent part in the life of Egypt is seen in an inscription on the tomb of Penno at Ibrim, in Nubia, in the reign of Rameses VI (*c.* 1150 B.C.), in which the boundaries and areas of five districts are given.[3]

Evidence of Egypto-Cretan Relations. Thus we see that before 1000 B.C. Egypt had developed enough knowledge of astronomy to devise an excellent calendar, and that she was in possession of a commercial system requiring extensive work in computation, of an elaborate scheme of leveling and surveying, of a considerable knowledge of what we would now consider as a kind of algebra, and of some ability in mensuration, especially as it related to granaries and to the use of grain products in the making of bread.

Recent excavations have shown the existence of a high degree of civilization in Crete in this period of progress in Egypt, and there is also evidence of amicable relations between these two countries in early times. Our knowledge of the subject is too limited, however, to determine whether it has any bearing upon the history of mathematics. The deciphering of the Cretan inscriptions still awaits the further efforts of scholars.

[1] Herodotus, II, 109.

[2] Breasted, *Anc. Records*, IV, 127 seq. See also S. Birch, *Zeitschrift für Aegyptische Sprache*, pp. 119 seq. (1872).

[3] Breasted, *Anc. Records*, IV, 233. For the Egyptian measures in common use at this period, and thus far identified as to equivalents, see *ibid.*, p. 88.

TOPICS FOR DISCUSSION

1. The countries in which mathematics flourished prior to 1000 B.C., and the reasons for this mathematical activity.

2. Reasons for supposing mathematics to have made some progress in the Late Stone Age, or even earlier.

3. Influences leading to an extension in the use of mathematics in the third millennium B.C.

4. Probable nature of the earliest mathematics of China, and the influences which developed the study of this science.

5. General nature of the early written mathematical works in China, with approximate dates.

6. The first traces of number mysticism in the East.

7. General period in which the *Nine Sections* was written. Nature of the work.

8. Probable nature of the early Hindu mathematics.

9. Influences that developed Babylonian mathematics and the method of recording the science.

10. General nature of Babylonian mathematics.

11. Evidence of early mathematics in Egypt. General nature of the work in the earliest periods.

12. Mathematics of the Feudal Age in Egypt.

13. The Ahmes Papyrus, its origin and general nature.

14. Evidences of development of commercial arithmetic between the time of Ahmes and 1000 B.C.

15. Types of problems in arithmetic, algebra, and mensuration that interested the ancient Egyptians.

16. Comparison of the mathematical progress and interests of China, India, Babylonia, and Egypt in early times.

17. A consideration of the reasons why this period was lacking in power to advance its mathematics.

18. A study of the evidence of mathematics in Crete and Cyprus before 1000 B.C., and the influence of this mathematics upon Greek science.

19. The evidences of interrelation of mathematical ideas in Mesopotamia, Egypt, and the islands of the Mediterranean Sea.

20. Mathematical and astronomical instruments of this period.

21. The degree of accuracy apparently secured by engineers before the year 1000 B.C.

CHAPTER III

THE PERIOD FROM 1000 B.C. TO 300 B.C.

1. The Occident in General

Geographical Limits. For our present purposes we may define the Occident of the period from 1000 B.C. to 300 B.C. as practically identical with Greece and her colonies. Whatever mathematics Rome had in this period was essentially Greek, and most of the Mediterranean world, aside from the hinterland of Phœnicia and Egypt, may therefore be conveniently classified as under the influence of the Hellenic civilization. Phœnicia contributed little that was not commercial, and the golden age of Egypt was already past.

Protected Regions. Philosophy, letters, mathematics, art, and all the finer products of the mind require peaceful surroundings for their development. It is for this reason that mathematics at this time flourished best on the protected islands of the Ægean Sea, on the Greek peninsula, and in the Greek towns of Southern Italy. In all these places invasion was difficult and the rewards of the invader were few. Commercial and intellectual communication with the rest of the world was possible, so that peace without stagnation was, relatively speaking, assured.

It would also be proper to include some mention of the mathematics of Mesopotamia, since this was quite as occidental as oriental; but aside from its use in astronomy the science was not sufficiently in evidence in Babylon at this time to demand our attention.

Chronological Limits. The reason for taking the lower arbitrary limit of 300 B.C. is that a new era in the history of mathematics begins with the founding of the Alexandrian School at about that time. This event led to a reshaping of

mathematics either through the efforts of scholars connected with the first great cosmopolitan university or through the works written by those who came under their influence.

2. The Greeks

Birth of Greek Arithmetic. Commercial arithmetic was well advanced in various neighboring states long before it was known in Greece. The merchants of the Phœnician coast (along and across which passed the routes of trade with the Orient), receiving inspiration from Babylon, early developed a fairly good business arithmetic, and in due time became the teachers of this art in Egypt, Asia Minor, and the Ægean isles. The recent excavations in the ancient palace of Knossos show that in early times the commercial arithmetic of Babylon reached even as far west as Crete, and future studies are likely to reveal much valuable information relating to this island. Indeed, in what is called the Early Minoan Period of Crete, Greece was still a forest, thinly peopled by a nomad race, for this was long before the warlike Dorians, about a thousand years before our era, made themselves masters of Peloponnesus and changed the whole tenor of Greek life. Thucydides describes the country at this early period as a theater of frequent migrations, when "each man cultivated his land only according to his immediate needs, with no thought of amassing wealth." Under such conditions, before the coining of money was known, only the most primitive arithmetic was demanded. A little counting and a little rude barter were all for which the ancient Greek civilization had created a need. Even at a much later period than the one we have described, Greece was little inclined to commerce. Her older cities were not generally seaports, and what little navigation she had was concerned with war and piracy rather than with the development of trade.

External Influences. It was only when the Greeks began to come into closer contact with other peoples that they showed any interest in arithmetic. Indeed, contrary to the idea that is commonly expressed, Greece always depended largely upon

external influences for her mathematics, and few who advanced this science in her schools were born within her continental area. But when we speak of the early efforts of Greece to put herself in contact with the external world through colonization, it must be understood that we are still ignorant as to how far peninsular Greece was then a colonizer and how far she herself was a colony, since her people lived as much along the Asiatic as along the European coast.

Miletus. We are told by Herodotus (*c.* 484–*c.* 425 B.C.) and Strabo (*c.* 66 B.C.–*c.* 24 A.D.), however, that Miletus, the greatest commercial town of the twelve forming the Ionian confederacy, was an Athenian colony, although there are good reasons for doubting the statement. Situated at a strategic point on the coast of Asia Minor, it in turn became a great colonizing center, and in the 7th century B.C. established no less than ninety towns along the shores of the Black Sea and the Mediterranean, even opening Egypt to her commercial settlements. This fact had a bearing upon the early science of the Greeks, since it was at Miletus that their mathematics had its beginning; and it was here, doubtless, that their commercial arithmetic first developed to any great extent. It was in Lydia, just east of here, that coins were first struck in the West, in the 7th century B.C., and Miletus at once recognized and adopted the new invention, anticipating Athens, indeed, by over half a century. The influence of this movement, particularly in relation to arithmetic, is evident. Without the aid of coins all business calculation must have been very cumbersome, money consisting of bars or ingots of metal that had to be weighed, and small currency being practically nonexistent save in the form of shells or trinkets. We can therefore determine fairly well the time and place of the beginning of any noteworthy business arithmetic among the Greeks, namely, about the 7th century B.C. and along the coast of Asia Minor.



ANCIENT COINS

Coins found in Asia Minor. They are among the earliest known, dating from about 550 B.C. From Breasted's *Ancient Times*


Logistic. At the period of which we have been speaking, the Greek science of numbers, the arithmetic proper, had not yet been invented. Only the art of calculating had made any appeal to these practical people. This branch of the subject went by the name of "logistic," and its beginnings must be sought in prehistoric times. Greek tradition states that it came from the Phœnicians, whose trading instincts are well known, and many comparatively recent writers have felt that this tradition had a foundation in fact. It must not be thought, however, that the interesting properties of numbers were entirely unrecognized before this time. Various curious relations had been the subject of discussion in the Orient for many centuries, and some knowledge of number mysticism had doubtless been acquired by the priestly caste in Greece long before logistic existed as a special subject of study.

Although by the tradesman in Miletus, and later in Corinth and other seaport towns, logistic must have been looked upon as important, it is probable that the ordinary Greek could neither multiply one number by another nor perform any other operation in what we now call arithmetic. There were doubtless schools at that time, for Herodotus (*c.* 450 B.C.) and Diodorus Siculus (1st century B.C.) both speak of them as then known, but logistic was looked upon as a technicality of trade, just as we may today look upon the use of a slide rule or a typewriter. A little later, however, it came more into favor, for Plato refers to it rather than to the theoretical part of the science when he says:

> Very unlike a divine man would be he who is unable to count, "one, two, three," or to distinguish odd and even numbers. . . . All freemen, I conceive, should learn as much of these branches of knowledge as every child in Egypt is taught when he learns his alphabet. In that country arithmetic games have actually been invented for the use of children, which they learn as a pleasure and amusement.

Furthermore, Plato recommends the use of apples and other objects in the presenting of the idea of number, quite as a modern teacher would employ them.

In spite of the extensive use of logistic among the Greek merchants, not a single treatise upon the subject remains. A Greek multiplication table, written on wax at about the beginning of our era, and hence somewhat later than the close of the period under discussion, is still preserved in the British Museum; and this, together with a few examples in addition, subtraction, and multiplication, and an abacus, are all that have come down to us that bears directly upon the practical



GREEK MULTIPLICATION TABLE ON A WAX TABLET

One of the few examples of the Greek logistic. This specimen is now in the British Museum and dates from about the beginning of the Christian era

computations of the Greeks. To these examples reference will be made when we come to consider the abacus and the various operations.

Arithmetic. Although the precise nature of the Greek logistic, the art of calculating, is very little known, fortunately the same cannot be said of the Greek arithmetic, the theory of numbers. As a subject for philosophers and by them committed to writing, it has come down to us as it was left by the later Greeks, and probably with its details little changed from the original form given to them in the earlier days. This topic, relating to the remote ancestor of our present number theory, will be considered, together with logistic, in Volume II.

Greek Geometry. Although both logistic and arithmetic developed in the Orient as well as in the Occident, geometry as a logical science is purely a product of the western civilization. On the other hand, intuitive geometry is universal, differing as a matter of course in degree of accomplishment in the various parts of the world. Egypt possibly knew the law of the square on the hypotenuse of a right-angled triangle long before Pythagoras, and there is reason to believe that China and India were also familiar with it; but the first proof of the theorem, and apparently the first idea of a geometric proof, are both due entirely to the Greeks. Indeed, we may say that all of our geometry, considered as a logical sequence of propositions, whether relating to two-dimensional or to three-dimensional space and whether limited to circles and straight lines or extended to include conic sections and higher plane curves, had its origin solely in the Greek civilization. So completely was the Greek mathematics given over to geometry that both arithmetic and the science that was much later known as algebra were treated almost entirely from the geometric standpoint. We shall therefore see that, although mathematics among the Greeks included geometry, arithmetic, logistic, music, and a kind of algebra, the central element was geometry.

Centers of Mathematical Activity. Mention has been made of Miletus, and before proceeding farther it is desirable to locate the other centers of mathematical activity in Greece and her colonies. The following cities and countries will be mentioned frequently, the numbers referring to the map on page 60:

Abdera, 16.
Alexandria, 41.
Amisus, 31.
Antinoopolis, 43.
Apameia, 35.
Aquitania, 2.
Athens, 13.
Byzantium, 28.
Cadiz, 1.
Chalcedon, 29.
Chalcis, 37.
Chios, 17.
Clazomenæ, 24.
Cnidus, 22.
Constantinople, 28.
Crete, 20.
Crotona, 9.
Cyprus, 34.
Cyrene, 11.
Cyzicus, 27.
Elea, 8.
Elis, 12.
Gades, 1.
Gerasa, 39.
Jerusalem, 40.
Laodicea, 32.
Larissa, 36.
Medma, 7.
Mendes, 42.
Miletus, 23.
Naples and Pompeii, 6.
Nicæa and Bithynia, 30.
Paros, 19.
Perga, 33.
Pergamum, 26.
Ptolemais, 44.
Rhodes, 21.
Rome, 3.
Samos, 18.
Sicily, 4.
Smyrna, 25.
Stageira, 14.
Syene, 45.
Syracuse and Messina, 5.
Tarentum, 10.
Thasos, 15.
Tyre, 38.



MATHEMATICAL-HISTORICAL MAP OF THE MEDITERRANEAN COUNTRIES IN CLASSICAL TIMES

This map shows the location of places most frequently mentioned in relation to Greek and Roman mathematics, with the names (page 61) of scholars connected with them. The numbers are arranged on the map from left to right. The dates following the names are merely approximate, and those before the Christian era are indicated by the letters B.C., as in the text. France and Spain are not shown, because they are mentioned only with respect to Cadiz (Gades) and Aquitania (in southern France), and to include them would reduce in size the more essential parts of the map. It will be observed that, as stated in the text, mathematics flourished best in territory that was protected by sea, by desert, or by mountainous regions

Cities and Countries by Map Numbers

1. *Cadiz* (Gades): Columella, 25. In southwestern Spain, not shown.
2. *Aquitania:* Victorius (Victorinus), 450. In southern France, not shown.
3. *Rome:* Varro, 60 B.C.; P. Nigidius Figulus, 60 B.C.; Vitruvius, 20 B.C.; Frontinus, 100; Menelaus, 100; Hyginus, 120; Balbus, 100; Domitius Ulpianus, 200; Nipsus, 180; Epaphroditus, 200; Sextus Julius Africanus, 220; Censorinus, 235; Serenus, 200; Porphyrius, 275.
4. *Sicily:* Diodorus Siculus, 1st century B.C.
5. *Syracuse* and *Messina:* Archimedes, 225 B.C.; Julius Firmicus Maternus, 340; Dicæarchus of Messina, 320 B.C.
6. *Naples* and *Pompeii:* Pliny, 75.
7. *Medma:* Philippus Medmæus, 350 B.C.
8. *Elea:* Zeno, 450 B.C.; Parmenides, 460 B.C.
9. *Crotona:* Pythagoras, 540 B.C.; Philolaus, 425 B.C.
10. *Tarentum:* Pythagoras, 540 B.C.; Philolaus (?), 425 B.C.; Archytas, 400 B.C.
11. *Cyrene:* Theodorus, 425 B.C.; Nicoteles, 250 B.C.; Eratosthenes, 230 B.C.; Synesius, 410.
12. *Elis:* Hippias, 425 B.C.
13. *Athens:* Solon, 600 B.C.; Agatharchus, 470 B.C.; Socrates, 425 B.C.; Meton, 432 B.C.; Phaeinus, 432 B.C.; Euctemon, 432 B.C.; Theætetus, 375 B.C.; Plato, 380 B.C.; Speusippus, 340 B.C.; Ptolemy, 150.
14. *Stageira:* Aristotle, 340 B.C.
15. *Thasos:* Leodamas, 380 B.C.
16. *Abdera:* Democritus, 400 B.C.
17. *Chios:* Œnopides, 465 B.C.; Hippocrates, 460 B.C.
18. *Samos:* Pythagoras, 540 B.C.; Conon, 260 B.C.; Aristarchus, 260 B.C.
19. *Paros:* Thymaridas, 380 B.C.
20. *Crete:* Early civilization, particularly at Knossos.
21. *Rhodes:* Eudemus, 335 B.C.; Geminus, 77 B.C.; Poseidonius, 100 B.C.
22. *Cnidus:* Eudoxus, 370 B.C.
23. *Miletus:* Thales, 600 B.C.; Anaximander, 575 B.C.; Anaximenes, 530 B.C.
24. *Clazomenæ:* Anaxagoras, 440 B.C.
25. *Smyrna:* Theon, 125.
26. *Pergamum:* Great library; parchment.
27. *Cyzicus:* Callippus, 325 B.C.
28. *Byzantium* (Constantinople): Proclus, 460; Psellus, 1075.
29. *Chalcedon:* Xenocrates, 350 B.C.; Proclus, 460.
30. *Nicæa* and *Bithynia:* Hipparchus, 140 B.C.; Theodosius, 50 B.C.; Sporus, 275.
31. *Amisus:* Dionysodorus, 50 B.C.
32. *Laodicea:* Anatolius, 280.
33. *Perga:* Apollonius, 225 B.C.
34. *Cyprus:* Early civilization.
35. *Apameia:* Poseidonius, 100 B.C.
36. *Larissa:* Domninus, 450.
37. *Chalcis:* Iamblichus, 325.
38. *Tyre:* Marinus, 150; Porphyrius, 275.
39. *Gerasa:* Nicomachus, 100.
40. *Jerusalem:* Religious.
41. *Alexandria:* Euclid, 300 B.C.; Eratosthenes, 230 B.C.; Apollonius, 225 B.C.; Aristarchus, 260 B.C.; Hypsicles, 180 B.C.; Heron, 50; Menelaus, 100; Ptolemy, 150; Diophantus, 275; Pappus, 300; Theon, 300; Hypatia, 400.
42. *Mendes:* Commercial.
43. *Antinoopolis* (Antinoe): Serenus, 50.
44. *Ptolemais:* Synesius, 410.
45. *Syene:* Eratosthenes, 230 B.C.

Early Greek Appreciation of Geometric Forms. Greece went through the same stages of appreciation of geometric forms that Egypt and Crete went through. This is seen in the use of crude parallels, then of the less crude and more elaborate forms, and finally of the more delicate forms found in the period just preceding the development of the highest type of Greek art. The use of these geometric forms was especially noteworthy just before the time of Thales, what is known as the geometric style in the decoration of vases having reached its climax in the 8th century B.C.




GREEK GEOMETRIC FORMS JUST PRECEDING THE TIME OF THALES

From a specimen of the 8th century B.C. in the Metropolitan Museum, New York

Greek Algebra. Algebra as a science distinct from arithmetic and geometry was invented long after Greece had ceased to be a center of civilization. Certain identities that we now express in algebraic form were well known to the Greeks, however, and were demonstrated with even greater rigor than is the case in our textbooks today, where the work is practically limited to rational numbers. For example, the Greeks proved that

$$(a+b)^2 = a^2 + 2ab + b^2,$$

although they had no algebraic shorthand by which to express the fact, and although they considered only lines and rectangles instead of numbers and products. In like manner they knew such other identities as



$$(a+b)(a-b)=a^2-b^2,$$

$$a(x+y+z)=ax+ay+az,$$

and $$(a-b)^2=a^2-2ab+b^2,$$

although they considered these also as geometric relations. They could complete the square of the binomial expression

$$a^2 \pm 2ab,$$


but, again, this was looked upon as the filling out of a geometric figure that was made up of a square increased or decreased by twice a rectangle. Greek algebra, as a form of arithmetic distinct from geometry, was developed some time after the period which we are now studying. When we come to consider the life and works of Diophantus (*c.* 275), for example, we shall see that the later Greeks made a remarkable advance in the analytic treatment of this subject. They developed a fairly good symbolism, and they considered algebra, under the name "arithmetic," as entirely distinct from the geometry of which we have been speaking.

3. Origins of Greek Mathematics

The Makers of Greek Mathematics. There are three important periods in the development of Greek mathematics, two of them within the chronological limits now being considered and one immediately following the later of these limits. The periods may be characterized as, first, the one subject to the influence of Pythagoras; second, the one dominated by Plato and his school; third, the one in which the Alexandrian School flourished in Grecian Egypt and extended its influence to Sicily, the Ægean Islands, and Palestine. We shall now consider the names of some of those who made the mathematics

of the first two of these important periods, and some of the influences which led them to undertake their epoch-making work.[1]

Thales. The first of the Greeks to take any scientific interest in mathematics in general, and in the union of astronomy, geometry, and the theory of numbers in particular, was Tha'les.[2] Before his time there had been the usual interest of early peoples in the mystery of the heavens, as witness the statement of the poet Archil'ochus[3] that a solar eclipse was observed some time before Thales was born. Not until the time of Thales, however, did the science of mathematics begin in the Greek civilization.



THALES

Ancient bust in the Capitoline Museum at Rome, not contemporary with Thales


Miletus was then, as we have seen, a trading and colonizing center, a city of wealth and influence. Herodotus (*c.* 450 B.C.) tells us that Thales was of Phœnician descent; but his mother, Cleobuline, bore a Greek name, while the name of his father, Examius, is Carian. The name of Thales himself was

[1] The most recent and elaborate works on this period are A. Mieli, *La Scienza Greca*, of which Volume I (Florence, 1916) deals with the history of Greek science before Aristotle; and Sir T. L. Heath, *A History of Greek Mathematics*, 2 vols. (Cambridge, 1921), of which the first volume covers the period from Thales to Euclid.

[2] Θαλῆς. Born at Miletus, *c.* 640 B.C.; died *c.* 546.

[3] Ἀρχίλοχος. Born *c.* 714 B.C.; died *c.* 676.

probably a common one. Indeed, we have in the Metropolitan Museum in New York at present a Cypriote vase of his period which bears the name in the form



—evidently that of the owner.[1] Considering the general recognition of the abilities of Thales, even during his lifetime, it is not impossible that a vase made in Cyprus may have been intended for him, but there is no further evidence that this was the case.

CYPRIOTE VASE WITH THE NAME OF THALES

The name *TA* + *LE* + *SE*, when viewed from above, has the appearance shown in the text. The vase is about contemporary with Thales of Miletus. From the Metropolitan Museum of Art, New York

Stories concerning Thales. Thales was a merchant in his younger days, a statesman in his middle life, and a mathematician, astronomer, and philosopher in his later years. In his mercantile ventures he seems to have been unusually successful, even in dealing with the shrewdest of the Greek trading races. Aristotle (*c.* 340 B.C.) tells us how he secured control of all the oil presses in Miletus and Chios in a year when olives promised to be plentiful, subletting them at his own rental when the season came. Plutarch (1st century) also testifies to his ingenuity in the following anecdote:[2]

Solon went, they say, to Thales at Miletus, and wondered that Thales took no care to get him a wife and children. To this Thales made no answer for the present; but a few days after procured a stranger to pretend that he had left Athens ten days earlier and

[1] Reading from right to left, the characters have the values *ta* + *le* + *se*, corresponding to the Greek Θα + λῆ + ς, Thales. The vase is part of the Cesnola collection found in Cyprus.

[2] In his life of Solon.

Solon inquiring what news there was, the man replied according as he was told: "None but a young man's funeral, which the whole city attended, for he was the son, they said, of an honorable man, the most virtuous of the citizens, who was not then at home, but had been traveling a long time." Solon replied, "What a miserable man is he! But what was his name?" "I have heard it," said the man, "but have now forgotten it, only there was great talk of his wisdom and justice." [After Solon had been drawn on to pronounce his own name and had learned that it was his own son,] Thales took his hand and, with a smile, said, "These things, Solon, keep me from marriage and rearing children, which are too great for even your constancy to support; however, be not concerned at the report, for it is a fiction."

Solon (*c.* 639–559 B.C.), it should be observed, was interested in astronomy and was the one who introduced a "leap month" into the Athenian calendar (594 B.C.).

Trade was then an honorable calling, and Thales seems to have traveled in Egypt on his commercial ventures, and early writers tell of his also visiting both Crete and Asia. He was not the only mathematician to have thus turned trade to profit, for Plutarch has this to say of him: "Some report that Thales and Hippocrates the mathematician traded, and that Plato defrayed the charges of his travels by selling oil in Egypt." In this way Thales may have accumulated the wealth that permitted him to indulge his taste for learning and to found the Ionian School. It was through this indulgence that he acquired such a reputation as to be enrolled as the first among the Seven Wise Men of Greece, and that he was esteemed as the father of Greek astronomy, geometry, and arithmetic.

Arithmetic of Thales. Of the nature of the arithmetic that Thales brought back from Egypt we have little direct knowledge. Iamblichus of Chalcis (*c.* 325 A.D.) tells us that he defined number as a system of units, and adds that this definition and that of unity came from Egypt. This is not much, but it is enough to show that Thales was interested in something besides the merely practical. It is probable that he knew many

other number relations, for the Ahmes papyrus contains some work in progressions, and such knowledge would hardly escape so careful an observer as Thales. It is, however, in his work in founding deductive geometry and in his capacity as a teacher of Pythagoras rather than as a discoverer of facts that Thales commands our attention.

Interest in Astronomy. He took much interest in astronomy, and Herodotus (I, 74) tells us that he even succeeded in predicting an eclipse. Some authorities suppose this eclipse to have occurred on May 28, 585 B.C., while others place it about twenty-five years earlier. He could have obtained certain information on this subject from a study of the Chaldean records, but whether this was his source of information we cannot say. At the present time we have numerous cuneiform tablets of the 7th century B.C. which record such prognostications. One of these reads: "To the king, my master, I have written that there was about to be an eclipse. The eclipse has now taken place. This is a sign of peace for the king, my master."

A man like Thales, possessed of an inquisitive mind, coming in contact with scholars from other lands, either on his travels or in the commercial center of Miletus, would lose no opportunity to secure information of this kind and to make use of it in his teaching. Doubtless his scientific training led him to discard the astrological notions of the Chaldeans but to retain whatever of astronomy came to his attention.

Geometry of Thales. In geometry he is credited with a few of the simplest propositions relating to plane figures. The list, according to the most reliable ancient writers, is as follows:

1. Any circle is bisected by its diameter.
2. The angles at the base of an isosceles triangle are equal
3. When two lines intersect, the vertical angles are equal.[1]
4. An angle in a semicircle is a right angle.[2]

[1] See Proclus, ed. Friedlein, pp. 157, 250, 299 (Leipzig, 1873).

[2] There is a doubt about his knowing this. It is inferred from a statement by Pam'phila (Παμφίλη, a woman historian, 1st century A.D.), but there is no early authority for the statement.

5. The sides of similar triangles are proportional.[1]

6. Two triangles are congruent if they have two angles and a side respectively equal (Euclid, I, 26).[2]

Importance of his Geometry. As propositions in geometry these may seem trivial, since they are intuitive statements; but their very simplicity leads us to believe that it was the fact that Thales was the first to prove them that led Eudemus (*c.* 335 B.C.) and other early writers to mention them. Up to this time geometry had been confined almost exclusively to the measurement of surfaces and solids, and the great contribution of Thales lay in suggesting a geometry of lines and in making the subject abstract. With him we first meet with the idea of a logical proof as applied to geometry, and it is for this reason that he is looked upon, and properly so, as one of the great founders of mathematical science. In the history of mathematics, as in the history of civilization in general, it is the setting forth of a great idea that counts. Without Thales there would not have been a Pythagoras—or such a Pythagoras; and without Pythagoras there would not have been a Plato—or such a Plato.

Philosophy of Thales. In philosophy he is said to have asserted that water is the origin of all things, that everything is filled with gods, that the soul is that which originates motion, and that matter is infinitely divisible; but his basis for belief in these assertions is not very satisfactory. Like most of his contemporaries, he left no written works.

Anaximander. At the death of Thales the leadership of the Ionian School passed over to Anaximan'der,[3] who is generally

[1] He used this proposition in measuring the height of a pyramid by means of the shadow of the pyramid and that of a staff. See Diogenes Laertius, *Vitæ Philosophorum*, ed. Cobet, p. 6 (Paris, 1878); Pliny, *Hist. Nat.*, XXXVI, 17; Plutarch, *Septem Sapientium Convivium*, ed. Didot, III, 174 (Paris, 1841). Pliny's statement that he measured the shadow at the time of day when the shadow "is equal in length to the body projecting it" is not very convincing. This would be too simple and is quite contrary to Plutarch's version.

[2] Eudemus (*c.* 335 B.C.), a disciple of Aristotle, refers this to Thales.

[3] Ἀναξίμανδρος. Born *c.* 611; died *c.* 547 B.C. J. Neuhaeuser, *Anaximander Milesivs* (Bonn, 1883).

thought to have been his pupil. Anaximander or some contemporary of his first brought into use in Greece the gnomon, an instrument resembling the sundial[1] and used for determining noon, the solstices, and the equinoxes. Aside from this, Anaximander seems to have had no interest in mathematics. It was about this time that the water clock (clep'sydra),[2] already known to the Assyrians, found its way into Greece, and very likely Anaximander's gnomon came also to be used for determining the time of day.

4. From Pythagoras to Plato

Pythagoras. Of all the interesting figures in the history of ancient mathematics Pythag'oras[3] ranks easily first, partly from the mystery surrounding his life, partly from his own mysticism, partly from the brotherhood which he established, and partly from the unquestioned ability of the man himself.

Early Life of Pythagoras. As with Euclid and Heron, of whom we shall presently speak, so with Pythagoras,—the date and the place of his birth are both unknown. He seems to have been born between the 50th and 52d Olympiads, to use the Greek system of chronology, or between 580 and 568 B.C. of our calendar. Although called a Samian, we are not certain that he was born on the island of Samos, for Suidas, a late medieval writer (*c.* 1000), says that he was born in Italy, and

[1] Really, the pointer which casts the shadow on the dial.

[2] Κλεψύδρα, from κλέπτειν (to hide) + ὕδωρ (water).

[3] Born at Samos (?), *c.* 572 B.C.; died at Tarentum (?), *c.* 501. W. Schultz, *Pythagoras und Heraklit*, Leipzig, 1905; A. Ed. Chaignet, *Pythagore et la philosophie pythagoricienne, contenant les fragments de Philolaüs et d'Archytas*, Paris, 1873; W. Bauer, *Der ältere Pythagoreismus*, Bern, 1897; W. W. Rouse Ball, "Pythagoras," in the *Math. Gazette*, London, January, 1915; G. J. Allman, *Greek Geometry from Thales to Euclid*, Dublin, 1889 (hereafter referred to as Allman, *Greek Geom.*); F. Cramer, *Dissertatio de Pythagora*, Prog., Sund, 1833; W. Lietzmann, *Der pythagoreische Lehrsatz*, Leipzig, 1912; Armand Delatte, "Études sur la littérature pythagoricienne," in the *Bibliothèque de l'École des hautes Études*, Vol. 217 (Paris, 1915). Of the early histories of Pythagoras the one best known is that of Iamblichus, *c.* 325. It first appeared in print at Franeker, 1598. Better editions by Ludolph Kuster (Amsterdam, 1707), and A. Nauck (Petrograd, 1884).

that as a child he migrated to Samos with his father.[1] Nevertheless the weight of authority favors his Samian birth, and a number of coins of the island, struck some centuries after his time, bear his name and figure, and this would hardly have been the case had he merely spent his boyhood there.[2] Various stories are told of his parentage, but we are quite uncertain whether his father was an engraver of seals or a merchant. At any rate he lived after Greece had enjoyed two centuries of commercial activity, and at the dawn of that golden age which began in Athens in the 6th century B.C. and closed for that city at the end of the 5th century B.C.



FIGURE OF PYTHAGORAS

A coin of Samos of the reign of Trajan (98–117), and therefore much later than Pythagoras. It shows the honor in which he was held and the claim of Samos as his native country

Period of Pythagoras. But in whatever land he was born, and in whatever year, and of whatever parentage, Pythagoras lived in stirring times and was himself one of the great makers of the civilization of his period. Samos was just becoming a center of Greek art and culture, Polyc'rates was just ascending the throne, and Anac'reon was beginning to write his famous lyrics in the Samian court. Pythagoras was therefore brought up amid scenes that could hardly fail to stimulate a youth of his native powers and urge him to a high intellectual life. Moreover, the spirit of the times was active in great works. Buddha was just promulgating his doctrine in India, and Confucius and Lao-Tze were laying the foundation for their philosophic cults in China; and, whether or not Pythagoras came into personal touch with the Far East, he lived when the world was ripe for great movements.

[1] See also M. Barbieri, *Notizie istoriche dei Mattematici e Filosofi di Regno di Napoli,* cap. ii (Naples, 1778), who (p. 25) thought that Samos was the modern Crepacore, a town in Southern Italy.

[2] One of these coins is shown in the illustration. There are also a few gems, of doubtful age, which are said to represent Pythagoras. See C. W. King, *Antique Gems and Rings,* I, 212, and II, plate XXXVIII, No. 1 (London, 1872).

The fact that arithmetic and geometry took such a notable step forward at this time was due in no small measure to the introduction of Egyptian papyrus into Greece. This event occurred about 650 B.C.,[1] and the invention of printing in the 15th century did not more surely effect a revolution in thought than did this introduction of writing material on the northern shores of the Mediterranean Sea just before the time of Thales.

Studies and Travels of Pythagoras. Our knowledge of the life of Pythagoras is very limited, the early writers having vied with each other in the invention of fables relating to his travels his miraculous powers, and his teachings. He seems to have sought out Thales and to have been his pupil. Tradition says that he was initiated by the master into the secrets of Zeus on Mount Ida, and was then told that if he would have further light he must seek it in Egypt. We now lose all definite knowledge of Pythagoras for a considerable period. Appuleius,[2] a Roman writer of about 150 A.D., asserts that he was captured by Cambyses the Persian,[3] that he learned from the Magi, and that he even sat at the feet of Zoroaster himself; but part of the story cannot be true, because Zoroaster probably died about the time that Pythagoras was born, and possibly much earlier than that, for the date is very uncertain. Isocrates, a writer of a century after Pythagoras, and Callimachus, librarian of the Alexandrian library, who lived in the 3d century B.C., both assert that he spent some years in Egypt. Pliny, writing in the 1st century of our era, says that Pythagoras was there in the time of Psammetichus,[4] and Strabo, about the beginning of the Christian era, states that he studied in Babylon. Others claim that he went as far east as India, but we have no definite proof of any of these statements.

[1] In the reign of King Psammetichus (Psammitichus) I, soon after 660 B.C., sometimes given as *c.* 640–610 and sometimes as 671–617. See T. Gomperz, *Les Penseurs de la Grèce*, French ed., p. 13 (Lausanne, 1904). This is the Psemtek of the monuments and the first king of the 26th Dynasty.

[2] Usually so spelled in ancient texts, but occasionally Apuleius.

[3] Reigned 529–522 B.C.

[4] That is, Psammetichus III, who reigned 526–525 B.C., when Pythagoras was about 46 years old.

Contact with the East. In spite of the assertion of various writers to the contrary, the evidence derived from the philosophy of Pythagoras points to his contact with the Orient. The mystery of the East appears in all his teachings.[1] His mysticism of numbers is quite like that found earlier in Babylon, and indeed his whole philosophy savors much more of the Indian than of the Greek civilization in which he was born. According to our best evidence the familiar proposition of geometry that bears his name was known, as already stated, in India, China, and Egypt (?) before his time, and all that can be claimed for him in relation to it is that he may have given the earliest general demonstration of its truth.

School of Crotona. When Pythagoras reappeared after his years of wandering, he sought out a favorable place for a school, and finally settled upon Crotona, a town on the southeastern coast of Italy, in a territory called by the Italic Greeks of that time Great Greece.[2] This town was a wealthy seaport, and it was to the young men of well-to-do families that Pythagoras made his appeal. Pretending to have the power of divination, given at all times to mysticism, and possessed in a remarkable degree of personal magnetism, he gathered about him some three hundred of the noble and wealthy young men of Magna Græcia and established a brotherhood that has ever since served as a model for all the secret societies in Europe and America. He divided his disciples into two groups, the hearers and the mathematicians, the latter having passed through a probationary period as members of the former group.

Oral Teaching of Pythagoras. Pythagoras never embodied his doctrines in any treatise. Like Thales, and also like those Oriental teachers from whom he probably learned, he transmitted his theories by word of mouth. This he did through the elect of his brotherhood, thus making known his doctrines freely to all who were deemed worthy to receive them. This

[1] E. W. Hopkins, *The Religions of India*, p. 559 (Boston, 1902); L. von Schroeder, *Indiens Literatur und Cultur*, pp. 718 seq.; *Reden und Aufsätze*, p. 168 (Leipzig, 1913); *Pythagoras und die Inder* (Leipzig, 1884).

[2] Ἡ μεγάλη Ἑλλάς, *Magna Graecia*.

method of imparting knowledge was not due merely to a spirit of mysticism, but was quite as dependent upon a lack of good writing material. Parchment had not as yet been invented, the wax tablet was serviceable only for brief epistles, the clay cylinders of Babylon were subject to similar limitations, and the fragile papyrus of Egypt was probably somewhat rare in Magna Græcia. Pythagoras therefore followed the custom of his time in passing his philosophy along by word of mouth, just as the ancients had transmitted to his generation the songs of Homer. Even in Plato's time there was no bookshop in all Athens where worthy manuscripts could be purchased, nor was there any when Euclid taught in Alexandria. Not until the time of Augustus was the book trade established, making possible the easy and certain transmission of knowledge, and not until fifteen hundred years later was printing known in Europe.

For the doctrines of Pythagoras we are indebted chiefly to Eudemus of Rhodes (*c.* 335 B.C.), whose works, though lost, are known to us through extracts preserved by later writers. We also know of the doctrines of Pythagoras through passages from a work by Philolaus of Crotona (who lived in the 5th century B.C.), from a statement by Archytas of Tarentum (*c.* 400 B.C.), a friend of Plato, and from a number of passages in the works of later writers.

Philosophy of Pythagoras. Pythagoras based his philosophy upon the postulate that number is the cause of the various qualities of matter. This led him to exalt arithmetic, as distinguished from logistic, out of all proportion to its real importance. It also led him to dwell upon the mystic properties of numbers and to consider arithmetic as one of the four degrees of wisdom,—arithmetic, music, geometry, and spherics (astronomy), these forming the quadrivium of the Middle Ages. Aristotle (384–322 B.C.) tells us that Pythagoras related the virtues to numbers, and Plutarch says that he believed that earth was produced from the regular hexahedron, fire from the pyramid, air from the octahedron, water from the icosahedron, and the heavenly sphere from the dodecahedron, in all of which the physical elements are related both to number and to form.

Philolaus probably voiced the teaching of the master when he asserted that five is the cause of color, six of cold, seven of health, and eight of love.

The Chinese say that five represents wind, and two represents earth, and these ideas are also claimed for the Pythagorean system.[1] Here again the resemblance between the mysticism of this school and that commonly found in the Far East leads to the belief that Pythagoras must have come into relations with the wise men of the Orient. Savoring of the East, too, is the description given by Suidas, a late medieval Greek compiler, of a ceremony called Pythagus, in which there is written something in blood on the face of a mirror, at the time of the full moon, the words then being read in the reflection in the circle of the moon; but there is no ancient authority for such a statement.[2]

Shakespeare refers to the acceptance by Pythagoras of the Hindu belief in the transmigration of souls, in these words:

> Thou almost mak'st me waver in my faith,
> To hold opinion with Pythagoras,
> That souls of animals infuse themselves
> Into the trunks of men.
>
> *Merchant of Venice*

Unity and Infinity. From various early writers we judge that Pythagoras asserted that unity is the essence of number, the origin of all things, the divine; that he had the idea of the limited and the unlimited; and that he held that from the latter came the ideas of time, space, and motion. Diogenes Laertius (2d century A.D.) says that he was interested in number, and that the part of mathematics "to which Pythagoras applied himself above all others was arithmetic";[3] and Aristox'enus[4] says that he esteemed this science above all others.

[1] J. Hager, *An Explanation of the Elementary Characters of the Chinese*, p. xv. London, 1801.

[2] J. C. Bulengerus, *De Lvdis privatis ac domesticis veterum*, p. 31. Lyons, 1627.

[3] Diogenes Laertius, VIII, i, 11, p. 207 ed. Cobet. Various references in connection with Greek mathematics may be found in Allman, *Greek Geom.*

[4] Ἀριστόξενος, a philosopher. Born at Tarentum, *c.* 350 B.C.

Geometry of Pythagoras. In the field of geometry Eudemus (*c.* 335 B.C.) informs us that Pythagoras "investigated his theorems from the immaterial and intellectual point of view," and that "he discovered the theory of irrational quantities and the construction of the mundane figures."[1] Favori'nus, a philosopher living in southern France *c.* 125, asserts that he employed definitions in his work in mathematics, this being the first trace that we have of such use.[2] In particular, he defined a point as "unity having position."[3] He or his school knew that the plane space about a point may be filled by six equilateral triangles, four squares, or three regular hexagons,—a fact which had doubtless been inferred as a result of the observation of mosaic pavements long before this time, but which no doubt he was able to prove. It is probable that Pythagoras proved the proposition relating to the sum of the angles of a triangle, that he constructed a polygon equivalent to one given polygon and similar to another, and that he could construct the five regular polyhedrons; and he may possibly have proved the theorem relating to the square on the hypotenuse. It seems likely that he taught that the earth is a sphere in space; at any rate, this theory was accepted by various later philosophers.[4]

Pythagoras on Music. Pythagoras is said to have discovered that the fifth and the octave of a note can be produced on the same string by stopping at $\frac{2}{3}$ and $\frac{1}{2}$ of its length, respectively, and it is thought that this harmony gave rise to the name of "harmonic proportion," since

$$1 : \tfrac{1}{2} = 1 - \tfrac{2}{3} : \tfrac{2}{3} - \tfrac{1}{2}.$$

Although he seems to have derived some knowledge of music from Egypt,[5] he is generally called the inventor of musical science, or the harmonic canon (a mere tradition), but we

[1] *I.e.*, of the five regular polyhedrons. Proclus (c. 412–485), ed. Friedlein, p. 65.

[2] Diogenes Laertius, VIII, i, 25, p. 215 ed. Cobet.

[3] Proclus, ed. Friedlein, p. 95.

[4] On doubts as to the Pythagorean Theorem see G. Junge, *Bibl. Math.*, VIII (3), 62, and H. Vogt, *ibid.*, IX (3), 19. On the astronomical question see P. Duhem, *Le Système du Monde* (Paris, 1913–1917).

[5] J. G. Wilkinson, *Manners and Customs of the Ancient Egyptians*, I, vi. London, 1878.

know nothing of the notes or of the system that he used.[1] With his love for music and number it is natural to believe that he must have taken great pride to himself for connecting the two in the harmonic proportion. He seems to have held that the



PYTHAGORAS THE MUSICIAN

From F. Gafurius, *Theorica Musice*, Milan, 1492. One of the first crude attempts to portray Pythagoras by means of a woodcut, and the first to portray him as a musician. In the same work he is also shown as a bell ringer


intervals between the heavenly bodies were determined by the laws of musical harmony, and hence arose the doctrine of the harmony of the spheres.

The influence of Pythagoras became so great that the government caused his brotherhood to be dispersed, although the members still spread the doctrines of the sect throughout Greece. Pythagoras died an exile from Crotona, possibly at Tarentum. Two centuries later, however, and during the disasters of the First Samnite War, 343 B.C., the Senate erected his statue in Rome, in response to an order of the Delphic Oracle to pay this honor to "the wisest and bravest of the Greeks," and the people learned to call him the preceptor of

[1] A. Baumgartner, *Geschichte der musikalischen Notation*, p. 11. Munich, 1856.

King Numa, while even the great Æmilian family was, in later years, proud to claim him as one of their honored ancestors.

We shall now consider a few of the other Greek mathematicians who attained prominence before the time of Plato. All of them were influenced by the doctrines of Pythagoras, and it is convenient and proper to consider them in close connection with the Pythagorean School, including in the discussion the members of the Eleatic School mentioned below.



MAP OF THE WORLD BY HECATÆUS, 517 B.C.

Showing the primitive ideas held at the time of Pythagoras. From Breasted's *Ancient Times*

Lesser Writers. Among the contemporaries of Pythagoras was Anaxim'enes of Miletus,[1] probably a pupil of Anaximander. Diogenes Laertius quotes two letters from him to Pythagoras, in one of which he speaks of Thales as his teacher; but his tastes were in the direction of philosophy rather than of mathematics. In this period there also flourished the geographer Hecatæ'us, whose map of the world serves to show how fragmentary was the knowledge then possessed even by the best scientists.

About the time of Pythagoras there also flourished Ameris'tus,[2] a geometer of some prominence and brother of the poet Stesich'orus. He is mentioned by Proclus (*c.* 412–485 A.D.), but nothing is known of his work.

[1] Ἀναξιμένης, born at Miletus, *c.* 585; died *c.* 528 B.C.

[2] The Mamercus of J. Gow, *History of Greek Mathematics*, p. 145 (Cambridge, 1884). Hereafter referred to as Gow, *Greek Math.*

Zeno of Elea.[1] About the time of the death of Pythagoras there was born at Elea[2] the philosopher Zeno, whose work on motion represented a noteworthy advance in the science, even though the mathematical feature is in evidence in only a single instance. It was from Elea that the Eleatic School of philosophers, one of the two great schools of southern Italy, derived its name. Zeno asserted that on account of the infinite divisibility of space through which an object must pass in moving, motion could not begin; that Achilles could not pass a tortoise, even though he went faster than the tortoise; that a moving object must be at once in motion and, because it occupies space, at rest; and that one space of time might, in different relations, be both long and short, reminding us of certain features of the modern doctrine of relativity. His argument with respect to Achilles and the tortoise may be thus expressed in modern units: If the tortoise has a mile the start and goes one tenth as fast as Achilles, when Achilles reaches the point where the tortoise was, the latter will be $\frac{1}{10}$ of a mile ahead; when Achilles has covered that distance, the tortoise will be $\frac{1}{100}$ of a mile ahead; and, similarly, whenever Achilles reaches a spot where the tortoise was, the latter will still be ahead, and so Achilles can never pass it.[3]

Anaxagoras. Among the noteworthy contemporaries of Zeno was Anaxag'oras,[4] the last of the celebrated philosophers of the Ionian School. He was a friend and teacher of Euripides, Pericles, and other great men of his time, but was condemned to death[5] at the age of seventy-two for being favorable to the Persian cause. Although his chief work was in philosophy, where his prime postulate was that "reason rules the world," he was interested in mathematics and wrote on the quadrature of

[1] Ζήνων (*Zenon*). Born at Elea, *c.* 496 B.C. He was living in the time of Pericles (died 429). Heath, *History*, I, 271.

[2] Ἐλέα or Ὑέλη (*Elea* or *Hyele*); Latin, *Velia*; in southern Italy.

[3] F. Cajori, "The History of Zeno's Arguments on Motion," *Amer. Math. Month.*, XXII, 1, 39, 77, 109, 143, 179, 215, 253, 293; "The Purpose of Zeno's Arguments on Motion," *Isis*, III, 7.

[4] Ἀναξαγόρας. Born at Clazomenæ, Ionia, *c.* 499; died *c.* 427 B.C.

[5] Ancient writers are not clear upon this point.

the circle and on perspective.[1] When banished from Athens he remarked, "It is not I who have lost the Athenians, but the Athenians who have lost me."

Agatharchus. About this period (470 B.C.) an Athenian artist, Agathar'chus,[2] applied stereometry to the theory of perspective. He is said to have painted the scenery for a tragedy which Æschylus produced. In his work on drawing he showed how to make use of the notion of projection upon a plane surface.

Socrates.[3] Although we do not commonly think of Soc'rates, the Athenian statesman and philosopher, as a mathematician, yet for his work on induction and for his insistence upon accurate definition he should be mentioned in connection with the early development of a logical geometry. As the teacher of Plato he assisted in the development of that great maker of philosophers and of those who based their mathematics upon sound logic. Socrates has left us no writings of his own, but we have the testimony of Plato, Euclid, and others that they were greatly his debtors. Probably no more noble tribute has been paid to him than that given by Dr. Jowett in his paraphrase of the words of Plato: "And he, Socrates, is a midwife, although this is a secret; he has inherited the art from his mother bold and bluff, and he ushers into light, not children, but the thoughts of men."[4] Xenophon[5] and Diogenes Laertius[6] tell us, however, that he felt that geometry and astronomy were useful merely for measuring fields and telling the time of day,—a view which, if really held by him, has been advanced by men of far less mentality in every generation since that time, and with the same empty results.

Œnopides of Chios.[7] Probably a Pythagorean and certainly one of the leading astronomers of his time, Œnop'ides is

[1] *Anaxagorae Fragmenta* (Leipzig, 1827); better edition by Schorn (Bonn, 1829). See also F. Breier, *Die Philosophie des Anaxagoras von Klazomenä nach Aristoteles* (Berlin, 1840).

[2] Ἀγάθαρχος.

[3] Σωκράτης. Born near Athens, 468 B.C.; died at Athens, 399.

[4] Jowett's *Plato*, IV, 123.

[5] *Memorabilia*, IV, 7.

[6] *Lives of the Philosophers*, II, 32.

[7] Οἰνοπίδης. Born in Chios; fl. *c.* 465 B.C.

thought to have learned the science of the stars and the obliquity of the ecliptic from the priests and temple astronomers of Egypt. He is said to have invented the cycle of 59 years for the return of the coincidence of the solar and lunar years, giving the length of the solar year as 365 days and somewhat less than 9 hours. Proclus (*c.* 460) attributes to him the discovery of two problems of Euclid, one (I, 12) referring to the drawing of a perpendicular to a given line from an external point, and the other (I, 23) referring to the making of an angle equal to a given angle. If this is really the case, it shows how slight had been the advance in demonstrative geometry, even in the century following the death of Pythagoras.

Democritus. Democ'ritus,[1] known to later generations as the Laughing Philosopher, inherited great wealth, spent his fortune in travel, met the learned men of many lands, was a man of remarkable diligence in study, and died in poverty. His works are lost, except for certain fragments.[2] One of his teachers in philosophy is said to have been Leucip'pus,[3] the founder of the atomic theory of the ancient philosophy which asserted that the original characteristics of matter are functions of quantity instead of quality, the primal elements being particles homogeneous in quality but heterogeneous in form. Archimedes tells us, in his work on *Method*, that Democritus was the first to show the relation between the volume of a cone and that of a cylinder of equal base and equal height, and similarly for the pyramid and prism. In spite of the manifest bearing of his work upon an infinitesimal calculus, it seems to have had no influence in this direction among the Greeks. It is said that Plato felt that all the writings of Democritus should be burned. At any rate he had so slight an opinion of the latter that he makes no mention of him in any of his works. Such

[1] Δημόκριτος. Born at Abdera, Thrace, *c.* 460 B.C.; died *c.* 357.

[2] F. W. A. Mullach (F. G. A. Mullachius), *Democriti operum fragmenta.* Berlin, 1843.

[3] Λεύκιππος. The date and place of his birth and of his death are unknown.

treatment at the hands of Plato was perhaps due to the boastful nature of Democritus, who speaks of himself in these words:

I have wandered over a larger part of the earth than any other man of my time, inquiring about things most remote; I have observed very many climates and lands and have listened to very many learned men; but no one has ever yet surpassed me in the construction of lines with demonstration; no, not even the Egyptian harpedonaptæ (Ἀρπεδονάπται), with whom I lived five years in all, in a foreign land.

These harpedonaptæ (literally, "rope stretchers") were the surveyors of ancient Egypt, and the quotation suggests that the logical demonstration of propositions was then practiced in that country as well as in Greece.

Parmenides of Elea. Parmen'ides of Elea[1] taught at Athens in the middle of the 5th century B.C., and among his theories of the universe was the one that the earth is a sphere. His work, however, was that of a philosopher rather than a mathematician. It was in his time that Herodotus (*c.* 450 B.C.) wrote his history, and it is in this work that the idea of a meridian first appears in any literature now extant. From this time on for several centuries the sphericity of the earth was accepted as valid by many philosophers. The theory was revived in the 12th century of our era and was strongly asserted by Roger Bacon (*c.* 1250).


Philola'us,[2] a distinguished Pythagorean, was born at Crotona, or possibly at Tarentum, and according to Plato was a contemporary of Socrates. Although Pythagoras handed down his doctrines by word of mouth, it is stated by Porphyr'ius (fl. *c.* 275) that Ly'sis, who was a prominent philosopher, and an obscure Pythagorean named Archip'pus, put into writing some of the doctrines of the school and transmitted them to their descendants as secret heirlooms. Philolaus, however, was the first to write a treatise on the teachings of Pythagoras and to

[1] Παρμενίδης. Born in Elea; fl. *c.* 460 B.C.
[2] Φιλόλαος. Fl. *c.* 425 B.C.

make it public. Judging from the fragments that have come down to us,[1] his interest was in philosophy rather than in mathematics, although he touches upon the latter field in his description of a gnomon.

Hip'pias of Elis,[2] known both as a statesman and as a philosopher, belonged to the sophists, a class of teachers who traveled from place to place and took money for their services,—a practice quite contrary to the ideas of earlier philosophers. He accumulated wealth by teaching and public speaking, and Plato speaks of him as a vain man, given to arrogance and boasting. He seems to have been possessed of a wide but superficial knowledge. His contribution to mathematics was confined to his invention of a simple device for trisecting any angle, this device being known as the quadratrix. Since it was studied and described at a later period by Deinostratus (Dinostratus, *c.* 350 B.C.), it generally bears the latter's name.[3]

Hippocrates.[4] Various stories are told of Hippoc'rates, among them one that he was an unsuccessful merchant, later becoming a Pythagorean philosopher with a special interest in mathematics. Aristotle speaks of him as skilled in geometry but as otherwise stupid and weak. He is mentioned by ancient writers as the first to arrange the propositions of geometry in a scientific fashion and as having published the secrets of Pythagoras in the



[1] A. Böckh, *Philolaos des Pythagoreers Lehren, nebst den Bruchstücken seines Werkes* (Berlin, 1819); W. R. Newbold, "Philolaus," in *Archiv für Geschichte der Philosophie*, XIX, 176.

[2] Ἱππίας. Born at Elis, on the west coast of Peloponnesus; fl. *c.* 425 B.C.

[3] Allman and Hankel do not believe that this is the Hippias to whom Proclus refers as the inventor of the quadratrix. There is room for the doubt that these eminent writers express, but Cantor, Montucla, and various other historians feel that the evidence is in favor of Hippias of Elis. See also Gow, *Greek Math.*, p. 162. The name "quadratrix" is due to the fact that the curve can also be used in the quadrature of the circle. On the application of this and other curves to the problems of geometry, see Volume II, Chapter V.

[4] Ἱπποκράτης. Born in Chios; fl. *c.* 460 B.C.

field of geometry. In his attempts at squaring the circle he discovered the first case of the quadrature of a curvilinear figure,[1] namely, the proof that the sum of the two shaded lunes here shown is equal to the shaded triangle. The proposition holds equally for any right-angled triangle, isosceles or not, although Hippocrates knew it only for the isosceles right-angled triangle. Proclus (*c.* 460) ascribes to him the method of reduction (*'απαγωγή*), the passing from one proposition to another that seems more simple, proving the latter, and then reversing the order. For example, Eratosthenes (*c.* 250 B.C.) tells us that Hippocrates showed that the duplication of a cube can be effected if two mean proportionals can be found between two given lines.[2]

Meton, Phaeinus, and Euctemon. That there was great interest in mathematical astronomy in Athens between the time of Pythagoras and that of Plato is seen in the work of the three astronomers Me'ton, Phaei'nus, and Eucte'mon.[3] It is not possible, however, to differentiate their contributions to the subject. The philosopher Theophras'tus[4] says that Phaeinus made astronomical observations on Lycabettus, at Athens, and that from these Meton constructed the cycle of 19 years, since known as the Metonic Cycle. The astronomer Ptolemy[5] says that Meton and Euctemon made observations at Athens and in other places. He adds that Meton made the length of the year to be $365\frac{1}{4}$ days $+\frac{1}{76}$ of a day, which is more than 30 minutes too long. Whether the 19-year cycle is really due to Meton, or was already known to Œnopides, or was obtained from Egyptian or other sources is, and is likely to remain, unknown.

[1] W. Lietzmann, *Der pythagoreische Lehrsatz* (Leipzig, 1912). For a further discussion see Volume II, Chapter V.

[2] That is, if $a : x = x : y = y : 2a$, then $x^2 = ay$, $x^4 = a^2y^2$, $y^2 = 2ax$, and hence $x^4 = 2a^3x$, or $x^3 = 2a^3$. F. Rudio, *Der Bericht des Simplicius über die Quadraturen des Antiphon und des Hippokrates*, Leipzig, 1907, with Greek and German text; P. Tannery, "Hippocrate de Chios et la quadrature des lunules," in the *Mémoires de Bordeaux* (1878); *La Géométrie Grecque*, p. 117 (Paris, 1887); "Le fragment d'Eudème sur la quadrature des lunules," *Mémoires scientifiques*, II, 46, 339 (Paris, 1912); Gow, *Greek Math.*, p. 164; M. Simon, *Archiv der Math.*, VIII (3), 269.

[3] *Μέτων, Φαεινός, Εὐκτήμων.* Fl. *c.* 432 B.C.

[4] *Θεόφραστος.* Fl. *c.* 250 B.C

[5] Born *c.* 85; died *c.* 165.

The Method of Exhaustion. Iamblichus (*c.* 325) mentions Bry'son,[1] or Bryso, as one of the youths whom Pythagoras instructed in his old age. If this is true, Bryson must have been born about 520 B.C., but it is commonly believed that he flourished about 450 B.C. He was formerly thought to have contributed to what is known as the method of exhaustion, a crude approach to the integral calculus whereby the area between a curvilinear figure (say a circle) and a rectilinear figure (say an inscribed regular polygon) could be approximately exhausted by increasing the number of sides of the latter. There is, however, no reliable ancient authority for connecting his name with the theory.[2] The method was effectively used by later writers, notably by Eudoxus and Archimedes, and was extended to include the mensuration of solids.

Antiphon and the Method of Exhaustion. Aristotle mentions a Greek sophist named An'tiphon,[3] or Antipho, whose attempts at the quadrature of the circle led him into this phase of geometry. Antiphon inscribed a regular polygon in a circle, doubled the number of sides, and continued doubling until, as he seems to have believed, the sides finally coincided with the circle. Since he could construct a square equivalent to any polygon, he could then, as he thought, construct a square equivalent to the circle; that is, he could "square the circle," thus finding its area. We have here another phase of the method of exhaustion, the area between the polygon and the circle being exhausted as the process of doubling the number of sides proceeds. It is one of the first steps in the development of an infinitesimal calculus applied to integration,—a type of mathematics that had to wait two thousand years for serious consideration.

Archytas. In Plato's time Archy'tas,[4] a distinguished Pythagorean philosopher, achieved a high reputation as a mathematician, a general, a statesman, a philanthropist, and an educator, and Cicero (106–43 B.C.) speaks of him as a friend of the great

[1] Βρύσων. [2] F. Rudio, *Bibl. Math.*, VII (3), 378.

[3] Ἀντιφῶν. Fl. *c.* 430 B.C.

[4] Ἀρχύτας. Born at Tarentum, *c.* 428 B.C.; died *c.* 347. See Allman, *Greek Geom.*, p. 102, for an excellent summary of his work.

master himself. Horace (65–8 B.C.) refers to his death by shipwreck in the Adriatic, speaking of him in these words:

> The scanty present of a little dust
> Near the Matinian shore confines thee, O Archytas,
> Measurer of the sea, the earth, and the innumerable sand.[1]

Archytas lived in Magna Græcia, then much more tranquil than Greece itself, disturbed as the latter was by the Peloponnesian War. It was because of these wars that many Pythagoreans returned to Crotona and Tarentum, the result being that scholarship again flourished in this part of Italy. Vitruvius[2] says that Archytas solved the problem of the duplication of the cube by means of cylindric sections. He was the first to apply mathematics in any noteworthy way to mechanics, and he also applied the science to music and even to metaphysics.[3]

Eudemus (*c.* 335 B.C.), speaking of his work in geometry, tells us that he was one of those who "enriched the science with original theorems and gave it a sound arrangement," and from another statement we infer that he knew and doubtless proved the following propositions:

1. If a perpendicular is drawn to the hypotenuse from the vertex of the right angle of a right-angled triangle, each side is the mean proportional between the hypotenuse and its adjacent segment.

2. The perpendicular is the mean proportional between the segments of the hypotenuse.

3. If the perpendicular from the vertex of a triangle is the mean proportional between the segments of the opposite side, the angle at the vertex is a right angle.

[1] Te maris et terrae numeroque carentis harenae
Mensorem cohibent, Archyta,
Pulveris exigui prope litus parva Matinum
Munera.

Carmen i, 28

[2] *Praefatio* to his *De Architectura*, ix.

[3] For a list of fragments attributed to him see J. A. Fabricius, *Bibliotheca Graeca*, 14 vols., I, 833 (Hamburg, 1705–1728). There is a later edition, Hamburg, 1700–1809. See also Gow, *Greek Math.*, p. 157.

4. If two chords intersect, the rectangle of the segments of one is equivalent to the rectangle of the segments of the other.

5. Angles in the same segment of a circle are equal.

6. If two planes are perpendicular to a third plane, their line of intersection is perpendicular to that plane and also to their lines of intersection with that plane.[1]

Theodorus of Cyrene.[2] Among those who assisted in preparing the way for scientific mathematics as distinguished from the intuitive form, Theodo'rus of Cyrene deserves at least brief mention. He was a Pythagorean philosopher, and Proclus (*c.* 460) says that he was a little younger than Anaxagoras, who was born *c.* 499 B.C. According to Appuleius (*c.* 150) and Diogenes Laertius (2d century), Plato went to Cyrene to study geometry under Theodorus, possibly learning from him the theory of irrationality, which, as we know, had received attention in the school of Pythagoras.

Theæte'tus[3] **of Athens** was a pupil of Theodorus and of Socrates and is described by Plato as a man of unusual brilliancy.[4] Although his works are lost, there are references in the writings of the ancient historians to show that he discovered a considerable part of elementary geometry and wrote upon solids. Euclid seems to have been indebted to him and to Eudoxus for some of the material used in writing the *Elements*.

With Theætetus may be said to have closed the period which began with Pythagoras and which prepared the way for Plato and his school. Pythagoras made scientific study popular with the leisure class,—or at least he created an influential group of scholars. Without the work of his school, supplemented by the contributions of such schools as the one at Elea, the world would not have been ready for Plato. The period just closing supplied the raw material, and we shall find that Plato furnished the tools for making good use of this supply.

[1] Allman, *Greek Geom.*, p. 114.

[2] Θεόδωρος. Fl. *c.* 425 B.C. Cyrene (Κυρήνη) was a city on the north coast of Africa.

[3] Θεαίτητος. Fl. *c.* 375 B.C.; died 368 B.C. H. Vogt, *Bibl. Math.*, XIII (3), 200.

[4] Allman, *Greek Geom.*, p. 206.

5. Influence of Plato and Aristotle

Plato. To few men can the words of Carlyle be more appropriately applied than to Plato: "In every epoch of the world the great event, parent of all others, is it not the arrival of a Thinker in the world!" For never in all her early history was Greece so desperately in need of men of soul as she was when Plato[1] began his life work. While he was still a young man (404 B.C.) Athens fell before the Spartan forces. The century in which were first produced the great tragedies of Æschylus, Sophocles, and Euripides, which saw the Acropolis adorned with the masterpieces of Ictinus, Phidias, and Callicrates, and which knew Athens under the reign of Pericles the Magnificent,—this century had passed away, and with it had gone forever that glory of the city that appealed to the masses,—the glory of arms, of the drama, of architecture, and of sculpture. The new century was to see a new Athens, dead to the present but filled with intellectual ambitions for the future. Three great names



PLATO

A fanciful portrait. From a drawing by Raphael in the Accademia at Venice. Inserted to show this artist's conception of the philosopher

[1] Πλάτων (*Platon*). Born at Athens, *c.* 430 B.C.; died *c.* 349.

of Athenian citizens has that future preserved, and by them was it powerfully influenced,—the names of Plato, Aristotle, and Demosthenes.

Plato's Studies. Of the first of these three great leaders of men Cicero has this to say:

> It is reported of Plato that he came into Italy to make himself acquainted with the Pythagoreans, and that when there he made the acquaintance, among others, of Archytas and Timæus[1] and learned from them all the tenets of the Pythagoreans.[2]

It is also said that Plato visited Egypt, partly, no doubt, for purposes of trade, but chiefly that he might acquire knowledge. It may have been from the priests along the Nile, but more likely through the Pythagoreans, that he came to appreciate so highly the value of geometry. At any rate, in later years he is said to have placed above the entrance to his school of philosophy (the Academy) the words, "Let no one ignorant of geometry enter my doors,"[3]—the oldest recorded entrance requirement of a college,—and to have spoken of God as the great geometer.[4]

Plato studied under Socrates and also under a certain Eucleides (Euclid) of Meg'ara,[5] a philosopher who has often been confused with Euclid the geometer (*c.* 300 B.C.). He traveled extensively, visiting not only Egypt and lower Italy but also Sicily and possibly Asia. He thus came in contact with the mathematics and philosophy of these various countries and returned to Athens filled with enthusiasm for an era of splendid thought in place of the era of splendid action which had characterized the century that had just closed.

[1] Not the historian, but a native of Locri. Probably no works of his are extant, although there is one doubtfully attributed to him.

[2] *Tusculan Disputations*, I, 17

[3] Μηδεὶς ἀγεωμέτρητος εἰσίτω μου τὴν στέγην.

[4] "God eternally geometrizes," Ἀεὶ θεὸς γεωμετρεῖ. This is not in Plato's works, but is stated by Plutarch as due to him. Plutarch, *Convivalium Disputationum libri novem*, viii, 2, ed. Didot (Paris, 1841).

[5] A Greek city.

Of his philosophy it is unnecessary to speak, since this has little bearing upon the problem in hand, but in the field of mathematics his great contribution was to the underlying principles of the science, including the method of analysis.

Plato's Interest in Arithmetic. In the study of numbers he, like all the ancient philosophers, was interested in arithmetic rather than logistic. In his *Republic* he says that the science has a double use, military and philosophical.

> For the man of war must learn the art of numbers or he will not know how to array his troops;[1] and the philosopher also, because he has to arise out of the sea of change and lay hold of true being, and therefore he must be an arithmetician. . . . Arithmetic has a very great and elevating effect, compelling the mind to reason about abstract number.

Mysticism of Numbers. One thing that particularly interested Plato was the mysticism of numbers. In his *Republic* (Book VIII) he speaks in an obscure fashion of a certain mystic number, but does not make it clear what this number is. He calls it "the lord of better and worse births," and subsequent writers have often tried to find exactly what he meant. One theory is that 60^4, or 12,960,000, is the Platonic number. This number played an important part in the mysticism of the Hindus and the Babylonians, and it is possible that Pythagoras found it on the banks of the Euphrates, if he really studied there, and that he took it with him to Crotona, passing it on to his disciples, who, in turn, told it to Plato and his followers.

Although Plato esteemed the science of numbers highly,[2] he gives us no information concerning the way it was taught in his school or what it included. We are about as ignorant of the subject as presented by him, and of the ground it covered, as we are of the ancient logistic.[3]

[1] Evidently referring to the square, heteromecic, and triangular numbers described in Volume II, Chapter I.

[2] *Laws*, V.

[3] P. Tannery, "L'éducation platonicienne: L'arithmétique," in the *Revue scientifique*, XI (1881), 287.

Plato on Geometry. More than any of his predecessors Plato appreciated the scientific possibilities of geometry, of which more will be said in Volume II of this work. By his teaching he laid the foundations of the science, insisting upon accurate definitions, clear assumptions, and logical proof. His opposition to the materialists, who saw in geometry only what was immediately useful to the artisan and the mechanic, is made clear by Plutarch (1st century) in his *Life of Marcellus*. Speaking of the use of mechanical appliances in geometry, Plutarch remarks upon "Plato's indignation at it and his invections against it as the mere corruption and annihilation of the one good of geometry, which was thus shamefully turning its back upon the unembodied objects of pure intelligence." That Plato should hold the view here indicated is not a cause for surprise. The world's thinkers have always held it. No man ever created a mathematical theory for practical purposes alone. The applications of mathematics have generally been an afterthought.[1]

Immediate Followers of Plato. Among the followers of Plato was his nephew, Speusip'pus,[2] who accompanied him on his third journey to Syracuse and succeeded him as head of the Academy (347–339 B.C.). He wrote upon Pythagorean numbers,—integers like 3, 4, 5, which represent the sides and hypotenuse of a right-angled triangle. He also wrote upon proportion. We get some information concerning him from an anonymous work of uncertain date called the *Theologumena*, which is also the title of a lost work by Nicomachus (*c.* 100 A.D.). In this work it is related that Speusippus was the son of Potone, the sister of Plato, and that he "ceased not to study with diligence" the lessons of the Pythagoreans, and especially of Philolaus. The work also states that he treated with rare elegance the subjects of linear, polygonal, plane, and solid numbers.[3]

[1] On the general subject of mathematics in Plato's time see B. Rothlauf *Die Mathematik zu Platons Zeit und Seine Beziehungen zu ihr* (Munich, 1878) Heath, *History*, I, 284.

[2] Σπεύσιππος.

[3] See Volume II, Chapter I.

Of the minor followers of Plato mention should be made of Leod'amas of Thasos,[1] who is referred to by Proclus (*c.* 460) and Diogenes Laertius (2d century) and is said to have made use of the analytic method of proof. There was also Philip'pus Medmæ'us,[2] an astronomer and geometer of Medma, or Mesma, in Magna Græcia, who, under the guidance of Plato, took up the study of mathematics.[3] Thymaridas, who devised a rule for solving simultaneous linear equations, seems to have lived about this time.

Eudox'us of Cnidus,[4] at one time a pupil of Plato, achieved eminence in astronomy, geometry, medicine, and law.[5] It is said that he introduced the study of spherics (mathematical astronomy) into Greece and made known the length of the year as he had found it given in Egypt. He was the first of the Greeks, so far as is known, to give a description of the constellations. Strabo asserts that the observatory of Eudoxus still existed at Cnidus in his time, that is, about the beginning of the Christian era. Seneca says that he brought from Egypt to Greece the theory of the motions of the planets; Aristotle records that he made separate spheres for the stars, sun, moon, and planets; and Archimedes says that he found the diameter of the sun to be nine times that of the earth and showed that a pyramid is one third of a prism of the same base and the same altitude, and similarly for a cone and cylinder. For the mensuration of the cone and cylinder he probably developed the method of exhaustion as a rigorous theory.[6] Vitruvius gives him credit for a new form of sundial called the spider's web,[7] which may, however, have been an astrolabe. Because of a note, possibly due to Proclus, he is often credited with having written a work on proportion which finally became Book V of Euclid, but for this statement there is no definite historical

[1] Λεωδάμας. Fl. *c.* 380 B.C. [2] Φίλιππος ὁ Μεδμαῖος Born *c.* 375 B.C.
[3] He is also known as Opuntius. H. Vogt, *Bibl. Math.*, XIII (3), 193, 195.
[4] Εὔδοξος. Born *c.* 408 B.C.; died *c.* 355. See Allman, *Greek Geom.*, p. 128; Gow, *Greek Math.*, p. 183; Heath, *History*, I, 322.
[5] Diogenes Laertius, VIII, 86. [6] See Allman, *Greek Geom.*, pp. 96, 139.
[7] Ἀράχνη. Part of the astrolabe resembles a web.

sanction.[1] Our principal knowledge of Eudoxus and his work comes from an astronomical poem written by Ara'tus,[2] and from a commentary of Hipparchus upon it.[3]

Menæch'mus[4] was a pupil of Eudoxus and a friend of Plato,[5] and possibly it is to him that we owe the first treatment of conics. It is said that Alexander the Great was his pupil and that he asked that geometry be made more simple for him; whereupon Menæchmus replied: "O King, through the country there are private and royal roads, but in geometry there is only one road for all."[6] The conic sections which Proclus (*c.* 460) says were considered by him were probably the "Menæchmian Triads" of Eratosthenes (*c.* 230 B.C.). It is said that he obtained them by cutting cones by planes perpendicular to an element,—the parabola from a right-angled cone, the hyperbola from an obtuse-angled cone, and the ellipse from an acute-angled cone. A friend of his, Theudius of Magnesia, wrote a textbook on geometry.

Deinos'tratus, or Dinostratus,[7] was a brother of Menæchmus. He is known chiefly for his study of the quadratix, a curve already invented by one Hippias, very likely Hippias of Elis. This curve enabled him to square a circle.[8]

Xenoc'rates,[9] a native of Chalcedon, was a friend of Plato and Aristotle and was prominent both as a philosopher and as a diplomat. Besides various works on philosophy and government he wrote on physics, geometry, arithmetic, and astrology.

[1] For a discussion of the matter, see Allman, *Greek Geom.*, p. 136; Sir T. L. Heath, *The Thirteen Books of Euclid's Elements*, 3 vols., II, 112 (Cambridge, 1908) (hereafter referred to as Heath, *Euclid*).

[2] Ἄρατος. Fl. *c.* 270 B.C. The poem was the *Phaenomena* (Φαινόμενα), and certain fragments were preserved by Hipparchus.

[3] Ἵππαρχος. Fl. *c.* 150 B.C. J. B. J. Delambre, *Histoire de l'astronomie ancienne*, I, 106 (Paris, 1817). This poem of Aratus was first printed at Venice in 1499.

[4] Μέναιχμος. Fl. 365–350 B.C. See *Bibl. Math.*, XIII (3), 194.

[5] See Allman, *Greek Geom.*, p. 153; Max C. P. Schmidt, "Die Fragmente des Mathematikers Menaechmus," in *Philologus*, XLII (1884), p. 77; Gow, *Greek Math.*, p. 185; Heath, *History*, II, 110.

[6] The story is due to Stobæus, a late Greek writer, *c.* 500. It is also related of Euclid and King Ptolemy.

[7] Δεινόστρατος. Fl. *c.* 350 B.C.

[8] The details are considered in Volume II, Chapter V. See also Allman, *Greek Geom*, p. 180; Gow, *Greek Math.*, p. 187.

[9] Ξενοκράτης. Born *c.* 396 B.C.; died 314.

Plutarch (1st century) tells us that he took the soul as a "self-moving number," and deified unity and duality,[1] speaking of the former as the first male existence, ruling in heaven, as father and Zeus, as uneven number and spirit; and duality as the first female, the mother of the gods, and the soul of the universe which reigns over the world,—all of which theory shows the Pythagorean influence. He also assumed the existence of indivisible lines and spoke of them as the elements of certain Platonic triangles, perhaps with some intuition of an infinitesimal calculus. He followed Speusippus as head of the Academy and wrote a history of geometry in five books, which, like his other works, is lost.

Ar'istotle[2] studied under Plato at Athens, and his diligence and brilliancy led the latter to call him the "intellect of the school."[3] He became one of the instructors of Alexander the Great, and later returned to Athens and founded the Peripatetic School of philosophy, probably so called from the place where he taught.[4] He was a voluminous writer, but although many of his works are extant the major part are lost. His interest in the mathematical sciences lay chiefly in their applications to physics. He speaks of mathematics as standing halfway between physics and metaphysics. He wrote two works of a mathematical nature, one on indivisible lines and the other on mechanical problems. Both have been edited and printed. We know that, contrary to the doctrines of the Pythagoreans, he advocated the separation of arithmetic and geometry. In his systematizing of logic he contributed indirectly to the great work of Euclid. To him, too, we owe the first known definition of continuity: "A thing is continuous when of any two successive parts the limits at which they touch are one and the same and are, as the word implies, held together."[5] Aristotle was also interested in the historical development of science, and this seems to have influenced the work of his disciples in

[1] Μονάς and δυάς.

[2] Ἀριστοτέλης. Born at Stageira (Stagira), the present Stavro, 384 B.C.; died at Athens, 322.

[3] Νοῦς τῆς διατριβῆς.

[4] Ὁ περίπατος.

[5] Gow, *Greek Math.*, p. 188.

gathering materials for the history of mathematics. Among those whose interests led them into this field was Theophras'tus,[1] a pupil of Plato as well as of Aristotle. He wrote on philosophy, oratory, poetry, botany, physics, politics, and mathematics, but his works are known chiefly from fragments.[2]

Eudemus. Eude'mus[3] of Rhodes, another disciple of Aristotle, who flourished *c.* 335 B.C., was also much interested in the history of mathematics. Most of his works are lost, but certain fragments remain and serve to throw considerable light upon the mathematics of the Aristotelian school. It seems to have been to his care that we are indebted for the preservation of certain works of Aristotle.

Dicæarchus. It is probable that Dicæar'chus[4] of Messina, a city just north of Syracuse, in Sicily, was also a disciple of Aristotle, although we know little of his life. He seems to have flourished *c.* 320 B.C. and to have died *c.* 285 B.C. His work in mathematics was connected chiefly with mensuration as applied to geography. There was another philosopher by the same name, a Pythagorean, whom Iamblichus (*c.* 325) quotes as having contributed to the history of mathematics, but his works are not extant.

Autolycus. Among the contemporaries of Aristotle should probably be included, although we are uncertain as to the date, the astronomer Autol'ycus.[5] Nothing is known of his personal history except that he wrote two treatises on astronomy, both of which are extant. These are the most ancient mathematical texts that have come down to us from the Greeks. The first is on the motion of the sphere, and the second is on the risings and settings of the fixed stars, and in each he shows considerable skill in geometry.

Aristæ'us,[6] known as Aristæus the Elder, is mentioned by Pappus, a mathematician of the 4th century, as one of the three geometers of the Greeks who were skilled in that branch of

[1] Θεόφραστος. Fl. *c.* 350 B.C.
[2] There are various editions of his works.
[3] Εὔδημος.
[4] Δικαίαρχος.
[5] Αὐτόλυκος. Fl. *c.* 330 B.C.
[6] Ἀρισταῖος. Fl. *c.* 320 B.C.

geometry which treats of analysis, the other two being Euclid and Apollonius. Pappus also relates that Aristæus wrote five books on solid loci,[1] supplementing five others on the elements of conics. Possibly these two works were the same. He also wrote on the five regular solids, and the 13th book of Euclid seems to have owed much to his skill. He was evidently one of those mathematicians of the 4th century B.C. who, inspired by Plato, helped to make possible the works of Euclid and Apollonius.

Callip'pus or Calippus,[2] an astronomer of Cyzicus, was a friend of both Eudoxus and Aristotle. Although not to be looked upon as a geometer, his astronomical observations deserve brief mention, being frequently referred to by Geminus and Ptolemy. The Callippic cycle of 76 years, 940 lunar months, or 27,759 days was such an improvement on the Metonic cycle of 19 years as to have been adopted by ancient astronomers. We have the testimony of Simplicius (6th century) that he was a pupil of Polemar'chus (4th century B.C.), who taught at Cyzicus, and that he lived for a time with Aristotle. Ptolemy tells us that he made astronomical observations on the shores of the Hellespont.

6. The Orient

Orient and Occident. The rise of mathematics in Greece, its remarkable development under the influence of such leaders as Thales, Pythagoras, and Plato, and its distinct characteristics, are such as to make it desirable to consider the Orient and the Occident separately from the time of Euclid until the two were joined by the new intellectual bond established by the Christian missions about the beginning of the 17th century. So little was accomplished in the Orient from 1000 B.C. to 300 B.C., however, that we may properly mention that little in the present chapter. Although each of these two great divisions of the world always influenced the other in developing a system of

[1] Τόποι στερεοί.

[2] Κάλλιππος or Κάλιππος. Fl. *c.* 325 B.C.

mathematics, the East has always been the East, and the West has always been the West. They have had many points in common, particularly in the application of mathematics to astronomy; but the development of a logical geometry, with all of its far-reaching results, is peculiar to the European peoples, while the less rigid and somewhat more poetic phases of mathematics have generally interested the Asiatic mind. Even the ancients recognized this difference, for Quintilian (*c.* 35–*c.* 96) remarks: "From of old there has been the famous division of Attic and Asiatic writers, the former being reckoned succinct and vigorous, the latter inflated and empty."[1]

China. It is an interesting fact that Egypt developed a worthy type of mathematics before 1000 B.C. and then stagnated, that Babylonia did the same, and that China followed a similar course. Was it that the world's vigor was concentrated in Greece? Had the older civilizations burned out? Or was there some subtle influence that subjected the original seats of mathematical thought to canonical expression instead of progressive action? Whatever the answer, between 1000 and 300 B.C. China produced no great classic in mathematics, unless possibly the *Nine Sections*[2] already mentioned, or the *Wu-ts'ao Suan-king* to be mentioned later, belongs to this period. It was rather in the impetus given to commercial calculation through the introduction of coins in the 7th century B.C., at about the same time as they appeared in Asia Minor, that China made her most noteworthy contribution to the progress of arithmetic. Knife money and spade money appeared *c.* 670 B.C., the coins representing such common articles of value as knives and spades. Circular coins were issued later and became the standard forms in the 3d century B.C. As to the methods used in calculating at this time, we are ignorant, but some mechanical means were probably employed in China as well as in other parts of the ancient world. About 542 B.C. the Chinese are known to have used in their calculations bamboo rods, in size and appearance somewhat like a new lead

[1] *Institutes of Oratory*, Bohn ed., XII, x, 16. [2] *K'iu-ch'ang Suan-shu.*

pencil. About 375 B.C. there appeared the earliest Chinese coins with weight or value inscribed upon them, and thus the monetary material for commercial arithmetic became fairly well perfected.

The Compass. As early as the 4th century B.C. there seems to have existed some kind of instrument for indicating the southern direction, probably the compass. In later literature the *chï-nan-kü*[1] (south-pointing chariot) is mentioned, but what it was is unknown.[2] In works of the 4th century it is ascribed to Huang-ti (2704–2595 B.C.) and is also mentioned as being in use in the reigns of Ki-li (1230–1185 B.C.) and his successor (1185–1135 B.C.)[3]

India. As already stated, we have no authentic records of India before the Mohammedan invasion (7th century), almost our only sources of information being the Vedic literature, the Buddhist sacred books, the heroic poems, such inscriptions as remain on monuments, and the metal land grants. Of these, the later Vedic literature, the heroic poems, and the Buddhist writings are all that give us any knowledge of the mathematics of the period from 1000 to 300 B.C. The Vedic writings probably extend down to about 800 B.C., although the *Vedāngas* ("Limbs for supporting the Veda") were written several centuries later. The dates of the *Śulvasūtra* period are unknown. Taking the opinions of various scholars and forming a rough estimate, we may put the ritualistic rules of the *Śulvasūtras* in the five centuries just preceding our era. The rules which have any mathematical interest relate indirectly to the proportions of altars in the temples. They include a statement about Pythagorean numbers, that is, numbers satisfying the relation $x^2 + y^2 = z^2$, and imply a statement of the Pythagorean Theorem itself. There is no reason for believing, however, that the Hindus had the slightest idea of the nature of a geometric proof. There is also evidence of a knowledge of irrationals and

[1] *Chï* means to point with the finger; *nan* means south; and *kü* means chariot.
[2] F. Hirth, *Ancient History of China*, p. 129.
[3] *Ibid.*, p. 135.

of an understanding of the uses of the gnomon.[1] The *Śulvasūtras* also state that the diagonal of a unit square is equal to

$$1+\frac{1}{3}+\frac{1}{3\cdot 4}+\frac{1}{3\cdot 4\cdot 34},$$

or 1.4142156. The area of the circle is asserted[2] to be

$$\left(\frac{13}{8}+\frac{1}{8\cdot 29}-\frac{1}{8\cdot 29\cdot 6}+\frac{1}{8\cdot 29\cdot 6\cdot 8}\right)^2 d^2.$$

Mathematics in the *Śulvasūtras*. The *Śulvasūtras* were changed more or less by such commentators as Apastamba, Baudhāyana, and Kātyāyana. The following statements from the Baudhāyana edition show the style:[3]

"The chord stretched across a square produces an area of twice the size."[4]

"The diagonal of an oblong produces by itself both the areas which the two sides of the oblong produce separately."[5]

The *Lalitavistara*, one of the sacred books of the Hindus, speaks of the arithmetical prowess of the Buddha.[6] Sir Edwin Arnold has put the statement in verse in his *Light of Asia*.[7]

[1] L. von Schroeder, *Pythagoras und die Inder*, Leipzig, 1884; and *Indiens Literatur und Cultur*, Leipzig, 1887; H. Vogt, "Haben die alten Inder den Pythagoreischen Lehrsatz und das Irrationale gekannt?" *Bibl. Math.*, VII (3), 6; A. Bürk, "Das Āpastamba-Śulba-Sūtra," *Zeitschrift der deutschen Morgenländischen Gesellschaft*, LV, 543; LVI, 327; M. Cantor, "Ueber die älteste Indische Mathematik," *Archiv der Math. und Physik*, VIII (3), 63; B. Levi, "Osservazioni e congetture sopra la geometria degli Indiani," *Bibl. Math.*, IX (3), 97; Smith and Karpinski, *The Hindu-Arabic Numerals*, p. 13 and bibliographical notes throughout (hereafter referred to as Smith-Karpinski), Boston, 1911; G. R. Kaye, *Indian Mathematics*, Calcutta, 1915.

[2] Thibaut in the *Journal of the Royal Asiatic Soc. of Bengal*, XLV (1875), p. 227; Dutt, *History of Civ. in Anc. India*, I, 271.

[3] The translation is Dr. G. Thibaut's. See his memoirs "On the Śulvasūtras," *Journ. Royal Asiat. Soc. Bengal*, XLIV (1874); "The Baudhāyana Śulvasūtra," *The Pandit*, 1875; "The Kātyāyana Śulvasūtra," *The Pandit*, 1882; G. Milhaud, "La géométrie d'Apastamba," *Revue générale d. sci.*, XXI, 512.

[4] That is, the square on a diagonal of a square is twice the original square.


[5] That is, the square on a diagonal of a rectangle is equal to the sum of the squares on the two sides; essentially the Pythagorean Theorem.

[6] The date of the birth of Buddha is often placed at 543 B.C., but in Burma, Siam, and Ceylon it is usually given as 80 years earlier, that is, 623 B.C.

[7] It is given in Smith-Karpinski, p. 16. See also the translation of the Mahābhārata in E. Arnold, *Indian Poetry*, London, 6th ed., 1891.

In all this there is nothing that is definite, but there is enough to show that mathematics was not limited to the meager needs of trade and that it was related, as with all thoughtful peoples, to the higher life.

Mesopotamia. Just before the period of which we are speaking the Arameans[1] established a flourishing dominion to the north of the Hebrew territory of Palestine. Their mercantile interests extended into the ancient cities of Assyria, as is proved by such bronze weights as the one shown on page 39.



HEBREWS PAYING TRIBUTE TO SHALMANESER III, KING OF ASSYRIA

This was about 850 B.C. The original is now in the British Museum. From Breasted's *Ancient Times*

By 1000 B.C. they had developed a system of alphabetic writing, and their bills of exchange were known in Mesopotamia, Persia, and India, as those of Babylon had been known before them. All through this early period the records of taxes show that this form of applied arithmetic was ever present.

In the 8th century B.C. the Assyrians subdued Mesopotamia and much of the territory to the west and became the dominating power in Western Asia. They maintained the first great army equipped with weapons of iron and by this means held a large territory in subjection. Militarism, however, eventually proved a weakness, and they in turn succumbed to

[1] Or Syrians, as they are often called.

the power of the Kaldi,—Semitic nomads already mentioned, —who came from the South and who, known to us as the Chaldeans, finally became the ruling power in Mesopotamia.

Contributions of Babylonia and Assyria. In the midst of all these changes two steps in the history of mathematics deserve special mention: (1) the Arameans brought the arithmetic of commerce to a higher standard, and (2) the Babylonians and Chaldeans extended the earlier work in astronomy. The science of astrology had by this time developed as a potent force in civilization, and astronomy had become recognized as the science par excellence. Ptolemy the astronomer (*c.* 150) refers to a Chaldean record of a lunar eclipse of 721 B.C. and to the division of the circle into 360°. The recognition of a zodiac of twelve signs, and the study of the courses of the planets, about 600 B.C., are further evidences of the interest of the Chaldean astronomers in this phase of applied mathematics.

As to astrology,—that daughter of astronomy who nursed her own mother, as Kepler writes of it,—there are various tablets of this period which show in what high esteem it was held. In general they are reports of the following kind to the king: "Two or three times of late we have searched for Mars but have not been able to see him. If the king, my master, asks me if this invisibility presages anything, I reply that it does not."[1]

Science reached its highest point in the reign of Nebuchadnezzar, which closed in 561 B.C. It is true that we have lists of the planets dating from 523 B.C. and from other years,[2] statements of the irregular insertion of intercalary days at about the same period, and a definite recognition of the leap year between 383 and 351 B.C.,[3] but the mathematics of Mesopotamia practically ceased to exist with the decay of the Chaldean power.

[1] Bigourdan, *L'Astronomie*, p. 29; R. C. Thompson, *The Reports of the Magicians and Astrologers of Nineveh and Babylon in the British Museum*, London, 1900.

[2] F. Hommel, "Die Babylonisch-Assyrischen Planetenlisten," in the *Hilprecht Anniversary Volume*, p. 170.

[3] F. H. Weissbach, "Zum Babylonischen Kalender," in the Hilprecht volume above cited, p. 282.

TOPICS FOR DISCUSSION

1. Influences favorable to the development of mathematics among the Greeks from 1000 B.C. to 300 B.C.

2. The nature of logistic and of arithmetic and the reasons for their treatment as unrelated subjects.

3. The advantages of the Greek method of treating arithmetic from the geometric standpoint, particularly in relation to the nature of irrational numbers.

4. The influence of Thales upon the subsequent development of mathematics in Greece.

5. The influences which contributed to the making of the character of Pythagoras.

6. The influence of Pythagoras upon mathematics in general, and particularly upon geometry and the theory of irrationals.

7. Music as a branch of ancient mathematics.

8. Beginnings of a kind of infinitesimal calculus in Greece, particularly with respect to the method of exhaustion.

9. Types of geometric propositions that attracted special attention in this period, thus showing the nature of geometry before the time of Euclid.

10. The influence of Plato upon mathematics in general and upon geometry in particular.

11. The influence of astronomy upon mathematics in Greece, particularly with reference to geometry and a primitive trigonometry.

12. The early steps in the invention of conic sections.

13. The study of higher plane curves among the Greeks in the period under discussion.

14. The influence of Aristotle upon mathematics in general, and particularly upon its applications.

15. Nature of mathematics in the Orient in this period.

16. General distinction between the mathematics of Greece and that of the East.

17. Mysticism of numbers as found in the Orient, in Mesopotamia, and in the West.

18. Early studies in the history of mathematics among the Greeks.

19. The recognition of the sphericity of the earth by various leading Greek philosophers.

20. The nature of the mathematics of the *Śulvasūtras*.

CHAPTER IV

THE PERIOD FROM 300 B.C. TO 500 A.D.

1. The School of Alexandria

Chronological and Geographical Considerations. The reason why the limitations of 300 B.C. and 500 A.D. are arbitrarily chosen for this chapter is that these dates mark approximately the period of influence of the greatest mathematical school of ancient times, the School of Alexandria. Moreover, the first of these dates is approximately that of Euclid, the world's greatest textbook writer, and the second is that of Boethius, whom Gibbon characterizes as "the last of the Romans whom Cato or Tully could have acknowledged for their countryman."

Within this period Greek civilization passed away, Rome rose and fell, and the ancient mathematics of the West descended from its most exalted to its most debased estate. We have, therefore, the most significant period of ancient mathematical history, at least in the matter of actual production, and we have the Mediterranean world, probably the most interesting of all ancient civilizations.

The School of Alexandria. The greatest mathematical center of ancient times was neither Crotona nor Athens, but Alexandria. Here it was, on the site of the ancient town of Rhacotis, in the Nile Delta, that Alexander the Great founded a city worthy to bear his name. Upon the death of the great Macedonian conqueror (323 B.C.) the vast domain which he had brought under his control was broken up. After the death of Antig'onus, his ablest general, the empire fell into three parts. Alexander's friend and counselor, and possibly his blood relative, Ptol'emy[1] So'ter (Ptolemy the Preserver), came into

[1] Πτολεμαῖος; Latin, *Ptolemaeus*.

possession of Egypt, Antigonus the younger laid claim to Macedonia, while Seleu'cus took for his part the provinces of Asia. Under Ptolemy's benevolent reign (323–283 B.C.) Alexandria became the center not only of the world's commerce but also of its literary and scientific activity.[1] Here was established the greatest of the world's ancient libraries and its first international university. Cardinal Newman, in speaking of these two features, says with poetic feeling that "as the first was the embalming of dead genius, so the second was the endowment of living." Here were trained more great mathematicians than in any other scientific center of the ancient world. With Alexandria are connected the names of Eu'clid. Archime'des, Apollo'nius, Eratos'thenes, Ptol'emy the astronomer, He'ron, Menela'us, Pap'pus, The'on, Hypa'tia, Diophantus, and, at least indirectly, Nicom'achus. Today, however, not the slightest trace remains of the famous library and museum, and even their exact locations are merely conjectural.

2. EUCLID


Euclid.[2] Of all the great names connected with Alexandria, that of Euclid is the best known. He was the most successful textbook writer that the world has ever known, over one thousand editions of his geometry having appeared in print since 1482,[3] and manuscripts of this work having dominated

[1] For a summary of the causes of its rise and a description of its library see W. Kroll, *Geschichte der klassischen Philologie*, p. 12 (Leipzig, 1908); hereafter referred to as Kroll, *Geschichte*.

[2] Εὐκλείδης. Fl. *c.* 300 B.C. The leading work upon Euclid and his *Elements* is that of Sir T. L. Heath, *The Thirteen Books of Euclid's Elements*, 3 vols., Cambridge, 1908. The best Greek and Latin edition of Euclid's works is Heiberg and Menge, *Euclidis Elementa*, Leipzig, 1883–1916. Of the many works and articles on the life of Euclid the following may be consulted to advantage: A. De Morgan, "Eucleides," in Smith's *Dict. of Greek and Roman Biog.*; T. Smith, *Euclid, his life and system*, New York, 1902; W. B. Frankland, *The Story of Euclid*, London, 1902; G. B. Biadego, "Euclide e il suo Secolo," in Boncompagni's *Bullettino*, V, 1; P. Tannery, *Pour l'histoire de la science hellène*, Appendix II (Paris, 1887); Gow, *Greek Math.*, p. 195; M. C. P. Schmidt, *Realistische Chrestomathie aus der Litteratur des klass. Altertums*, I, 1 (Leipzig, 1900), hereafter referred to as Schmidt, *Chrestomathie*; Pauly-Wissowa's *Real-Encyclopädie*, Vol. VI (Stuttgart, 1909), has an extensive article on Euclid.

[3] P. Riccardi, *Saggio di una Bibliografia Euclidea*, p. 4. Bologna, 1887.

the teaching of the subject for eighteen hundred years preceding that time. He is the only man to whom there ever




PAGE FROM A TRANSLATION OF EUCLID'S ELEMENTS

This manuscript was written *c.* 1294. The page relates to the propositions on the theory of numbers as given in Book IX of the *Elements*. The first line gives Proposition 28 as usually numbered in modern editions

came or ever can come again the glory of having successfully incorporated in his own writings all the essential parts of the accumulated mathematical knowledge of his time.

Of the life of Euclid nothing definite is known. A recent writer expresses the belief that the evidence indicates that he was born as early as 365 B.C. and that he wrote the *Elements* when he was about forty years old,[1] but we have no precise information as to his birthplace, the dates of his birth and death, or even his nationality. It was formerly asserted that he was born at Meg'ara, a Greek city, but it is now known that Euclid of Megara was a philosopher who lived a century before Euclid of Alexandria. The Euclid in whom we are interested may have been a Greek or he may have been an Egyptian who came to the Greek colonial city of Alexandria to learn and to teach. There is some reason for believing that he studied in Athens, but in the way of exact information we have nothing concerning him. Under any circumstances, the period of his real influence upon mathematics begins about 300 B.C.



PARCHMENT ROLL

Upon such a roll a "book" of Euclid was written

The Books of Euclid. As was the custom in the days when all treatises were written on long strips of parchment or papyrus, the separate parts of his work were rolled up and called volumes, from a Latin root meaning to roll. Because of the difficulty of handling large rolls, they were cut into smaller rolls known as *biblia* (*βιβλία*, bibles), a word meaning books. Hence we have the books of Homer, the books of geometry, the books of the Bible, and so on. Euclid's greatest work is known as the *Elements*,[2] and in the books relating to geometry there was arranged that mass of material treating of circles, rectilinear figures, and ratios which had accumulated during the two centuries following the death of Pythagoras. No doubt there were also many propositions that were original with Euclid; but the feature which made his treatise famous, and which

[1] H. Vogt, *Bibl. Math.*, XIII (3), 193; but see Heath, *History*, I, 354.

[2] Στοιχεῖα. So great was Euclid's fame that he was known to the Greeks as ὁ στοιχειωτής, "teacher of the *Elements*."

accounts for the fact that it is the oldest scientific textbook still in actual use, is found in its simple but logical sequence of theorems and problems. It has been said of Shakespeare that he "took the stillborn children of lesser men's brains and breathed on them the breath of life,"—and so it was with Euclid.

Contents of the *Elements*. The various books of the *Elements* treated of the following topics, respectively: I, Congruence, parallels, the Pythagorean Theorem; II, Identities which we would now treat algebraically, like $(a+b)^2=a^2+2ab+b^2$, but which were then treated geometrically; areas; the Golden Section; III, Circles; IV, Inscribed and circumscribed polygons; V, Proportion treated geometrically; in part, a geometric way of solving fractional algebraic equations; VI, Similarity of polygons; VII–IX, Arithmetic (the ancient theory of numbers) treated geometrically; X, Incommensurable magnitudes; XI–XIII, Solid geometry. We have in this text the earliest extant evidence of a systematic arrangement of definitions, axioms, postulates, and propositions. Euclid differs from most of our modern writers on geometry in his greater seriousness of purpose, in his desire to be more rigorous, and in the following details of treatment: He has no intuitive geometry as an introduction to the logical; he uses no algebra as such; he demonstrates the correctness of his constructions before using them, whereas we commonly assume the possibility of constructing figures and postpone our proofs relating to constructions until we have a fair body of theorems; he does not fear to treat of incommensurable magnitudes in a perfectly logical manner; and he has no exercises of any kind.

Euclid's Other Works. Euclid wrote a number of other works. Among them are the *Phænomena*,[1] dealing with the celestial sphere and containing twenty-five geometric propositions; the *Data*; possibly a treatise on music;[2] and works on optics,[3]

[1] Φαινόμενα.

[2] Εἰσαγωγὴ Ἁρμονική or else the Κατατομὴ Κανόνος, both doubtful. See Heath *Euclid*, I, 17.

[3] Ὀπτικά.

porisms,[1] and catoptrics.[2] He also wrote a work on divisions of figures, touching upon questions which arise, for example, in surveying.[3]

Immediate Effect of Euclid's Work. The natural effect of Euclid's work on geometry was to give rise to the feeling that elementary geometry had attained to perfection and that the next step in the progress of mathematics must be in the direction of some kind of higher geometry or else in the field of mensuration. As a result, mathematics pursued both courses, at first with little effect, as was the case with the predecessors of Euclid, and then, when another genius appeared, with great rapidity.

Minor Writers. There were, for example, at first such minor writers as Co'non[4] of Samos, who, influenced by his observations of the coiled basketry work of the Egyptians, may have invented the spiral of which Archimedes developed the properties. He is also mentioned by Apollonius (*c.* 225 B.C.) as having studied the number of points of intersection of two conics. There was also Nicot'eles[5] of Cyrene, possibly a student in Alexandria, of whom Apollonius speaks as his predecessor in the study of conics. Still another writer of influence appeared in the person of the astronomer Aristar'chus,[6] a native of Samos but a teacher at Alexandria. It was he who first showed how to find, by means of the Pythagorean triangle, the relative distances of the sun and the moon from the earth, and for nearly two thousand years no better plan was known. His instruments of observation were such as to make his result far from being even approximately correct.[7] His greatest glory, however, lies

[1] Relating to methods of solution. See M. Breton, in *Journal de Math. pures et appliquées*, XX (1), III (2).

[2] Κατοπτρικά.

[3] Περὶ Διαιρέσεων βιβλίον. R. C. Archibald, *Euclid's Book on Divisions of Figures*, Cambridge, 1915.

[4] Κόνων. Fl. *c.* 260 B.C.

[5] Νεικοτέλης or Νικοτέλης. Fl. *c.* 250 B.C.

[6] Ἀρίσταρχος. Born *c.* 310 B.C.; died *c.* 230. See Sir T. L. Heath, *Aristarchus of Samos*, Oxford, 1913; *The Copernicus of Antiquity*, London, 1920.

[7] Carl Snyder, *The World Machine*, chap. vii, "Aristarchus and the distance and grandeur of the sun" (London, 1907); Bigourdan, *L'Astronomie*, 252.

in the fact that he was the first to place the sun in the center of the universe, asserting that the earth and the other planets revolved about it, thus anticipating Copernicus by seventeen centuries. In the field of arithmetic he found $\sqrt{2}$, possibly by a method analogous to that of continued fractions.[1]

There are also extant various papyri of the Ptolemaic period containing information about the financial problems of Egypt. These problems relate chiefly to taxes and the cost of various commodities, but they add nothing to our information as to methods of calculation in ancient times.[2]

Men like Conon were merely the usual heralds calling out the approach of genius. The three men whose advent they heralded were Archimedes, Apollonius, and Heron. Before speaking of Archimedes, however, reference should be made to a scholar whose interests were so scattered as to make his contributions to pure mathematics of relatively little importance. This man was the poet, librarian, arithmetician, and geographer Eratos'thenes.

3. Eratosthenes and Archimedes

Eratosthenes[3] lived some years after Euclid, and was one of the greatest scholars of Alexandria.[4] His admirers exaggerated his attainments by calling him "the second Plato,"[5] and some have thought that his nickname "Beta" signified that he was the second of the wise men of antiquity, the Greek letter *beta* standing for two. Others have said that he was called by this name because his room in the university bore the number two. But whether or not his followers ranked him second among the wise men of Greece, we are justified in calling him the first prominent geographer of antiquity. He was educated at Athens

[1] P. Tannery, *Mémoires de Bordeaux*, V (2), 237; IV (3), 79.


[2] For a bibliography, a list of the papyri, and a summary of the information available, see H. Maspero, *Les Finances de l'Égypte sous les Lagides*, Paris, 1905.

[3] Ἐρατοσθένης. Born at Cyrene, *c.* 274 B.C.; died *c.* 194.

[4] Schmidt, *Chrestomathie*, I, 29, 114. See also *Bibl. Math.*, XIII (3), 193.

[5] B. Baldi, *Cronica de Matematici*, p. 29 (Urbino, 1707), hereafter referred to as Baldi *Cronica*; Heath, *History*, II, 104.

and is known to have taught at Alexandria after *c.* 240 B.C., to have been librarian of the university, and to have been a poet of some merit. His contribution to arithmetic was his sieve,[1] a method of sifting out the composite numbers in the natural series, leaving only primes. This he did by writing all the odd numbers and then canceling the successive multiples of each, one after the other, thus: 3, 5, 7, ~~9~~, 11, 13, ~~15~~, 17, 19, ~~21~~, 23, ~~25~~, ~~27~~, 29, 31, ~~33~~, ~~35~~, 37, ~~39~~, . . .[2] Prime numbers



MAP OF THE WORLD ACCORDING TO ERATOSTHENES

This shows the knowledge of the geography of the world in the 3d century B.C. and should be compared with the map of Hecatæus (517 B.C.) shown on page 77. From Breasted's *Ancient Times*

have been studied from that time until the present, but no general formula is yet known for detecting all of them. For example, we do not yet know whether there are an infinite number of primes of the form $x^2 + 1$, whether $2 = x - y$ has an infinite number of prime solutions, or whether a prime number can always be found between n^2 and $(n+1)^2$.

Earth Measure. Of the mathematical achievements of the Greek astronomers none is more interesting than the measurement of the circumference and diameter of the earth by Eratosthenes, the first noteworthy step in the science of

[1] Κόσκινον. It was called the *Cribrum Arithmeticum* by Latin writers
[2] M. C. P. Schmidt, *loc. cit.*, I, 114 (Greek text).

geodesy.[1] Learning that the sun at noonday was exactly in the zenith at Syene[2] when it was 7° 12′ south of the zenith at Alexandria, he decided that Alexandria was 7° 12′ north of Syene on the earth's surface. Since the distance was known to be 5000 stadia, and since 7° 12′ = $\frac{1}{50}$ of 360°, he judged the circumference of the earth to be 50 × 5000 stadia, or 250,000 stadia. This result he altered to 252,000 stadia so as to have




DIAGRAM SHOWING METHOD USED BY ERATOSTHENES IN MEASURING THE EARTH

Eratosthenes found that, when the sun was directly over Syene, at the First Cataract, it was 7° 12′ south of Alexandria. From this he computed the circumference of the earth. From Breasted's *Ancient Times*

700 stadia, a more convenient number, to a degree, and from this he computed the diameter to be the equivalent of 7850 miles, in our system of measure, which is only 50 miles less than the polar diameter as we know it.[3]

Eratosthenes also stated that the distance between the tropics is $\frac{11}{83}$ of the circumference, which makes the obliquity of the

[1] Carl Snyder, *The World Machine*, chap. vi, "Eratosthenes and the earliest measures of the earth" (London, 1907); Schmidt, "Erdmessung des Eratosthenes," Greek text, *Chrestomathie*, II, 105; see also I, 29.

[2] Sy e'ne, Συήνη, the modern Assouan (Arabic from *al* + *Syene*) at the first cataract of the Nile.

[3] The problem of earth measure is more fully treated in Volume II, Chapter V.

ecliptic[1] 23° 51′ 20″. Plutarch tells us that he found the sun to be 804,000,000 stadia from the earth, and the moon to be 780,000 stadia,—results which are remarkably close when we consider the instruments then in use. That the knowledge of geography had increased in the preceding 250 years may be seen by comparing his map with that of Hecatæus (517 B.C.).[2]

In one of his letters Eratosthenes also discussed the problem of the duplication of the cube.

Archime'des[3] was a friend of Eratosthenes and, if the testimony of Plutarch is accepted, was related to King Hiero. Leibniz praised his genius by saying that those who knew his works and those of Apollonius marveled less at the discoveries of the greatest modern scholars.[4] These words are justified, for Archimedes anticipated by nearly two thousand years some of the ideas of Newton and his contemporaries, and in the application of mathematics to mechanics he had no equal in ancient times. One of the Italian historians of mathematics[5] uses the happy phrase that he had "a genius more divine than human," and Pliny calls him "the god of mathematics," a phrase which one of his French translators felicitously renders as "the Homer of geometry."

It is related that Archimedes set fire to the besieging ships in the harbor of Syracuse by the aid of burning mirrors, and there is nothing improbable in the idea that he may have

[1] This term seems first to have been used by Ambrosius Aurelius Theodosius Macrobius, a grammarian of *c.* 400.


[2] On the general subject of the history of mathematical geography consult S. Günther, *Studien zur Geschichte der math. und physik. Geographie*, with extensive series of bibliographies (Halle a. S., 1879).

[3] Ἀρχιμήδης. Born at Syracuse, the modern Siracusa, Sicily, 287 B.C.; died at Syracuse, 212 B.C. Sir T. L. Heath, *Archimedes*, Cambridge, 1897 (hereafter referred to as Heath, *Archimedes*); German translation by Kliem, Berlin, 1914; P. Midolo, *Archimede e il suo tempo*, Syracuse, 1912; C. Snyder, *The World Machine*, chap. x, "Archimedes and the first ideas of gravitation," London, 1907; Schmidt, *Chrestomathie*, III, 64, 100; Heath, *History*, II, 16.

[4] "Qui Archimedem et Apollonium intelligit, recentiorum summorum virorum inventa parcius mirabitur," *Archimedis Opera* (Geneva, 1768), V, 460. The definitive edition is that of Heiberg, Leipzig, 1880–1915.

[5] Baldi, *Cronica*, p. 26: "Hebbe ingegno più divino, che humano."

made them at least untenable by the soldiers.[1] A glance at the map, and particularly at the lesser harbor then in use, and a consideration of the fact that the ships were then hardly larger than our pleasure yachts of today, and that they were all anchored close to the rocky shore, will show that the task was not so great as one might at first suppose. At a time when the breeze was blowing in from the sea, escape would have been difficult, even with oars.



ARCHIMEDES


Conjectural portrait bas-relief in the Capitoline Museum at Rome. Date uncertain

Archimedes and Mechanics. Plutarch, in his life of Marcellus, relates this incident to illustrate the genius of Archimedes in mechanics:

> Archimedes . . . had stated that, given the force, any given weight could be moved; and even boasted . . . that, if there were another earth, by going into it he could remove this one. Hiero being struck with amazement at this, . . . [Archimedes] fixed accordingly upon a ship of burden . . . which could not be drawn out of the dock without great labor and many men; and loading her with many passengers and a full freight, sitting himself the while far off with no great endeavor, but holding the head of the pulley in his hand and drawing the cords by degrees, he drew the ship in a straight line as smoothly and evenly as if she had been in the sea.

[1] On the subject of burning mirrors in Greek literature see Sir T. L. Heath, in *Bibl. Math.*, VII (3), 225.

The Sand Reckoner. Archimedes saw the defects of the Greek number system, and in his *Sand Reckoner*[1] he suggested an elaborate scheme of numeration, arranging the numbers in octads, or the eighth powers of ten. In this work he



MAP OF ANCIENT SYRACUSE

Showing the general situation at the time of the Third Peloponnesian War and continuing until the time of Archimedes. On the river Anapus here shown papyrus still grows luxuriantly. From Breasted's *Ancient Times*

recognized, in substance, that $a^m a^n = a^{m+n}$, a law that is the basis of our present operations by logarithms.

Other Mathematical Activities. Among his many activities was the summation $\sum_1^n n^2$, the first example of the systematic treatment of higher series of any kind. By the intersection of

[1] Ψαμμίτης (*psammites*); Latin, *arenarius* or *harenarius*.

conics he was able to solve cubic equations which we should now write in the form $x^3 \mp ax^2 \pm b^2c = 0$. He also succeeded in squaring a parabola,[1] that is, in finding the area of a segment, showing that it is two thirds of a circumscribed parallelogram. In the measure of a circle he showed that $3\frac{1}{7} > \pi > 3\frac{10}{71}$. In his work in mensuration Archimedes included the sphere, cylinder, and cone, the rules concerning the two latter having already been known to Menæchmus. He also studied ellipsoids and paraboloids of rotation. In his treatise on the mensuration of circles and round bodies he was aided by the method of exhaustion which had been developed by Menæchmus and others. In the study of specific gravity and the center of gravity of planes and solids he was a pioneer, and in the study of hydrostatics he was unequaled in the Greek period. He is also known for his study of spirals, possibly led thereto by his friend Conon. In general, he stands out as one of the greatest mathematicians and physicists in all history.

Method of Archimedes. In 1906 Professor Heiberg, who had already edited the works of Archimedes, discovered in Constantinople a manuscript on certain geometric solutions derived from mechanics.[2] This is especially interesting from the fact that it sets forth the method taken by Archimedes in deriving geometric truths from principles of mechanics. Some idea of the working of his mind may be obtained from the following:

After I had thus perceived that a sphere is four times as large as the cone whose base is the largest circle of the sphere and whose altitude is equal to the radius, it occurred to me that the surface of a sphere is four times as great as its largest circle, in which I proceeded from the idea that just as a circle is equal to a triangle whose base is the periphery of the circle and whose altitude is equal to the radius, so a sphere is equal to a cone whose base is the same as the surface of the sphere and whose altitude is equal to the radius of the sphere.[3]

[1] See Volume II, Chapter X.

[2] Translated by Lydia G. Robinson (Chicago, 1909) and by Sir T. L. Heath (Cambridge, 1912).

[3] For his method with respect to the parabola, see Volume II, Chapter X.

Death of Archimedes. Of the death of Archimedes at the siege of Syracuse under Marcellus (212 B.C.), Plutarch has this interesting record:

Nothing afflicted Marcellus so much as the death of Archimedes who was then, as fate would have it, intent upon working out some problem by a diagram, and having fixed both his mind and his eyes upon the subject of his speculation, he did not notice the entry of the Romans nor that the city was taken. In this transport of study and contemplation a soldier unexpectedly came up to him and commanded him to go to Marcellus. When he declined to do this before he had completed his problem, the enraged soldier drew his sword and ran him through. Others write that a Roman soldier ran towards him with a drawn sword and threatened to kill him, whereupon Archimedes . . . earnestly besought him to stay his hand that he might not leave his work incomplete; but the soldier, unmoved by his entreaty, instantly slew him. Others again relate that Archimedes was carrying to Marcellus some mathematical instruments, dials, spheres, and angles, by which the size of the sun might be measured . . . , and some soldiers . . . thinking that he carried gold in a vessel, slew him. Certain it is that his death brought great affliction to Marcellus; that he ever after regarded the one who killed him as a murderer; and that he sought for the kindred of Archimedes and honored them with signal favors.

Discovery of the Tomb of Archimedes. In his *Tusculan Disputations* (V, 23) Cicero relates that he himself discovered the tomb of Archimedes "when the Syracusans knew nothing of it and even denied that there was any such thing remaining." He relates the incident as follows:

I remembered some verses which I had been informed were engraved on his monument, and these set forth that on the top of the tomb there was placed a sphere with a cylinder. When I had carefully examined all the monuments . . . I observed a small column standing out a little above the briers, with the figure of a sphere and a cylinder upon it. . . . When we could get at it and were come near to the front of the pedestal, I found the inscription, though the latter parts of all the verses were effaced almost half away. Thus one of the noblest cities of Greece, and one which at

one time likewise had been very celebrated for learning, had known nothing of the monument of its greatest genius, if it had not been discovered to them by a native of Arpinum.[1]

Of his works that have come down to us, those which are of chief interest in the history of mathematics are the ones on the quadrature of the parabola, on the sphere and the cylinder, on the measure of a circle, on spirals, conoids, and spheroids, and on notation. Archimedes seems also to have been interested in astronomy,[2] although no work of his upon this subject is extant.

4 Apollonius and his Successors

Apollo'nius of Per'ga[3] was known as "the great geometer" because of his work on conic sections. He was educated in Alexandria, and since he died under Ptolemy IV (Philop'ator, reigned 222–205 B.C.) he very likely knew Eratosthenes. He improved on the numeration system of Archimedes by using 10^4 as the base. This number, the myriad,[4] had long been in use in the Orient, and was the base of all great systems of numeration in the East as well as in Europe for many centuries. His chief work was on the conic sections, to which he gave the names ellipse, parabola, and hyperbola.[5]

This work[6] consisted of eight books, the first four of which have come down to us in Greek and the next three in Arabic, the last book being lost. In the first book Apollonius shows how the three conics are produced from the same cone. He uses a kind of coordinate system, the diameter serving for what we call the x-axis, and the perpendicular at the vertex serving for the y-axis. Books I–IV probably contain little that was

[1] Translated by C. D. Yonge. London, 1891.

[2] Livy (XXIV, 34) speaks of him as "unicus spectator coeli siderumque."

[3] Apollonius Pergæus, Ἀπολλώνιος. Fl. *c.* 225 B.C.; born at Perga, in Pamphylia, on the south coast of Asia Minor; Heath, *History*, II, 126.

[4] Μύριοι, ten thousand.

[5] Ἔλλειψις, παραβολή, ὑπερβολή.

[6] Sir T. L. Heath, *Apollonius of Perga* (Cambridge, 1896); J. L. Heiberg. edition of his works (Leipzig, 1891); E. Halley, *Apollonii Pergaei Conicorum libri octo . . .* (Oxford, 1710), and his *De Sectione Rationis libri duo* (Oxford, 1706); G. Eneström, *Bibl. Math.*, XI (3), 7.

not already known, but he arranged the material anew, as Euclid had arranged systematically many propositions that his own predecessors had known. Books V–VII seem to contain the discoveries which he himself had made. Book V treats of normals to a curve; Book VI, of the equality and similarity of conics; and Book VII, of diameters and rectilinear figures described upon these diameters. In general, his propositions are those which we now treat by analytic geometry, his method being synthetic and analogous to that of Euclid with respect to the circle and rectilinear figures.

Apollonius wrote various other works on geometry, including one on plane loci.[1] Ptolemy speaks of him as having been also a contributor to astronomy, but he probably confused him with another Apollonius who lived a little earlier.

In the works of Apollonius Greek mathematics reached its culminating point. Without Euclid as a guide, Apollonius could never have reached the summit; together, they dominated geometry for two thousand years.

Minor Writers. After the death of Apollonius no great writers on mathematics appeared for about two centuries. Greek civilization was receding. War was taking its toll. Arithmetic seemed for a time to sink into a comatose state after the slight attempts of Eratosthenes, and elementary geometry seemed to die with Euclid and Apollonius.

There was some indication at this time of the coming birth of a geometry of higher plane curves, just as the centuries immediately preceding Euclid and Apollonius foretold the appearance of these masters. The Greek civilization, however, had not strength to fulfill the momentary promise. Many generations had to come and go before another people, living far to the north, speaking a new language and making use of new symbols and of a new method, brought to light the theory.

[1] For his geometric works see a convenient list in Gow, *Greek Math.*, pp. 246, 261. See also various restorations of his lost works, such as Woepcke's in the *Mémoires présentés par divers savants à l'Académie des Sciences*, XIV; reprint (Paris, 1856); G. Eneström, *Bibl. Math.*, XI (3), 7, 8. Since we are concerned at present with the elementary field, the reader who wishes to consider the general history of conics should consult the *Encyklopädie*, III (ii), 1.

Perseus. Of those who treated of these curves, one of the first was Per'seus,[1] who lived *c.* 150 B.C. He wrote on sections of the anchor ring,[2] but his works are known only by references of later writers.

Nicomedes. Among the minor geometers of this period one of the best known was Nicome'des,[3] who flourished *c.* 180 B.C. and who invented a curve called the conchoid (mussel-shaped), by which the trisection of an angle is easily effected.[4]

Diocles. A probable contemporary of his, Di'ocles,[5] invented the cissoid,[6] by which the duplication of a cube can be accomplished.[7] He also studied[8] the problem of Archimedes, to cut a sphere by a plane in such a way that the volumes of the segments shall have a given ratio.

At about the same time, say 180 B.C., Zenodo'rus[9] wrote upon isoperimetry,[10] but most of his writings are lost.

A little later, Poseido'nius[11] taught in Rhodes, acquired a high reputation as a cosmographer and geometer, and had the honor of claiming Cicero and Pompey as his pupils. His measurements of the distance to the sun and of the circumference of the earth, known to us through the works of Cleome'des (*c.* 40 B.C.), were far from being as accurate as those of Eratosthenes, but his results seem to have been more generally accepted by ancient geographers.

Contributions of the Astronomers. Owing largely to the influence of the Egyptian and Chaldean priest-astronomers, whose achievements had attracted more and more attention on the part of the Greeks as intercourse became more free,

[1] Περσεύς. Heath, *History*, II, 203.

[2] Σπεῖρα, a torus, or ring-shaped solid of revolution (in a special case, an anchor ring), a solid already studied by Eudoxus.

[3] Νικομήδης. His birthplace is unknown. Heath, *History*, II, 199.

[4] This method and the use of other important curves are considered in Volume II, Chapter V.

[5] Διοκλῆς.

[6] "Ivy-shaped" curve.

[7] Such cases are more fully discussed in Volume II, Chapter V.

[8] In his Περὶ πυρείων.

[9] Ζηνόδωρος.

[10] Fourteen of his propositions have been preserved by Pappus (V, Pt. I) and Theon of Alexandria (*Comment. Almagest.*).

[11] Ποσειδώνιος, born at Apameia, in Syria, *c.* 135 B.C.; died *c.* 44. Sometimes called the Apamean. The name is also spelled Posidonius.

the 2d century B.C. was noteworthy for its advance in the study of the stars. In this century two names stand out prominently, not merely for their work as observers but because of their mathematical attainments.

Hypsicles. The first of these astronomers was Hyp'sicles[1] of Alexandria, who may have written the so-called fourteenth book of Euclid's *Elements*, containing seven propositions on regular polyhedrons. He was also interested in polygonal numbers,[2] in progressions, and in certain indeterminate equations. His prime interest, however, was in astronomy, and about his time there begins, among the Greeks, the division of the circle into 360° and the definite, scientific use of sexagesimal fractions, which the Babylonians had already suggested.

Hipparchus. About this time Hippar'chus,[3] working chiefly in Rhodes, wrote a famous work on astronomy in which were set forth the basic principles of the science. For this work he needed to measure angles and distances on a sphere, and hence he developed a kind of spherical trigonometry. Plane trigonometry had as yet taken only rudimentary form, and, so far as we know, there were no tables of functions. Hipparchus worked out a table of chords, that is, of double sines of half the angle, and thus was definitely begun the science of trigonometry. With him also began the theory of stereographic projection, a phase of geometry which Agatharchus (470 B.C.) had already put in practice. Hipparchus used it for the purpose of representing the projection of the celestial sphere upon the plane of the equator. He left a catalogue of 850 fixed stars, a number which Ptolemy (*c.* 150) increased to 1022 and which was not further materially increased until modern times.[4]

[1] Ὑψικλῆς. Fl. *c.* 180 B.C.. De Morgan places him *c.* 160 A.D. on general rumor, but asserts that he could not have lived before 550 A.D. and places Diophantus even later! The Arab writers say that he was born in Ascalon. See Smith's *Dict. of Greek and Roman Biog.*, II, 541. [2] See Volume II, Chapter I.

[3] Ἵππαρχος. Born at Nicæa, in Bithynia, Asia Minor, *c.* 180 B.C.; died *c.* 125.

[4] F. Boll, "Die Sternkataloge des Hipparch und des Ptolemaios," *Bibl. Math.* II (3), 185. This article disputes the usual assertion that Hipparchus listed 1022 stars, and asserts that he knew only about 850, the rest being catalogued by Ptolemy. See also Heath, *History*, II, 255.

Mathematics of Rome. The next two or three centuries witnessed the rise of the Roman military power and the consequent suppression of intellectual ideals. Art, philosophy, science, politics, ethics, and mathematics had all sunk to a low level. In literature, however, Rome made progress, although Vergil took Homer as his model, and Cicero followed in the footsteps of Demosthenes. Even as early as the 6th century B.C. Etruscan art had become wholly Greek in its technique and in its use of Greek mythology and customs. Rome simply followed in the same lines, not merely in art but in letters and science as well. In mathematics she showed no originality and possessed no high ideals. The science was worth to her precisely what it would fetch in the coin of the realm and no more. Rome created a goddess Numeraria, but she favored the acquisition of wealth rather than the creation of men. Money is everything only when man is nothing.[1] Whenever a genius like Heron of Alexandria, for example, arose in Greco-Roman territory, his interests were usually in the applications of the science as already developed, not in extending the boundaries. As to Rome herself, it is noteworthy how many of her scholars and literary men were born outside of Italy. Spain furnished the two Senecas, Lucanus, Martial, Quintilian, and probably Hyginus; France, Favorinus and Domitius Afer; Palestine, Josephus; Egypt, Philo; and Greece, Plutarch and Epictetus.[2] If Pythagoras and Archimedes may be ranked as dwellers in Italy, they were essentially Greek, and after the death of the latter, exact science may be said to have taken her departure. Cicero lamented this attitude of the Latin mind, contrasting the high honor in which geometry was held among the Greeks with the lack of appreciation on the part of the Romans.[3]

In this period there flourished Marcus Terentius Var'ro (116–28 B.C.), whom Quintilian called the most learned of the

[1] "L'argent n'est tout que dans les siècles où les hommes ne sont rien." Libri, *Histoire*, I, p. xiv.

[2] Libri, *Histoire*, I, p. 53.

[3] "In summo honore apud Graecos geometria fuit; itaque nihil mathematicis illustrius; at nos ratiocinandi metiendique utilitate huius artis terminavimus modum." *Tusculanarum Disputationum Libri V*, I, 2.

Romans. St. Augustine said of him that he had read so much that we wonder that he had time to write anything, and that he had written so much that we can scarcely believe that anyone could find time to read it all. Of such a dilettante nothing very scientific could be expected, and his one extant work[1] certainly has no great merit. In his *Disciplinarum Libri* he treated of arithmetic, and he wrote a *Mensuralia* or *De Mensuris* which related to practical mensuration, but, so far as we know, his works were mere compilations.[2] He is the one of the few pre-Christian mathematicians of whom we have a contemporary portrait, his profile appearing on a coin struck when he was the proquæstor of Pompey.

Geminus. Among those who showed any interest in the history of mathematics at this time the best known is Gem'inus,[3] who was a native of Rhodes but may have written in Rome. He is said to have divided mathematics into two groups, the pure group, including arithmetic (in the ancient sense) and geometry, and the applied group, including mechanics, astronomy, optics, geodesy, canonics, and logistic. Proclus, who lived in the 5th century A.D., tells us that he wrote a geometry which treated of spirals, conchoids, and cissoids. Only one of his works is extant, the *Phænomena*,[4] a treatise on astronomy. Proclus has numerous historical notes based upon the works of Geminus, these notes being found mostly in fragments that remain of the latter's *Arrangement of Mathematics*.

Minor Sources. Another source of information in the history of mathematics was written by a Sicilian who flourished a little later than this, having apparently been living in the year 8 B.C. Diodo'rus, usually called Diodorus Siculus, was born in Agyrium on the island of Sicily. He wrote forty books on history, and while his style is not good and his facts are ill-sorted, his

[1] *De Re Rustica Libri III.*

[2] Montucla, *Histoire*, I (2), 488, tells us that a MS. of his arithmetic was extant as late as the close of the 16th century, but it is now lost.

[3] Γεμῖνος or Γεμεῖνος. Fl. *c.* 77 B.C. M. C. P. Schmidt, "Was schrieb Geminos?" *Philologus*, XLV, 63; *Chrestomathie*, I, 45; C. Tittel, *De Gemini Studiis Mathematicis*, Leipzig, 1895; Heath, *History*, II, 222.

[4] Εἰσαγωγὴ εἰς τὰ Φαινόμενα. It was first printed, in Greek and Latin, in 1590.

works give us considerable information on the nature of the mathematics studied in the classical period, particularly in the schools of Egypt.

Still another writer to whom we are indebted for numerous bits of knowledge relating to the ancient mathematics is Strabo[1] the geographer. His second book deals with mathematical geography and passes certain criticisms on the map of the world prepared by Eratosthenes.

A little later than Geminus there lived P. Nigid'ius Fig'ulus,[2] a Pythagorean philosopher, who was highly esteemed in his time. He was known to his contemporaries as a philosopher, statesman, mathematician, and astrologer, but his contributions had little influence. In the field of mathematical astronomy he wrote *De Sphaera Barbarica et Graecanica*, but only fragments of his works have come down to us.

Probably contemporary with Figulus (but we are not sure of the dates) there was a certain geometer named Dionysodo'rus,[3] who lived in Asia Minor, probably in Ami'sus.[4] He is known for a solution of the problem proposed by Archimedes and already discussed by Diocles,—to cut a sphere by a plane in a given ratio.[5] He also invented a new type of conic sundial.

Cæsar the Mathematician. It is also proper to speak of the contributions of Julius Cæsar (100–44 B.C.) to the reform of the calendar (46 B.C.), a work undertaken with the help of Sosig'enes[6] of Alexandria, an astronomer of whom almost nothing further is known. Cæsar himself was well versed in astronomy and wrote a poem on the subject and a work *De Astris*, neither of which is extant. He also planned extensive surveys of the empire.

About 40 B.C. the Greek astronomer Cleome'des[7] seems to have flourished and to have composed a treatise on the circular

[1] Born *c.* 66 B.C.; died *c.* 24 A.D.

[2] Fl. *c.* 60 B.C.; died in exile 44 B.C.

[3] Διονυσόδωρος. Fl. *c.* 50 B.C.

[4] Αμισός.

[5] It is preserved in Eutocius's commentary (*c.* 560) on II, 5, of the work of Archimedes on the sphere and cylinder. The method employed is that of the intersection of a parabola and a hyperbola.

[6] Σωσιγένης.

[7] Κλεομήδης.

theory of the heavenly bodies.[1] It was said three centuries ago that manuscripts of his treatises on arithmetic and the sphere were still in existence,[2] but they have since been lost.

Vitruvius. Of the Romans who made extensive practical use of mathematics, none is more prominent than Marcus Vitru'vius Pollio, commonly known as Vitruvius. Although the dates are uncertain, it is thought that his great work on architecture[3] was written between 20 and 14 B.C. In Book IX he treats of various types of sundials, and throughout the work he shows his early training as an engineer. He also has something to say on perspective, the ancient science of optics.

Referring to the same general line of applied mathematics, Lucius Junius Moderatus Columel'la (*c.* 25 A.D.) of Gades (Cadiz) wrote on agriculture[4] and included in his treatise a certain amount of information on astronomy, the calendar, and the art of surveying.

Although the name of Gaius Plin'ius Secundus,[5] commonly known as Pliny, is connected chiefly with his *Natural History*, a work in thirty-seven books, it should be recalled that he incorporated a certain amount of mathematics in his treatise. Book II contains a brief account of astronomy and is particularly valuable because of its historical information. Our knowledge of the practical use of the Roman numerals is enriched by his frequent reference to them in this work.

Frontinus. Next to Vitruvius, the most prominent of the Roman writers who made any practical use of mathematics was Sextus Julius Fronti'nus (*c.* 40–106), general, superintendent of water supply, and author of a work on war[6] and of one on

[1] Κυκλικῆς θεωρίας μετεώρων βιβλία δύο, first printed in Latin at Venice, 1498; in Greek, at Paris, 1539.

[2] Baldi, *Cronica*, p. 43. The date of Cleomedes is often given as a century later, but since he mentions no writer later than Poseidonius (died *c.* 44 B.C.), it is probable that he lived in the 1st century B.C.


[3] *De Architectura Libri X* (first printed at Rome *c.* 1486); *The Ten Books on Architecture*, translated by M. H. Morgan, Cambridge, Massachusetts, 1914.

[4] *De Re Rustica.*

[5] Born at Como, 23; died at the destruction of Pompeii, 79. The first name often appears as Caius.

[6] *Strategematicon Libri IV.* There is an edition by Gundermann, Leipzig, 1888.

aqueducts.[1] Some appreciation of the engineering works of this period may be formed from a consideration of the aqueduct of Claudius, which was constructed in the 1st century A.D. There are also preserved certain other books, generally believed to have been written by Frontinus, setting forth the principles of land surveying as commonly practiced by the Romans.[2]



THE AQUEDUCT OF CLAUDIUS

Constructed about the time of Frontinus, or probably just before his period of activity. From Breasted's *Ancient Times*

Hyginus. Among those who made use of mathematics in the work of surveying, Hygi'nus[3] (*c.* 120), known as Gromaticus (the surveyor), is one of the most prominent. The *gromatici* were those who used the *groma*,[4] an instrument employed in measuring and laying out land, and Hyginus was well known as a writer on the subject, although the fragments of his works

[1] *De Aquaeductibus urbis Romae Libri II.* First printed in Rome *c.* 1490; there is a recent edition in English by Herschel, London, 2d ed., 1913.

[2] His writings are collected in the so-called *Codex Arcerianus*. See K. Lachmann and A. Rudorff, *Gromatici Veteres*, being Vol. I of F. Blume, K. Lachmann, and A. Rudorff, *Die Schriften der Römischen Feldmesser*, 2 vols. (Berlin, 1848).

[3] Also spelled Hygenus and Higinus.

[4] Also spelled *gruma*. It is from γνώμων (*gnomon*), the shaft set up for the ancient shadow-reckoning, for sundials, and for general astronomical purposes. See Lachmann and Rudorff, *loc. cit.*, I, 108.

extant show no mathematical contributions to the science. There was an earlier Hyginus,[1] who wrote a work of no merit on astronomy,[2] and who is sometimes confused with his more prominent namesake, the surveyor.

The Roman surveyor Balbus (*c.* 100) was very likely contemporary with Hyginus Gromaticus, but his contributions[3] were unimportant.

Theodosius. There lived about this time, and certainly in the reign of Trajan (98–117), the mathematician and astronomer Theodo'sius.[4] He seems, on the testimony of Suidas, to have been a native of Tripoli, on the Phœnician coast. He wrote several works, the most important being his treatise on the sphere.[5] While this work possessed but little merit, it was translated into the Arabic along with most of the other Greek works on astronomy, and its brevity gave it considerable standing in Arabian schools. He is often confused with a Theodosius of Bithynia, who lived *c.* 50 B.C. and wrote on the sundial.

5. Period of Menelaus

Heron, or He'ro, of Alexandria[6] represented the applications of mathematics more completely than any other writer of about the beginning of our era. He seems to have been an Egyptian,

[1] Gaius Julius Hyginus, a friend of Ovid and therefore living in the 1st century B.C.

[2] *Poeticon astronomicon Libri IIII.*

[3] *Expositio et ratio omnivm formarvm.* See Lachmann and Rudorff, *loc. cit.*, I, 91.

[4] Θεοδόσιος. Fl. *c.* 100.

[5] Σφαιρικὰ ἐν βιβλίοις τρισίν. It was first printed, in Latin, at Paris, in 1529. See A. A. Björnbo, "Wann lebte Theodosios?" *Abhandlungen, zur Geschichte der Mathematik*, hereafter referred to as *Abhandlungen*, Leipzig, *v. d.*, XIV, 64.

[6] Ἥρων. Fl. *c.* 50 A.D. This date is based upon the careful researches of Wilhelm Schmidt, *Heronis Alexandrini Opera quae supersunt omnia*, Leipzig, 1899–1914. He places Heron in the 1st century A.D. It was formerly thought that he lived under the Ptolemies Philadel'phus and Euer'getes (283–222 B.C.), and it was also asserted that he flourished *c.* 100 B.C. See also R. Meier, *De Heronis aetate*, Leipzig, 1905; *Abhandlungen*, VIII, 195; T. H. Martin, "Recherches sur la vie et les ouvrages d'Héron d'Alexandrie" in *Mémoires présentés par divers savants à l'Académie des Inscriptions*, IV (1) (Paris, 1854); F. Hultsch, *Heronis Alexandrini geometricorum et stereometricorum reliquiae*, Berlin, 1864. Heath, *History*, II, 298, states that the evidence at present favors the 3d century A.D., but at best the date is very uncertain.

his style not being that of a Greek. He invented the pneumatic device commonly known as Heron's Fountain, a simple form of the steam engine, and various other machines, showing much ingenuity in all his numerous activities. He wrote on pneumatics, dioptrics, and mechanics, but from the standpoint of mathematics his work on mensuration is the most interesting. In this he treats of land surveying, probably summarizing the methods in use by the Egyptians. As is the case with many of the Greek scholars, some of his works are lost. His formula for the area of a triangle, $A = \sqrt{s(s-a)(s-b)(s-c)}$, is well known. It appears in the geodesy,[1] which is contained in his metrics,[2] but the proof is given (possibly an interpolation) in his dioptrics.[3] In his geometry may be found the first definite use of the trigonometric rule which we express by the formula $c = \frac{n}{4} \cdot \cot \frac{180^\circ}{n}$, where n is the number of sides of a regular polygon of area A and side s, and where $c = A/s^2$. He computed c for $n = 3, 4, \cdots, 12$, but his method is unknown. He was able to solve the equation which we write in the form $ax^2 + bx = c$, so that the general quadratic as we know it today was thus fully mastered by the Greek mathematicians.

About this time there lived Sere'nus of Antinoop'olis.[4] He was the author of a treatise on the *Section of the Cylinder*, containing thirty-three propositions, and of one on the *Section of the Cone*, with sixty-nine propositions. The latter has considerable work on maxima and minima. He also employed the principle of a harmonic pencil of rays.

Menelaus. Of those who, in the period of decay of Greek mathematics, showed any evidence of genius, Menela'us[5] was one of the most prominent. He was a native of Alexandria and wrote

[1] Γεωδαισία. [2] Μετρικά.

[3] Περὶ διόπτρας. On the formula in the Middle Ages, see G. Enestrom, *Bibl. Math.*, V (3), 311.

[4] Antin'oe, Ἀντινόεια, a city on the eastern bank of the Nile. He is often called Serenus of Antissa. See Cantor, *Geschichte*, I, chap. 20. The date of Serenus is quite uncertain. J. L. Heiberg, who edited his *Opuscula* (Leipzig, 1895), is inclined (p. xvii) to place him in the 4th century.

[5] Μενέλαος. Fl. *c.* 100. Heath, *History*, II, 260.

a treatise on the sphere,[1] particularly with respect to the geometric properties of spherical triangles. He is known to have made astronomical observations in Rome in the year 98. Besides his treatise on the sphere he also wrote six books on the calculation of chords. One of his most important theorems states that if the three lines forming a triangle are cut by a transversal, the product of the lengths of three segments which have no common extremity is equal to the products of the other three. This appears as a lemma to a similar proposition relating to spherical triangles, "the chords of three segments doubled" replacing "three segments." The proposition was often known in the Middle Ages as the *regula sex quantitatum* because of the six segments involved. He also knew the invariant property of the anharmonic ratio of the line segments formed by a transversal cutting four concurrent lines,—a property the discovery of which was formerly attributed to Pappus,[2] who flourished about two centuries later.

Nicomachus. The best known of the Greek writers on arithmetic, although not the greatest arithmetician, was Nicom'achus[3] of Gerasa, his birthplace being probably the modern Jerash, a town situated about fifty-six miles northeast of Jerusalem. Since he mentions Thrasyl'lus,[4] who lived under Tiberius (reigned 14–37), but says nothing of the work of Theon of Smyrna, who lived under Hadrian (reigned 117–138), and since his work was translated from Greek into Latin by Appuleius, who lived in the time of Antoninus Pius (reigned 138–161), we are safe in asserting that he lived about the close of the first century.

Nicomachus wrote a treatise on music and a work in two books on arithmetic.[5] The arithmetic as it has come down to us may be only a compendium of a larger work which has

[1] The Latin title, by which it is best known, is *Sphaericorum Libri III.* There are editions by Maurolycus (1558), Mersenne (1644), and later writers.

[2] See the *Abhandlungen*, XIV, 96, 99.

[3] Νικόμαχος Γ*ερασηνός*, or Γ*ερασινός*. Fl. *c.* 100. Heath, *History*, I, 97.

[4] Probably Thrasyllus of Rhodes, died *c.* 36.

[5] This was first printed in 1538, at Paris. The best edition is that of Hoche, Leipzig, 1866.

long since been lost. Some such work seems to have been known to Boethius (*c.* 510) and to have been used by him in compiling his own treatise on the subject.

The Works of Nicomachus. Nicomachus belonged to the Neopythagoreans, a sect of philosophers then flourishing in Alexandria and trying to revive the teachings of Pythagoras. It is therefore quite possible that Nicomachus made the journey from Gerasa to Alexandria to study their doctrines.[1] At any rate there is a considerable amount of the Pythagorean theory of numbers in the tiresome treatment that he accords to arithmetic. The period was one of intellectual decadence, and had he not happened to summarize the ancient teachings in a field that had not been entered by writers of the first rank, we should never have heard of him.[2] His arithmetic[3] was rather an introduction to the philosophy of the subject than a scholarly treatment of the science itself. For lack of anything better it was adopted as a textbook in the few remaining schools of philosophy, and Boethius did much to perpetuate its influence.[4] In the *Philop'atris*,[5] probably a spurious dialogue inserted among the genuine works of Lucia'nus,[6] perhaps as late as the 10th century,[7] it is said of a certain man that "he reckons like Nicomachus of Gerasa."[8] The remark is ludicrous, and very likely was so intended, because there is no evidence that Nicomachus could reckon with any skill whatever, his interest being rather in the theory of numbers, which. as we have seen, was quite distinct from logistic.

[1] On the rise of other intellectual centers, however, as Alexandria began to lose prestige, see Kroll, *Geschichte*, p. 32.

[2] P. Tannery, *Revue philosophique*, XI, 289.

[3] *Introductionis Arithmeticae Libri duo*; Greek, Ἀριθμητικῆς εἰσαγωγῆς βιβλία β. There are various editions in Latin and Greek. For a summary of the work in English see G. Johnson, *The Arithmetical Philosophy of Nicomachus of Gerasa* (Lancaster, Pennsylvania, 1916); Heath, *History*, I, 97, and II, 238.

[4] It was also known in Hebrew, at least in paraphrase, in 1317. See M. Steinschneider, "Die Mathematik bei den Juden," *Bibl. Math.*, XI (2), 79.

[5] Φιλόπατρις.

[6] Λουκιανός. A humorous Greek writer of the 2d century.

[7] For discussion, see M. C. P. Schmidt, *Chrestomathie*, III, 19.

[8] Καὶ γὰρ ἀριθμέει ὡς Νικόμαχος ὁ Γερασηνός.

Nicomachus mentions the sieve of Eratosthenes and often cites the Pythagorean doctrines. He gives an extended treatment of figurate numbers, and in his work appears an early form of the Greek multiplication table. Extensive multiplication tables are found in the Babylonian tablets, but no earlier Greek example is known, unless it be the one on the ancient wax tablet mentioned on page 58. The medieval name, *mensa Pythagorica*, may mean that a certain form of the multiplication table, mentioned in Volume II, came from the Neopythagoreans.

Another work of Nicomachus, the *Theologumena*,[1] has been lost, the extant work by that name being a later compilation.

The'on[2] **of Smyrna**, so called to distinguish him from Theon of Alexandria, who is mentioned later, lived in the time of Hadrian (reigned 117–138). He was interested in arithmetic and astronomy, and was the author of a work[3] which is commonly known in the Latin translation as the *Expositio*. Of this work, which set forth the mathematics necessary for the reading of Plato, two books are extant, one on arithmetic and one on astronomy, and very likely these are all that he wrote. The former resembles the work of Nicomachus but is less systematic.[4]

Marinus of Tyre. Mari'nus[5] of Tyre, a Greek scientist, who lived *c.* 150, may properly be called the founder of ancient mathematical geography. Apparently with greater success than Hipparchus (*c.* 150 B.C.) he definitely located places by reference to two coordinates, namely, latitude and longitude, and his maps set a new standard which the astronomer Ptolemy recognized a little later. The maps themselves, however, have not come down to us. He established the prime meridian

[1] Θεολογούμενα ἀριθμητικά.

[2] Θέων. Fl. *c.* 125. Heath, *History*, II, 238.

[3] Τῶν κατὰ τὸ μαθηματικὸν χρησίμων εἰς τὴν τοῦ Πλάτωνος ἀνάγνωσιν (βιβλία). The best Greek edition is E. Hiller, *Theonis Smyrnaei Philosophi Platonici Expositio* . . . (Leipzig, 1878). There is a French translation by J. Dupuis (Paris, 1892).


[4] On his astronomy see the edition by T. H. Martin, *Theonis Smyrnaei Platonici Liber de Astronomia* (Paris, 1849); J. B. Biot, review in the *Journal des Savants* (April, 1850).

[5] Μαρῖνος.

through the *Fortunatae Insulae*,[1] and this meridian was adopted by Ptolemy. At a later date the meridian was more definitely located through Ferro,[2] one of the Canary Islands, and this position was recognized until modern times.

6. Ptolemy and his Successors

Ptol'emy, or Claudius Ptolemæus,[3] whose period of greatest activity was *c.* 140–160, did for astronomy what Euclid did for plane geometry, Apollonius for conics, and Nicomachus for



PTOLEMY'S MAP OF THE WORLD

This shows the great growth in the knowledge of geography from the time of Eratosthenes. See page 109. From Breasted's *Ancient Times*

arithmetic. He brought together in a single treatise the discoveries of his predecessors, arranging the material systematically, and, like the first two mentioned, was possessed of such genius as to make his work a standard of excellence for many

[1] Αἱ τῶν Μακάρων νῆσοι, Islands of the Blessed, probably including the Canary, Madeira, and Azores groups. It was here, in what Milton calls the "thrice happy isles," that Hesiod and Pindar placed Elysium.

[2] Ancient *Pluvialia*, the Πλουϊτάλα of Ptolemy.

[3] Πτολεμαῖος Κλαύδιος. Born *c.* 85; died *c.* 165. Heath, *History*, II, 273.

centuries. As to his life we know only that he taught in Athens and Alexandria. His greatest work, commonly known as the *Almagest*,[1] contains much information on the history of ancient astronomy. He also wrote on the planisphere, on music, and on applied mathematics. There is a question as to the genuineness of a work on optics that is often attributed to him. In the *Almagest* there is a summary of the computations of Eratosthenes, Poseidonius, and others as to the size of the earth, the position of certain places, and the size of islands and countries. In the application of mathematics to astronomy and geography Ptolemy stands preëminent among Greek scholars. He extended the use of sexagesimal fractions and elaborated the table of chords already used by Hipparchus. He also wrote a treatise on the postulate of parallels and a work of an astrological nature which is generally known in English as the *Tetrabiblos*.[2]

Minor Writers. Among the minor writers who came after Ptolemy there was the jurist Domi'tius Ulpia'nus (*c.* 170–228), a prolific contributor to the law and the compiler of the first table of mortality of which we have any knowledge.

Probably in the same period (*c.* 180) there lived the Roman surveyor Marcus Junius Nip'sus, but his contributions to the science relate chiefly to mensuration and are unimportant.[3] At

[1] The original title is usually given as Μαθηματικῆς συντάξεως βιβλίον πρῶτον; but on this question see J. L. Heiberg, *Ptolemaei Opera*, II, p. cxl (Leipzig, 1898–1907). Since he wrote another σύνταξις, the Arabs seem to have called the greater work *al* μεγάλη, and afterwards *al* μεγίστη (Smith's *Dict. of Greek and Roman Biog.*, III, 570). From μεγίστη, with the Arabic *al* (the), the Arabs made the word which has come to us as Almagest, so that to speak of "the almagest" is like speaking of "the the-greatest." The work was first printed, in an abridged form prepared by Regiomontanus, at Venice in 1496; the first complete edition appeared in Venice in 1515. For the latest work on the subject see C. H. F. Peters and E. B. Knobel, *Ptolemy's Catalogue of the Stars, a Revision of the Almagest*, Washington, 1915.

[2] Τετράβιβλος σύνταξις. The first printed edition appeared at Venice, 1484; first Greek edition, Nürnberg, 1535. It is also known by the Latin name, *Quadripartitum*.

[3] They are given in the *Codex Arcerianus* under the following titles: *fluminis uaratio, limitis repositio, uarationis repositio, lapides etc., podismus.* See Lachmann and Rudorff, *Gromatici Veteres*, I, 285.

about the same time (*c.* 200) there flourished another Roman surveyor named Epaphrodi'tus, who wrote not only on surveying but also on the theory of numbers.[1] He showed that if r is the radius of the circle inscribed in a right-angled triangle of sides a, b, and hypotenuse c, then $2r = a + b - c$. It is proper to refer, chiefly for the sake of showing the low estate to which learning had fallen, to the *Chronicon* of Sextus Julius Africa'nus (*c.* 220), a considerable part of which work is lost, but the extant portion of which contains information of value on the history of the calendar, and also to another work attributed to him, in which some notes appear on the history of other branches of mathematics.

Among the lesser Roman geometers and astronomers there was Censori'nus (*c.* 235), who wrote a book (238) entitled *De die natali*, a work primarily on astrology but containing a limited treatment of chronology, astronomy, and computation. It has been stated that he also wrote a geometry, although the work, if it ever existed, is lost.

We are also told by early writers of the interest taken in mathematics by the wealthy Roman dilettante Quintus Sammonicus Sere'nus (died 212). He was a prolific writer and his works include medicine, mathematics, and other sciences, but in general they merely show the debased state of learning. He is not to be confused with Serenus of Antinoopolis, already mentioned.

A little later (*c.* 275) Spo'rus[2] of Nicæ'a wrote a work from which we derive certain information relating to the history of early mathematics, particularly with reference to duplicating the cube and squaring the circle. He may have been the teacher of Pappus, who is usually put a century later.[3]

[1] V. Mortet, "Un Nouveau Texte des Traités d'Arpentage et de Géométrie d'Epaphroditus et de Vitruvius Rufus," *Notices et Extraits des Manuscrits de la Bibl. Nat.*, XXXV (1896), p. 510.

[2] Probably the same as Porus of Nicæa. The date is very uncertain; it is often given as of the 2d century.

[3] P. Tannery, *Mémoires de Bordeaux*, V (2), 211, and *Mémoires scientifiques*, Paris, 1912, I, 178, thinks he was the teacher of Pappus, or possibly one of his older pupils. The dates are so uncertain as to allow of either possibility.

It is possible that Metrodo'rus,[1] the compiler of the arithmetical epigrams in the Greek Anthology,[2] flourished about 325, but the date *c.* 500 is more probable. These epigrams were puzzle problems, like the one about the pipes filling the cistern, which we should now solve by algebra. For a long time such problems have interested students of arithmetic and algebra, and will doubtless continue to do so for all time to come. Sir Thomas Heath believes that their use dates back at least to the 5th century B.C.

7. Diophantus and his Successors

Diophan'tus[3] of Alexandria was one of the greatest mathematicians of the Greek civilization. That he flourished about the middle of the 3d century seems now fairly certain, although various other dates have from time to time been given. Psellus (11th century) says that Diophantus and Anato'lius[4] wrote on Egyptian computation and that "the very learned Anatolius collected the most essential parts of the doctrine . . ., dedicating his work to Diophantus." Very likely, therefore, Anatolius may have studied under Diophantus. Since he became bishop of Laodicea *c.* 280, he doubtless wrote this work some time before that date, and so Diophantus, who seems to have been the elder, probably flourished *c.* 250–275.[5]

[1] Μητρόδωρος. [2] English translation by W. R. Paton, London, 1918, p. 25.

[3] Διόφαντος. Also written Diophantes, Diophantis, and Diophantos. Fl. *c.* 250–275. There were several writers by this name. Sir T. L. Heath, *Diophantus of Alexandria*, 2d ed. (Cambridge, 1910). On the text see also Tannery's edition of his *Opera Omnia* (Leipzig, 1893, 1895). The first Latin edition of his works was that of Xylander (Wilhelm Holzmann), Basel, 1575; the second, that of Bachet (Paris, 1621), contained the Greek text; the third was that of Bachet with Fermat's notes, Toulouse, 1670. Stevin published a French translation of the first four books in his *Arithmétique*, Leyden, 1585, with editions in 1625 and 1634.

[4] Ἀνατόλιος. *Bibl. Math.*, IV (3), 306. Some fragments of his works are given in J. A. Fabricius, *Bibliotheca Graeca*, III, 275 (Hamburg, 1716). His computus was published by J. P. Migne, *Patrologia Graeca*, Vol. X (Paris, 1857).

[5] Heath, *Diophantus*, 2d. ed., p. 1. Tannery confirms this by an ingenious study of the price of wine at this time, finding that it conforms to that which Diophantus gives. See his *Mémoires scientifiques*, I, 62 (Paris, 1912).

All that is known of his life is given in a curious problem in the Greek Anthology, probably dating from the 5th century. The problem states that his boyhood lasted $\frac{1}{6}$ of his life, his beard grew after $\frac{1}{12}$ more, after $\frac{1}{7}$ more he married, 5 years later his son was born, the son lived to half his father's age, and the father died 4 years after his son. While the statement is obscure at one point, it is generally thought to mean that Diophantus married at 33 and died at 84.

Works of Diophantus. Diophantus wrote three works: (1) *Arithmetica*, originally in thirteen books, of which six are extant[1]; (2) a tract *De polygonis numeris*[2] of which a portion is extant; (3) a number of propositions under the title of *porisms*. Of these, the work of greatest importance is the *Arithmetica*. This work relates, as the title indicates, to the theory of numbers as distinct from computation, and covers much that is now included in algebra. The equations of the first degree are determinate and are so framed as to give positive values for the unknowns. In solving determinate quadratic equations Diophantus used only one root, even where both are positive. He solved a single special case of a cubic equation, but it is thought that further work on such equations may have been given in the lost books. His indeterminate quadratic equations are generally of the types $Ax^2 + C = y^2$ and $Bx + C = y^2$. His simultaneous quadratics relate only to special cases.[3]

Diophantus introduced a better algebraic symbolism than had been known before his time. In general he anticipated by several centuries the progress of algebra, as this progress appears in the works of other writers; and his work, while known to the Arabs, was not really appreciated until its discovery in Europe in the 16th century. He stands out in the history of science as one of the great unexplained geniuses. We do not know what teachers inspired him, we do not know

[1] Heath, p. 16, lists altogether twenty-five MSS., each containing more or less of the works of Diophantus. See also the Tannery edition, I, xxii.

[2] Περὶ πολυγώνων ἀριθμῶν. See the Tannery edition, I, 450; Heath, *Diophantus*, 2d ed., p. 247.

[3] Heath, *Diophantus*, 2d ed., p. 93.

the books he read, and we cannot explain how it happened that he appeared like a giant in a century of pigmies. Perhaps Seneca's statement that "no age is shut against great genius"[1] is the only explanation to be expected.

Lesser Writers. Not far from this time there also flourished the Neoplatonist Porphy'rius,[2] originally known as Malchus[3] the Tyrian and commonly spoken of as Porphyry. He wrote on the life of Pythagoras[4] and a work on the music of Ptolemy. He resided in Athens and Rome, spent some time in Sicily, and is known chiefly for his philosophical works and his antagonism to Christianity. His tomb, or one traditionally designated as his, is still pointed out in Constantinople.

One of the pupils of Anatolius and Porphyrius was Iam'blichus,[5] the author of several works, including one on arithmetic. He wrote a commentary on Nicomachus,[6] and we are indebted to him for considerable information relating to the latter, to Pythagoras,[7] and to other Greek writers. To him is due the theorem that if a number equal to the sum of the three integers $3n$, $3n-1$, $3n-2$ is taken, and if the separate digits of this number are added, and the digits of this result, and so on, the final sum is 6.

About 340 Julius Fir'micus Mater'nus, a Sicilian, wrote a work entitled *Eight Books on Mathematics*[8] but concerned

[1] "Nullum saeculum magnis ingeniis clausum est."

[2] Πορφύριος. Born in Syria, 232 or 233; died *c.* 300.

[3] From *Melekh*, the Hebrew for "king"; in the Greek of that period, Μάλχος. The name was changed, according to tradition, to Porphyrius (wearer of the purple).

[4] Πυθαγόρου βίος, possibly a fragment of his history of the philosophers.

[5] Ἰάμβλιχος. Born at Chalcis, Cœlesyria, *c.* 283; died *c.* 330.

[6] It appeared in various editions in the 16th century. The title-page of the 1668 edition begins: *Jamblichus Chalcidensis ex Coele-Syria in Nicomachi Geraseni Arithmeticam introductionem* (Arnheim, 1668).

[7] Περὶ Πυθαγόρου αἱρέσεως, of which four books are extant, the first containing the life of Pythagoras. The latter was published in Greek and Latin, at Franeker, in 1598. There have been other editions. See *Bibl. Math.*, VIII (3), 309.

[8] *Iulii Firmici Materni Junioris Siculi V. C. Matheseos Libri VIII.* It was first printed at Venice in 1497. The definitive edition is that of Kroll and Skutsch, Leipzig, 1897–1913. L. Thorndike, "A Roman Astrologer as a Historical Source: Julius Firmicus Maternus," *Classical Philology*, VIII, 415.

exclusively with judicial astrology according to the precepts of the Babylonians and Egyptians. Such works have little place in a history of mathematics except as they show from time to time the tendencies of the devotees of the science.

There are also various other isolated cases of mathematical interest in this period of general decay of scholarship, as in the constructing of an astrolabe by Syne'sius of Cyrene (*c.* 378–*c.* 430), the poet and orator, a pupil of Hypa'tia. He became bishop of Ptolemais in 410.

About 390 The'on of Alexandria, known as Theon the Younger, father of the learned Hypatia, edited Euclid's *Elements* and the great work of Ptolemy, wrote various scientific treatises, and set forth a method for finding square roots by the aid of sexagesimal fractions. Manuscripts of his edition of Euclid have been helpful to modern writers in determining the accurate text of the *Elements*.

A little later (*c.* 450) Domni'nus[1] of Larissa, in Syria, wrote on arithmetic, philosophy, and optics. He followed the geometric, deductive method of Euclid rather than the inductive method of Nicomachus, and seems to have had access to some important work that is now lost on the theory of numbers.

Pap'pus[2] of Alexandria, a late Greek geometer, flourished probably in the 3d century, although the date is uncertain. Suidas (*c.* 10th century), not a very careful writer, however, places him in the reign of Theodosius (379–395), but others believe him to have lived two centuries earlier. Of his greatest work, the *Mathematical Collections*,[3] only the last six of the eight books that it originally contained have come down to us. The third book treats of proportion, inscribed solids, and the duplication of the cube; the fourth, of spirals and of such other higher plane curves as the quadratrix; the fifth, of maximum and isoperimetric figures; the sixth, of the sphere; the

[1] Δομνῖνος. P. Tannery, Darboux *Bulletin*, VIII (2), 288.

[2] Πάππος. Fl. *c.* 300. Heath, *History*, II, 355.

[3] Μαθηματικῶν συναγωγῶν βιβλία. The text of this work in Greek and Latin was published with notes by Hultsch, Berlin, 3 vols., in 1876–1878. There was a Latin edition published at Pesaro in 1588, reprinted without change at Venice in 1589 and at Pesaro in 1602. See also *Bibl. Math.*, XII (3), 252.

seventh, of analysis and its history among the Greeks; and the eighth, of mechanics. Two well-known theorems bear his name, one on the generation of a solid by the revolution of a plane figure about an axis, later known as Guldin's Theorem, and the other a generalization of the Pythagorean Theorem. He also knew the doctrine of the involution of points and the constancy of anharmonic ratios in the case of a transversal cutting a pencil, the latter having already been known to Menelaus.

Hypa'tia[1] of Alexandria was the first woman who took any noteworthy position in mathematics, and on this account and because of her martyrdom she has occupied an unduly exalted place in history. She was the daughter and pupil of Theon, and such were her attainments that she was called upon, so tradition says, to preside over the Neoplatonic School at Alexandria. Much that passes for history in her case seems to be fiction, as the statement of Suidas (*c.* 10th century) that she married Isidorus of Gaza, the Neoplatonist. It seems certain, however, that she was slain in one of the city brawls between followers of rival sects. Suidas says that she wrote a commentary on an astronomical table of a certain Diophantus, possibly the algebraist, and one on the conics of Apollonius. Her works, however, are all lost.[2]

Pro'clus, surnamed the Successor[3] because he was looked upon as the successor of Plato in the field of philosophy,[4]

[1] Ὑπατία. Born at Alexandria, *c.* 370; died at Alexandria, 415.

[2] For the romantic side of her life, see J. Toland, *Hypatia, or the history of a most beautiful, most vertuous, most learned . . . lady*, London, 1720; C. Kingsley, *Hypatia*, London, 1853; F. Mauthner, *Hypatia, Roman aus dem Altertum*, 2d ed., Stuttgart, 1892. For a critical study, see R. Hoche, "Hypatia, die Tochter Theons," in *Philologus*, XV (1860), 435; S. Wolf, *Hypatia, die Philosophin von Alexandria*, Vienna, 1870; W. A. Meyer, *Hypatia von Alexandria*, Heidelberg, 1886. See also Heath, *History*, II, 528.

[3] Πρόκλος Διάδοχος. Born at Byzantium, *c.* 412; died 485. A certain Marinus, not to be confused with Marinus of Tyre, gives his birth as February 8, 412. The name also appears as Proculus. The best of the partial editions of his works is that of Cousin, *Procli Opera*, 6 vols., Paris, 1820–1827; 2d ed., 1864. The best edition of his commentary on Euclid I is that of G. Friedlein, *Procli Diadochi in primum Euclidis Elementorum librum*, Leipzig, 1873. His *Institutio Physica*, edited by A. Ritzenstein, was published at Leipzig in 1912.

[4] Or because he succeeded Syrianus, the philosopher, at Alexandria.

studied at Alexandria and taught at Athens. He was a prolific writer and his works include a paraphrase of difficult passages from Ptolemy, a work on astronomy, a commentary on Euclid I, and a brief treatise on astrology. He also shows evidence of a study of certain higher plane curves. His works are valuable sources of information on the history of Greek geometry. For information concerning his life we are indebted to Mari'nus,[1] of Flavia Neapolis in Palestine (the old Sichem), who succeeded him in 485.[2] This Marinus, very likely a Jewish scholar,[3] also wrote an introduction to the *Data* of Euclid.

At about this time Victo'rius[4] of Aquitania (457) wrote a *Canon Paschalis*, one of the first of the *Computi*,—books on the finding of the date for Easter. He suggested beginning our era at the time of the first full moon after the death of Christ. He also wrote a *calculus*, that is, a practical arithmetic. In this he gave considerable attention to fractions and to tables for the multiplication of large numbers.

The name of Capella might, for chronological reasons, be included in this chapter, but on account of the relation of his work to that of writers of the 6th century it is considered in Chapter V.

8. The Orient

China. The period from 300 B.C. to 500 A.D. was one of mathematical activity in China, and some slight but noteworthy trace remains of an interest in numbers in Japan.[5] At the beginning of this period the event of greatest concern in the history of Chinese mathematics was the burning of all books[6] (213 B.C.), as already mentioned in Chapter II, by order of the emperor Shï Huang-ti,[7] founder of the Ch'in (Ts'in)

[1] Μαρῖνος. As stated above, he must not be confused with the astronomer already mentioned.

[2] His life of Proclus was first printed at Zürich in 1559.

[3] S. Krauss, *Jewish Quarterly Review*, 1897, p. 518.

[4] Often written Victorinus. It is thought that he was born in Limoges.

[5] Smith and Mikami, *History of Japanese Mathematics*, chap. i (Chicago, 1914); hereafter referred to as Smith-Mikami.

[6] An exception was made of books on medicine, agriculture, and divination.

[7] She Huang-ti, "the First Emperor," born 259 B.C.; died 210 or 211.

Dynasty (221 B.C.), who wished to appear in the eyes of posterity as the creator of a new era of learning. The penalty for not burning the books was branding and four years' service on the Great Wall. The records say that four hundred and sixty scholars protested against this odious law and were buried alive as an example to others. How many of the ancient classics survived, or how many were faithfully transmitted by means of copies made from memory, we do not know, but it is probable that Chinese scholars will in due time apply the methods of textual criticism to the determination of this point.

About this time, and probably just after the burning of the books, there lived the learned Ch'ang Ts'ang (*c.* 250–152 B.C.), a statesman of highest rank, who wrote (176 B.C.) a new *K'iu-ch'ang Suan-shu* (*Arithmetic in Nine Sections*),[1] basing it upon fragments of the earlier work of the same name. The nine chapters or sections have already been given (page 32).

Ch'ang Ts'ang gave the area of a segment of a circle as $\frac{1}{2}(c + a)a$, where c is the chord and a is the altitude of the segment. Among his problems is that of finding the height of the trunk of a tree, the upper part of which was 10 feet high but has fallen over and reaches the ground 3 feet from the base. The rule for the area of the segment of a circle is later found in the work of the Hindu Mahāvīra (*c.* 850), and the problem about the tree is found in various Hindu mathematical works after the time of Āryabhaṭa (*c.* 510).

Minor Chinese Writers and Events. The period following the burning of the books was, as might have been expected from the need thus created, one of considerable intellectual activity. In this respect, but from a wholly different cause, it was not unlike the century following the impetus given to learning by Plato. Ch'êng Kiang Chen (also known as Chŭn Shuen), who died in 200 B.C., wrote on knotted cords which perhaps, like the Peruvian *quipu*, were for keeping accounts.

[1] K. L. Biernatzki, "Die Arithmetik der Chinesen," Crelle's *Journal*, LII, 11; A. Wylie, *Chinese Researches*, Part III (Shanghai, 1897). These writers put the date *c.* 100 B.C., but Ch'ang Ts'ang appears to have died in 152 B.C., upwards of 100 years old, and to have written the work in 176 B.C. See Mikami, *China*, p. 9.

Then as always in Chinese history the regulation of the calendar occupied the attention of scholars. Thus it is recorded that *c.* 104 B.C. the emperor reëstablished official astronomy and a new calendar was devised.[1] It is also worthy of note, as bearing upon the arithmetic of commerce, that about this time (135 B.C.) coinage became a government prerogative.[2] The Chinese annals of this period also speak of the efforts of the emperor[3] to open up communication with the region about the river Oxus, all such efforts having relation to the unsolved problem of the transmission of mathematical knowledge between the East and the West. The famous Chinese general Ch'ang K'ién went to the countries of the Jaxartes and the Oxus in the 2d century B.C., and about 100 B.C. an envoy was sent as far west as Lake Baikal.[4] This intercourse between the East and the West was maintained for several centuries. For example, an Aramaic manuscript of the 1st century (*c.* 1–20), the earliest known specimen of rag paper, has been found on the Chinese border.[5] That China had intercourse with India is evident from the fact that the records show such relations as early as 218 B.C. and that the name Sin-du appears in the Chinese annals of about 120 B.C. It is also well established that China was known in the West at this period. Ptolemy the astronomer (*c.* 150) speaks of the country under the name of Thin, and in 166 Marcus Aurelius sent an embassy to the emperor's court.

[1] J. B. Biot, *Études sur l'astronomie Indienne et sur l'astronomie Chinoise*, p. 299. Paris, 1862.

[2] H. B. Morse, "Currency in China," from the *Journal of the North-China Branch of the Royal Asiatic Society*, XXXVIII; reprint, p. 2.

[3] Wu-tí (140–87 B.C.). On the general subject of the relations of China with the West see S. W. Williams, *A History of China*, p. 58 (New York, 1897); F. Hirth, "The Story of Chang K'ién," *Journal of the Amer. Oriental Soc.*, XXXVII, 89, 185, 186; T. W. Kingsmill, "The Intercourse of China with Central and Western Asia in the 2d Century B.C.," *Journal of the China Branch of the Royal Asiat. Soc.*, XIV (N.S.), 1; Hirth and Rockhill, *Chau Ju-Kua: His Work on the Chinese and Arab Trade in the twelfth and thirteenth centuries* (Petrograd, 1911), the preface to which considers the whole question from earliest times to the 13th century.

[4] E. Bretschneider, *Mediaeval Researches*, I, 32. London, 1910.

[5] M. A. Stein, *Ruins of the Desert of Cathay*, II, 114. London, 1912.

It is probable that this continued interchange of thought is one of the causes of the frequent changes in the calendar and of the study of the related geometric figure of the circle. About 25 A.D. there lived a well-known philosopher and astronomer named Liu Hsiao, who was of the Imperial house of the Han Dynasty.[1] He was one of the most prominent of the "circle squarers" of his day. His son, Liu Hsing,[2] devised a new calendar,[3] thus using his time to better advantage than the father. A few years later (*c.* 75 A.D.) Pan Ku wrote a work[4] in which the use of the bamboo rods, a primitive form of abacus, is mentioned. At about this time Ch'ang Höng (78–139), chief astrologer and minister under the emperor An-tí, constructed an armillary sphere and wrote on astronomy and geometry. He gave $\sqrt{10}$ as the value of π, this being one of the earliest uses of this approximation.[5] Perhaps contemporary with him, although we are uncertain, there lived Ch'ang ch'un-ch'ing, who wrote a commentary on the *Chóu-peï*. About 190 there flourished Ts'ai Yung,[6] one of the numerous experts on the calendar, but his works are lost. He was sentenced to death for political reasons, but the sentence was commuted to having his hair pulled out. His convivial habits gave him the name of Drunken Dragon.

Wu-ts'ao Suan-king. Possibly about the beginning of the Christian era, for the date is so uncertain[7] that we are not safe in fixing the time even within the limits of several centuries, there was written one of the best-known but least worthy Chinese classics on mathematics, the *Wu-ts'ao Suan-king*.[8] The author seems to have been Sun-tzï,[9] but

[1] This dynasty lasted from 206 B.C. to 25 A.D.

[2] Biot (p. 305) transliterates the name as Lieou-hin.

[3] The *San-t'ung* calendar, devised in the year 66.

[4] The *Han Shu*. Pan Ku died in 92.

[5] On account of the unreliability of early Chinese texts, all such statements are open to some doubt.

[6] Born 133; died 192.

[7] Mikami, *loc. cit.*, p. 37, says in the former (beginning *c.* 206 B.C.) or later (*c.* 25–220 A.D.) Han Dynasty.

[8] *Arithmetic Classic in Five Books.*

[9] Also given as Sun Tsze, Sun Tsü, Suentse, Sun Wu tsze, and Sun Tsu Yeh Ch'i-sun. The work is also known as the *Sun-tzï Suan-king*. Père Vanhée puts the date as probably the 1st century A.D., while Biernatzki (p. 21) says that Sun-tzï may have lived 220 B.C.

even as to this we are uncertain. The work is obscurely written and is not so accurate in its statements as the *Nine Sections*. It relates chiefly to the mensuration of areas. A single problem will serve to show its nature:

"There is a quadrangular field of which the eastern side is 35 paces, the western side 45 paces, the southern side 25 paces, and the northern side 15 paces. Required the area of the field."

Evidently a solution is impossible through lack of sufficient data; but the author assumes that he may take one fourth the product of the sums of the pairs of opposite sides,[1] such approximations as this being not uncommon all through the East in these early times.

Liu Hui. The best-known Chinese mathematician of the 3d century was Liu Hui.[2] In 263 he wrote the *Sea Island Arithmetic Classic*,[3] a work which probably took its name from the first problem that it contains, this problem beginning with the statement, "There is a sea island that is to be measured." The work is concerned with the mensuration of heights and distances, the rules seeming to show some familiarity with the manipulation of algebraic formulas.

Liu Hui also wrote a commentary on the *Nine Sections*, and it seems to have been in the performing of this task that he accumulated the materials for his "Sea Island" work.

Minor Chinese Writers from 200 to 500. Of the minor writers of the 3d century mention may properly be made of Wang Pi (*c.* 225–249), the leading authority on the mysticism of the *I-king*[4]; of Wang Fan (229–267), the astronomer, who asserted that $\pi = \frac{142}{45}$; of Siu Yo (*c.* 250), who wrote the *Omissions noted in the Art of Numbers*[5]; of Li Ping, the great

[1] Mikami, *China*, p. 38.

[2] Also transliterated Lew Hui, Lew Hwuy, and Lieou Hoei.

[3] *Hai-tau Suan-king*. Wylie says that this title first appeared in an edition prepared in the 8th century.

[4] See page 25.

[5] *Shu-shu-ki-yi*, or *Chou-chou-ki-yi*. There are many commentaries on this work. See A. Vissière, *Recherches sur l'origine de l'abaque Chinois . . .*, p. 22 (Paris, 1892).

irrigation engineer of the 3d century;[1] of L... ...h[2] (*c.* 289), who is possibly the one who gave the so-called "Chih's value of π," that is, $\pi = 3\frac{1}{8}$; and Hsü Yüeh, who wrote a commentary on Siu Yo's work above mentioned.

The 5th century is more interesting because of the evidence that we have of intercourse between China and the rest of the world than because of any definite contributions to mathematics. A few names of mathematicians are known,[3] but it was the visit of the Buddhist missionaries and pilgrims from India that is significant. The result of this visit was the translation of an arithmetic and of various astronomical works of the Brahmans, which stimulated the activity of Chinese scholars in these fields. This interchange of thought was not new, for Buddhism was transmitted from India to China at least as early as the year 65. In 399 a Chinese Buddhist, Fa-hién, went to India, and after his return in 414 he devoted his life to the translation of Hindu works. Since religion was closely related to astronomy, and astronomy to mathematics, the influence of this interchange of religious thought must have been stimulating to the science of China. Moreover, after about the year 450 there are many references in the Chinese annals to the people of Po-ssï (Persia), and thereafter many embassies passed between the two countries.

Among the mathematicians of this period whose names have come down to us is P'i Yen-tsung (*c.* 400–*c.* 450), who is said to have computed a noteworthy value of π which has since been lost. There is also Tsu Ch'ung-chih (430–501), an expert in mechanics, who revived the knowledge of the "south-pointing vehicle" and constructed a motor boat, all details of which are lost. He gave $\frac{22}{7}$ as an "inaccurate value" of π, and $\frac{355}{113}$ as the "accurate value," and he also showed that π lies between our present decimal forms 3.1415926 and 3.1415927. About the year 450 a new calendar was devised by Ho' Ch'êng-t'ien,

[1] H. K. Richardson, *Asia*, XIX, 441.

[2] There was another mathematician of the same name (1311–1375), who devised a new official calendar.

[3] For example, Tun Ch'üan (*c.* 425), who wrote the *San-töng-shu*, and Wang Jong, an arithmetician.

and at about the same time one Wu, a geometer, gave the equivalent of 3.1432 + as the value of π. These details have little significance except as they show the nature of the scientific interests of China during this long period.

Japan in Earliest Times. Prior to the year 500 Japan seems to have made no progress either in literature or in science. There is a tradition that Chinese ideograms made their way through Korea and into Japan in the year 284. There is also reference to the *Jindai monji*, or "letters of the era of the gods," in early times, possibly a kind of system of cabala with numerical values assigned to the letters, but nothing is definitely known upon the subject. A tradition also exists that in 660 B.C. the Japanese had a system of numeration extending to very high powers of ten. In this system the special name *yorozu* was used for 10,000, corresponding to the Greek myriad already mentioned, and this may possibly be some slight evidence of the early interrelations between the East and the West.[1]

Of the rest of Japanese mathematics in the early periods we know only that there was a system of measures and that, as among all other ancient peoples of any intellectual standing, a calendar existed.

India. The noteworthy contribution of India in this period was probably the Hindu numeral system, which will be discussed later.[2] A second event of importance in the history of mathematics in India, and one which chronologically precedes the writing of the numerals, was the invasion of this country by the army of Alexander the Great (327 B.C.) and the sending of Greek ambassadors to reside in Indian courts. How much influence this event had upon the science and particularly upon the astronomy of the Hindus it is difficult at present to say. It is worthy of note, however, that the later Hindu writers used such Greek adaptations as *jāmitra* (from the Greek διάμετρος), *kendra* (κέντρον), and *dramma* (δραχμή).[3]

[1] For discussion and bibliography see Smith-Mikami, p. 4.

[2] See Volume II, Chapter II.

[3] G. R. Kaye, *Indian Mathematics*, p. 26 (Calcutta, 1915) (hereafter referred to as Kaye, *Indian Math.*); H. T. Colebrooke, *Algebra with Arithmetic and*

Just before the beginning of the Christian era there were numerous invasions from the north that interfered seriously with the spread of Greek science, and in the 4th century A.D. there appeared at least one work which definitely sought to replace the astronomy of Greece by the ancient science of India.

The first important work on astronomy produced in India, so far as now known, was the *Sūrya Siddhānta*,[1] probably written about the beginning of the 5th century, although known to us only in later manuscripts. The ritualistic mathematical formulas of the *Śulvasūtras* now gave place to the mathematics of the stars. This change was possibly due to the influence of Greek scholars whose works might still have been appreciated by the descendants of the ancient Greeks who settled in India after Alexander's time. Varāhamihira, who will be mentioned later, speaks of five *Siddhāntas*, but places the *Sūrya Siddhānta* at the head. Among the five is the *Paulisa Siddhānta*, probably of about the same period. This contains an excellent summary of early Hindu trigonometry, the rules, expressed in modern symbolism, being as follows:

$$\sin 30^\circ = \tfrac{1}{2}, \qquad \pi = \sqrt{10},$$

$$\sin 60^\circ = \sqrt{1 - \tfrac{1}{4}}, \quad \sin^2\phi = \left(\frac{\sin 2\phi}{2}\right)^2 + \left(\frac{1 - \sin(90^\circ - 2\phi)}{2}\right)^2.$$

There is also included in this work a table of sines which was apparently derived from Ptolemy's table of chords.

The absence of an authentic Hindu chronology and of a careful study of the effect of the Greek civilization upon the sciences in India renders difficult a satisfactory assessment of her mathematical achievements in this period.

Mensuration, from the Sanscrît, p. lxxx (London, 1817) (hereafter referred to as Colebrooke, *Āryabhaṭa*, or *Brahmagupta*, or *Bhāskara*, according to the part of the work considered, and with the modern spellings as here).

[1] E. Burgess, "The Sūrya Siddhānta," in the *Journ. of the Am. Oriental Soc.*, VI (New Haven, 1860); G. R. Kaye, "Ancient Hindu Spherical Astronomy," *Journ. and Proc. of the Asiatic Soc. of Bengal*, Vol. XV; Bapu Deva Sastri and L. Wilkinson, *The Sūrya Siddhānta and the Siddhānta Śiromani* (Calcutta, 1861). Albêrûnî, the Arab writer on India (*c.* 1000), speaks of the work as "the Siddhānta of the sun, composed by Lāṭa."

Decay of Civilization in Mesopotamia. For about two thousand five hundred years before the period now under consideration Mesopotamia had maintained a high civilization. Assyria, Sumeria, Babylonia, and Chaldea had contributed in a large way to the world's commercial machinery, to its science, to its laws, and to its art. Mathematics, medicine, religion, sculpture, architecture, literature, and the science of government are all indebted to the genius of those who dwelt in the lands bordering upon or in the vicinity of the Two Rivers.

With the close of the 6th century B.C., however, there came a change that was disastrous to the native civilization of this region. The Persian conquest of 539 B.C. and the subsequent coming of the Parthians, the Greeks, and the Romans, each of whom held in subjection some or all of the territory of Mesopotamia, left little of her ancient glory. Trajan, hoping to repeat the conquests of Alexander, visited Babylon early in the 2d century A.D., and "saw nothing worthy of such fame, but only heaps of rubbish, stones, and ruins," and this was symbolic of the decay of a civilization which had perhaps exerted a greater influence upon the world than any that had existed prior to the rise of Greece.

Astrology continued to retain its power over the mass of people, as it does in a large part of Asia today. This is shown by tablets of the 2d century B.C., in which reports are made to the king with respect to predictions as to the positions of the planets. If superstition affected the court, much more would it have affected the people at large.

In all the records of this region only a single name stands out that is worthy of mention in the history of the mathematics of this period, and this only in connection with a sister science. About 250 B.C. Berosus (probably Bar Oseas, that is, the son of Oseas), a Chaldean, founded a school on the island of Cos, and introduced into Greece the astronomy and the astrological beliefs of his people, constructing a sundial and probably other instruments.[1]

[1] A. Wittstein, "Bemerkung zu einer Stelle im Almagest," *Zeitschrift für Math.*, XXXII (Hl. Abt.) (Leipzig, 1887), 201.

TOPICS FOR DISCUSSION

1. The School of Alexandria, its rise, its influence, the great scholars connected with it, and its decay.

2. Euclid, his life, his works, and his influence.

3. The work of Eratosthenes, particularly with respect to geodesy.

4. The life, the works, and the influence of Archimedes.

5. Apollonius and his contribution to the study of conics.

6. The mathematical contributions of the Greek astronomers.

7. Mathematics in the Roman civilization. Causes of the disregard for the science.

8. The life and works of Heron. His influence upon the development of applied mathematics as compared with that of Archimedes.

9. The work of Nicomachus compared with the works of Euclid and Apollonius.

10. The work of Claudius Ptolemæus, or Ptolemy.

11. The life and works of Diophantus.

12. The decay of Greek geometry, with a special consideration of the work of Menelaus, Hypatia, Proclus, and Pappus.

13. Causes and probable effects of the burning of the books in China in 213 B.C.

14. The period in which the *Nine Sections* was written and the general nature of this work.

15. The knotted cords of China and the general subject of knotted cords in the keeping of records and in religious ceremonial.

16. Efforts at opening communications between the East and West at this period, and the probable effect of these efforts on science in general and mathematics in particular.

17. The periods and nature of the *Arithmetic Classic in Five Books* and the *Sea Island Classic*.

18. Influx of Hindu learning into China in this period and the probable effect of this intercourse on the mathematics of both China and India.

19. The invasion of India by Alexander the Great and its effect upon the mathematics of the East.

20. The nature of the *Sūrya Siddhānta* and the bearing of this work upon the mathematics of India.

21. Causes of the decay of mathematics in Mesopotamia in the five centuries after the time of Alexander.

CHAPTER V

THE PERIOD FROM 500 TO 1000

1. China

Intercourse with India and the West. The five centuries extending from 500 to 1000 saw the general trend of mathematics to the West rather than in the opposite direction. Europe was intellectually dormant, drugged with a new narcotic, while most of the East was, as always, superstitious but inquisitive. On this account it is proper to consider first the work of this period as it appears in the Orient. Even in the Dark Ages, however, the West influenced the East, passing traces of the later Greek culture on to the intellectual centers of China and probably to those of India.

In so far as this intercourse was commercial it influenced the art of calculation, while the travel of pilgrims and the movements of armies resulted in the exchange of a knowledge of both astronomical and abstract mathematics. Moreover, the priest, whose leisure allowed time for the study of mathematics, was often an astronomer, and he or the professional astrologer was looked upon as a natural attendant at court or a necessary adjunct to the general's staff. Where the army went, there went a knowledge of mathematics. Astrologers of one country thus consulted with those of another. The itinerant tradesman, the pilgrim, and the army were the means of the exchange of ideas in all ancient times, just as books and periodicals are the corresponding media in our day.

Evidences of this Intercourse. Of the many evidences of intercourse that we have in this period, a few may be mentioned simply as typical. In 518 Hui-sing, a Buddhist pilgrim, visited India; sometime in the 7th century a Sanskrit

calendar[1] was translated into Chinese; in 615 an Arab[2] embassy visited China; in 618 a Hindu astronomer[3] was employed by the Chinese Bureau of Astronomy to devise a new calendar; in 629 Hüan-tsang[4] went to India and after his return in 645 he devoted his life to the translation of Hindu works, of which he had brought no less than 657 from India; in 636, so the Chinese records assert, a Roman priest whom these records speak of as A-lo-pen came to the capital of China; and at the end of the 7th century Buddhist pilgrims sailed from Canton to Java and Sumatra. In the 8th century Arab ambassadors visited China several times, in particular in 713, 726, 756, and later; in 719 an ambassador[5] was sent from Rome to the Chinese court; between 713 and 825, foreign ships of large tonnage visited Canton, and an important customhouse is known to have existed there at that time; and about 775 the geographer Kia Tan (730–805) wrote[6] the itinerary of a voyage by sea from Canton to Persia. About 800, when Bagdad was rapidly becoming the center of the mathematical world, the Chinese received an embassy from A-lun (Harun al-Rashid). In records of the Tang Dynasty (618–907) there are numerous references to the Arabs (Ta-shi), and until the 12th century the intercourse between the Chinese and these people is frequently mentioned. Mas'ûdî (died at Cairo, 956), the famous Arab geographer and historian, visited India, Ceylon, and China in 915, and his *Meadows of Gold*, in which he mentions these countries, is well known. With such evidences as these we have a simple answer to the question as to whether it is probable that China knew of the status of Western mathematics before her own period of remarkable activity, and whether, on the other hand, the West could have known anything of Oriental progress. The answer is that it would have been very strange if each had not been the case.

[1] The *Chiu-chi-li*, as it was called in Chinese. The translator was Chü-t'an Hsi-ta.

[2] That is, if the name *Ta-shi* is taken, as usual, to mean "Arab."

[3] In Chinese, Chü-t'an Chüan.

[4] Original name was Ch'ön I. See Giles, *Biog. Dict.*, No. 801.

[5] Called in Chinese by the name T'u-huo-lo.

[6] In the *T'ang shu*.

The Sixth Century. The 6th century is an important one in the history of Chinese mathematics, owing to the appearance of several works of considerable merit. The earliest of the prominent writers was probably the learned Buddhist Ch'ön Luan,[1] who seems to have been living in 535, but who devised a calendar in the second half of the century.[2] He wrote the *Arithmetic in the five classics*,[3] in which he included various problems of the standard type that had appeared in earlier works. He also wrote commentaries on several of the earlier treatises.[4]

Probably about the same time as Ch'ön Luan there lived Ch'ang K'iu-kien[5] (*c.* 575), whose arithmetic[6] in three books is nearly all extant. The work is devoted chiefly to fractions, and it seems quite clear that the author knew the modern rule of division by multiplying by the reciprocal of the divisor. It also treats of arithmetic progression, the Rule of Three, mensuration, and indeterminate linear equations.

Another contemporary of Ch'ön Luan was the arithmetician Hsia-hou Yang[7] (*c.* 550), the author of a treatise that is still extant.[8] This work includes, as was the custom in most cases of the kind, some problems in mensuration as well as a treatment of certain processes of arithmetic. The arithmetic problems all relate to multiplication, division, and percentage.

In this century there also flourished a geometer by the name of Men (*c.* 575), of whom little is known, but who is said to have given 3.14 as the value of π.

Seventh to the Tenth Century. The most prominent Chinese mathematician of the 7th century was Wang Hs'iao-t'ung,[9]

[1] Given by Père Vanhée as Tsen Loan and by Biernatzki (p. 12), who puts him early in the 7th century, as Tschin Lwan. On all these names Mikami's work has been freely used.

[2] In the reign of Wu-tí, of the Chou monarchy (557–581), in the Chin Dynasty.

[3] *Wu-king Suan-shu.*

[4] For example, on the *Chóu-peï* and the *K'iu-ch'ang Suan-shu.*

[5] Biernatzki (p. 12) transliterates the name as Tschang Kiu Kihn and gives the date as early in the 7th century.

[6] *Ch'ang K'iu-kien Suan-king.*

[7] Biernatzki gives the name as Hea Hau yang. The date is uncertain, but he probably lived in the period from *c.* 550 to *c.* 600.

[8] The *Hsia-hou Yang Suan-king* (*Arithmetic Classic of Hsia-hou Yang*).

[9] Also written Wang Hiao-t'ong and Wang Heau tung.

known to have been living in 623 and in 626. He was an expert on the calendar and was one of the first of the Chinese to write on cubic equations. His work,[1] most of which is extant, contains twenty problems on mensuration, and in some of these problems the cubic equation enters. No method of solving such equations, however, is given.

The 8th century saw no work of importance in mathematics. In 727 I-hsing devised a new calendar,[2] and two centuries later (*c.* 925) there appeared an astrological treatise of some merit,[3] but neither contained any mathematics beyond such as was needed in the work on the calendar. The Dark Ages of the West had spread over the East as well.



SHŌTOKU TAISHI, *c.* 600

From a bronze of the 18th century, showing the prince with a *soroban*, a chronological impossibility

2. JAPAN

Beginnings of Japanese Mathematics.[4] Although Chinese influence had begun to show itself in the intellectual development of Japan before 500, it was not until the Buddhist missionaries began to appear, in 522,[5] that any very pronounced results were noticed. Indeed, it was not until 552 that Buddhism was really introduced, and not until two years later that two scholars, learned in matters pertaining to the calendar,[6] crossed over from Korea and brought to Japan the Chinese system of chronology. Not far from the year 600 a Korean priest, Kanroku, presented to the empress a set of books on astrology and the calendar, and Prince Shōtoku Taishi showed so much interest in calculation that tradition thereafter made him the father of Japanese arithmetic.

[1] *Ch'i-ku Suan-king.*
[2] The *T'ai-yen* calendar.
[3] The *K'ai-yüan Chan-king.*
[4] See Smith-Mikami.
[5] The first to come was Szŭ-ma Ta, known in Japanese as Shiba Tatsu.
[6] These were Wang Pao-san and Wang Pao-liang. See Smith-Mikami, p. 8.

Chinese Influence in Japan. From now on for many generations Japan came completely under Chinese influence in all her intellectual life. The Chinese system of measures was adopted, a school of arithmetic was founded (*c.* 670), an observatory was established at about the same time, and in 701 a university system was inaugurated. Nine Chinese works were specified for students of mathematics,[1] and these seem to have been the classics which influenced the Japanese study of mathematics for several centuries.



TENJIN, PATRON OF MATHEMATICS, *c.* 890

From a bronze. The portrait is found also in early paintings of the Japanese

Aside from Shōtoku Taishi the man whose name stands out most prominently in the history of Japanese mathematics in this period is Tenjin,[2] counselor and teacher at the imperial court (*c.* 890) and a great patron of science and letters.

Altogether the era was one of preparation, contributing nothing new to what China had already developed. Indeed it was not until the 17th century that Japan really awoke to her possibilities in the field of mathematics.

3. India

General Nature of the Work. In the period from 500 to 1000 there were four or five mathematicians of prominence in India. These were the two Āryabhaṭas,[3] Varāhamihira the astronomer, Brahmagupta, and Mahāvīrācārya. In the works of all these writers there is such a mixture of the brilliant and the

[1] These were (1) *Chóu-peï Suan-king*, (2) *Sun-tzï Suan-king*, (3) *Liu-chang*, (4) *San-k'ai Chung-ch'a*, (5) *Wu-ts'ao Suan-shu*, (6) *Hai-tau Suan-shu*, (7) *Kiu-szu*, (8) *Kiu-ch'ang*, (9) *Kiu-shu*, of which the third, fourth, and seventh are lost.

[2] His name was Michizane, but after his death he was canonized as Tenjin, "Heaven man."

[3] For rules for pronouncing Hindu names, see page xxi.

commonplace as to make a judgment of their qualities depend largely upon the personal sympathies of the student. Albêrûnî (*c.* 1000), the Arab historian, speaks of this peculiarity of their writings in these words:

I can only compare their mathematical and astronomical literature . . . to a mixture of pearl shells and sour dates, or of pearls and dung, or of costly crystals and common pebbles. Both kinds of things are equal in their eyes, since they cannot raise themselves to the methods of a strictly scientific deduction.[1]

Āryabhaṭa.[2] The first of the great writers whose name has come down to us is the elder Āryabhaṭa,[3] born at Kusumapura (Kousambhipura), the City of Flowers,[4] a small town on the Jumma just above its confluence with the Ganges.[5] The place is not far from the present Patna (Patnā), called by the Mohammedans Azimabad, by the ancient Buddhists Pāṭaliputra (Pātoliputra), and by Megasthenes, the Syrian ambassador,

[1] Albêrûnî's *India*, translated by E. C. Sachau, 2 vols., I, 25 (London, 1910); hereafter referred to as Albêrûnî's *India*. On the relation of Greek and Hindu arithmetic see H. G. Zeuthen, *Bibl. Math.*, V (3), 97. On the relation of India to the West in general, see H. G. Rawlinson, *Intercourse between India and the Western World from the Earliest Times to the Fall of Rome* (Cambridge, 1916). For extreme Hindu claims, see Benoy Kumar Sarkar, *Hindu Achievements in Exact Science* (New York, 1918).


[2] Born 475 or 476; died *c.* 550.

[3] Sir M. Monier-Williams, *Indian Wisdom*, 4th ed., p. 175 (London, 1893) (hereafter referred to as Monier-Williams, *Indian Wisdom*); J. Garrett, *Classical Dictionary of India*, p. 767 (Madras, 1871); C. M. Whish, "On the Alphabetical Notation of the Hindus," *Trans. of the Literary Society of Madras* (London, 1827); L. Rodet, "Leçons de Calcul d'Âryabhaṭa," *Journal Asiatique*, XIII (7), 393; L. Rodet, "Sur la véritable signification de la notation numérique inventée par Âryabhaṭa," *ibid.*, XVI, p. 440. Certain fragments of his works were published by H. Kern in the *Journal of the Royal Asiatic Society*, XX (1863), 371. See also G. R. Kaye, *Indian Math.*, p. 11, and "Āryabhata," *Journ. and Proc. of the Asiatic Soc. of Bengal*, IV (N.S.), p. 111 (hereafter referred to as Kaye, *Āryabhaṭa*); an article on "Ancient Hindu Spherical Astronomy," *ibid.*, XV; and an article in *Scientia*, XXV, 1, all claiming Greek origin for most of the Hindu work.

[4] The term is also applied to Pāṭaliputra. If we may trust to the rather obscure statements of Albêrûnî, it was the younger Āryabhaṭa, however, who was born at Kusumapura. See the mention of him later.

[5] Sir E. Clive Bayley, *Journal of the Royal Asiatic Society*, XV (N.S.), 21.

Palibothra.[1] Because of this geographic proximity Āryabhaṭa is often said to have been born at Pāṭaliputra. A tradition says that the city was originally called Pāṭaliputraka, being founded by Putraka, the knight of the magic cup and staff and slippers, who married the princess Pāṭali.[2] The tradition further asserts that Buddha, toward the close of his life, crossed the Ganges at this point and prophesied the future greatness of the city.[3] By the beginning of the 5th century, and nearly a century before the birth of Āryabhaṭa, it had lost some of its ancient prestige, for Fa-hién, the Chinese traveler already mentioned, describes (*c.* 400) the ruins of the royal palace which Aśoka commissioned the genii to build, although he speaks of the remarkable hospitals and other institutions still to be found there.[4] Āryabhaṭa evidently wrote there or at



MATHEMATICAL-HISTORICAL MAP OF INDIA

At Delhi, Jaipur, and Benares are interesting relics of native observatories; Patna is approximately the birthplace of Āryabhaṭa (*c.* 475); Ujjain was the leading mathematical center of ancient India and is known particularly for Varāhamihira (*c.* 505), Brahmagupta (*c.* 628), and Bhāskara (*c.* 1150); about 75 miles from Poona are the Nānā Ghāt inscriptions with early numerals; it was at Mysore that Mahāvīra (*c.* 850) lived

[1] E. Reclus, *Asia*, American ed., III, 222. [2] J. Garrett, *loc. cit.*, p. 770.

[3] R. W. Fraser, *A Literary History of India*, p. 143 (N. Y., 1898); E. W. Hopkins, *Religions of India*, pp. 5, 311 (Boston, 1898).

[4] Dutt, *Hist. of Civ. in Anc. India*, II, 58 (London, 1893). On the sojourn of Megasthenes, 306–298 B.C., see Fraser, *loc. cit.*, p. 175. On its importance at about the time of Āryabhaṭa, see the inscriptions of Chandragupta II in

Kusumapura, for he says in one of his works: "Having paid homage to Brahma, to Earth, to the Moon, to Mercury, to Venus, to the Sun, to Mars, to Jupiter, to Saturn, and to the constellations, Āryabhaṭa, in the City of Flowers, sets forth the science venerable."[1]

It was probably because Āryabhaṭa lived so far from Ujjain, the ancient center of mathematics and astronomy, that his works were so little known among Hindu scholars of the centuries immediately following.

Āryabhata's Work. His work, often called the *Āryabhaṭīya*[2] or *Āryabhaṭīyam*, consists of the *Gītikā* or *Daśagītikā*, a collection of astronomical tables, and the *Āryāṣṭasata*, which includes the *Ganita*, a note on arithmetic[3]; the *Kālakriyā*, on time and its measure; and the *Gola*, on the sphere.

The arithmetic carries numeration by tens as far as 10^{18}, treats of plane and solid numbers, and gives a rule for square root. It contains a rule for summing an arithmetic series after the pth term, which may be expressed in modern symbols thus:

$$s = n\left[a + \left(\frac{n-1}{2} + p\right)d\right].$$

It also has a rule which we express by the formula

$$n = \frac{1}{2}\left(1 + \frac{-2a \pm \sqrt{(d-2a)^2 + 8sd}}{d}\right).$$

The rest of the work shows a knowledge of the quadratic equation and of the indeterminate linear equation.

J. F. Fleet, *Corpus Inscriptionum Indicarum*, III, pl. iv, A, B (London, 1888). A century and a half later (629–645) the Chinese pilgrim Hüan-tsang remarked, "Although it has long been deserted, its foundation walls still survive." See Fraser, *loc. cit.*, p. 248.

[1] Rodet, *loc. cit.*, p. 396. For a slightly different translation see Kaye, *Āryabhata*, p. 116.

[2] Monier-Williams, *Indian Wisdom*, p. 175; Mrs. Manning, *Ancient and Mediaeval India*, 2 vols. (London, 1869), largely from the *Journal of the Royal Asiatic Soc.*, I (N.S.), 392, and XX, 371.

[3] Rodet, "Leçons," *loc. cit.*, p. 395. He translates the second part, p. 396 See also Kaye, *Āryabhaṭa*, p. 111; *Bibl. Math.*, XIII (3), 203.

Among the rules relating to areas is one for the isosceles triangle, and this will serve to show the imperfect form of statement used by Āryabhaṭa: "The area produced by a trilateral is the product of the perpendicular that bisects the base, and half the base." The formula for the volume of a sphere is very inaccurate, being $\pi r^2 \sqrt{\pi r^2}$, which would make π equal to $\frac{16}{9}$, possibly an error for the $(\frac{16}{9})^2$ of Ahmes.

The rule for finding the value of π is given as follows: "Add four to one hundred, multiply by eight, and add again sixty-two thousand; the result is the approximate value of the circumference when the diameter is twenty thousand." This makes π equal to $\frac{62832}{20000}$, or 3.1416.[1] Āryabhaṭa also gives a rule for finding sines, and the *Gītikā* has a brief table of these functions.

His work is also noteworthy as containing one of the earliest attempts at a general solution of a linear indeterminate equation by the method of continued fractions.[2]

As stated above, the Āryabhaṭa here mentioned is known as the elder of the two mathematicians of the same name. This fact appears in the work of Albêrûnî[3] and has been the subject of comment by recent writers.[4] The date of the younger Āryabhaṭa is unknown, nor is it possible as yet to differentiate clearly between the works of the two. He seems, from the meager authorities now known, to have been born at Kusumapura.

Varāhamihira (*c.* 505). Among the astronomers of India[5] two appear with the name of Varāhamihira, one living *c.* 200 and the other *c.* 505.[6] The latter of these scholars is the most celebrated of all the writers on astronomy in early India. He wrote several works, of which the *Pañca Siddhāntikā*, treating of astrology and astronomy, is the best known. It includes the computation necessary for finding the position of

[1] Kaye, in his *Āryabhaṭa*, questions whether this is from the works of the elder Āryabhaṭa, the one of whom we are speaking. He thinks it is due to the younger one mentioned below.

[2] Kaye, *Indian Math.*, p. 12.

[3] *India*, II, 305, 327.

[4] See summary by Kaye, *Āryabhaṭa*, p. 113.

[5] A list with dates is given in Colebrooke, *loc. cit.*, p. xxxiii.

[6] The date is quite uncertain. Varāhamihira is said by some Oriental authorities to have died *c.* 587.

a planet, shows an advanced state of mathematical astronomy, but is chiefly valuable in the history of mathematics because of the description that it gives of the five *Siddhāntas* which had been written just before this time.[1] He urged his people to appreciate the work of the Greeks, saying: "The Greeks, though impure, must be honored, since they were trained in the sciences and therein excelled others. What then, are we to say of a Brahman if he combines with his purity the height of science?"[2]

Varāhamihira taught the sphericity of the earth, and in this respect he was followed by most of the other Hindu astronomers of the Middle Ages.[3] Two of his works were translated into Arabic by Albêrûnî (*c.* 1000).[4]

Brahmagupta. The most prominent of the Hindu mathematicians of the 7th century was Brahmagupta,[5] whose period of activity has been fixed as *c.* 628, both from astronomical data and from the testimony of various Hindu writers.[6] He lived and worked in the great astronomical center of Hindu science, Ujjain or Ujjayinī, a town in the state of Gwalior, Central India, said to have been the viceregal seat of Aśoka during his father's reign at Patna. Varāhamihira also carried on his work at the observatory in Ujjain.

When he was only thirty years old, Brahmagupta wrote an astronomical work in twenty-one chapters entitled *Brahmasiddhānta*,[7] which includes as special chapters the *Gaṇitād'hāya*[8]

[1] G. Thibaut and Sudharkar Dvivedi, *The Pañcha-siddhāntikā of Varaha Mihira*. Benares, 1889.

[2] Albêrûnî's *India*, I, 23.

[3] Albêrûnî, *loc. cit.*, I, 266.

[4] For a list of his works, see Albêrûnî, *loc. cit.*, I, xxxix. For the influence of the Greeks upon his work and upon Hindu astronomy in general, see Colebrooke, *loc. cit.*, p. lxxx.

[5] Colebrook, *loc. cit.* Albêrûnî (*c.* 1000) speaks of him as "the son of Jishṇu, from the town of Bhillamāla." Sūryadāsa, a commentator on Bhāskara, also speaks of him as the son of Jishṇu.

[6] Colebrooke (*loc. cit.*, p. xxxv) makes the date 581 or 582, from Brahmagupta's reference to the position of the star *Chitrá* (*Spica Virginis*). The Hindu astronomers make it *c.* 628. He seems to have been born *c.* 598.

[7] Also called the *Brāhma-sphuṭa-sidd'hānta*, "Brahma correct system," possibly a revision. Albêrûnî (*c.* 1000) gives twenty-four chapters, with the title of each. See his *India*, I, 154; II, 303.

[8] *Lectures on Arithmetic.*

and the *Kuṭakhādyaka*.[1] The former begins by a definition of a *gaṇaca*, that is, a calculator who is competent to study astronomy: "He who distinctly and severally knows addition and the rest of the twenty logistics and the eight determinations, including measurement by shadow, is a *gaṇaca*."[2]

Nature of Brahmagupta's Arithmetic. The arithmetic includes work with integers and fractions, progressions, barter, Rule of Three, simple interest, the mensuration of plane figures, and problems on volumes and on shadow reckoning (a primitive plane trigonometry applied by him to the sundial). The mensuration is often faulty, as where Brahmagupta states a rule which would give the area of an equilateral triangle of side 12 as 6×12, or 72; that of the isosceles triangle 10, 13, 13 as 5×13, or 65; and that of the triangle 13, 14, 15 as $7 \times \frac{1}{2} \times (13 + 15)$, or 98. He also states that the area of any quadrilateral whose sides are a, b, c, d is

$$\sqrt{(s-a)(s-b)(s-c)(s-d)},$$

where $s = \frac{1}{2}(a+b+c+d)$, a formula that is true only for cyclic quadrilaterals. His rule for the quadrilateral is as follows: "Half the sum of the sides set down four times, and severally lessened by the sides, being multiplied together, the square root of the product is the exact area."[3] He uses 3 as the "practical value" of π and $\sqrt{10}$ as the "neat value."

Brahmagupta's Algebra. The *Kuṭakhādyaka* applies algebra to astronomical calculations. For example, "One who tells, when given positions of the planets, which occur on certain lunar days or on days of other denomination of measure, will recur on a given day of the week, is versed in the pulverizer."[4]

[1] *Lectures on Indeterminate Equations.* The *kuṭaka* (*kuṭṭaka*, *cuṭacā*) is defined by Colebrooke (*loc. cit.*, p. vii) as "a problem subservient to the general method of resolution of indeterminate problems of the first degree." The word means "pulverizer" and is used as a name for algebra. *Ibid.*, p. 325; J. Taylor, *Lilawati*, p. 129 (Bombay, 1816); hereafter referred to as Taylor, *Lilawati*, with this spelling. The word *khādyaka* means "sweetmeat," such fanciful names being common in the East.

[2] Colebrooke, *loc. cit.*, p. 277.

[3] *Ibid.*, p. 295.

[4] Question 7 of the Colebrooke translation.


In his chapter on computation[1] Brahmagupta gives the usual rules for negative numbers. He also has a chapter on quadratic equations, the rule[2] for solving an equation of the type $x^2 + px - q = 0$ being substantially a statement of the formula

$$x = \frac{\sqrt{p^2 + 4q} - p}{2},$$

which evidently gives one root correctly.

In the case of simultaneous equations of the first degree the unknowns are spoken of as "colors," and the problems are chiefly astronomical. Indeed, Brahmagupta was the first Indian writer, so far as we know, who applied algebra to astronomy to any great extent. While the fanciful problems so often found in Indian works are generally wanting, a commentator has supplied various examples to illustrate certain of his rules. Two such problems are as follows:

On the top of a certain hill live two ascetics. One of them, being a wizard, travels through the air. Springing from the summit of the mountain he ascends to a certain elevation and proceeds by an oblique descent diagonally to a neighboring town. The other, walking down the hill, goes by land to the same town. Their journeys are equal. I desire to know the distance of the town from the hill, and how high the wizard rose.



The commentator takes the case here shown, and finds x to be 8.

A bamboo 18 cubits high was broken by the wind. Its tip touched the ground 6 cubits from the root. Tell the lengths of the segments of the bamboo.[3]

Indeterminate Equations. It is indicative of the state of algebra at this time that Brahmagupta was interested in the solution of indeterminate equations. Āryabhaṭa had already

[1] *Shaṭ-triṅśat-paricarman.* [2] Page 346 of the Colebrooke translation.
[3] From his arithmetic. For an early Chinese version, see page 139.

considered the question of the integral solutions of $ax \pm by = c$, but Brahmagupta actually gave as the results

$$x = \pm cq - bt,$$
$$y = \mp cp + at,$$

where t is zero or any integer and p/q is the penultimate convergent of a/b.[1] He also considered the so-called Pell Equation of the form

$$Du^2 + 1 = t^2,$$

but the solution was first effected, so far as we know, by Bhāskara in the 12th century.

For the sides of the right-angled triangle Brahmagupta gave the two sets of values

$$2mn, \quad m^2 - n^2, \quad m^2 + n^2,$$

and

$$\sqrt{m}, \quad \frac{1}{2}\left(\frac{m}{n} - n\right), \quad \frac{1}{2}\left(\frac{m}{n} + n\right),$$

values which he probably obtained from Greek sources.

Brahmagupta was accused of propagating falsehoods relating to science for the purpose of pleasing the bigoted priests and ignorant rabble of his country, hoping thus to avoid the fate that befell Socrates,[2] all of which shows that he was a man of recognized importance in his day.

Progress retarded in India. From this time to the year 1000 learning seems to have made but little progress in northern India. In the 8th century the Rajput dynasty succeeded the high-minded Valabhis, and for two hundred years the history of this part of India is a blank. Not a piece of literature of any value remains, nor any work of art or of industry.[3] The abode of mathematics now moved northward and is found for two or three centuries in Persia and in the other lands which had been brought under Moslem rule. In southern India, however, there must have been some encouragement of mathematics, as will be seen from the great work of Mahāvīra.

[1] Kaye, *Indian Math.*, p. 16.
[2] See Sachau's note in his translation of Albêrûnî's *India*, II, 304.
[3] Dutt, *History of Civ. in Anc. India*, II, 162.

Mahāvīra. The third of the great Hindu writers of this period is Mahāvīrācārya, Mahāvīra the Learned, who wrote the *Gaṇita-Sāra-Sangraha.*[1] This writer probably lived at the court of one of the old Rāshṭrakūta monarchs who ruled over what is now the kingdom of Mysore, and whose name is given as Amoghavarsha Nirpatuṅga. This king ascended the throne in the first half of the 9th century, so that we may roughly fix the date of the treatise in question as *c.* 850, or between the dates of Brahmagupta and Bhāskara,[2] though nearer to the former.

The work begins, as is not unusual with Oriental treatises, with a salutation of a religious nature. In this case the words are addressed to the author's patron saint, the founder of the religious sect of the Jainas (Jinas), a contemporary of Buddha:

> Salutation to Mahāvīra, the Lord of the Jinas, the protector [of the faithful], whose four infinite attributes, worthy to be esteemed in [all] the three worlds are unsurpassable [in excellence].
>
> I bow to that highly glorious Lord of the Jinas, by whom, as forming the shining lamp of the knowledge of numbers, the whole of the universe has been made to shine.

Mahāvīra's Sources. In general it may be said that Mahāvīra seems to have known the work of Brahmagupta. It would have been strange if this had not been so, for the *Brāhma-sphuṭa-siddhānta* was probably recognized in his time as one of the standard authorities. Mahāvīra seems to have made the effort to improve upon the work of his predecessor, and certainly did so in his classification of the operations, in the statement of rules, and in the nature and number of problems. As a result his work became well known in southern India, although there is no definite proof that Bhāskara (*c.* 1150), living in Ujjain, far to the north, was familiar with it.

Mahāvīra's Work. The work itself consists of nine chapters. The first is introductory and relates chiefly to the measures

[1] M. Raṅgācārya, *The Gaṇita-Sāra-Sangraha of Mahāvīrācārya*, Sanskrit and English, Madras, 1912; hereafter referred to as *Mahāvīra. Gaṇita-Sāra* means "Compendium of Calculation."

[2] He lived *c.* 1150. See page 275.

used, the names of the operations, numeration, negatives, and zero. Eight operations with numbers are given, addition (except in series) and subtraction (even with fractions) being omitted as if presupposed. One interesting feature is the law relating to zero, which is stated thus: "A number multiplied by zero is zero, and that [number] remains unchanged when it is divided by, combined with, [or] diminished by zero." That is, the law given by Bhāskara for dividing by zero is not here recognized, division by zero being looked upon as of no effect. The law of multiplication by negative numbers is stated, and the imaginary number is thus disposed of: "As in the nature of things a negative [quantity] is not a square [quantity], it has therefore no square root."

In his arithmetic operations he first treats of multiplication. He then considers in order the topics of division, squaring, square root, cubing, cube root, and the summation of series. In his work in series he includes some treatment of arithmetic and geometric progressions and of *Vyutkalita*, that is, the summation of a series after a certain number of initial terms (*ista*) have been cut off, a theory which, as we have seen (p. 155), occupied the attention of Āryabhaṭa.

The most noteworthy feature in his treatment of fractions is that relating to the inverted divisor, the rule being set forth as follows: "After making the denominator of the divisor its numerator [and *vice versa*], the operation to be conducted then is as in the multiplication [of fractions]." It is curious that this device, which from another source we know to have been used in the East, became a lost art until again adopted in Europe in the 16th century.

His method of approach to the subject of quadratic and radical equations is through fanciful problems of which the following is a type:

> One fourth of a herd of camels was seen in the forest; twice the square root [of that herd] had gone on to mountain slopes; and three times five camels [were] however, [found] to remain on the bank of a river. What is the [numerical] measure of that herd of camels?

This evidently requires the finding of the positive root of the equation $\frac{1}{4}x + 2\sqrt{x} + 15 = x$, or, in general, the solution of an equation of the type $x - (bx + c\sqrt{x} + a) = 0$, the rule for which is given. The chapter also contains various other types of equations involving some knowledge of radical quantities.

A single example will also suffice to show the nature of his indeterminate problems:

Into the bright and refreshing outskirts of a forest, which were full of numerous trees with their branches bent down with the weight of flowers and fruits, trees such as jambu trees, lime trees, plantains, areca palms, jack trees, date palms, hintala trees, palmyras, punnāga trees, and mango trees—[into the outskirts], the various quarters whereof were filled with the many sounds of crowds of parrots and cuckoos found near springs containing lotuses with bees roaming about them—[into such forest outskirts] a number of weary travelers entered with joy. [There were] sixty-three [numerically equal] heaps of plantain fruits put together and combined with seven [more] of those same fruits, and these were equally distributed among twenty-three travelers so as to have no remainder. You tell me now the numerical measure of a heap of plantains.

Mahāvīra's Treatment of Areas. His work in the measurement of areas is somewhat like the corresponding chapter in Brahmagupta's treatise, although it is distinctly in advance of the latter. Mahāvīra makes the same mistake as Brahmagupta with respect to the formula for the area of a trapezium (trapezoid) in that he does not limit it to a cyclic figure. The same error enters into his formula for the diagonal of a quadrilateral, which he gives as

$$\sqrt{\frac{(ac+bd)(ab+cd)}{ad+bc}} \quad \text{or} \quad \sqrt{\frac{(ac+bd)(ad+bc)}{ab+cd}}.$$

For the Pythagorean triangle Mahāvīra gives rules similar to those of Brahmagupta. For π he uses $\sqrt{10}$, a common value all through the East and also in medieval Europe. He was the only Hindu scholar of the native school who made any serious attempt to treat of the ellipse, but his work was inaccurate.

His rule for the sphere is interesting, the approximate value being given as $\frac{9}{2}(\frac{1}{2}d)^3$, and the accurate value as $\frac{9}{10}\cdot\frac{9}{2}(\frac{1}{2}d)^3$, which means that π must be taken as $3.03\frac{3}{4}$.

All things considered, the work of Mahāvīra is perhaps the most noteworthy of the Hindu contributions to mathematics, possibly excepting that of Bhāskara, who lived three centuries later. Mahāvīra may have known the works of Chinese scholars, for the value that he gives for the area of the segment of a circle, $\frac{1}{2}(c+a)a$, was given six centuries earlier by Ch'ang Ts'ang, but in any case he was a man of scientific attainments.

Bakhshālī Manuscript. Another work that stands out with some prominence in this period is the Bakhshālī manuscript.[1] This work, of uncertain origin and date, contains material relating to both arithmetic and algebra. It was formerly referred to the early part of our era and then to the 8th or 9th century, but it gives evidence of having been written even after the latter period, and possibly it is not even of Hindu origin. The nature of the work may be inferred from a single problem:

A merchant pays duty on certain goods at three different places. At the first he gives $\frac{1}{3}$ of the goods, at the second $\frac{1}{4}$ [of the remainder] and at the third $\frac{1}{5}$ [of the remainder]. The total duty is 24. What was the original amount of the goods?[2]


4. Persia and Arabia

Persia. We are apt to think that the rise of learning in the lands conquered by the Mohammedans was due solely to Arab influence, but this is not the case. In Persia, for example, Khosrú the Holy,[3] a generous patron of science, invited to his

[1] R. Hoernle, "The Bakhshālī Manuscript," *Indian Antiquary*, Vol. XVIII (1888); G. R. Kaye, "Notes on Indian Mathematics," in *Journ. and Proc. of the Asiatic Soc. of Bengal*, III (2), 501 (hereafter referred to as Kaye, *Notes*); and "The Bakhshālī Manuscript," *ibid.*, VIII (2), 349 (hereafter referred to as Kaye, *Bakhshālī*).

[2] The answer is 40, which necessitates the bracketed words.

[3] Khosrú I, Anôschirvân. He was a contemporary of Justinian, who was crowned emperor in Constantinople in 527. See W. S. W. Vaux, *Persia*, p. 169 (London, 1875); T. Nöldeke, *Aufsätze zur persischen Geschichte*, p. 113 (Leipzig, 1887).



BAKHSHĀLĪ MANUSCRIPT

A portion of this manuscript of which the date is still unsettled. It may be of the 10th century. From Kaye's *Indian Mathematics*

court scholars from Greece and encouraged the influx of Western culture. In his reign Aristotle and Plato were translated and doubtless the works of the Greek mathematicians were made known.

Christian Scholars in Mesopotamia. At about the time of the rise of the Mohammedan power there were various Christian centers of learning in the regions over which the Arabs were soon to hold sway. These were found in the monasteries which were scattered throughout the Near East. Of the scholars who taught in these retreats, the most learned one of the 7th century was Severus Sebokht,[1] a titular bishop who lived in the convent of Kenneshre on the Euphrates in the time of the patriarch Athanasius Gammala (who died in 631) and his successor John. He distinguished himself in the studies of philosophy, mathematics, and theology, and in his time the convent of Kenneshre became the chief seat of Greek learning in western Syria. He wrote on astronomy, the astrolabe, and geography. In one of the fragments of his works which have come down to us, of date 662, he directly refers to the Hindu numerals. He seems to have been hurt by the arrogance of certain Greek scholars who looked down on the Syrians, and in defending the latter he claims for them the invention of astronomy. He asserts the fact that the Greeks were merely the pupils of the Chaldeans of Babylon, and he claims that these same Chaldeans were the very Syrians whom his opponents condemn. He closes his argument by saying that science is universal and is accessible to any nation or to any individual who takes the pains to search for it. It is not, therefore, a monopoly of the Greeks, but is international.

Sebokht on our Numerals. It is in this connection that he mentions the Hindus by way of illustration, using the following words:

[1] J. Ginsburg, "New Light on our Numerals," *Bulletin of the Am. Math. Soc.*, XXIII (2), 366, from which extracts have been freely made. Attention of English readers was first called to this writer's mathematical works by Professor Karpinski, *Science* (U.S.). June, 1912. See also E. R. Turner, *Popular Sci. Mo.*, December, 1912

I will omit all discussion of the science of the Hindus, a people not the same as the Syrians; their subtle discoveries in this science of astronomy, discoveries that are more ingenious than those of the Greeks and the Babylonians; their valuable methods of calculation; and their computing that surpasses description. I wish only to say that this computation is done by means of nine signs. If those who believe, because they speak Greek, that they have reached the limits of science should know these things they would be convinced that there are also others who know something.

Bagdad. It was at Bagdad, on the Tigris River, that mathematics had its greatest encouragement under the Mohammedan ascendancy. Built upon the ruins of an ancient town by the caliph[1] al-Mansûr (712–774/5), one of the Abbasides,[2] Bagdad[3] became the intellectual center of the Mohammedan world, a second Alexandria in its fostering of learning. In al-Mansûr's reign (*c.* 766) a work mentioned as the *Sindhind* is said to have been brought to his court by an Indian scholar named Kankah (Mankah?), the Hindu astronomy and mathematics being thus made known to the scholars of Bagdad. This work may have been the *Sūrya Siddhānta* or it may have been some other work bearing the title *Siddhānta*, this name being nearer to *Sindhind* than any other Sanskrit word likely to be meant. It is generally believed, however, that it was the *Brahmasiddhānta* of Brahmagupta, whose works are known to have been brought to Bagdad at this time.[4]

To the court of the Caliphs there also came, so the story goes, a Persian by the name of Ya'qûb ibn Ṭâriq (died 796). He is said to have written (775) on the sphere (mathematical astronomy) and the calendar, and to have edited, and probably to have assisted in translating, the works of Brahmagupta

[1] Calif, from *Khalīfah*, successor (of the Prophet).

[2] Ab bās'ides or Ab'ba sides, the (at least pretended) descendants of Abbâs, uncle and adviser of Mohammed. Al-Mansûr reigned from 753/4 to 774/5. For rules for pronouncing Arabic names, see page xx.

[3] Persian *Bagadata*, "God-given"; in Arabic, Dār al-Salām, "Abode of Peace." Also spelled Baghdad.

[4] Sachau's preface to his translation of Albêrûnî's *India*, I, xxxi. As to the absence of Arabic records to prove that any embassy came from India at this time, see *ibid.*, II, 313.

above mentioned. To the same court there came (and the records of this fact are somewhat more trustworthy) the astronomer Abû Yaḥyâ,[1] and there he translated the *Tetrabiblos* of Ptolemy, thus assisting to begin the great movement that led to the introduction of the classics of Greek mathematics into the court of the Caliphs.

About the same time al-Fazârî[2] (died 777), working also at Bagdad, wrote on astrology and the calendar. He was the first Moslem, so far as is known, to construct astrolabes and to write on mathematical instruments. His famous contemporary, Jeber,[3] the greatest alchemist of the Arabs, also wrote on the astrolabe and possibly on mathematics.[4]

It was in this reign that another al-Fazârî,[5] son of the one already mentioned, a man of unusual scholarship, particularly in the field of astronomy, was asked by the caliph to translate the *Siddhānta* brought to Bagdad by Kankah. It was on this translation that Mohammed ibn Mûsâ al-Khowârizmî (*c.* 825) based his astronomical tables.

Harun al-Rashid. Harun al-Rashid,[6] well known to us from the *Arabian Nights Tales*, was a great patron of learning. Under his influence several of the Greek classics in science, including part of Euclid's works, were translated into Arabic. Indeed, it is to the Arabic versions that medieval Europe was indebted for its first knowledge of Euclid's *Elements*. In his reign there was a second influx of Hindu learning into Bagdad, especially in the line of medicine and astrology.

[1] Abû Yaḥyâ al-Baṭrîq, who died about 796–806.

[2] Ibrâhîm ibn Ḥabîb ibn Soleimân ibn Samora ibn Jundab, Abû Isḥâq al-Fazârî.

[3] Jâbir ibn Ḥaiyân al-Ṣûfî, Abû 'Abdallâh (died *c.* 777), one of two prominent scholars known by the name of Geber in the Middle Ages.

[4] The matter is discussed briefly in H. Suter, "Die Mathematiker und Astronomen der Araber und ihre Werke," in Volume X of the *Abhandlungen*. The transliteration of Arabic names is taken from Suter's list, with the change of *el* to *al*, of *ǧ* to *j*, of *š* to *sh*, of *ch* to *kh*, of *w* to *v*, and of *j* to *y* as in English. While this is not always desirable in the case of *sh*, as in Isḥâq, it is much simpler for the general reader. With respect to all these names the student should consult Suter's work.

[5] Mohammed ibn Ibrâhîm ibn Ḥabîb, Abû 'Abdallâh al-Fazârî. He died between 796 and 806.

[6] Hârûn al-Rashîd, Aaron the Just. He reigned from 786 to 808/9.

Al-Mâmûn. Harun al-Rashid's son, al-Mâmûn (reigned 809–833), was also a great patron of learning; indeed, he was more than a mere patron, for he erected an observatory at Bagdad and himself took observations there. He is also credited with supervising two geodetic surveys in Mesopotamia for the purpose of determining the length of a degree of the meridian. Under his direction the translation of the Greek classics continued, the *Almagest* of Ptolemy being put into Arabic and the translation of the *Elements* of Euclid being completed. In order to show the great activity among the Arabs in the field of mathematics, and the general nature of the work accomplished, a brief list of names will be given, with notes that are necessarily condensed, although it is apparent that most of the names are unfamiliar and most of the details will pass from the reader's mind.

It is evident that astronomy was the science of this period that did most to bring mathematics into a favorable light at court. Linked up with astrology on the one hand and with mathematics on the other, it introduced just enough of superstition through the former to help establish the latter science

Writers in al-Mâmûn's Reign. Among those who, in al-Mâmûn's remarkable reign, wrote upon mathematical astronomy, thus assisting to advance the study of trigonometry, the following scholars deserve special mention, not so much for their genius as for their spirit: al-Tabarî,[1] who wrote a commentary on Ptolemy's *Tetrabiblos*; al-Nehâvendî,[2] who prepared a set of astronomical tables; al-Mervarrûḏî,[3] who made astronomical observations at Damascus and Bagdad (*c.* 830); al-Aṣṭorlâbî,[4] who lived in Bagdad (*c.* 830), wrote on astronomy and geodesy, and was celebrated as a maker of astrolabes and other astronomical instruments; Messahala,[5] a Jewish

[1] Omar ibn al-Farrukhân, Abû Ḥafṣ al-Tabarî, died *c.* 815.

[2] Aḥmed ibn Mohammed al-Nehâvendî, died *c.* 835–845.

[3] Khâlid ibn 'Abdelmelik al-Mervarrûḏî. [4] 'Alî ibn 'Isâ al-Aṣṭorlâbî

[5] Mâ-shâ'-allâh ibn Aṯarî. The spelling in the text is the one commonly used in the West. The text of one of his MSS. was published by W. W. Skeat in his edition of Chaucer's *Astrolabe*, London, 1872. His chief work was done just before al-Mâmûn's reign.

astrologer, who wrote (*c.* 800) a treatise on the astrolabe that seems to have influenced the later works of Rabbi ben Ezra (*c.* 1150) and Chaucer (*c.* 1400); and Alfraganus[1] (*c.* 833), to use his European name, who wrote on sundials, astronomy, and the *Almagest*.

Mohammed ibn Mûsâ al-Khowârizmî. The greatest mathematician at the court of al-Mâmûn was Mohammed ibn Mûsâ al-Khowârizmî,[2] Abû 'Abdallâh (died between 835 and 845), a native of Khwarezm, the country in which is now the city of Khiva. Although an astronomer and the author of several astronomical tables and of works on dials, the astrolabe, and chronology, he is best known for having written the first work bearing the name "algebra," a treatise based upon Greek models.[3] He also wrote on arithmetic, this work being translated into Latin by Robert of Chester or by Adelard of Bath under the title *Algoritmi de numero Indorum*, whence such words as *algorism* and *augrim*,[4] derived from al-Khowârizmî. The title of the algebra was *'ilm al-jabr wa'l muqabalah*, "the science of reduction and cancellation."[5] After al-Mâmûn's death mathematics continued to flourish in Bagdad for about a century and a half, although, as might be expected, with somewhat less encouragement.[6]

[1] Mohammed ibn Ketîr al-Farġânî. The European translators also used such forms as Alfergani and Alfragan. Johannes Hispalensis translated his version of the *Almagest* into Latin, and it was printed at Ferrara in 1493 and again, with a preface by Melanchthon, at Nürnberg in 1537.

[2] This transliteration is more familiar to English readers than is Suter's el-Chowârezmî or Chwârezmî. The name means Mohammed son of Moses, the Khwarezmite. C. Huart, *History of Arab Literature*, pp. 131, 292, 207 (London, 1903), says that there were two others by the name of al-Khowârizmî, one (the geographer) of 935–993 (or 1002), and the other of *c.* 1036, but Huart seems to have been confused in this matter. The name also appears as al-Khowârazmî and as al-Khowâruzmî.

[3] L. C. Karpinski, *Robert of Chester's Latin Translation of the Algebra of al-Khowarizmi*, New York, 1915.

[4] So Chaucer speaks, in the *Canterbury Tales*, of "augrim stones."

[5] See Volume II, Chapter VI.

[6] Another Mohammed of Bagdad wrote a work on the division of surfaces. On the relation of this work to Euclid's book on the divisions of figures, see the careful study by Professor R. C. Archibald in his *Euclid's Book on Divisions of Figures*, pp. 1–8 (Cambridge, 1915).

Other Scholars of Bagdad. Almâhânî[1] (*c.* 860), as he is commonly called, an astronomer of high standing, is perhaps best known for having written upon the familiar problem of Archimedes relating to the cutting of a sphere into segments having a given ratio of volume. In his stereometric solution of the cubic equation involved in this problem he made use of the sine of a trihedral angle. He also wrote commentaries on Books V and X of Euclid's *Elements* and on the work of Archimedes on the sphere and cylinder.

Alchindi[2] (*c.* 860), to use the name by which he was generally known in medieval Europe, was commonly called "the philosopher of the Arabs." He wrote on a large variety of topics, including astronomy, astrology, optics, and number. Gherardo of Cremona (*c.* 1150) translated his work on optics into Latin.

About 870 there lived in Bagdad three scholars known as the Benî Mûsâ (sons of Moses) or the Three Brothers.[3] They were the sons of Mûsâ ibn Shâkir, a reformed robber who had finally devoted himself to geometry and astronomy in al-Mâmûn's court. Of these brothers, Mohammed, Aḥmed, and al-Ḥasan, the first-named was the most celebrated, but all three gave attention to securing the best scientific works of the Greeks and to having them translated. They wrote on medicine, conics, geometry, mensuration, the trisection of an angle, and other scientific subjects. They used the conchoid in the trisection problem and the string fastened to the foci in the construction of an ellipse.

At this period there worked for a time in Bagdad the celebrated Ṯâbit ibn Qorra[4] (826–901), a physician of prominence, but better known for his work in philosophy and mathematics, and particularly for the claim that he was successful in applying

[1] Mohammed ibn 'Îsâ, Abû 'Abdallâh al-Mâhânî, of Bagdad, died probably between 874 and 884.

[2] Ya'qûb ibn Isḥâq ibn al-Ṣabbâḥ al-Kindî, Abû Yûsuf, died *c.* 873/4.

[3] M. Curtze, "Liber Trium Fratrum de Geometria," in *Nova Acta der K. Leop.-Carol. Deutschen Akad. der Naturforscher*, XLIX, No. 2 (Halle, 1885).

[4] Ṯabit ibn Qorra ibn Mervân, Abû-Ḥasan, al-Ḥarrânî, a native of Ḥarrân in Mesopotamia, where he also spent some of his later years.

algebra to geometry. He revised the translation of Euclid's *Elements* made by Isḥâq ibn Ḥonein, a renowned physician (died 910), and the translation of the so-called "middle books," that is, of those books written between the time of Euclid and that of Ptolemy.[1] He also wrote extensively on astronomy, the *Almagest*, conics, elementary geometry, Euclid, magic squares, amicable numbers, and astrology. Gherardo of Cremona (*c.* 1150) and Johannes Hispalensis (*c.* 1140) translated certain of his works. He had a son,[2] a physician, who also followed in his father's steps, writing on astronomy and geometry, and revising one of the translations of Archimedes from the Syriac into Arabic.

At about this time an Egyptian—Aḥmed ibn Yûsuf[3]—wrote on proportion and astronomy and discussed the *figura cata*, that is, the proposition of Menelaus relating to the segments of the sides of a triangle cut by a transversal.

Christian and Jewish Scholars in Bagdad. To Bagdad there also came at this time various Jewish and Christian writers, their names being commonly given in Arabic form. Among these were Sahl ibn Bishr,[4] an astrologer, who had already gained considerable reputation in Khorâsân. He wrote a work on algebra. Part of his writings appeared in print in Venice (1493) and part in Basel (1533). There was also Abû'l-Ṭaiyib,[5] who gave up his Jewish religion and adopted the faith of Islam. He compiled a set of astronomical tables and seems to have written on trigonometry. Among the Christians there was Qosṭâ ibn Lûqâ al-Ba'albekî[6] (died *c.* 912/3), a

[1] L. M. L. Nix, *Das fünfte Buch der Conica des Apollonius von Perga*, with Arabic text and German translation. Leipzig, 1889.


[2] Sinân ibn Ṯâbit ibn Qorra, Abû Sa'îd, died 943. See Suter's list, *Abhandlungen*, X, 51.

[3] Aḥmed ibn Yûsuf ibn Ibrâhîm, Abû Ja'far, al-Miṣrî (died *c.* 912/3). Al-Miṣrî means the Egyptian, and the name is applied to other writers as well. There is some doubt as to his works. He was the son of Yûsuf ibn Ibrâhîm ibn al-Dâya, who was known as "the Arithmetician" and lived in Damascus, Bagdad, and Egypt.

[4] Sahl ibn Bishr ibn Ḥabîb ibn Hânî (or Hâyâ), Abû 'Oṯmân (*c.* 850).

[5] Sind ibn 'Alî, Abû'l-Ṯaiyib (*c.* 850).

[6] Kosta, son of Luke, from Baalbek, known to early Europeans as Kusta ben Luca.



PYTHAGOREAN THEOREM IN TÂBIT IBN QORRA'S TRANSLATION OF EUCLID

The translation was made by Isḥâq ibn Ḥonein (died 910) but was revised by Tâbit ibn Qorra, *c.* 890. This manuscript was written in 1350

physician, who translated the *Spherics* of Theodosius and parts of Aristarchus, Autolycus,[1] Hypsicles, Heron, and Diophantus, and who wrote a geometry in catechism form. There was also a Greek Christian, Nazîf ibn Jumn (or Jemen), known as al-Qass (the priest), who translated Euclid X; and another of the same faith, al-Jorjânî,[2] a physician, who wrote a compendium of the *Almagest*.

It is possible that it was about this time and in this region that the anonymous Hebrew work entitled *Mishnath ha-Middoth (Theory of Measures)* was written, but the place and date are quite unknown. It is primarily on the measurement of geometric solids, and some of its features recall the work of al-Khowârizmî on mensuration.[3]

Later Writers. After the reigns of the first three caliphs of Bagdad the science of astronomy still continued to be the antechamber of mathematics. Thus we find such writers in this field as al-Mervazî,[4] who wrote extensively on astronomy and astronomical instruments; Albumasar[5] (died 886), the most celebrated of the Arab writers on astrology, who was led by this science to the study of astronomy; Aḥmed ibn al-Ṭaiyib[6] (*c.* 890), of Persian origin, a pupil of Alchindi's, who wrote on algebra and arithmetic as well as on astrology and music; and al-Dînavarî,[7] who wrote on algebra, astronomy, and the Hindu methods of computation. There was also the

[1] A Greek astronomer who lived *c.* 360 B.C. The others have already been mentioned.

[2] 'Îsâ ibn Yaḥyâ al-Masîḥî, Abû Sahl, al-Jorjânî, died *c.* 1009/10. Al-Masîḥî means a believer in the Messiah, a Christian. The Suter list does not give the place where either of the last two lived.

[3] M. Steinschneider, *Festschrift Zunz* (Berlin, 1864); H. Shapiro, *Abhandlungen*, with translation and commentary, III, 3; F. Rosen, *The Algebra of Mohammed ben Musa*, p. 70 (London, 1831).

[4] Aḥmed ibn 'Abdallâh al-Mervazî, a native of Merv (probably died between 864 and 874), known as Ḥabash al-Ḥâsib ("Ḥabash the computer").

[5] As he was commonly known in medieval Europe. His name was Ja'far ibn Mohammed ibn 'Omar al-Balkhî (from Balkh, in Khorâsân), Abû Ma'shar.

[6] Aḥmed ibn Mohammed ibn Mervân, Abû'l-'Abbâs, al-Sarakhsî, known as Aḥmed ibn al-Ṭaiyib.

[7] Aḥmed ibn Dâ'ûd, Abû Ḥanîfa, al-Dînavarî (died 895). He lived most of the time in Dînavar, his native place.

well-known scholar Albategnius[1] (died 929), as he was called in Europe, who was justly esteemed for his astronomical writings[2] and tables. Among the many other scholars of this period there may be mentioned Rhases (died 932),[3] to use his European name, a celebrated physician who wrote on geometry and astronomy; a grandson of Ṯâbit ibn Qorra,[4] also a physician, who wrote on conics, dialing, and elementary geometry; al-Fârrâbî,[5] a native of Fârâb in Turkestan, who wrote a commentary on Euclid and was a philosopher of high standing; Ibn Yûnis,[6] who, next to al-Battânî, was the most celebrated astronomer among the Arabs; and al-Ḥarrânî,[7] who wrote a commentary on Euclid.

The 10th century saw several writers of somewhat higher attainments, among whom the best-known was Abû'l-Wefâ[8] (940–998), celebrated for his improvements in trigonometry, his introduction of the tangent (*umbra versa*), and his computation of tables of sines and tangents for every 10′; it is also very likely that he is entitled to credit for the use of secants and cosecants. He was also prominent as a writer on arithmetic, algebra, geometry, and astronomy.

Among the other writers of this period who are worthy of special mention were al-Ḥaiṯam of Baṣra,[9] who wrote on algebra, astronomy, geometry, gnomonics, and optics; Abû Ja'far al-Khâzin (died between 961 and 971), who attempted

[1] Mohammed ibn Jâbir ibn Sinân, Abû 'Abdallâh, al-Battânî, a native of Battan, in Mesopotamia. He is also known as al-Raqqî, from the fact that he made his observations at Raqqa on the Euphrates.

[2] Translated by Robert of Chester (*c.* 1140) or Robertus Retinensis, referred to later. The work was printed in 1537.

[3] Mohammed ibn Zakarîyâ al-Râzî, Abû Bekr.

[4] Ibrâhîm ibn Sinân Ṯâbit ibn Qorra, Abû Isḥâq, son of the Sinân already mentioned. Born 908/9; died 946.

[5] Mohammed ibn Mohammed ibn Ṭarkhân ibn Auzlag, Abû Naṣr, al-Fârrâbî; died at Damascus, 950/1.

[6] 'Alî ibn Abî Sa'îd 'Abderraḥmân ibn Aḥmed ibn Yûnis (or Yûnos), Abû'l-Ḥasan, al-Ṣadafî; died 1009.

[7] Ibrâhîm ibn Hilâl ibn Ibrâhîm ibn Zahrûn, Abû Isḥâq, al-Ḥarrânî. Born 923; died at Bagdad, 994.

[8] Mohammed ibn Mohammed ibn Yaḥyâ ibn Ismâ'îl ibn al-'Abbas, Abû'l-Wefâ al-Bûzjânî.

[9] Al-Ḥasan ibn al-Ḥasan ibn al-Ḥaiṯam, Abû 'Alî, *c.* 965–1039.

the solution of the cubic equation by the aid of conics and who wrote on Euclid and astronomy; and Kûshyâr ibn Lebbân,[1] who wrote on arithmetic, trigonometry, and astronomy.

Al-Nairîzî[2] (died *c.* 922/3) was one of the notable 10th century writers on Euclid. He was interested in astronomy and geometry, writing commentaries on both Ptolemy and Euclid, but it is the commentary on the *Elements*, translated into Latin by Gherardo of Cremona, that is best known.

As a type of the lesser commentators on Euclid in the 10th century there may be mentioned al-Ḥasan ibn 'Obeidallâh,[3] who wrote a commentary on the difficult parts of the *Elements*.

Translators into Arabic. Among the noteworthy translators of this period were al-Hajjâj,[4] who made two translations of at least six books of Euclid's *Elements*, and also translated Ptolemy's *Almagest*; al-Jauharî,[5] who made astronomical observations at Bagdad and Damascus (*c.* 830) and wrote a commentary on the *Elements* of Euclid; Ḥonein ibn Isḥâq,[6] who translated various Greek works, possibly including Ptolemy's *Tetrabiblos*, and who wrote on astronomy, but was more celebrated as a physician and a philosopher; and his son Isḥâq,[7] who was a physician and translated Euclid's *Elements* and *Data*, the *Almagest*, Archimedes on the sphere and cylinder, and probably the *Spherics* of Menelaus. Somewhat less well known, but worthy of mention, are al-Arjânî,[8] who wrote a commentary (*c.* 850) on Euclid X; al-Ḥimṣî,[9] who translated the first four books of Apollonius; and Sa'îd ibn Ya'qûb,[10] a physician, who translated parts of Euclid and of Pappus.

[1] Kûshyâr ibn Lebbân ibn Bâshahrî al-Jîlî, Abû'l-Ḥasan, *c.* 971–*c.* 1029.

[2] Al-Faḍl ibn Ḥâtim al-Nairîzî, Abû'l-'Abbâs.

[3] Al-Ḥasan ibn 'Obeidallâh ibn Soleimân ibn Vahb, Abû Mohammed, *c.* 925.

[4] Al-Hajjâj ibn Yûsuf ibn Maṭar, *c.* 786–*c.* 835.

[5] Al-'Abbâs ibn Sa'îd al-Jauharî.

[6] Ḥonein ibn Isḥâq, al-'Ibâdî Abû Zeid; born 809/10; died at Bagdad, 873.

[7] Isḥâq ibn Ḥonein ibn Isḥâq al-'Ibâdî, Abû Ya'qûb, died 910.

[8] Ibn Râhiweih al-Arjânî, or Arrajânî, according to Steinschneider the same as Isḥâq ibn Ibrâhîm ibn Makhlad al-Mervazî, who died at Nishâpûr in 852/3.

[9] Hilâl ibn Abî Hilâl al-Ḥimṣî, died 883/4. Al-Ḥimṣî means "from Emessa," in Syria.

[10] Sa'îd ibn Ya'qûb al-Dimishqî, Abû 'Otmân. He was living in 915.

Abû Kâmil. Between 850 and 930 there lived in Egypt Abû Kâmil,[1] who is known for several works but especially for his treatise on the pentagon and decagon[2] and for his arithmetic and algebra.[3] No writer of his time showed more genius than he in the treatment of equations and in their application to the solution of geometric problems.

About the same time there lived Abû'l-Faradsh Mohammed ibn Isḥâq, known as Ibn Abî Ya'qûb al-Nadîm, whose *Kitâb al-Fihrist* (*Book of Lists*), written *c.* 987, is a collection of brief biographies of various prominent mathematicians, both Greek and Mohammedan.[4]

Close of the Golden Age of Bagdad. In a general way it may be said that the Golden Age of Arabian mathematics was confined largely to the 9th and 10th centuries; that the world owes a great debt to Arab scholars for preserving and transmitting to posterity the classics of Greek mathematics; and that their work was chiefly that of transmission, although they developed considerable originality in algebra and showed some genius in their work in trigonometry.

5. The Christian West

The Dark Ages. The period from 500 to 1000 extends from about the time of the fall of Rome (455)[5] to the first reawakening of Europe under Pope Sylvester II (Gerbert). It includes the so-called Dark Ages, the period of the slow civilizing of the northern races, of the development of monastic schools, of the work of Charlemagne, and of the contact with Oriental civilization, chiefly through the Moors in Spain. In mathematics it was the era of the development of the Christian calendar in the West, and of little else. The barbarian had to be civilized, to assimilate slowly the Roman culture which he would have destroyed, and to receive a better religion. The Roman schools

[1] Abû Kâmil Shojâ ibn Aslam ibn Mohammed ibn Shojâ.

[2] H. Suter, *Bibl. Math.*, X (3), 15, 33.

[3] L. C. Karpinski, *Amer. Math. Month.*, XXI, 37, and *Bibl. Math.*, XII (3), 40. See also H. Suter, *Bibl. Math.*, XI (3), 100.

[4] Suter's translation appeared in the *Abhandlungen* (VI, 1) in 1892.

[5] The barbarians entered the city first in 410. The final fall is often given as 476.

had to be supplanted by those of the cathedral and the monastery, and all the mathematics required was limited to the needs of trade, to the keeping of accounts, and to the fixing of dates for Church festivals. In those parts of Europe less subject to Northern influence, such as Marseilles, Arles, and Narbonne, the needs of commerce were still such as to render necessary the arithmetic of exchange in the training of the merchant's apprentice. These cities maintained in this period their trade with Italy, Constantinople, and the Orient, sending dyes, cereals, pottery, and salt to the East, and importing silk from China, pearls from India, and even papyrus rolls from Egypt.[1]

Boethius. Anicius Manlius Severinus Boe'thius,[2] a Roman citizen, a member of the distinguished family of the Anicii, statesman, philosopher, mathematician, man of letters, and founder of the medieval scholasticism, lived at the opening of the period now under discussion. Persecuted for his uprightness, executed for his fearlessness, accepted by the Church as a martyr, his reputation and scholarship gave his books on mathematics high standing in the monastic schools for many centuries.

His greatest work, written while he was in prison, is the *Consolation of Philosophy*.[3] His mathematical works are an arithmetic,[4] a geometry,[5] and a work on music,[6] a subject then

[1] A. Rambaud, *Histoire de la Civilisation Française*, 12th ed., I, 115. Paris, 1911.

[2] Born at Rome *c.* 475; died at Pavia, 524. The more nearly correct Latin form is Boetius.

[3] *De consolatione philosophiae.*

[4] *Boetii de institutione arithmetica libri duo*, ed. Friedlein (Leipzig, 1867); hereafter referred to as *Boethius, ed. Friedlein.* The earliest manuscripts used in this edition are three of the 10th century, a fact worth noting in view of questions as to interpolations discussed later. Readers of Boethius and other Latin writers will find assistance in B. Veratti, "Sopra la Terminologia Matematica degli Scrittori Latini," *Memorie della R. Accad. . . . di Modena*, Vol. V.

[5] *Boetii quae fertur geometria*, in the Friedlein edition cited above. The earliest manuscript used in this edition is one of the 10th century. There is serious doubt as to whether Boethius wrote the *Ars Geometriae* attributed to him. See Tannery, *La Géométrie Grecque*, 128; H. Weissenborn, "Die Boetius-Frage," in the *Abhandlungen* II, 185.

[6] *Boetii di institutione musica libri quinque*, in the Friedlein edition cited above. The earliest manuscript used in this edition was mostly of the 9th century, although Books IV and V were earlier and some parts were missing.

ranked as part of mathematics. The arithmetic was based on the work of Nicomachus, and the geometry on the *Elements* of Euclid. Neither showed any originality in the domain of mathematics, but each was sufficiently successful in its presentation of the subject treated to permit of the general use of



FROM A DRAWING BY RAPHAEL

Fanciful sketches of Ptolemy and Boethius, now in the Accademia in Venice

these books in those monastic schools that had advanced far enough to demand courses in the theory of numbers and in demonstrative geometry.

Minor Writers. It is natural to expect that among the first Christian scholars few would be found with any interest in mathematics or the natural sciences. Their religious faith was too intense, their persecutions too real, and their lives too precarious to permit of speculations in these fields. The names of a few Christians have already been mentioned, but their contributions to mathematics were insignificant. With the

close of the 5th century, however, Christianity had become powerful enough to permit of the development of an intellectual class with interests outside of religious faith, and in this class we find the names of several scholars who showed some knowledge of the mathematics of the classical period.

Among these writers was Magnus Aurelius Cassiodo'rus,[1] a descendant of an ancient Roman family.[2] He was a statesman of distinction and was honored both by the last of the Roman rulers and by their Ostrogothic successors. He founded a monastery at Vivarium, and passed his last years within its walls. He insisted upon a high standard of scholarship for the clergy, and his writings show that he himself possessed, within the limits which conditions then imposed, that which he demanded for others. Cassiodorus wrote *De artibus ac disciplinis liberalium literarum*, a trivial sort of compendium of the seven liberal arts,—grammar, rhetoric, and dialectic composing the trivium, and arithmetic, geometry, astronomy, and music composing the quadrivium.[3] This work was widely used in the schools of the Middle Ages,[4] and nothing could better show the low state of learning than this feeble attempt at scholarship. There is also doubtfully assigned to him a *Computus Paschalis sive de indicationibus cyclis solis et lunae*, written in 562, one of the first treatises on the Christian calendar. The plan for the adoption of the Christian era, however, was worked out by Dionysius Exiguus, a Roman abbot, *c.* 525.

[1] Born at Scylaceum (Squillace), *c.* 470; died *c.* 564, a date sometimes given as 585. The name is also spelled Cassiodorius.

[2] For a popular but vivid account of his achievements see M. Crawford, *Rulers of the South*, II, 9.


[3] A common medieval verse reads:

> Gram loquitur, Dia verba docet, Rhet verba colorat,
> Mus canit, Ar numerat, Ge ponderat, As colit astra.

Petrus Pictaviensis, in a verse to Peter of Cluny, writes:

> Musicus, astrologus, arithmeticus, et geometra,
> Grammaticus, rhetor, et dialecticus est.

[4] The first collected edition of his works was published at Paris in 1584 and 1598.



MANUSCRIPT OF THE *ARITHMETICA* OF BOETHIUS

This MS., now in Mr. Plimpton's library, was written *c.* 1294. The scribe has used modern instead of Roman numerals

A little before the time of Cassiodorus there flourished Martianus Mineus Felix Capel'la,[1] author of an encyclopedia known as the *Nuptials of Philology and Mercury*.[2] It is a medley of prose and verse, one part of the work being on geometry and another on arithmetic. In connection with the latter Capella discusses various classes of numbers and the supposed mysteries of the smaller numbers. The book is even more arid than that of Cassiodorus, the only redeeming feature being the statement that Mercury and Venus revolve about the sun instead of the earth.[3]

Before the close of the 5th century there was born in Damascus a Syrian who took his name, Damas'cius,[4] from his birthplace. He was the last of the important Neoplatonists and was a disciple of the Marinus who succeeded Proclus (*c.* 485). In 510 he became director of the school at Athens. When Justinian closed the heathen schools of philosophy in that city (529), Damascius went to Persia, but returned five years later (534). His works were mostly philosophical, but his name has doubtfully been connected with a fifteenth book to be added to Euclid's *Elements*.

Almost the last of the Greeks to show any appreciation of mathematics before the medieval period fairly began was Euto'cius[5] of Ascalon. He wrote commentaries on the first four books of the conics of Apollonius. He also wrote on certain works of Archimedes,—the sphere and cylinder, the quadrature of the circle, and the work on equilibrium; and on the *Almagest* of Ptolemy, this last commentary being lost. These writings of Eutocius are of little value except as they supply certain information relating to Greek mathematics.

[1] Born possibly at Carthage, *c.* 420; died *c.* 490. See E. Narducci, in Boncompagni's *Bullettino*, XV, 505, with biography and bibliography. The name might properly have been given in Chapter IV, but Capella is more closely related to Boethius and Cassiodorus than to the last of the Greeks.

[2] The first edition appeared at Vincenza in 1499, *Opus Martiani Capelle de Nuptijs Philologie & Mercurij libri duo.*

[3] In the *De Astronomia*, the chapter entitled *Tellus quod non sit centrum omnibus planetis.* Fol. 333 of the 1592 edition.

[4] Δαμάσκιος.

[5] Εὐτόκιος. Fl. *c.* 560.

In the 6th century there seems also to have been written the *Codex Arcerianus*,[1] so called from the fact that it belonged at one time (1566–1604) to one Johannes Arcerius in Grö-ningen. While it relates largely to legal matters of a rural nature, it contains considerable information concerning the Roman surveyors.

There is little else to say for the century. It represents the lowest point on the curve of intellectual progress in Europe. The ecclesiastical element was unable to overcome the general ignorance of the masses, and aside from a faint light in the Irish monasteries, Europe was in darkness.

Isido'rus. The centuries immediately following the death of Boethius saw little interest in the literature and science of the classical period. Even as eminent a man as St. Ouen (*c.* 609–683) spoke of the works of Homer and Vergil as the trifling songs of impious poets[2] and made two distinct personages of Tullius and Cicero; while Gregory of Tours (538–594) uttered the lament: "Unhappy our days, for the study of letters is dead in our midst, and there is to be found no man able to record the history of these times." So debased was civiliza-tion that the few who stood for even the remnants of the old Latin cult resorted to doggerel verse, as Capella had done, or diluted their learning in the form of encyclopedias.

Prominent among those who developed the latter plan was Isidorus of Seville,[3] historian, grammarian, orator, theologian, bishop, and general scholar, as well as one of the most re-markable statesmen of the Middle Ages. St. Martin, in his funeral oration, describes him as "generous in his giving, affable in his entertaining, sober in his affections, free in his sentiments, equitable in his judgments, indefatigable in his ministrations," and celebrated for his integrity. A man of fortunate birth, he was helped by his family connections to begin a career of such remarkable success, relative to that of

[1] Mommsen puts it *c.* 450, and Cantor (*Die Römischen Agrimensoren*, p. 95 (Leipzig, 1875); hereafter referred to as Cantor, *Agrimensoren*) thinks it not later than the 7th century. See also Cantor, *Geschichte*, I, chap. 26.

[2] "Sceleratorum neniae poetarum." [3] Born at Seville, *c.* 570; died April 4, 636.

any contemporary, that the Council of Toledo (653), a few years after his death, could truthfully speak of him as "the extraordinary doctor, the latest ornament of the Catholic Church, the most learned man of the latter ages, always to be named with reverence." Since he was the most learned man of his time, it would be expected that his encyclopedia of the trivium and the quadrivium, the seven liberal arts, would contain some mathematics of merit. This work, called by him the *Origines* but often known as the *Etymologies*, consists of twenty books, the third one being on mathematics. The treatment, however, is trivial, the arithmetic being simply a brief condensation of Boethius, and the rest of the work being of as little scientific value.

Bede the Venerable. It was about a century after Isidorus that there was born at Monkton in Northumberland one of the greatest of the Church scholars of the Middle Ages, Baeda (*c.* 673–735), commonly known as Beda Venerabilis, the Venerable Bede,[1] and called by Burke "the father of English learning."

Of him Hallam[2] remarked that he "surpasses every other name of our ancient literary annals; and, though little more than a diligent compiler from older writers, may perhaps be reckoned superior to any man the world (so low had the East sunk like the West) then possessed." Four years before his death he prepared a list of the thirty-seven works which he had written up to that time, and added these words: "I have spent my whole life in the same monastery, and while attentive to the rule of my order and the service of the Church, my

[1] G. F. Browne, *The Venerable Bede* (London, 1880). For a discussion of his scientific works, see J. A. Giles, *Miscellaneous Works of the Venerable Bede*, VI, pp. v, 123 (London, 1843); J. Mabillon, "Ven. Bedae elogium historicum," in the *Opera Omnia* of Bede (Paris, 1862); K. Werner, *Beda der Ehrwürdige und seine Zeit* (Vienna, 2d ed., 1881). Bede was buried at Jarrow, but his remains were moved to Durham *c.* 1050 and his tomb may now be seen in the Galilee Chapel of the Cathedral. A good setting for the study of the education of the period may be found in F. P. Barnard, *Companion to English History (Middle Ages)*, p. 303 (Oxford, 1902).

[2] *Literature of Europe*, Chapter I, § 7 (London, n.d.).

constant pleasure lay in learning, or teaching, or writing."[1] Taught by Aldhelm and by John of Beverley, two of the heirs to the intellectual and spiritual treasure which Augustine bequeathed to Canterbury,[2] he was also a disciple of Archbishop Theodore of Tarsus and Abbot Adrian, two pioneers in bringing a high grade of scholarship to the monasteries, and thus he was well prepared to render service to the world and to lead a life "consecrated in noiseless activity to God."[3]

In mathematics his interests were in the ancient number theory, the ecclesiastical calendar, and the finger symbolism of number, and his writings include these and other mathematical subjects.[4] To him we are indebted for the best work on the calendar written during the Dark Ages, and for the best work up to his time on digital notation.[5] Certain mathematical recreations have also been attributed to him, but the evidence concerning their authorship is not conclusive.

Alcuin of York. The next great European scholar in mathematics was Al'cuin (735–804). Born in the year of Bede's death, less of a scholar than the latter but more of a man of action, he attained prominence in the State as well as in the Church. He studied in Italy,[6] taught at York,[7] was called (782) by Charlemagne[8] to assist him in his ambitious project for the education of his people, and became abbot of St. Martin

[1] "Semper aut discere aut docere aut scribere dulce habui," words worthy of the one whom Green, the historian, speaks of as "the first great English scholar."

[2] W. F. Hook, *Lives of the Archbishops of Canterbury*, I, 42 (London, 1860); A. Neander, *Church History*, 5th American ed., III, 12 (Boston, 1855); J. E. G. de Montgomery, *State Intervention in English Education*, p. 6 (Cambridge, 1902). See page 187, note 2.

[3] Neander. For a description of Bede's death see *loc. cit.*, p. 153.

[4] *De numeris, De temporum ratione, De numerorum divisione, De circulis sphaerae et polo, De astrolabio.*

[5] His *De temporibus* comes down only to 701/2. His *De temporum ratione* comes down to 726. This second work contains his *De Indigitatione sive de computo per gestum digitorum* and his *De ratione unciarum.*

[6] As shown by one of his letters (XV); see Libri, *Histoire*, I, 89.

[7] On the nature of the schools, see W. W. Capes, *The English Church in the 14th and 15th centuries*, p. 332 (London, 1900).

[8] Who addressed him as "Carissime in Christo praeceptor." Charlemagne reigned as king or emperor from 768 to 814.

of Tours. He wrote on arithmetic, geometry, and astronomy,[1] and his name is connected with a certain collection of puzzle problems[2] which has influenced the writers of textbooks for a thousand years. It is uncertain how much he may have had to do with this set of mathematical recreations, and considerable doubt has been thrown upon his connection with them through recent studies of a certain manuscript at Leyden.[3] This manuscript dates from the first part of the 11th century and is thought to have been written by, or at least inspired by, a monk named Adémar or Aymar, of the ancient house of Chabanais, who was born in 988 and who died on his way to the Holy Land in 1030. He had considerable reputation as a historian and a controversialist[4] and seems to have collected a large amount of material with no scientific care. These problems were very likely part of the medieval versions of Æsop's *Fables*, collections which, although probably begun by Æsop, in Samos, in the 7th century B.C., were modified by Babrius about the 3d century, and were still further corrupted in the Middle Ages. While problems attributed to Alcuin are found here, and probably interested Adémar as they did hundreds of others, there seems to be no good reason to believe that Alcuin may not have collected them from the medieval versions attached to the *Fables*. Certain it is that letters of Alcuin show that he wrote a set of puzzle problems, although there is no direct evidence that this is the one.[5] It would have been in keeping with his ideas to compile a book that should be amusing enough to relieve education of the drudgery of the time.[6]

[1] For his life and works, see G. F. Browne, *Alcuin of York* (London, 1908); C. J. B. Gaskoin, *Alcuin; his life and his work* (London, 1904); R. B. Page, *The Letters of Alcuin* (New York, 1909); A. F. West, *Alcuin and the rise of the Christian Schools* (New York, 1912).

[2] *Propositiones ad acuendos juvenes.*

[3] Cod. Vossianus Lat. oct. 15, edited by G. Thiele and published at Leyden in 1905.

[4] J. Lair, *L'Histoire d'Adémar*, Paris, 1899.

[5] It has been published in the works of Bede as well as in those of Alcuin. The oldest MS. of the work, written early in the 11th century, is now in Karlsruhe. In this are the words: "Dilectissimo fratri siguulfo presbytero alcuinus salutem," but naturally these words are not absolutely conclusive evidence.

[6] When "sub virga degere" meant school life and "pueri subiugales" meant pupils. T. Ziegler, *Geschichte der Pädagogik*, p. 29 (Munich, 1895).

In the collection is to be found, for example, the problem of the hare and hound, already ancient but made the more mysterious by the cipher title,

"De cursu cbnks bc fugb lepprks,"
for "De cursu canis ac fuga leporis."[1]

The continued private wars among petty lords in the 10th and 11th centuries made France a poor field for mathematical or other intellectual progress, and hence these two centuries produced little that was noteworthy.

Decay of British Learning. After the death of Alcuin the brilliant era that started in Great Britain with St. Au'gustine of Canterbury[2] (died *c.* 604 or 613) closed as suddenly as it began. The ravages of the Danes put an end to that feeling of security which makes for intellectual development, and when Alfred (848–900) came to the throne (871) he could only lament, "There was a time when people came to this island for instruction, but now we must obtain it abroad if we desire it." When Aethelstan,[3] the grandson of Alfred, came to the throne (925), however, he showed great interest in the fostering of learning, and in a poem written in the 14th century reference is made to the introduction of Euclid into England in the reign of this powerful ruler:

Thys grete clerkys name wes clept Euclyde,
Hys name hyt spradde ful wondur wide. . . .
The clerk Euclyde on thys wyse hyt fonde,
Thys crafte of gemetry yn Egypte londe;
Yn Egypte he tawȝhte hyt ful wyde,
Yn dyvers londe on every syde . . .
Thys craft com ynto Englond as y ȝow say
Yn tyme of good kynge Adelstonus day.[4]

[1] Cantor, *Agrimensoren*, 139, 142.

[2] Not to be confused with the greater St. Augustine of Hippo (354–430).

[3] Athelston, Ethelstan, Adelstan, Adelston, Edelstan, and other spellings Born *c.* 895; died 941.

[4] The MS. is in the British Museum (Bib. Reg. 17A, I. p. 32), and was published by J. O. Halliwell, *The Early History of Freemasonry in England* (London, 1840).

Jewish Activity. Probably about the time of Alcuin a Jewish mathematician, Jacob ben Nissim, wrote a work entitled *Sefer Jezira*,[1] which, like various Hebrew writings on mathematics, contains some material on the theory of numbers.

Hrabanus Maurus. Alcuin's most famous pupil was Magnentius Hrabanus Maurus,[2] "Primus praeceptor Germaniae," abbot of the monastery at Fulda (822), and archbishop of Mainz (847). In his younger days he traveled extensively[3] and wrote a worthy treatise on the calendar, based on Bede's work and showing a commendable knowledge of astronomy, a science which included most of the mathematics of his time.[4]

One of his contemporaries, Walafried Strabus[5] (*c.* 806–849), is known to have taught mathematics at Reichenau, near Constance, but he left no works upon the subject.

Remigius of Auxerre. A second great pupil of Alcuin's and a witness to the beneficent influence of the Church in France, was Remi'gius[6] of Auxerre, a Benedictine monk who did much for the schools at Rheims and who founded a school at Paris out of which the university is thought by some to have developed.[7] He wrote a commentary on the arithmetic of Capella,[8] not an important contribution to mathematics, but typical of a period given to useless disputation and empty sophistry.

[1] *Book of Creation.* See M. Steinschneider, "Miscellen zur Gesch. der Math.," in *Bibl. Math.*, III (2), 35; IX (2), 23. An Arabic commentary is known to have been written upon it in the 10th century. The question of the authorship of the *Sefer Jezira* is still unsettled. The work relates chiefly to number mysticism.

[2] Born *c.* 776; died 856. The name appears also as Rabanus Maurus. The date of his birth is also given as *c.* 784.

[3] Ego quidem, cum in locis Sidonis aliquoties demoratus sim." See Neander, *Church History*, III, 457.

[4] He also wrote an encyclopedia, *De universo libri XXII, sive etymologiarum opus.* On his life see J. N. Bach, *Hrabanus Maurus der Schöpfer des deutschen Schulwesens* (Fulda, 1835); D. Türnau, *Rabanus Maurus, der praeceptor Germaniae* (Munich, 1900).

[5] Walafrid Strabo. Cantor, *Geschichte*, I (2), 792.

[6] The name comes from Remy, Remi, *i.e.*, Rheims. Died *c.* 908.

[7] It is also said to have developed from a school of dialectics opened by William of Champeaux, *c.* 1100 to 1110.

[8] The Vatican codex was published in Boncompagni's *Bullettino*, XV, 572.

Hrotsvitha. A certain amount of light is thrown upon the barren field of monastic mathematics of this period by the story of the learned nun Hrotsvitha[1] of the Benedictine abbey of Gandersheim, in Saxony. She wrote several plays and in these she shows a knowledge of the Greek language and of either Greek or Boethian arithmetic. In the *Sapientia* the emperor Hadrian demands the ages of the three daughters of Wisdom (*Sapientia*), namely, of Faith, Hope, and Charity. Wisdom then says that the age of Charity is a defective evenly even number; that of Hope a defective evenly odd one; and that of Faith an oddly even redundant one. Upon Hadrian's remarking, "What a difficult and tangled question has been raised about the mere ages of these girls!" Wisdom replied, "In this is to be praised the great wisdom of the Creator and the marvelous knowledge of the Author of the universe."[2] Hrotsvitha incidentally speaks of three perfect numbers besides 6, namely, 28, 496, and 8128.[3]

Other Writers of the Tenth Century. In the 10th century there may also have been written a treatise on the abacus by

[1] Born *c.* 932; died *c.* 1002. *Hrosvithae Opera*, edited by Winterfeld (Berlin, 1902) (in the *Scriptores rerum Germanicarum*); *Hrotsvithae Opera*, edited by Strecker (Leipzig, 1906); all but one of her works, edited by Conrad Celtes, and with engravings by Dürer, were published at Nürnberg in 1501; there was also an edition by Schurzfleisch (Wittenberg, 1707), and a complete edition by Barack (Nürnberg, 1858). See also Ch. Magnin, *Théatre de Hrotsvitha religieuse Allemande du X^e siècle* (Paris, 1845); E. R. A. Köpke, *Die älteste deutsche Dichterin* (Berlin, 1869), with a refutation of a charge made by Aschbach (1867) that Celtes had forged the works.

The old historian Henricus Bodo referred to her in saying, "Rara avis in Saxonia visa est." The name appears in various other forms such as Roswitha and Hrotsuit. In the Munich MS., apparently contemporary, it appears as Hrotsvith and Hrotsuitha.

[2] A few of the sentences will show the style of the original:

Sapientia. Placetne vobis, O filiae, ut hunc stultum arithmetica fatigem disputatione?

Fides. Placet, mater, . . .

Sapientia. O Imperator, si aetatem inquiris parvularum, Caritas imminutum pariter parem mensurnorum [=annorum] complevit numerum; Spes autem aeque imminutum, sed pariter imparem; Fides vero superfluum impariter parem.

[3] "XXVIII, CCCCXCVI, VIII millia CXXVIII perfecti dicuntur." A perfect number is one that is equal to the sum of its aliquot parts, that is, of its factors and unity; for example, $6 = 1 + 2 + 3$.

Odo of Cluny (879–*c.* 942), although it may be the work of a 12th century writer[1]; but in general the period was a barren one. Only one other writer is worthy of mention, Abbo of Fleury (945–1003), a native of Orléans, who wrote on Easter reckoning,[2] on astronomy, and on the arithmetic of Boethius. His chief title to remembrance, however, is the fact that he was a teacher of Gerbert, the most learned man of his time, whose life and works are considered in the next chapter.

Another example of the ecclesiastical scholar is seen in the case of Bernward, who became Bishop of Hildesheim in 993[3] and who wrote a work on mathematics which was devoted chiefly to the Boethian theory of numbers. A manuscript of this work, possibly the original, is still extant at Hildesheim.

6. The Christian East

Egypt and Constantinople. The eastern countries touching upon the Mediterranean did little for mathematics for a period of five centuries after the fall of Rome. Even the brilliant reign of Justinian (527–565), "the Lawgiver of Civilization," was not able to remove the fears of a barbarian invasion, nor to suppress the disastrous feuds between the Blues and the Greens in Byzantium. Add to this the great fire of 532 and the terrible pestilence of ten years later, and it will be seen that the banks of the Bosporus were not the place for an intellectual revival.

Decay of Alexandria. In Alexandria the chance of progress in the arts and sciences seemed to die out with the fall of Rome, and with the rise of Mohammedanism as a world power the last hope of any revival of the city's ancient glories definitely disappeared. Eighty years after the death of Mohammed his followers had conquered all of northern Africa and had established themselves firmly in Spain. In 642 the great library

[1] S. Günther, *Geschichte der Mathematik*, I, 244 (Leipzig, 1908) (hereafter referred to as Günther, *Geschichte*); Cantor, *Geschichte*, I (3), 843; Th. Martin, "Les Signes Numéraux," *Annali di Mat. pura ed applic.*, V, 50, and reprint, Rome, p. 78 (1864).

[2] *Liber in calculum paschalem.*

[3] H. Düker, *Der liber mathematicalis des Heiligen Bernward.* Hildesheim, 1875.

of Alexandria was destroyed by fire, probably the most serious loss that ever befell any great institution of learning.

Nevertheless a few names appear in the Christian East. Anthemius,[1] an assistant architect in the building of St. Sophia, wrote on conics, and a century later (*c.* 610) Stephen of Alexandria wrote on mathematics and astronomy and taught in Constantinople. In Alexandria, just before the Mohammedan invasion, Asclepias of Tralles (*c.* 635[2]) wrote a commentary on Nicomachus, and Joannes Philop'onus (*c.* 640[3]), known also as Joannes Grammat'icus, did the same and also wrote upon the astrolabe.[4]

Toward the close of the 19th century there was found at Akhmim,[5] in Upper Egypt, a Greek papyrus which seems to have been written about the 7th or 8th century. In this there are tables of unit fractions similar to those found in the Ahmes papyrus, but the work shows no advance over its predecessor of more than two thousand years earlier. Science had long been dead in Egypt except in that part which came under the influence of Alexandria.

School of Cairo. In the early part of the 10th century the Fatimites, a branch of the Mohammedan ruling class, drove their rivals for power out of the city which they thereupon called al-Kâhira, the Victrix,—the modern Cairo. Here they proceeded to establish a school which they ventured to hope would rival that of ancient Alexandria, and which indeed became a center of astronomical activity. With it were connected the names of Ibn Yûnis (p. 175) and al-Ḥaiṯam (p. 175), but it was short-lived, the caliphate of Egypt being destroyed by Saladin in 1171.

[1] Died at Constantinople, 534.

[2] Possibly a century earlier.

[3] The date is very uncertain, being possibly a century too late.

[4] *De vsv astrolabii ejusque constructione libellus.* It was published by H. Hase, *Rheinisches Museum für Philologie*, VI, 127 (Bonn, 1839). His work on Nicomachus was edited by Hoche, Leipzig, 1864, and Wesel, 1867.

[5] Or Ekhmim, the site of the ancient Chemmis or Panopolis. It became a great religious center under the Christians of the early Middle Ages. Nestorius (5th century), the patriarch of Constantinople, was deprived of his honors and banished to Akhmim for heresy. See also Heath, *History*, II, 543.

It is probable that the Jewish scholar Sa'adia ben Joseph[1] studied at Cairo during this period. He wrote on the division of inheritances and on the calendar. He taught in Babylon, where he doubtless met with Isaac ben Salom, who wrote on the Hindu arithmetic and on astronomy.

7. Spain

Oriental Civilization in the West. After the burning of the Alexandrian library (642) the Mohammedans continued their conquests, sweeping along the north coast of Africa and finally entering Spain in 711, defeating the Visigothic king, and establishing themselves for a sojourn of eight hundred years. Bringing with them the Oriental faith in astrology, their primary interest in mathematics was related chiefly to astronomy, trigonometry, and the conics; possessed of esoteric tastes, the mysteries of numbers and of gematria[2] appealed to them; coming into constant relations with the Jews, the cabala doubtless impressed them; inspired by the intellectual brilliancy of Bagdad, the classics of the Greeks found place in their schools. By the time the intellectual supremacy of Bagdad was seriously threatened in the East, Cordova was becoming the intellectual center of Islam in the West. Alhakem II, who reigned from 961 to 976, established a considerable library there, and about the close of the 10th century al-Majrîṭî,[3] a native scholar, wrote on amicable numbers, astronomy, and geometry.

Even in the 10th century the activity in the field of mathematics was not great. The first writer of note was Muslim ibn Aḥmed al-Leiṭî, Abû 'Obeida, also called Sâḥib al-Qible (died 907/8), a native of Cordova and a writer on astronomy and arithmetic. About the same time Cordova produced Salhab ibn 'Abdessalâm al-Faraḍî, Abû'l-'Abbâs (died 922/3), an arithmetician of some note.

[1] In Arabic, Sa'îd ibn Yûsuf al-Fayyumî. He died in 941. The Hebrew name Sa'adia Gaon means Sa'adia the Genius (Great).

[2] Largely concerned with the evaluating of names by the numerical value of the letters.

[3] Abû'l-Qâsim Maslama ibn Aḥmed al-Majrîṭî, died 1007/8.

TOPICS FOR DISCUSSION

1. Intercourse between China and other countries, and its possible influence upon mathematics.

2. Progress of Chinese mathematics from 500 to 1000.

3. Nature and sources of early Japanese mathematics.

4. General nature of Hindu mathematics from 500 to 1000.

5. The work of the two Āryabhatas.

6. Brahmagupta and the School of Ujjain.

7. The work of Mahāvīra compared with that of Brahmagupta.

8. The Bagdad School, its rise and its relation to the Hindu and Greek learning.

9. The nature of the contributions of the Persian and Arab mathematicians of the ninth and tenth centuries.

10. The life and works of Mohammed ibn Mûsâ al-Khowârizmî.

11. Causes of the decay of eastern Arabic mathematics.

12. Indebtedness of medieval Europe to Oriental mathematics in the Middle Ages.

13. Causes of the low state of mathematics in Europe during the greater part of the Middle Ages.

14. Boethius as a mathematician.

15. The life and mathematical works of Bede.

16. The life, influence, and mathematical works of Alcuin.

17. Evidences of an interest in the Greek theory of numbers in the Middle Ages.

18. The influence of mathematics in the Middle Ages upon the science at present.

19. The mathematics of the quadrivium.

20. The nature of the encyclopedias produced by the Church scholars of the Middle Ages.

21. Mathematical recreations in the Middle Ages.

22. Nature of the mathematics studied in the British Isles in the early part of the Middle Ages.

23. The Church schools as preservers of mathematical knowledge in the early part of the Middle Ages.

24. The mathematical contributions of the Mohammedans of the ninth and tenth centuries in Spain.

25. The relation of medieval astrology to astronomy and also to mathematics in general.

CHAPTER VI

THE OCCIDENT FROM 1000 TO 1500

1. Christian Europe from 1000 to 1200

Religious and Political Influences. Just how much influence the passing of the first Christian millennium had upon the common people it is difficult to say. Historians pay much less attention to the "terreur de l'an Mil" than was formerly the case. It is not probable that many educated persons took literally the biblical remark relating to the period of a thousand years, but it is certain that it was so taken by some. At any rate, the passing of this milestone saw the Christian world aroused to new interests.

Then, too, there were the crusades (1095–*c.* 1270), which have been called "the first Renaissance," and which did for a civilization that had long been dormant one thing which the World War did for the civilization of the 20th century,—it let one part of the race know more of what other parts were doing and thinking and hoping. It was war, but it was in general beyond the boundaries of intellectual Europe.

There was also the potent influence in Europe of a foreign and highly developed civilization in her midst,—the Saracen supremacy in Spain; and it was the Saracen scholars who made known to Latin scholars the best of the Greek and Oriental civilizations.

Moreover, Europe was seeing the folly of her private wars, the "Truce of God" was beginning to make its power felt, and the blessings of peace were once more settling upon France and her neighbors, rendering intellectual pursuits possible.

To these influences there should be added that of the Norman Conquest, which, without prolonged warfare, awakened and united England, and showed her what the Continent had for her in the way of science and art.

As a result of such influences Europe entered upon a new era, one in which cathedral building,[1] church reform, renewed attention to art, political experiment, and scientific achievement played great parts.

Gerbert. Nevertheless, the period was still dominated by the spirit of the earlier centuries of the Middle Ages, "when faith overpowered intelligence" and "authority became the enemy of investigation," when "scholars degenerated into schoolmen" and "science lost itself in the morasses of alchemy or astrology and became anathema to the faithful."[2] This is seen in the attitude of the learned world toward that remarkable churchman and scholar, Gerbert,[3] one of the greatest popes that ever added lustre to the Church and to the city of Rome. Elevated to the papal throne, he reigned under the name of Sylvester II from 999 until 1003. He was born of humble parents,[4] but his natural brilliancy led to his call to study under the monks at Aurillac, and particularly under such a worthy scholar as Abbo of Fleury, and to his being sent to Spain (967) to perfect his education.[5] About 970 he went to Italy, where he was presented to the pope and by him to the emperor, returning to

[1] "It was as though the world had arisen and tossed aside the worn-out garments of ancient time, and wished to apparel itself in a white robe of churches." —Raoul Glaber (985–*c.* 1046).

[2] W. C. Abbott, *The Expansion of Europe*, I, chap. i, New York, 1918.

[3] Born near Aurillac, in Auvergne, *c.* 950; died at Rome, May 12, 1003. The name is pronounced zhĕr-bâr.

[4] "Obscuro loco natum," as an old chronicle states.

[5] For bibliography and for a more elaborate sketch, see Smith-Karpinski, p. 110 seq. See also Cantor, *Geschichte*, II, chap. 39; J. Havet, *Lettres de Gerbert* (983–997), Paris, 1889; N. Bubnov, *Gerberti postea Silvestri II papae opera Mathematica*, Berlin, 1899; A. Olleris, *Œuvres de Gerbert*, Paris, 1867; F. Picavet, *Gerbert, un pape philosophe, d'après l'histoire et d'après la légende*, Paris, 1897; H. Weissenborn, *Gerbert. Beiträge zur Kenntnis der Math. des Mittelalters*, Berlin, 1888; C. F. Hock, *Gerberto o sia Silvestro II Papa ed il suo secolo*, Milan, 1846; A. Nagl, *Gerbert und die Rechenkunst des X. Jahrh.*, Vienna, 1888; G. Friedlein, "Die Entwickelung des Rechnens mit Columnen," *Zeitschrift für Mathematik und Physik*, X, Hl. Abt., 241 (hereafter referred to as *Zeitschrift* (Hl. Abt.)), and *Gerbert, die Geometrie des Boethius*, Erlangen, 1861; K. Werner, *Gerbert von Aurillac*, Vienna, 1878; B. Carrara, *Memorie dell' Accad. d. Nuovi Lincei*, XXVI, 195; K. Schultess, *Papst Silvester II. (Gerbert) als Lehrer und Staatsmann*, s. l. a.

France in 972. He held various offices in the Church, and in 999 was elected to the papacy. He was a man of great learning, was "accused—our learning's fate—of wizardry," combated error, aroused new interest in mathematics, acquired a knowledge of the Hindu-Arabic numerals, gave some attention to the study of astrology (a subject then looked upon as a worthy science), and wrote on arithmetic,[1] geometry,[2] and other mathematical subjects, and probably on the astrolabe.[3]

Minor Church Writers. Contemporary with Gerbert, but living a life as humble as Gerbert's was magnificent, was an English monk of the abbey of Ramsey, Byrhtferth[4] by name. He traveled in France and studied under Abbo of Fleury. Returning to England he found waiting for him at Ramsey a group of students to whom he proceeded to teach astronomy, the calendar, and the principles of mathematics.[5] Times were not propitious for study, however. For three centuries in England (1000–1300) there was an average of a famine every fourteen years, and life was hard. Perhaps the need for the conquest of mind over matter, which such calamities set forth, was one of the influences that made possible the later thinkers of England.

On the Continent, St. Gall was one of the chief centers of monastic learning at this time, and here the well-known scholar Notker Labeo[6] (*c.* 950–1022) translated parts of the encyclopedia of Capella and possibly some of the arithmetic of Boethius, besides writing a computus.[7]

[1] *Regulae de numerorum abaci rationibus; Scholium ad Boethii arithmeticam.*

[2] *Gerberti Isagoge Geometriae.* Some doubt has been expressed as to his authorship, but he probably compiled the work.

[3] *Gerberti Liber de astrolabio*, placed by Bubnov with other works among the *Opera Dubia.*

[4] Or Bridferth. Fl. *c.* 1000.

[5] *De temporum ratione, De natura rerum, De indigitatione, De ratione unciarum, De principiis mathematicis*, the extant MSS. being merely notes of his lectures. The Anglo-Saxon text of his *Handboc* was published by F. Kluge in *Anglia*, VIII, 298. See also the Cologne edition (1612) of Bede's works. There are two other works doubtfully attributed to him.

[6] "Notker the Thick-lipped," so called to distinguish him from earlier scholars of the same family.

[7] A. A. Björnbo in the *Reallexikon der Germanischen Altertumskunde*, IV, 465. Strasburg, 1916.

Of the mathematical pupils of Gerbert the most prominent was Bernelinus of Paris, who wrote an arithmetic[1] in which he explained the use of Gerbert's counters, but concerning his life nothing further is known.[2]

A little later (*c.* 1028) Guido of Arezzo (Aretinus), a Benedictine monk from Pomposa, near Ferrara, wrote on arithmetic,[3] and at about the same time (*c.* 1066) Franco of Liège did the same and, what was not so common at this time, wrote on the quadrature of the circle.[4] Among his contemporaries was Wilhelm, abbot of Hirschau (1026–1091), who taught mathematics and astronomy.

Hermannus Contractus. The most prominent of the successors of Gerbert in the 11th century was Hermannus (1013–1054), son of the Swabian Count Wolverad. His limbs having been painfully contracted from childhood, he is known in history as Hermannus Contractus.[5] Educated in the monastic school at Reichenau, he afterwards joined the Benedictine order, became a lecturer on mathematics, and gathered about him a large number of pupils. He wrote on the astrolabe,[6] the abacus, and the number game of rithmomachia.[7]

Psellus. The period of intellectual activity in the West had very little counterpart in Constantinople. Life was still stagnant there. In the 11th century only a single name stands out as representing any interest whatever in mathematics in the eastern capital,—that of Michael Constantine Psellus[8] (1020–1110), a Greek writer who studied at Athens, became a zealous Neoplatonist, and returned to Constantinople to teach philosophy.

[1] *Liber Abaci.* [2] Gerbert's *Œuvres*, ed. Olleris, p. 357 (Paris, 1867).

[3] B. Baldi, Boncompagni's *Bullettino*, XIX, 590.

[4] *Abhandlungen*, IV, 135.

[5] Treutlein, Boncompagni's *Bullettino*, X, 589, where his *Abacus* is published; Günther, *Math. Unterrichts*, p. 47; Baldi, *Cronica*, p. 70. He is also known as Hermann the Lame.

[6] There is a beautifully written MS. of this work, 12th century, in the British Museum (22,790), first published by Pez in Volume III of his *Thesaurus Anecdotorum* and republished by Migne in Volume CXLIII of *Patrologiae cursus completus*. [7] See page 198.

[8] Ψέλλος, called also Psellus the Younger, there having been another Psellus who taught philosophy *c.* 870. Heath, *History*, II, 545.

He lived during the reigns of several rulers, consulted by the emperors and honored by them with the title of Prince of Philosophers.[1] An introduction to the study of Nicomachus and Euclid is attributed to him, but the authorship is doubtful. Partly because of the fact that he was almost the last of the Greek writers on mathematics, partly because his works were easily read, and partly because of his reputation for learning in general, he is one of the few scholars of his time whose mathematical contributions attracted any attention in the Renaissance period. His leading works on mathematics[2] were published at least thirteen times in the 16th century. The fact that he takes $\sqrt{8}$ as the value of π shows how little he merited his reputation as a scientist.

Rithmomachia. In speaking of the 11th century mention should be made of the number game of rithmomachia.[3] One of the earliest treatises on the subject is due to Fortolfus, a monk, who lived probably at the close of the 11th century,[4] and the indications are that it was not known before that century, although it is occasionally attributed to Boethius and even to Pythagoras. There is a manuscript in the Vatican library on the subject, under the title "Ritmachya," written in 1077 by a monk known as Benedictus Accolytus, and the game is also referred to in a medieval poem *De Vetula*.[5] Among the early

[1] Φιλοσόφων ὕπατος.

[2] *Sapientissimi Pselli opus dilucidum in quattuor Mathematicas disciplinas, Arithmeticam, Musicam, Geometriam, & Astronomiam*, edited by Archbishop Arsenius, Venice, 1532. This was the first edition, the text in Greek. The *Compendium Mathematicum*, containing various works, appeared at Leyden in 1647, but numerous others still remain unedited.


[3] Literally, "combat of numbers." The word is spelled in various ways, rithmimachia, ritmachya, richomachie, and rhythmimachia being among the most common forms.

A work on the subject by Boissière, a French mathematician of the 16th century, is entitled *Nobilissimvs et antiqvissimus ludus Pythagoreus (qui Rythmomachia nominatur)*, Paris, 1556. See *Rara Arithmetica*, pp. 12, 63, 271, 340.

[4] R. Peiper, "Fortolfi Rythmimachia," *Abhandlungen*, III, 167, 198.

[5] "O utinam ludus sciretur Rythmimachiae!
ludus Arithmeticae folium, flos fructus et eius
gloria laus et honor."
Abhandlungen, III, 222

writers who were interested in the subject were Hermannus Contractus (1013–1054), as already stated, and both Jordanus Nemorarius (died *c.* 1236) and Nicole Oresme (*c.* 1323–1382).



RITHMOMACHIA

From a work published at Paris in 1496. The middle portion of the board is omitted. The part on Rithmomachia may be due to Bishop Shirwood of Durham (died 1494), but is usually ascribed to Faber Stapulensis (1455–1536)

The game is based on the Greek theory of numbers as set forth by Nicomachus. It was played upon a double chess-board, rectangular in form, one side having eight squares and

the other sixteen. The pieces were triangles, squares, circles, and pyramids, each possessing a certain value. These pieces were arranged as shown in the illustration (p. 199) from a work of 1496. The numbers were not taken at random, but the plan on which they were arranged is too elaborate for description in this work. Suffice it to say that when we form the triangles we have $81 = 72 + \frac{1}{8}$ of 72, $72 = 64 + \frac{1}{8}$ of 64, $6 = 4 + \frac{1}{2}$ of 4, and $9 = 6 + \frac{1}{2}$ of 6; that in the case of the square pieces, $45 = 25 + 20$ and $15 = 9 + 6$; that the pyramids are superposed squares such that $91 = 6^2 + 5^2 + 4^2 + 3^2 + 2^2 + 1^2$ and $190 = 8^2 + 7^2 + 6^2 + 5^2 + 4^2$. In the case of the squares there is a formula $s = \left(\frac{2n+1}{n+1}\right) s'$, in which the meaning of each letter may be found by looking at the illustration, where $s = 25$, $s' = 15$, $n = 2$; or $s = 81$, $s' = 45$, $n = 4$. The play is very complicated, and for our purposes we may say that the climax of the game was reached in the *Victoria praestantissima*, in which it was necessary to get four numbers in a row, embodying all three of the common progressions,—arithmetic, geometric, and harmonic, the only possible solutions with these pieces being six in number. It will be seen that the game requires such familiarity with the Greek number theory as to make it available only for the élite in mathematics in the Middle Ages. Its popularity is attested by at least three manuscripts of the 11th century and three of the 12th and 13th centuries, besides several printed treatises on the subject, all going to show that there were more scholars in number theory than we should think from the meager list of names that have come down to us.[1]

A Century of Translators. The 12th century was to Christian Europe what the 9th century was to the eastern Mohammedan world, a period of translations. In the case of Bagdad, these translations were from the Greek into Arabic; in the case of

[1] For a description of the game, see D. E. Smith, "Number Games and Number Rhymes," in *Teachers College Record*, XIII (New York), 385, together with a history of "The Great Number Game of Dice." The article on rithmomachia may also be found in the *Amer. Math. Month.*, April, 1911. See also E. Wappler, *Zeitschrift* (Hl. Abt.), XXXVII, 1.

Christian Europe, from the Arabic into Latin. The reasons for this desire to know the science of the East are not difficult to find. The causes already mentioned in connection with the 11th century were even more potent a hundred years later, and the advancement of Moorish Spain in the arts and sciences was already causing intellectual unrest in the higher class of Church schools in France, Italy, and England. The result of this unrest was an influx of students into Spain, an acquiring of some knowledge of Arabic on the part of various scholars, and a strong desire to know and to make known the science of the East. Just as Bagdad never translated the Greek literature, but sought diligently to know Greek science, so Europe gave little attention to Arab letters, but devoted great care to those works on astronomy, arithmetic, trigonometry, optics, astrology, geometry, and medicine that had acquired reputation in the capital of the caliphs. Even the *Elements* of Euclid became known to the scholars of the Latin Church chiefly through its Arabic translation instead of through the original Greek.

Italian and French Translators. In the 12th century Italy and France produced two or three prominent scholars whose knowledge of Arabic and taste for mathematics led them to make known to the Latin world various classics of the Mohammedan and Greek civilizations.

The first of these translators was Plato of Tivoli, or Plato Tiburtinus,[1] who lived *c.* 1120. He translated the astronomy of Albategnius (al-Battânî), the *Spherics* of Theodosius, the *Liber Embadorum* of Abraham bar Chiia (*c.* 1120), and various works on astrology.

About this time Sicily was also active in the translation of Greek and Arabic works.[2] Among the treatises thus brought to the attention of scholars was Ptolemy's *Almagest*, which was turned into Latin by an unknown translator, *c.* 1160,[3]

[1] B. Boncompagni, *Delle versioni fatte da Platone Tiburtino*, Rome, 1851.

[2] C. H. Haskins and D. P. Lockwood, "The Sicilian Translators of the Twelfth Century . . .," in the *Harvard Studies in Classical Philology*, XXI, 75.

[3] There is in the Vatican a MS. of this translation, written *c.* 1300. It is this that was used by Professors Haskins and Lockwood in the work above cited.

from a Greek manuscript which had formerly been brought from Constantinople to Palermo by a Sicilian scholar.

Some years later Gherardo Cremonense, or Gherardo of Cremona (1114–1187),[1] studied in Italy and then in Spain, learning Arabic in Toledo. With him, as with many other scientists in the Middle Ages and even later, astrology formed a nexus joining medicine and mathematics, his interests therefore lying in all three lines. He translated various mathematical and astronomical works from the Arabic, including Euclid's *Elements* and *Data*, the *Spherics* of Theodosius, a work by Menelaus, and Ptolemy's *Almagest*,[2] "for the love" of which book he journeyed to Toledo.[3] In his translation is found one of the early uses of the word *sinus* for a half chord, this being the first of our modern names for the trigonometric functions.[4] There was a younger Gherardo of Cremona who lived in the 13th century, called da Sabbionetta, who wrote on astronomy.[5]

Among the Italian and French translators there may properly be included Rudolph of Bruges, since most of his work was done under French influence. About his time (*c.* 1143) Hermann of Carinthia translated Ptolemy's *Planisphere*.[6]

[1] Apparently a native of Cremona in Lombardy, although certain Spanish writers have claimed him for Carmona in Andalusia. The name appears in English as Gerard and in Latin as Girardus, with variants. B. Boncompagni, *Della vita e delle opere di Gherardo Cremonense*, Rome, 1851. A considerable amount of information relating to such early Italian mathematicians is given in B. Veratti, *De' Matematici Italiani anteriori all' Invenzione della Stampa*, a pamphlet with bibliographical notes, Modena, 1860.

[2] The translation was finished in 1175, as an old MS. asserts. This was about fifteen years after the Sicilian translation, a work of which Gherardo was apparently ignorant. See also Rose, in *Hermes*, VIII, 332. It was printed in Venice in 1515. On the question of his translations see A. A. Björnbo, *Bibl. Math.*, VI (3), 239.

[3] Amore tamen almagesti, quem apud latinos minime reperiit, Toletum perrexit."

[4] On the question of priority and of the use of the term by Plato of Tivoli, see A. Braunmühl, *Geschichte der Trigonometrie*, I, 49 (Leipzig, 1900, 1903); hereafter referred to as Braunmühl, *Geschichte*. The term was probably first used in Robert of Chester's revision of the tables of al-Khowârizmî. See also *Bibl. Math.*, I (3), 521.

[5] His *Theorica planetarum* was printed at Ferrara in 1472.

[6] This was printed in 1507. See M. Chasles, *Aperçu historique sur l'origine et développement des méthodes en géométrie*, Paris, 1837; 2d ed., 1875, hereafter referred to as Chasles, *Aperçu*; 3d ed., 1889. See also *Bibl. Math.*, IV (3), 130.

English Translators. England produced two or more translators of prominence in the 12th century, and Ireland seems to have produced at least one. Of these the best known is Adelard[1] of Bath (*c.* 1120), a British scholar who studied at Toledo (1130), at Tours, at Laon, and also in the East, and who journeyed through Greece, Asia Minor, Egypt, and possibly Arabia, bringing back numerous mathematical works.[2] He is credited with a knowledge of Greek and was one of the first to translate Euclid into Latin, but he seems to have made this translation from the Arabic.[3] Either he or Campanus seems to have determined the sum of the angles of a stellar polygon, a figure then attracting considerable interest, possibly because of its use in astrology. He probably translated the astronomical tables of al-Khowârizmî, and he is said to have written a commentary on the arithmetic of this author and to have composed a work entitled *Regulae abaci.*[4] Adelard was by no means the first to bring Euclid's name into England, for, as we have seen (p. 187), it was probably known to British scholars in the 10th century.

A few years after Adelard's sojourn in Toledo two other English scholars who were interested in mathematics went to Spain to pursue their studies. The first of these was Robert of Chester (*c.* 1140),[5] who translated al-Khowârizmî's algebra[6] into Latin and prepared several astronomical tables. He was archdeacon of Pampeluna, in northern Spain, and seems also to have traveled in Italy and Greece. He was the first to translate the Koran into Latin (1143).

[1] The older English form was Aethelhard. See C. H. Haskins, "Adelard of Bath," *English Historical Review*, p. 491 (1911).

[2] F. Woepcke, *Journal Asiatique*, I (6), 518.

[3] From certain similarities in the different manuscripts there seems to have been an unknown scholar whose version was consulted by Adelard and various other early translators.

[4] Boncompagni's *Bullettino*, XIV, 1.

[5] Robertus Retinensis, Robertus Ketensis, Robert de Ketene, Robert de Retines, Robertus Cataneus, and other variants. He is known to have been in Spain in 1141 and seems to have been studying at Barcelona with Plato of Tivoli in 1136. As already stated, he translated the astronomy of Albategnius.

[6] For reference to the translation see page 170, note 3.

The second of these English scholars was Daniel Morley,[1] who studied at Oxford in 1180. He went to Paris and thence to Toledo,[2] and wrote on astronomy and mathematics,[3] quoting freely from Arabic authors. That such men were compelled to go abroad for their mathematics at this time is apparent from the records of the work done in the schools of London, this work being chiefly of the nature of grammar and disputation.[4] That they should go to Spain was quite natural, not merely for linguistic reasons but because of the close ties that existed between Castile and England, owing to the marriage of Alfonso VIII (1158–1214) to Lenora, daughter of Henry II.

Other Scholars. One of Adelard's pupils, N. O'Creat,[5] wrote a work on multiplication and division which shows his indebtedness to Arab writers on mathematics. Of O'Creat himself, however, nothing further is known, but the name suggests the country of his birth. The work contains a rule of Nicomachus for squaring a number by using the formula $a^2=(a-b)(a+b)+b^2$, thus: $109^2 = 100 \cdot 118 + 81 = 11{,}881$. He used the Roman numerals, but with both o and a character like the Greek τ for zero.[6]

[1] Daniel of Merlai, Merlac, Marlach. In Latin, Morleius, Merlacus. A MS. in the British Museum (J. O. Halliwell, *Rara Mathematica*, London, 1838–1839, 2d ed., 1841, p. 84; hereafter referred to as Halliwell, *Rara Math.*) begins, "Philosophia magistri Danielis de Merlai ad Johannem Norwicansem episcopum." See also C. Singer in *Isis*, III (1920), 263.

[2] See A. à Wood, *Historia et Antiquitates Vniversitatis Oxoniensis*, I, 56 (Oxford, 1674); hereafter referred to as Wood, *Historia Oxon.*

[3] Probably *De principiis mathematicis.*

[4] See the *Descriptio nobilissimae civitatis Londoniae* written by Fitzstephen (died *c.* 1190), prefixed to his *Life of Becket*, and published in John Leland's *Itinerary* (London, 1770); J. Stow, *Survey of London*, p. 703 (London, 1633).

[5] Probably the same as Joh. Ocreatus. See N. Bubnov, *Gerberti . . . Opera Mathematica*, p. 174, n. 7 (Berlin, 1899); C. Henry, *Abhandlungen*, III, 129. O'Creat begins his work with these words: "N. O. Creati liber de multiplicatione et divisione numerorum ad Adelardum Bathoniensem magistrum suum." There is a 13th century MS. of the work in the Bibliot. nat. in Paris. The prologue begins, "Prologus N. Ocreati in Helceph, ad Adelardum Batensem magistrum suum." On the uncertain meaning of Helceph, see Henry, *loc. cit.*

[6] Possibly from the medieval τζίφρα, from theca (teca), or from τήκειν (to come to naught).

About the year 1125 Radulph of Laon (died 1133) wrote on arithmetic, and a little before this time (*c.* 1090) Gerland, prior of St. Paul, of Besançon,[1] wrote a computus and a brief work on the abacus.[2]

Early in the 12th century there was an astrologer, geometrician, and abacist by the name of Walcherus, a native of Lorraine, who attained considerable prominence in England and wrote a work on astronomy.[3] Such names are of interest simply as they bear witness to the nature of mathematics in the Church schools of the time.

2. Oriental Civilization in the West

Spain. After the year 1000 numerous Moorish scholars appeared in Spain and contributed to the literature of arithmetic and astronomy, and occasionally to that of algebra. A list of a few of the most prominent of these scholars will serve to show their range of interest and achievement in the general field of mathematics.

Ibn al-Ṣaffâr,[4] a native of Cordova, wrote on astronomical tables and instruments. A little later (*c.* 1050) Ibn al-Zarqâla,[5] probably a native of Cordova, wrote on astronomy and astrology, and prepared a set of tables.[6]

[1] An ancient record speaks of "Gerlandus vel Garlandus Prior S. Pauli, annis, 1131, 1132." A document of 1134 records: "Huius praefatae concordiae testes sunt . . . Garlandus magister . . . anno . . . M.C.XXXIIII." He is also mentioned in 1148: "Magistrum quoque Jarlandum Bisuntinum & magistrum Theodericum Carnotensem [i.e., of Chartres], duoa fama & gloria doctores nostri temporis excellentissimos." He is again mentioned in a letter written to him in 1157: "Gerlando scientia trivii, quadriviique onerato & honorato." See Boncompagni's *Bullettino*, X, 654.

[2] See Boncompagni's *Bullettino*, X, 653; Cantor, *Geschichte*, II (2), 843.

[3] In the south aisle of the chancel of the old priory at Great Malvern may still be seen his tomb with this inscription, in part: Philosophvs · dignvs · bonvs · astrologvs · Lotheringvs | vir · pivs · ac · hvmilis · monachvs · prior · hvjvs . . . | . . . geometricvs · ac · abacista :— | Doctor · Walchervs . . . MCXXV · ✠ He should not be confused with the Walcherus of Lorraine who became Bishop of Durham and was murdered in 1075.

[4] Aḥmed ibn 'Abdallâh ibn 'Omar al-Gâfiqî, Abû'l-Qâsim (died 1035).

[5] Ibrâhîm ibn Yaḥyâ al-Naqqâsh, Abû Isḥâq.

[6] Schoner translated one of his works in 1534.

In the latter part of the century Abû'l-Ṣalt,[1] a Spanish physician from Denia, wrote on geometry and astronomy, and Jabir ibn Aflah (died between 1140 and 1150), commonly known as Geber, flourished at Seville and wrote on astronomy, spherical trigonometry, and the transversal theorem of Menelaus.[2] He is often confounded with an alchemist of similar name.

Jewish Scholars of the Eleventh Century. The most learned scholars in Spain at the close of the 11th century, however, were not Mohammedans. The Jewish race, which may conveniently be mentioned in connection with the Oriental civilization in Spain, was generally accorded better treatment under Saracen than under Christian rule, although it had flourished somewhat in Italy before this time. Through the encouragement received from the Moors the Jews contributed in no small degree to the advance of mathematics in Spain, and to them the Christians were indebted for their first knowledge of the Arabic works on the subject.[3] The first of their prominent scholars in this century was Abraham bar Chiia[4] (Abraham Judæus), commonly known as Savasor'da[5] (*c.* 1070–*c.* 1136), a native of Barcelona. He wrote on astronomy, but is chiefly known for an encyclopedia which included arithmetic, geometry, and mathematical geography.[6] Of this only fragments are now extant. He also wrote a work entitled *Liber Embadorum*,[7] treating of geometry but containing numerous definitions used in the theory of numbers. In this he accuses the French

[1] Omeiya ibn 'Abdel'azîz ibn Abî'l-Ṣalt, Abû'l Ṣalt (1067/8–1133/4).

[2] His astronomy was translated by Gherardo of Cremona and was printed in 1534.

[3] Libri, *Histoire*, I, 154 n.

[4] Or Chijja, Chiya.

[5] From *Sahib al-Shorta*, "Chief of the Guards." The transliteration to Savasorda is due to Plato of Tivoli (*c.* 1120).

[6] *Iesode ha-Tebuna u-Migdal ha-Emuna.*

[7] This is one of the sources of Fibonacci's geometry. See M. Curtze, *Abhandlungen* XII, where the Latin and German translations are given. The title is medieval Latin from the Greek ἐμβαδόν, an area or surface. For his works and for the contemporary Jewish writers see J. Bensaude, *L'Astronomie Nautique au Portugal*, p. 52 (Bern, 1912); hereafter referred to as Bensaude, *Astron. Portug.* See *Bibl. Math.*, 1896, p. 36.

Jews of being ignorant of geometry and therefore weak in their arithmetic. This work was translated from Hebrew into Latin by Plato of Tivoli.

Rabbi ben Ezra. The second great Hebrew scholar of the period was Abraham ben Ezra.[1] He wrote on the theory of numbers, the calendar, magic squares, astronomy, and the astrolabe, was much interested in the cabala, and is justly ranked as the most learned Jew of his time.[2] He traveled extensively, going at least as far as Egypt to the east and as far as London (1158) to the north. Besides his contributions to astronomy, the calendar, and allied subjects, he wrote three or four works on number: (1) *Sefer ha-Echad*;[3] (2) *Sefer ha-Mispar*,[4] chiefly on arithmetic; (3) *Liber augmenti et diminutionis vocatus numeratio divinationis*, known only in Latin translation and possibly not due to him;[5] (4) *Ta 'hbula*, containing the Josephus Problem, possibly a separate work, and probably due to him. Of these the *Sefer ha-Mispar* is the only one of importance. It is based on the Hindu arithmetic but uses Hebrew letters for the numerals, with a zero as in algorism. He employed the check of casting out nines, as several of his predecessors had done. The following is an example of his rules: "Whoever would know how great the sum of the numbers is

[1] Born at Toledo, between 1093 and 1096; died at Rome or Rouen, 1167. This is the Rabbi ben Ezra of Browning's poem. He is sometimes confused with Abraham ben Chiia, probably because each was called Abraham Judæus. On his life see M. Steinschneider, *Bibl. Math.*, IX (2), 43; *Abhandlungen*, III, 57; Bensaude, *loc. cit.*, p. 52.

[2] Attention was called to him as a mathematician by O. Terquem, *Journal des mathématiques pures et appl.*, VI, 275. Since then his work has been studied by Luzzato, Rodet, and Steinschneider. See also Smith and Ginsburg in the *Amer. Math. Month.*, XXV, 99.

[3] *Book of Unity*. This has twice been published: Bamberg, 1856; Odessa, 1867.

[4] *Book of Number*. M. Silberberg published a German translation at Frankfort a. M. in 1895. See also *Bibl. Math.*, IX (2), 91.

[5] In favor of his authorship, Cantor, *Geschichte*, I (3), 730; against it, G. Wertheim, *Bibl. Math.*, II (3), 143. See also P. Tannery, *Bibl. Math.*, II (3), 45. The *Liber augmenti . . . divinationis* was published by Libri. His *Liber de nativitatibus* (Venice, 1485) was, however, the first of his works to appear from the press.

which follow one another in a series to a certain number, multiply this by its half increased by $\frac{1}{2}$. The product is the sum."[1]

Although highly esteemed by Jews and Christians alike, his fate was not altogether a happy one, and in his struggle against adversity he voices his lament in words like these:

> Were candles my trade it would always be noon;
> Were I dealing in shrouds Death would leave us alone.

In connection with the Jewish activity of this period there should also be recalled the name of one Hasan, a judge, who may have written in the 10th century, but whose country is unknown, and of Yehuda ben Rakufial, who seems to have been a physician in Spain. Both of these men wrote on the Jewish calendar, and the former is referred to by Rabbi ben Ezra.

Twelfth Century in Spain. The 12th century was even more favorable than its predecessors to the study of mathematics in Spain. The first of the Arab writers was Aver'roës (*c.* 1126–1198/9), as he was commonly called in the Middle Ages,[2] who wrote on astronomy and trigonometry. His most prominent scientific contemporary was Avenpace, as he was called by the Christians,[3] who lived at Seville and Granada *c.* 1140 and wrote on geometry.

Jewish Writers of the Twelfth Century. In this as in the preceding century, however, it was the Hebrew scholar who made the greatest contributions to the advance of mathematics. Aside from Rabbi ben Ezra, two of these scholars are deserving of special mention: Maimonides[4] (1135–1204), a native of Cordova, physician to the sultan, and an astronomer of

[1] That is, $s = n\left(\frac{n}{2} + \frac{1}{2}\right)$. *Sefer ha-Mispar*, ed. Silberberg, p. 24 (Frankfort a. M., 1895).

[2] His name was Mohammed ibn Aḥmed ibn Mohammed ibn Roshd, Abû Velîd.

[3] His name was Mohammed ibn Yaḥyâ ibn al-Ṣâig, Abû Bekr, also known as Ibn Bâjje and as Ibn Ṣâig. The name "Avenpace" (also spelled Avempace) is a Spanish form and, as such, is pronounced ah vän pä'thā.

[4] Rabbi Moses ben Maimun. He became rabbi of Cairo in 1177.

prominence,[1] and Johannes Hispalensis[2] (fl. *c.* 1140), who professed Christianity and wrote on arithmetic and astrology (1142) and translated various Arabic works on mathematics into Latin.[3]

In the same century there were various Jewish scholars of less prominence, such as Samuel ben Abbas,[4] who wrote on arithmetic,[5] the Hindu numerals and their use,[6] algebra, and geometry. There was also an unknown English Jew who wrote a work called by English historians *Mathematum Rudimenta quaedam.*

Jewish Writers of the Thirteenth Century. The 13th century saw various translations made from the Arabic into Hebrew, and several of the translators are known. Among these was Moses ben Tibbon,[7] whose father and grandfather were celebrated as translators of philosophical and scientific works from the Arabic into Hebrew. He was actively at work about the middle of the century and translated (1259) the astronomy[8] of Alpetra'gius (*c.* 1200) and probably, as stated on page 210, the arithmetic of al-Ḥaṣṣar.

The other Jewish scholars of this period also showed their chief scientific interest in astronomy. Jehuda ben Salomon Kohen of Toledo (died 1247), for example, wrote upon Ptolemy's *Almagest*, although he also prepared a brief extract from Euclid

[1] There is a Jewish calendar of his among the manuscripts in the Bodleian Library. Parts of his works on the calendar were printed at Paris in 1849, at Leipzig in 1859, and at Berlin in 1881.

[2] John of Seville, John of Luna. As in many such cases, the first name is often written Joannes. The full name is also written Johannes Hispanensis or Johannes de Hispania. The date of his death may have been 1153.

[3] At least some were translated into Spanish and were then put into Latin by Domenico Gondisalvi. His *Alghoarismi de Practica Arismetrice* was published by B. Boncompagni, Rome, 1857. It is based on Arab sources, but is not a translation. His translations include works by Alfraganus (al-Fargânî, *c.* 833), Abû 'Alî al-Chaiyât (a prominent astrologer, died 835), and Ṯâbit ibn Qorra (*c.* 875). His works were published at Nürnberg in 1548.

[4] M. Steinschneider, *Bibl. Math.*, X (2), 81. He adopted the Mohammedan faith, his Arabic name being Samû'îl ibn Yaḥyâ ibn 'Abbâs al-Maġrebî al-Andalusî. He died in 1174/5.

[5] *Al-Tab'sira.*

[6] *Al-Qiwâmî*, probably named after a patron, Qiwâm ed-din Yaḥyâ.

[7] *Bibl. Math.*, X (2), 112.

[8] *Kitâb al-hei'a.*

and wrote a commentary upon it, and Isaac ben Sid, of Toledo (died 1256), edited the Alfonsine Tables (see page 228) just before his death.

About the middle of the 13th century there was born in Cordova another descendant of the celebrated Tibbon family,—Jacob ben Machir, known as Prophatius. He lived in Montpellier, wrote on a quadrant which he had invented (the *quadrans Israelis* or *quadrans Judaicus*),[1] translated from the Arabic into Hebrew the *Elements* and *Data* of Euclid and the *Sphere* of Menelaus,[2] and composed a work on the almanac.

Arab Writers of the Twelfth Century. Of the writers on arithmetic among the western Arabs of the 12th century one of the best known was Abû Bekr Mohammed ibn ʿAbdallâh,[3] commonly known as al-Ḥaṣṣâr.[4] His work was so well received that, as already stated, it was translated into Hebrew by Moses ben Tibbon[5] (1259). The work is evidently Western, since it uses the ġobâr numerals.

Arab Writers of the Thirteenth Century and Later. Early in the 13th century Alpetragius,[6] as the Christians called him, lived in Spain, probably in Seville, and wrote on astronomy (*c.* 1200). His theory of planetary motion, which gives him a place in the list of mathematical writers, was translated into Latin by Michael Scott.

Contemporary with Alpetragius there was a certain Ibn al-Kâtib[7] (died 1210/11), who wrote two works which included a little discussion of arithmetic, geometry, and architecture.

[1] There is a good MS. of the work in the Columbia University Library, apparently of the 15th century. The work has been several times translated into Latin.

[2] Boncompagni's *Bullettino*, IX, 595; *Bibl. Math.*, XI (2), 35. The name also appears as Propatius. He died *c.* 1308.

[3] A Gotha MS. gives the name as Abû Zakarîyâ Mohammed ibn ʿAbdallâh ibn ʿAiyâsh. See *Bibl. Math.*, XIII (2), 87.

[4] That is, the Computer; but Suter thinks this a family name. See H. Suter, "Das Rechenbuch des Abû Zakarîjâ el-Ḥaṣṣâr," in *Bibl. Math.*, II, (3), 12, and III (2), 109. See also *ibid.*, XIII (2), 87.

[5] Probably. The Vatican MS. has still to be studied critically.

[6] Nûr ed-dîn al-Betrûjî, Abû Isḥâq.

[7] Mohammed ibn ʿAbderraḥmân, Abû ʿAbdallâh.

Of the scholars born in northern Africa in the 13th century, and geographically closely related to the Spanish civilization, the best known is Albanna,, or Ibn al-Banna.[1] From the fact that he is also known as al-Marrâkushî we infer that he was a native of Morocco.[2] He wrote on astronomy, mensuration, algebra, the astrolabe, and proportion. His best-known work is the *Talchîs*, a treatise on arithmetic.[3]

There was also a Mohammedan scholar of Seville, known as Ibn Bedr[4] or Abenbêder, who wrote a compendium of algebra about this time.[5] The date is uncertain, but there is a commentary upon it in verse which was written in 1311/12.

The last of the great Moorish arithmeticians of Spain was al-Qalasâdî,[6] a native of Baza, a town near Granada. He wrote extensively on arithmetic and seems to have had some originality in the treatment of the theory of numbers. He introduced a new radical sign and a sign of equality, and proposed a system of ascending continued fractions.[7]

3. Christian Europe from 1200 to 1300

General Activity of the Thirteenth Century. Whatever may be thought of the mathematics of the 13th century, it is certain that the century itself represents the real awakening of the world after a long period of intellectual torpor. The centuries

[1] That is, Son of the Architect. His full name is Aḥmed ibn Mohammed ibn 'Oṯmân al-Azdî, Abû'l-'Abbâs. Born *c.* 1258; died in Morocco *c.* 1339.

[2] A. Marre, *Atti dell' Accademia Pontificia dei Nuovi Lincei*, XIX (hereafter referred to as *Atti Pontif.*); M. Steinschneider, Boncompagni's *Bullettino*, X, 313. Suter's list omits the occasional name al-Marrâkushî, and places the date of his birth as *c.* 1258 or later, although it is sometimes given as early as 1252. The father seems to have belonged to a Granada family.

[3] Discussed in Cantor, *Geschichte*, I (3), 806.

[4] Mohammed ibn 'Omar, Abû 'Abdallâh.

[5] José A. Sánchez Pérez, *Compendio de Álgebra de Abenbéder*, Arabic text and Spanish translation, Madrid, 1916.

[6] Ali ibn Mohammed ibn Mohammed ibn 'Alî al-Qoreshî al-Basṭî, Abû'l-Ḥasan. The name "al-Qalasâdî" means the Upright, or Versed in the Law. Suter gives the place and date of his death as Tunis, 1486.

[7] Woepcke in the *Journal Asiatique*, 1854, II, 358, and 1863, I, 58. See also the *Atti Pontif.*, XII, 230, 399.

immediately preceding had produced writers on mathematics in Europe, but they had produced no mathematicians. But now a Spirit of the Times was abroad. The Far East felt its influence, and hence the remarkable revival and development of algebra in China; India felt it, and hence the appreciation of the merit of Bhāskara, now a generation dead; and all of intellectual Europe felt it as never before. It was not a century of great beacon lights, but it was one in which lanterns were hung in all the thoroughfares of the West, promises of the great illumination that was to come with the period of the Renaissance.[1]

Rise of the Universities. The most potent influence in the development of the world's mathematical knowledge has, of course, been the universities, and it is from the 13th century that we trace the rise of these institutions in the modern sense of the term. The earliest medieval universities grew out of the cathedral or Church schools and hence their date of beginning is necessarily obscure. In most cases the years in which they received official privileges from some sovereign, civil or ecclesiastical, are known, however, and are commonly taken as the dates of foundation. In some cases there are two dates, one of the receipt of the privilege from the State and the other that from the Church, the latter giving to the holders of degrees a right to teach. Thus Paris had a charter from the State in 1200 and its degrees were recognized by the pope in 1283. The corresponding dates for Oxford were 1214 and 1296; and for Cambridge, 1231 and 1318. The University of Padua was founded in 1222, and that of Naples in 1224.[2] The 14th and 15th centuries saw a number of other universities established, but we may look upon the 13th century as the one which laid the foundation for this type of higher education, although the mathematics taught was still very meager.[3]

[1] J. J. Walsh, *The Thirteenth, Greatest of Centuries*, New York, 1907.

[2] Some of these dates are uncertain, but they are approximately as stated.

[3] H. Suter, "Die Mathematik auf den Universitäten des Mittelalters," *Festschrift der Kantonschule in Zürich* (Zürich, 1887), p. 39; hereafter referred to as Suter, *Univ. Mittelalt.*



THE TOWER OF KNOWLEDGE

Illustrating the educational system of the Middle Ages. From the *Margarita phylosophica*, 1503

Medieval Curriculum. The student began his study of grammar with Donatus and Priscian, and took his logic from Aristotle and his rhetoric from Cicero. He then entered upon his mathematical studies, such as they were,—arithmetic according to Boethius, music according to Pythagoras, geometry according to Euclid, and astronomy according to Ptolemy. The goal for those who were preparing for church activities was the metaphysics and theology of Peter Lombard (*c.* 1150). This progress was illustrated in a tower of knowledge given by Gregorius Reisch in his *Margarita phylosophica* (1503), as shown on page 213.



LEONARDO FIBONACCI

Modern engraving. The portrait is not based on authentic sources

Leonardo Fibonacci. The first great mathematician of the 13th century, and indeed the greatest and most productive mathematician of all the Middle Ages, was Leonardo Fibonacci, known also as Leonardo Pisano or Leonardo of Pisa.[1]

[1] Born at Pisa, *c.* 1170; died *c.* 1250. On his life and works see B. Boncompagni, *Scritti di Leonardo Pisano*, 2 vols., Rome, 1857–1862 (hereafter referred to as Boncompagni, *Scritti Fibonacci*); *Della vita e delle opere di Leonardo Pisano*, Rome, 1852; *Intorno ad alcune opere di Leonardo Pisano*, Rome, 1854; and *Tre scritti inediti di Leonardo Pisano*, Florence, 1854 (hereafter referred to as Boncompagni, *Tre Scritti*); Cantor, *Geschichte*, II, chaps. xli, xlii; Libri, *Histoire*, I, 156; E. Lucas, "Recherches sur plusieurs ouvrages de Léonard de Pise et sur diverses questions d'arithmétique supérieure," Boncompagni's *Bullettino*, X, 129; G. Loria, "Leonardo Fibonacci," *Gli Scienziati Italiani* (Rome, 1919), p. 4, with excellent bibliography; G. B. Guglielmini, *Elogio di Lionardo Pisano*, Bologna, 1813; F. Bonaini, *Memoria unica sincrona di Leonardo Fibonacci*, Pisa, 1858 (republished in 1867), and also in the *Giornale Arcadico*, Vol. CXCVII (N. S., LII), and in an article by G. Milanesi, *Documento inedito e sconosciuto intorno a Lionardo Fibonacci*, Rome, 1867; V. A. Le Besgue, "Notes sur les opuscules de Léonard de Pise," Boncompagni's *Bullettino*, IX, 583; O. Terquem,

At the time of Leonardo's birth, Pisa ranked with Venice and Genoa as one of the greatest commercial centers of Italy. These towns had large warehouses where goods could be stored and duty paid in all important ports of the Mediterranean, the head of such an establishment being a man of considerable prominence. It was such a position that the father[1] of Leonardo held at Bugia[2] on the northern coast of Africa, and in this town Leonardo received his early education from a Moorish schoolmaster.[3] As a young man he traveled about the Mediterranean, visiting Egypt, Syria, Greece, Sicily, and southern France,[4] meeting with scholars and becoming acquainted with the various arithmetic systems in use among the merchants of different lands. All the systems of computation he counted as poor, however, compared with the one that used our modern numerals.[5] He therefore wrote a work in 1202, *Liber Abaci*,[6]

"Sur Léonard Bonacci de Pise et sur trois écrits . . . ," *Annali di Sci. Mat.*, Vol. VII (reprint, Rome, 1856); M. Lazzarini, "Leonardo Fibonacci," *Bullettino di Bibliogr. di. Sci. Mat.*, VI, 98, and VII, 1; P. Cossali, *Scritti inediti*, ed. Boncompagni, p. 342 (Rome, 1857); Libri, *Histoire*, II, 21.

[1] Guglielmo Bonaccio. But the name "Fibonacci" is thought by Boncompagni and Milanesi to be a family name like Johnson, the form "filius Bonacci" being merely a Latin translation. An ancient document of 1226 has "Leonardo bigollo quondam Guilielmi," in which the Latin form "Bonaccius" does not appear, but in which the grandfather has this name. It seems more reasonable, however, to think that when Leonardo himself wrote "filius Bonacci," "filius Bonaccij," and "filius Bonacii," he knew what the words would mean to Latin readers. Leonardo speaks of his father as being "in duan a bugee,"—in the custom house at Bugia.

[2] Modern Bougie, whence France imported her wax candles (*bougies*). Little of its ancient splendor remains except the Moorish gate (Bab-el-Bahr, "sea-gate") in the old ramparts.

[3] "Vbi ex mirabili magisterio in arte per nouem figuris indorum introductus." *Liber Abaci*, p. 1.

[4] ". . . apud egyptum, syriam, greciam, siciliam et prouinciam."

[5] "Sed hoc totum etiam et algorismum atque arcus pictagore quasi errorem computavi respectu modi indorum," as it appears in the Florentine MS. published by B. Boncompagni, Rome, 1857.

Early writers attributed to him the introduction of these numerals into Italy; thus ". . . e questi fù il primo, che portò nell' Italia i carrateri dei numeri conforme testifica Luigi Colliado." *Aritmetica di Onofrio Pvgliesi Sbernia Palermitano*, p. 12 (Palermo, 1670).

[6] "Incipit liber Abaci Compositus a leonardo filio Bonacij Pisano In Anno M° cc° ij°." This is the title as it appears in the first line of Boncompagni,

in which he gave a satisfactory treatment of arithmetic and elementary algebra. The work is divided into fifteen chapters, and the following brief statement of the contents will serve to show its general scope: 1. Reading and writing of numbers in the Hindu-Arabic system;[1] 2. Multiplication of integers;[2] 3. Addition of integers; 4. Subtraction of integers;[3] 5. Division of integers; 6. Multiplication of integers by fractions; 7. Further work with fractions; 8. Prices of goods; 9. Barter; 10. Partnership; 11. Alligation; 12. Solutions of problems; 13. Rule of False Position;[4] 14. Square and cube roots; 15. Geometry and algebra, the former being devoted to problems in mensuration.

Possibly it was his indulgence in travel that caused him to write his name occasionally as Leonardo Bigollo,[5] since in Tuscany *bigollo* meant a traveler. The word also means blockhead, and it has been thought that he had been so called by the professors of his day because he was not a product of their schools, and that he retaliated by adopting the name simply to show the learned world what a blockhead could do. It would be human to hope that the latter explanation is the correct one, just as it is human to rejoice that the son of a provincial official became the greatest medieval mathematician. Such a remarkable career as Fibonacci's warns us, as Froude so

Scritti Fiboracci, I, 1, from the Codex Magliabechianus. The spelling of *abacus* varies in this and other MSS., often appearing as *abbacus*. B. Boncompagni, *Intorno ad alcune opere di Leonardo Pisano*, p. 1 (Rome, 1854).

[1] "Nouem figure indorum he sunt

9 8 7 6 5 4 3 2 1

Cvm his itaque nouem figuris, et cum hoc signo O, quod arabice zephirum appellatur, scribitur quilibet numerus." P. 2.

[2] "Incipit capitulum secundum de multiplicatione integrorum numerorum."

[3] "Incipit capitulum quartum de extractione minorum numerorum de maioribus."

[4] "De regulis elchatayn . . . Elchataieym quidem arabice, latine duarum falsarum posicionum regula interpretatur." See Volume II, Chapter VI.

[5] "Incipit flos Leonardi bigolli pisani . . . ." See Boncompagni, *Tre Scritti*, 1. The word *bigolli* also appears as *pigolli*. See also F. Bonaini, *Iscrizione . . . a onore di Leonardo Fibonacci . . .*, Pisa, 1858; 2d ed., 1867.

truly said, "that we should draw no horoscope; that we should expect little, for what we expect will not come to pass."[1]

In the same years and in the same regions in which Leonardo was bringing new light into the science of mathematics, St. Francis, humblest of the followers of Christ, was bringing new light into the souls of men. Each was one of the world's geniuses, and for a genius there is no human explanation.

Fibonacci's Other Works. Leonardo also wrote three other works, the *Practica geometriae*[2] (1220), the *Liber quadratorum*[3] (1225), and the *Flos*,[4] besides which there is extant a letter of his to Theodorus, philosopher to Frederick II, relating to indeterminate analysis and to geometry. These works treat of the theory of numbers in a way that shows that Leonardo was a mathematician of remarkable ability, considering the time in which he lived. His name attaches to the series 0, 1, 1, 2, 3, 5, 8, 13, . . ., in which $u_n = u_{n-1} + u_{n-2}$, where $u_{n-1} : u_n \to \frac{1}{2}(\sqrt{5} - 1)$.

So far as the schools were concerned, Leonardo's works were like a voice crying in the wilderness. It is probably within the bounds of truth to say that not a professor in the University of Paris, to select what was soon to become the greatest intellectual center of the world, could have made anything whatever out of the fine reasoning of the *Liber Quadratorum* or could have comprehended what the *Flos* was meant to convey to the

[1] "Un brevet d'apothicaire n'empêcha pas Dante d'être le plus grand poète de l'Italie, et ce fut un petit marchand de Pise qui donna l'algèbre aux Chrétiens." Libri, *Histoire*, I, xvi.

[2] On his knowledge of Euclid see G. Eneström, *Bibl. Math.*, V (3), 414.

[3] R. B. McClenon, "Leonardo of Pisa and his Liber Quadratorum," *Amer. Math. Month.*, XXVI, 1. There is a question about the date 1225, although it is given in the MS.

[4] "Incipit pratica geometrie composita a Leonardo pisano de filiis bonaccij anno M°. cc°. xx°."

"Incipit liber quadratorum compositus a leonardo pisano Anni. M. CC. XXV."

"Incipit flos Leonardi bigolli pisani super solutionibus quarumdam questionum ad numerum et ad geometriam uel ad utrumque pertinentium."

These titles are from the Boncompagni editions. *Flos* is a fanciful title,—blossom or flower.

mind. Since the course of study was concerned with little that was scientific,[1] mathematics had no standing there or in the schools of Italy.

Campanus. Roger Bacon speaks highly of a certain Master Nicholas who lived about this time, but concerning whom we know nothing further, and also of Master Campanus de Novaria.[2] The latter is Johannes Campanus[3] (fl. *c.* 1260), sometime chaplain to Urban IV, who reigned as pope from 1261 to 1264. It was he who prepared the translation of Euclid's *Elements* that was used in most of the early printed editions, but which seems to have depended upon at least three earlier translations from the Arabic. Campanus also wrote a *Tractatus de Sphaera*, a *Theoria Planetarum*, a *Calendarium*, a work *De Computo Ecclesiastico*, a work on perspective, and a memoir *De Quadratura Circuli* which seems lost but which was mentioned a century later by Albert of Saxony (*c.* 1370). Contemporary writers have little to say of his life. He held relatively minor positions in the Church, and it is probable that in his later years he was a canon in Paris.[4]

In the appendix to his translation of Euclid he showed how to compute the sum of the angles of a stellar pentagon. It is not improbable that the figures used by the astrologers of this period account for the interest developed by various writers in the study of stellar polygons in general. Campanus also considered the trisection problem, the irrationality of the Golden Section (not yet known by this name), and the angle between a circle and a tangent. He was, therefore, not merely a compiler of translated material but a man genuinely interested in geometry.[5]


[1] The oldest statutes now extant (1215) record: "Non legant in festivis diebus, nisi Philosophos et rhetoricas et quadrivalia et barbarismum et ethicam, si placet." See Suter, *Univ. Mittelalt.*, p. 56.

[2] Of Novara, near Milan.

[3] Giovanni Campano. Cantor, *Geschichte*, II (2), 90; C. S. Peirce, in *Science*, XIII (N.S.) (New York), 809.

[4] Pacioli (1509), in his *De diuina proportione* (I, 4) speaks of him as "el gran philosopho Campan, nostro famosissimo mathematico."

[5] B. Baldi, Boncompagni's *Bullettino*, XIX, 591.



FIRST PAGE OF THE CAMPANUS EUCLID

From the MS. probably given by Campanus to Pope Urban IV, at that time Jacques Pantaléon, Patriarch of Jerusalem. Now in the library of Mr. Plimpton

Other Italian Writers. In the 13th century Italian mathematics consisted almost entirely of astronomy, and of the works bearing upon this subject the most popular one produced was the *Tractatus Sphaerae* of Bartolomeo da Parma, who is known to have been teaching mathematics at Bologna in 1297. He also wrote on geometry and astrology.[1]

Whether Guglielmo de Lunis belonged to the 13th century is uncertain,[2] but he translated an algebra from the Arabic.[3]

Pietro d'Abano[4] (*c.* 1250–*c.* 1316), a professor of medicine[5] at Padua, wrote an *Astrolabium planum*,[6] his interest in the astrolabe being doubtless due to its applications to astrology.

British Scholars.[7] Fibonacci was not a product of the universities, but he speaks of his master[8] as one who had studied in the universities of Oxford and Paris, and no doubt, in his mature years at least, he learned from him. This man was "the wizard" Michael Scott,[9] who had not only studied at the

[1] E. Narducci, Boncompagni's *Bullettino*, XVII, 1, 43, 165.

[2] *Bibl. Math.*, XII (3), 270.

[3] An algebra MS. of the 15th century treats of "la regola de Algebra amucabale . . . secondo ghuglielmo de lunis." See *Rara Arithmetica*, p. 463; *Bibl. Math.*, IV (2), 96, and V (2), 32, 118.

[4] Petrus Aponensis. The dates are sometimes given as 1253–*c.* 1319.

[5] On the relations of mathematics to medicine two or three centuries later, see the author's article on "Medicine and Mathematics in the Sixteenth Century," in *Annals of Medical History*, I, 125.

[6] Published in Venice in 1502.

[7] Cantor, *Geschichte*, II, chap. xlvi.

[8] He dedicates to him the second edition of his *Liber Abaci* in these words: "Scripsistis mihi domine mi magister Michael Scotte, summe philosophe, ut librum de numero, quem dudum composui, uobis transcriberem."

[9] Spelled also Scot. Born possibly at Balwearie, Scotland, *c.* 1175; died *c.* 1234. He was also called Michael Mathematicus.

> "In these fair climes it was my lot
> To meet the wondrous Michael Scott;
> A wizard of such dreaded fame,
> That when, in Salamanca's cave,
> Him listed his magic wand to wave,
> The bells would ring in Notre Dame!"
> Scott, *Lay of the Last Minstrel*, II, xiii

> "That other, round the loins
> So slender of his shape, was Michael Scot,
> Practised in every slight of magic wile."
> Dante, *Inferno*, XX, Cary translation

universities mentioned, but had learned Arabic and made astronomical observations at Toledo. He was later appointed astrologer to Frederick II and seems to have been employed by this ruler to make known to scholars, through translations from the Arabic, the newly discovered Greek texts.

Upon those who, unlike Michael Scott, studied chiefly in England at this time, some influence may have been exerted through the arrival at Oxford in 1224 of the first of the Franciscans. These men[1] were not learned in the sciences, but they came from the intellectual centers of Southern Europe and could not have been ignorant of what scholars were doing beyond the Alps and the Pyrenees. Their presence in one of the university centers was especially significant.

Sacrobosco. The second of the prominent British scholars of this century was Sacrobosco,[2] who was educated at Oxford and entered the University of Paris *c.* 1230. He afterwards taught mathematics and philosophy in Paris and died there *c.* 1256.[3] He was buried in the Cloister Sodalium Mathurinalium, his astrolabe being placed on his tomb.[4]

Sacrobosco wrote the most popular work on the sphere that had appeared up to that time, and did much, through his

[1] A list is given by Wood, *Historia Oxon.*, I, 67–77, and in A. G. Little, *The Grey Friars in Oxford*, chap. i and p. 176 (Oxford, 1892).

[2] Born at Halifax, Yorkshire, *c.* 1200; died at Paris, *c.* 1256. The name appears in various forms, such as Johannes de Sacrobosco, John of Halifax, John of Holywood, Sacro Bosco, Sacrobusto. Sacrobosco is the Latin for Holywood (Holyfax, Halifax). Widman (1489) writes the name Iohānē vō sacrobusto, and Pacioli (1494) writes it Giouā de sacro busco. J. Aubrey, *Brief Lives*, ed. Clark, I, 408 (Oxford, 1898) (hereafter referred to as Aubrey, *Brief Lives*), says that "Dr. [John] Pell is positive that his name was Holybushe."

[3] The date of his death was formerly given definitely as 1256 on the authority of G. J. Vossius, *De Vniversae Mathesios Natvra & Constitvtione Liber*, p. 179 (Amsterdam, 1650). P. Tannery has shown that the obscure verse from which Vossius obtained this date refers to the completion of his *Compotus*, and moreover that the date should be read 1244 instead of 1256. The verse is

"M Xristi bis C quarto deno quater anno." See *Bibl. Math.*, XIII (2), 32.

[4] J. C. Heilbronner, *Historia Matheseos Universae*, p. 471 (Leipzig, 1742); hereafter referred to as Heilbronner, *Historia*. Wood (*Historia Oxon.*, I, 85) speaks of his teaching there: "Joh. de Sacro bosco. Claruit apud Parisienses in Mathesi & in Philosophia."

Tractatus de Arte Numerandi[1] or *Algorismus*, to make the Hindu-Arabic arithmetic known to European scholars. These books were widely used for three hundred years, and continued in use until the close of the 16th century.[2]

The third of the prominent British scholars of this period was Robert Grosseteste, or Greathead (died October 9, 1253), at one time a student at Paris, later a student and teacher at Oxford, and finally bishop of London.[3] His interest was chiefly in the applications of mathematics to physics and astronomy,[4] but he also wrote a *Praxis geometriae* and a work on Euclid's *Optics*.[5]

Among the Oxford men of this period was John of Basingstoke (died 1252),[6] who learned Greek in Athens (1240) and took back to England some knowledge of the numeral systems and possibly of the mathematics of classical times.

Roger Bacon. The most prominent scholar in England in the 13th century, however, was Roger Bacon (1214–1294), a man of erudition and of prophetic vision. His works show a knowledge of Euclid's *Elements* and *Optics*, of Ptolemy's *Almagest* and *Optics*, of Theodosius on the *Sphere*, of parts of the works of Hipparchus, Apollonius, and Archimedes, and of the works of various Arab writers. He was familiar with the writings of Aristotle and with some of the commentaries upon them. Of

[1] Printed in Halliwell, *Rara Math.*, 1. For early editions, see the *Rara Arithmetica*. An edition by M. Curtze appeared in 1897.

[2] Suter, *Univ. Mittelalt.*, p. 67; *Bibl. Math.*, XI (2), 97; P. Riccardi, *Bibl. Math.*, VIII (2), 73.

[3] The variants of his name, as given by Wood (*Historia Oxon.*, I, 81), are interesting. They include such forms as Grossum caput, Groshedius, Grouthede, Grokede, and Groschede. He was also known as Robertus Lincolniensis and Rupartus Lincolniensis.

[4] *Theorica planetarum*, *De astrolabio*, *De cometis*, *De sphaera coelesti*, *De computo*, *Praxis geometriae*, and a *Calendarium*.

[5] See also L. Baur, "Der Einfluss des Robert Grosseteste auf die wissenschaftliche Richtung des Roger Bacon," in A. G. Little, *Roger Bacon Essays*, p. 33 (Oxford, 1914) (hereafter referred to as Little, *Bacon*), and *Die philosophischen Werke des Robert Grosseteste*, Münster, 1912.

[6] Under this date Matthew Paris records: "Obiit magister Johannes de Basingestokes, archidiaconus Legrecestriae, vir in trivio et quadrivio ad plenum eruditus."



FROM A MANUSCRIPT OF SACROBOSCO

This MS. was written in Germany, *c.* 1442. It shows distinctly the numerals as they then appeared. Now in the library of Mr. Plimpton

algebra he knew little except the name.[1] Mathematics as then understood was little more than astronomy, and for the work of most of his contemporaries in this field he had a profound contempt. This contempt was even more pronounced with respect to teaching, in which he asserted that an enormous amount of time was wasted. He stated that he had devoted forty years to study, and that the entire ground could have been covered in from three to six months.[2] The teachers at Paris he could only characterize by their "four defects,—infinite and puerile vanity, ineffable falsity, voluminous superfluity, and the omission of all that is worthy." The charge was untrue, as are most epigrams of the kind, for Bacon was given to dipping his pen in vitriol. It is no wonder that his contemporaries generally hated him. Although spoken of in later times as *doctissimus mathematicus* he contributed nothing to pure mathematics, and his chief work in applied mathematics was a calendar which the world was not ready to appreciate.[3]

John Peckham. Of Bacon's influence upon his pupils, in the direction of mathematics, we have little evidence. There is some reason for thinking that he was possibly the one who inspired John Peckham[4] to take up the study of the science. At any rate Peckham was a scientist of repute and his *Perspectiva communis* was looked upon as a classic for three hundred years.[5] He became archbishop of Canterbury in 1279.


[1] In his *Opus majus*, he says: "Algebra quae est negotiatio, et almochabala quae est census."

[2] "Multum laboravi in scientiis et lingua, et posui jam quadriginta annos postquam didici primo alphabetum . . . et tamen certus sum quod infra quartam anni, aut dimidium anni, ego docerem ore meo hominem sollicitum et confidentem, quicquid scio de potestate scientiarum et linguarum." *Opus Tertium*, cap. xx.

[3] Besides his published works by S. Jebb (1733, 1750), J. S. Brewer (1859), J. H. Bridges (1897), and Robert Steele, consult E. Charles, *Roger Bacon, sa vie, ses ouvrages, ses doctrines* . . . (Paris, 1861), and Little, *Bacon*.

[4] Born in Kent, probably some time before 1240; died at Mortlake, December 8, 1292. The name also appears as Peachamus, Peccamus, and Pithsanus, with various other modifications.

[5] Facio Cardano (1444–1524) edited it under the title *Prospectiua coīs d. Iohānis archiepiscopi Cātauriēsis*, and it was printed s. l. a. (but 1482, at Milan). There are various editions of this work.



FIRST PAGE OF THE EARLIEST FRENCH ALGORISM

Written *c.* 1275, and now in the Library of Ste. Geneviève, in Paris

French Scholars. France produced no mathematicians of importance in the 13th century. During a considerable part of the time her great university was a place of rioting rather than a seat of learning. In spite of this fact, however, several respectable scholars appeared, one of the first being Alexandre de Villedieu[1] (*c.* 1225), a Franciscan monk from Bretagne. He wrote *De Sphaera*, *De Computo Ecclesiastico*, and *De Arte Numerandi*, and taught in Paris. He is best known, however, for his *Carmen de algorismo*,[2] a little arithmetic in Latin verse that probably did more to make known the new Hindu-Arabic numerals than any other work of the century.[3] A little later (*c.* 1275) the first algorism in the French language was written.

Among the contemporaries of Alexandre de Villedieu there was Vincent de Beauvais[4] (*c.* 1250), a Dominican, whose encyclopedia, the *Speculum Majus*,[5] written for Louis IX ("Saint Louis"), includes the quadrivium, the subject being very poorly treated.

Roger Bacon mentions as one of the greatest mathematicians of his time (*c.* 1265) a certain Petrus de Maharncuria,[6] but all that is known of him is that he wrote a work on the magnet.

German Writers. Of the German mathematicians of the 13th century only three deserve special mention. Of these the first in order of time and of mathematical ability was Jordanus Nemorarius,[7] who studied at Paris and wrote an *Arithmetica decem libris demonstrata*[8] and possibly an *Algorismus demonstratus*.[9] He also wrote a work on mathematical astronomy,—*Tractatus de sphaera*; one on geometry,—*De triangulis*; and one of the leading books of the Middle Ages on algebra,—*Tractatus*

[1] De Villa Dei, or De Villa Dei Dolensis.

[2] *Song on Algorism*, that is, on al-Khowârizmî's arithmetic methods.

[3] It is printed in full in J. O. Halliwell, *Rara Mathematica* (London, 1838), 2d ed. (1841), p. 73.

[4] Vincentius Bellovacensis. [5] His *Opera* appeared at Venice in 1494.

[6] Also called Peter de Maharn-Curia and Petrus Peregrinus. Maharncuria seems to have been Maricourt in Picardy. Boncompagni's *Bullettino*, I, 1.

[7] Also known as Jordanus de Saxonia and Jordan of Namur. He was born at Borgentreich in the diocese of Paderborn and died in 1236 or 1237. Cantor, *Geschichte*, II, chap. xliii. [8] Published at Paris in 1496.

[9] As to the doubt upon this point see G. Enestrom, *Bibl. Math.*, V (3), 9.

de numeris datis. The *Arithmetica*[1] is on the theory of numbers as set forth in treatises like that of Boethius, and is the least original of his works. The one noteworthy feature of the book is the use of letters to represent general numbers. This is already found to a certain extent in the works of earlier writers, including Aristotle and Diophantus, but Jordanus uses letters quite as they are used today, letting *b*, for example, represent any number whatsoever.[2] The *Tractatus de sphaera* was for a long time a classic and several editions were printed. The *De triangulis*[3] is a work in four books containing seventy-two propositions of the usual type, together with propositions on such topics as the center of gravity of a triangle, curved surfaces, and similar arcs.

The *Tractatus de numeris datis* is a system of algebraic rules. The problems[4] generally relate to a *numerus datus*, a given number,[5] which has to be divided in some stated manner, as in many of the problems in our current algebras.[6]

He also wrote a work entitled *De Ponderibus Propositiones XIII*, which was printed at Nürnberg in 1533 and contains a brief treatment of statics. He is the Jordanus de Saxonia who, in 1222, became general of the Dominican order.[7]

[1] On the MSS. in the library of the Royal Society and in Oxford, both of which differ from the first printed edition, see J. O. Halliwell, *A Catalogue of Miscellaneous Manuscripts preserved in . . . the Royal Society* (London, 1840), and J. Wallis, *Algebra*, p. 13 (Oxford, 1693).

[2] Cantor, *Geschichte*, II (2), 56; Eneström, *Bibl. Math.*, VII (3), 85.

[3] "Jordani Nemorarii Geometria vel de Triangulis libri IV," in the *Mittheilungen des Coppernicusvereins*, Heft VI (Thorn, 1887).

[4] See P. Treutlein, *Abhandlungen*, II, 135.

[5] "Numerus datus est cuius quantitas nota est."

[6] One of his first problems is practically this: To separate a given number into two parts such that the sum of the squares of the parts shall be another given number. *E.g.*, $x+y=10$, $x^2+y^2=58$, whence $x=7$, $y=3$. *Abhandlungen*, II, 136 (4).

[7] One Oxford MS. distinctly calls him Jordanus de Saxonia. Nicolas Trivet, an English chronicler of the 14th century, under the year 1222, states: "Hoc anno in Capitulo Fratrum Praedicatorum generali tertio, quod Parisiis celebratum est, successor beati Dominici in Magisterio Ordinis Fratrum Praedicatorum factus est frater Iordanus, natione Teutonicus, Dioecesis Moguntinae, qui cum Parisiis in scientiis saecularibus et praecipue in Mathematicis magnus haberetur. . . ."

One of the greatest of the German scholars of this period was Albertus Magnus,[1] Count of Bollstädt, a Dominican priest, and Bishop of Regensburg. He studied at Padua and taught at Bologna, Strasburg, Freiburg, Cologne, and Paris. So versatile was he that he was called "Doctor Universalis." His interests were chiefly in philosophy and physics, but his works include material on astronomy and some reference to Pythagorean arithmetic.[2] Claude Fleury, who wrote an ecclesiastical history in 1691, remarked that he could see nothing great in him but his volumes.

At the close of the century (*c.* 1270) Witelo or Vitello,[3] probably from Thüringen but possibly from Poland,[4] wrote on perspective (optics) and astronomy, a fact which shows the interest in this phase of applied mathematics in Poland in the 13th century.

Other Thirteenth-Century Writers. Of the other scholars of this century whose works touched upon mathematics the most prominent was Alfonso X, King of Castile (1223–1284), known as el Sabio (the Wise). He was an astronomer of merit and his name appears in the Alfonsine Tables,—planetary tables which improved upon the imperfect ones left by Ptolemy. Work upon them began in 1248 and was completed in 1254.[5] Tycho Brahe is said to have deplored the waste of money involved in their compilation, although they unquestionably stimulated the study of mathematical astronomy.

[1] Born at Lauingen, Swabia, 1193 or 1205; died at Cologne, 1280. Albertus Teutonicus, de Colonia, or Ratisbonensis. See Cantor, *Geschichte*, II (2), 86, and Dixon's translation (London, 1876) of his biography by Sighart (Regensburg, 1857).

[2] The first edition of his *Opera Omnia* appeared in Leyden in 1651. The best edition is that of Paris, 1890.

[3] In the oldest MSS. the name appears as Witelo. The forms Vitello and Vitellius are later. There are many variants, such as Witilo, Witulo, Widilo, Wito, and Vitellion.

[4] He speaks of his country, saying: "In nostra terra, scilicet Poloniae habitabili . . . ," and of himself as "Thuringo-polonus" and as "Filius Thuringorum et Polonorum," so that possibly his mother was a Pole.

[5] The date of completion is sometimes placed later than this. The tables were first printed in Venice in 1483.

Another prominent writer of the period was Arnaldo de Villa Nova[1] (*c.* 1235–*c.* 1313), who taught at Paris, Barcelona, and Montpellier. While known principally as a physician and for his twenty works on alchemy, he wrote a *Computus Ecclesiasticus & Astronomicus*,[2] probably being led to a study of the subject through its relation to astrology.[3]

Roger Bacon, in his condemnation of most of his contemporaries, speaks of "the notorious William Fleming who is now in such reputation, whereas it is well known to all the literati at Paris that he is ignorant of the sciences in the original Greek, to which he makes such pretensions." This Flemish writer was William of Moerbecke,[4] chaplain to Clement IV and Gregory X. Among his translations were the catoptrics of Heron and the writings of Archimedes on floating bodies.[5] It is thought that Tartaglia[6] took his translation of Archimedes from this writer.[7] He also wrote on perspective.

Byzantine Writers. In the 13th century the only writer of note in the Near East was Georgios Pachymeres,[8] who may for convenience be classified as European, although born in Asia Minor. He wrote on the *Four Mathematical Sciences*,[9] that is, on arithmetic, music, geometry, and astronomy. The work is important only as showing that interest in learning had not

[1] Arnauld de Villeneuve, Arnald Bachuone, Arnoldus Villanovanus. He was probably born at Villa Nova, Catalonia, but possibly at Villeneuve in Southern France.

[2] Printed at Venice in 1501. With respect to the edition see the *Rara Arithmetica*, p. 73.

[3] On the geometric figures used in astrology the *Ars Magna* of Raymundus Lullus (*c.* 1235–1315), a writer of this period, may be consulted.

[4] Guilielmus Brabantinus or Flemingus. He died *c.* 1281 as archbishop of Corinth.

[5] *De iis quae in humido vehuntur.* See also J. L. Heiberg, *Zeitschrift* (Hl. Abt.), XXXIV, 1–84; XXXV (Hl. Abt.), 41–48, 48–58, 81–100, and later volumes.

[6] See page 297. [7] Cantor, *Geschichte*, II (2), 514.

[8] Born at Nicæa, in Bithynia, about 60 miles from Constantinople, in 1242; died *c.* 1316.

[9] Περὶ τῶν τεσσάρων μαθημάτων Παχυμεροῦς μεγάλου διδασκάλου. There are various MSS. extant. See E. Narducci, "Di un Codice Archetipo e Sconosciuto dell' opera di Giorgio Pachimere," *Rendiconti della R. Accad. dei Lincei*, VII, 194.

wholly died out in the period between the capture of Nicæa by the Crusaders in 1097 and its downfall before the Turkish invaders in 1330.

Otherwise there is little known of the mathematics of Constantinople in the 13th century. There is evidence, however, to show that her scholars were using Greek numerical characters even as late as the 15th century, augmented by a symbol for zero resembling our inverted *h*. Their problems were trivial, chiefly relating to mensuration.[1] Although they used the Greek forms, they were acquainted with the numerical system of the Arabs and spoke of it as Hindu in origin, but they were not familiar with the numerals which we commonly call Arabic.

4. Christian Europe from 1300 to 1400

General Activity of the Fourteenth Century. After the brilliant beginning of a renaissance of learning in the 13th century, it would naturally be expected that the 14th century would see a notable revival of science and letters. To understand why this expectation was not fully realized, it is necessary to consider the peculiar conditions by which Europe was confronted. As for Italy, this country was at last fully awakened to the beauties of ancient literature, and so Dante (1265–1321), taking Vergil as his master, produced the *Divina Commedia*, the "Epic of Medievalism"; Petrarch (1304–1374) made a notable collection of manuscripts of the ancient classics and started a movement that resulted in a new appreciation of the literature of Greece and Rome; and Boccaccio (1313–1375) showed great zeal in the attempt to collect and study the works of the ancients. In this search Constantinople was drawn upon for Greek manuscripts, with the result that Italian scholars were interested anew in the study of science and letters. Furthermore, the Florentine republic had just become practically a government by the merchant class, owing to modifications in the constitution between 1282 and 1292,—a fact that must have had much to do with the great prominence

[1] J. L. Heiberg, "Byzantinische Analekten," *Abhandlungen*, IX, 161.

of Florentine arithmetic in the schools of the 14th century. A general accumulation of wealth must also have followed, which would naturally tend to foster the arts and sciences. All this was promising, and the result would probably have been the hastening of the period popularly known as the Renaissance, had it not been for two deterring factors.

The first of these factors was the Hundred Years' War (say 1338–1453, although also given as 1328–1491), which overturned the economical and political systems of the two most advanced countries of Europe north of the Alps. The battle of Crécy (1346) struck at something besides feudalism.

The second deterring factor was the terrible ravaging of the Black Death (1347–1349), by which from a third to a half of the population of Europe is thought to have been swept away.

As to the universities in the 14th century, they did little for mathematics. Those of Italy were behind their contemporaries in Paris, England, and Germany, the statutes of 1387 making no mention whatever of the subject.[1] In England, Merton College, Oxford, was the mathematical center[2] and made some pretense at work in this science, while Paris had lectures on algorism, astronomy, and geometry, such as they were. In the newly founded University of Erfurt (1392), which may be taken as a German type, an elementary knowledge of mathematics was offered but apparently was not required.[3]

Italian Writers. No Italian writer of the 14th century stands out as showing any real genius in mathematics, as a brief list will bear witness. Cecco d' Ascoli (1257–1327), also known as Francesco di Simone Stabili and as Francesco degli Stabili, a native of Ascoli in Romagna, was professor of philosophy at Bologna and Rome, wrote a commentary on the *Sphaera* of Sacrobosco,[4] and did much to bring into high repute once more

[1] Suter, *Univ. Mittelalt.*, p. 75. [2] *Ibid.*, p. 83.

[3] W. Hellmann, *Ueber die Anfänge des math. Unterrichts an den Erfurter . . . Schulen*, I, 4. Erfurt, 1896.

[4] On the early Bologna mathematicians in general, consult Silvestro Gherardi, *Di alcuni materiali per la Storia della Facoltà Matematica . . . di Bologna*, p. 17 (Bologna, 1846); hereafter referred to as Gherardi, *Facoltà Mat. Bologna.*

the ancient belief in astrology, a subject which perhaps reached its greatest popularity in this century.

Andalò di Negro (*c.* 1260–*c.* 1340), a native of Genoa, had considerable reputation as a mathematician and astronomer, writing several works on the astrolabe, a book on the planets, and a *Tractatus de sphaera.* There is also ascribed to him a practical arithmetic.[1]

Barlaam (*c.* 1290–*c.* 1348), a native of Seminara in Calabria, Italy, bishop of Geraci, studied in Constantinople and wrote on computing, astronomy,[2] the science of numbers,[3] algebra,[4] and Book II of Euclid.

Paolo Dagomari,[5] a native of Prato, in Tuscany, was prominent in Florence as an arithmetician and astronomer. His *Trattato d'Abbaco, d'Astronomia, e di segreti naturali e medioinali* contained a little commercial arithmetic and gave him a reputation more extended than scientific.[6] That he wrote on algebra is asserted by at least one later writer.[7] He may have been the Paolo Pisano who is said to have lived about this time.

A more worthy writer on mathematics appeared in the person of Rafaele Canacci (*c.* 1380) of Florence, author of an algebra[8] with a number of historical notes.

[1] C. de Simony, Boncompagni's *Bullettino*, VII, 313, 330.

[2] *Libri V logisticae astronomicae.* See also B. Baldi, *Cronica di Matematici*, p. 85 (Urbino, 1707), and Boncompagni's *Bullettino*, XIX, 598. Barlaam's (Barlaamo's) given name may have been Bernardo, but this is uncertain. He is occasionally known, from his birthplace, as Calabro.

[3] *Arithmetica demonstratio eorum quae in secundo libro elementorum (Euclidis) sunt.* It was printed at Strasburg in 1564.

[4] Λογιστικη, *sive arithmeticae, algebraicae libri VI.* It was printed at Strasburg in 1572.


[5] Born at Prato, *c.* 1281; died at Florence, 1365 or 1374. Known also as Paolo dell' Abaco, Paolo Astrologico, Pagolo Astrologo, Paoli il Geometra, Paolo Geometra, and Paolo Arismetra. F. Villani (fl. 1404), in *Le Vite d' Uomini illustri Fiorentini*, 2d ed., Florence, 1826, speaks of him as "geometra grandissimo, e peritissimo aritmetico . . . diligentissimo osservatore delle Stelle e del movimento de' cieli." For a résumé of his work see D. Martines, *Origine e progressi dell' aritmetica*, p. 59 (Messina, 1865); hereafter referred to as Martines, *Origine aritmet.*

[6] For a description of the work, see *Rara Arithmetica*, p. 435.

[7] See *Rara Arithmetica*, p. 463, with reference to "*m°*. paolo fiorj che circha al. 1360. duro."

[8] See *Rara Arithmetica*, p. 459.

There was also a Master Biagio[1] of Parma (died 1416) who wrote an arithmetic and an algebra, but neither has been published. He taught astrology and philosophy at Paris, Pavia, Bologna, Padua, Venice, and Parma, wrote a commentary on Oresme's *De latitudinibus formarum*,[2] and wrote on statics and perspective. The famous educator Vittorino da Feltre (1378–1446, but the dates are doubtful), born in poverty, worked as



FROM DAGOMARI'S TRATTATO D'ABBACO

From an Italian MS. of *c.* 1339. Notice also the early per cent sign, p 100 lano (per 100 the year), and the sign for *lb.* The latter is possibly the origin of the dollar sign

a scullery boy in Biagio's house so as to learn geometry from him, and in turn became one of the best teachers of mathematics of his time.[3]

Toward the close of the century Antonio Biliotti[4] (*c.* 1383) of Florence taught mathematics in Bologna, but he left no works on the subject. Altogether the mathematical output of Italy in this century was not encouraging.

Constantinople. Constantinople was at this time experiencing an intellectual revival similar to the one seen in Italy before the coming of the Black Death. Prominent among her scholars was Maximus Planudes (*c.* 1340), a Greek monk, at one time

[1] Also Biagio da Parma and Pelacani. There was also a "m°. biagio che circha al. 1340. anj morj," as a MS. of *c.* 1440 asserts, although this 1340 may be wrong and the two may be the same. See *Rara Arithmetica*, p. 463.

[2] See page 199. The students of his day in Paris had a phrase "aut diabolus est, aut Blasius Parmensis."

[3] W. H. Woodward, *Vittorino da Feltre*, Cambridge, 1892.

[4] Also called Antonio dall' Abaco.

(1327) ambassador to Venice, who wrote on Diophantus and who also wrote an arithmetic based upon the Hindu-Arabic numerals.[1] He was a man of industry but of no genius, and his arithmetic is of value chiefly as showing the influence of Bagdad upon the mathematical thought of Constantinople. It sets forth the system of notation by the "nine figures received from the Hindus" together with the zero, and is the first of the Greek works to give any attention to modern methods of calculation. Planudes is also deserving of credit as a translator of various Latin classics into Greek.[2]

Among the minor contemporaries of Planudes there was Joannes Pedias'imus (*c.* 1330), also called Galenus, who was keeper of the seal of the patriarch of Constantinople. He wrote a work on geometry in which he attempted to pattern after the style of Heron of Alexandria, and also wrote upon the duplication of the cube, and upon arithmetic.[3] In general, however, his work was literary and philosophical. Among his contemporaries he was known as the "Chief of Philosophers."

There lived in Constantinople a little later than Joannes Pediasimus the celebrated grammarian Manuel Moschopou'lus,[4] a native of Crete. The dates are uncertain, but he seems to have lived *c.* 1300. Although there were two men of the same name, this one and his nephew, it seems from a manuscript of a work by Nicholas Rhabdas, referred to below, that this is the one who wrote a treatise on magic squares, the earliest contribution to the subject in the Mediterranean countries.[5]

[1] He called his work *Ψηφοφορία κατ' Ἰνδούς* (*Indian Arithmetic*). There is a Greek edition by C. I. Gerhardt (Eisleben, 1865), and a German translation by H. Wäschke, *Das Rechenbuch des Maximus Planudes* (Halle, 1878) (hereafter referred to as Wäschke, *Planudes*). See also Heath, *History*, II, 546.

[2] Kroll, *Geschichte*, p. 70.

[3] Boncompagni's *Bullettino*, III, 303.

[4] Or Emanuel. In Greek the name appears both as Μανουήλ and as Ἐμανουήλ Μοσχόπουλος. Heath, *History*, II, 549.

[5] S. Günther, *Vermischte Untersuchungen zur Geschichte der math. Wissenschaften*, p. 195 (Leipzig, 1876) (hereafter referred to as Günther, *Vermischte Untersuch.*); P. Tannery, "Manuel Moschopoulos et Nicolas Rhabdas," *Bulletin des Sciences math. et astr.*, VIII (2), September 2, 1884.

About this time Nicholas Rhabdas[1] (*c.* 1341), a Greek "arithmetician and geometer"[2] from Smyrna, wrote from Constantinople two letters on arithmetic, and particularly on finger reckoning,[3] a subject first treated of with any completeness by Bede. He also edited a work of Planudes on the Hindu arithmetic,[4] possibly during the lifetime of the latter. With him there flickered out what once had been a great beacon light,—the mathematics of the Greeks,—and at the same time any real appreciation of the language itself almost ceased to exist. Petrarch began to study classical Greek in 1342, with the aid of a monk who had lived in Constantinople, and a learned scholar, Manuel Chrysoloras, lectured upon it in Florence from 1397 to 1400; but it was not until the 16th century that mathematical works of the Greeks began to be known again in the original tongue.

English Writers.[5] England produced several mathematicians of more than ordinary ability in this century, all but one of them doing his real work before the years of the pestilence.

Richard of Wallingford (born c. 1292; died 1336) lectured on the liberal arts at Oxford and wrote on trigonometry[6] and arithmetic.[7] He seems to have been one of the best-known mathematicians of his time.[8] It was no doubt his influence

[1] Nicholas Smyrnaeus, Rhabda, Artabasda, Artabasdes. In one manuscript the name appears as Nicolas Artavasdan. See P. Tannery, "Notice sur les deux lettres arithmétiques de Nicolas Rhabdas," in *Notices et extraits des manuscrits de la Bibliothèque nationale*, Paris, XXXII (1886), 121.

[2] As he describes himself,— "*ἀριθμητικοῦ καὶ γεωμέτρου.*"

[3] "*Ἔκφρασις τ.ῦ δακτυλικοῦ μέτρου.* For a review of this arithmetic see *Bibl. Math.*, I (2), 28. It has been printed several times, as by N. Caussinus. *Eloquentia sacra et humana*, Paris, 1636, and by Morellus, *Nic. Smyrnaei Artabasdae, graeci mathematici,* "*Ἔκφρασις numerorum notationis per gestum digitorum*, Paris, 1614. Heath, *History*, II, 550.

[4] Published by Gerhardt, Eisleben, 1865. [5] Cantor, *Geschichte*, II, chap. xlvi.

[6] *Quadripartitum de sinibus demonstratis*; *De sinibus et arcubus in circulo inveniendo*, *De chorda et arcu*, and *De chorda et versa*. The word *sinibus* often appears in the MSS. as *sinubus*, and *arcibus* commonly as *arcubus*. See Montucla *Histoire*, I (1), 529; Cantor, *Geschichte*, II (2), 101.

[7] *De rebus arithmeticis* and *De computo*.

[8] One of the medieval writers speaks of him as "in mathesi omnium sui temporis primus."

that led John Manduith[1] (fl. *c.* 1320) to follow in his footsteps and lecture on trigonometry[2] and astronomy at Oxford.

The most prominent of the English mathematicians of the 14th century was Thomas Bradwardine,[3] known as the "Doctor Profundus." He was professor of theology at Oxford, chancellor of St. Paul's cathedral, and an upholder of liberalism, and he died as archbishop of Canterbury. He wrote four works on mathematics. In his *Arithmetica Speculatiua*[4] he followed the Boethian model, the work relating solely to the theory of numbers. His other works were a *Tractatus de proportionibus*, *Geometria speculativa*, and *De quadratura circuli*. His geometry includes some work on stellar polygons,[5] isoperimetric figures, ratio and proportion, irrationals, and loci in space.

About this time there flourished in England a Cistercian monk by the name of Richard Suiceth[6] (*c.* 1345), probably a native of Glastonbury, in Somersetshire. He was educated at Merton College, Oxford, and wrote an obscure work on mathematics.[7] It treats of a subject just beginning to attract attention in England and in France, *De latitudinibus formarum*.[8]

In this period there also lived a well-known writer, Walter Burley,[9] whose work on the lives of the philosophers and

[1] Mandwith, Manduit.

[2] *De chorda et arcu recto et verso, et umbris*, showing that he was acquainted with the use of tangents.

[3] Born at Hertfield (Hartfield), Chichester, *c.* 1290; died at Lambeth Palace, London, August 26, 1349. The name appears in such forms as Bragwardin, Brandnardinus, Bredwardyn, Bradwardyn, de Bradwardina, and de Bredwardina. Pacioli (*Sūma*, 1494, fol. 68, *r.*) calls him Tomas beduardin.

[4] Printed at Paris in 1495.

[5] ". . . figuris angulorum egredientibus."

[6] The first name may possibly have been Roger or Raymund, and the last name appears in such forms as Suisset, Suicetus, Swincetus, Swineshead, and Suineshevedus, a word derived from the Cistercian cloister, Vinshed, on the Holy Island off the coast of Northumberland.

[7] *Opus aureum calculationum . . . Per . . . Iohanē de Cipro . . . emēdat*[9] *et explicit*, s. l. a., but Pavia, *c.* 1480; also Pavia, 1498. It may have been this work, which also went by the name "calculator," that led to his being called "calculator acutissimus" by one of the early writers.

[8] See Volume II, Chapter V.

[9] Born at Oxford, *c.* 1275; died *c.* 1357. The dates are very uncertain. Cantor, following Prantl, gives his death as 1337. The Latin spelling, Gualterus Burlaeus, and the late English Burleigh are also used.

poets[1] contains biographical notes on such prominent Greek mathematicians as Pythagoras, Plato, and Ptolemy.

There also flourished in the latter part of the 14th century a celebrated English mathematician and physician,—Simon Bredon.[2] He wrote on astronomy,[3] arithmetic,[4] the calculation of chords,[5] geometry,[6] and other related subjects, and various manuscripts of these works still exist.[7] He was one of the earliest European scholars to pay much attention to trigonometry.

Among the minor writers of the period there were William Reade,[8] of Merton College, who had considerable reputation as a mathematician, and who prepared some astronomical tables, and Walter Bryte,[9] who is said to have written on arithmetic,[10] astronomy, and surgery.

In the 14th century there was written an interesting but anonymous manuscript on the mensuration of heights and distances,[11] beginning: "Nowe sues here a Tretis of Geometri wherby you may knowe the heghte, depnes, and the brede of mostwhat erthely thynges." It is a practical work on shadow reckoning and surveying, using the compass, staff, and quadrant. Like many such works it is divided into three parts, very likely due to the Christian idea of the Trinity.[12]

[1] *De Vita et moribus Philosophorum et Poetarum.* The first printed edition was s. l. a., but Cologne, *c.* 1467. There were at least fourteen editions printed before 1501.

[2] Born at Winchcomb; living in 1386. The name appears as Bridonus and Biridanus.

[3] *In demonstratione Almagesti.*

[4] *Arithmetica theorica.*

[5] *Calculationes chordarum*, and *Tabulae chordarum.*

[6] *Quadratura circuli per Campanum et Simon Bredon.*

[7] See Suter, *Univ. Mittelalt.*, 84.

[8] Reede, Rede. He died in 1385 as bishop of Chichester.

[9] Brithus, Brit, Brytte. Possibly identical with Walter Brute, a lay follower of Wycliffe.

[10] *Tractatus algorismalis, De rebus mathematicis.* There is much doubt as to the authorship of each of the works assigned to him.

[11] Halliwell, *Rara Math.*, 56, where the complete text is given. The MS. is Bib. Sloan. 213. xiv, fol. 120, in the British Museum.

[12] "This tretis es departed in thre. þat es to say. hegh mesure. playne mesure. and depe mesure."

Another anonymous manuscript of considerable interest, *The Crafte of Nombryng*, was written *c.* 1300. It is one of the first works on algorism to appear in the English language.[1]

French Writers.[2] In spite of the calamities of war and plague, France did some noteworthy work in mathematics in the 14th century, not only through those born within her own boundaries but through scholars from other lands who found in Paris a more congenial intellectual atmosphere than they could find elsewhere.

Among those whom France could claim by adoption was Petrus Philomenus de Dacia,[3] a native of Denmark,[4] rector of the University of Paris (1326 or 1327), author of works on algorism[5] and the church calendar,[6] and a compiler of certain tables.[7] Still another of the adopted sons of France was Johannes Saxoniensis, or Johann Danck, who carried on his astronomical work in Paris. He left various writings on astronomy.[8]

Of those born in France, the first in point of time was Johannes de Lineriis or Jean de Lignères (*c.* 1300–1350), professor of mathematics at Paris, who adapted the Alfonsine Tables to the meridian of that city.[9] Joannes de Muris, or Jean de Meurs,[10] was a contemporary of his who studied at the Sorbonne and taught there. He wrote (1321) on arithmetic, astronomy,

[1] This manuscript is described more fully in Volume II.

[2] Cantor, *Geschichte*, II, chap. xlvii.

[3] There is an Easter computation of 1300 attributed to him.

[4] Whence "de Dacia."

[5] *Commentum super algorismum prosaicum Johannis de Sacro Bosco.* M. Curtze, *Petri Philomeni de Dacia in Algorismum Vulgarem Johannis de Sacrobosco Commentarius*, Copenhagen, 1897.

[6] *Computus ecclesiasticus.*

[7] His *Tabula ad inveniendam propositionem cujusvis numeri* contains a multiplication table to 49 × 49. See *Bibl. Math.*, IV (2), 32.

[8] *De astrolabio* and on the Alfonsine Tables. See Boncompagni's *Bullettino*, XII, 352.

[9] His nationality is not certain, nor is it clear whether he is the same as Johannes de Liveriis (or Liverius); whose work on fractions was printed at Paris in 1483, and who may have been a Sicilian. See *Rara Arithmetica*, p. 13; M. Steinschneider, Boncompagni's *Bullettino*, XII, 345, 352, 420.

[10] Born in Normandy *c.* 1290; died after 1360. The name also appears as Johannes de Murs or de Muria. L. C. Karpinski, *Bibl. Math.*, XIII (3), 99.

and music. Of his works on arithmetic,[1]—the *Canones tabula proportionum, Arithmetica communis ex diui Seuerini Boetij, Tractatus de mensurandi ratione, De numeris eorumque divisione*, and *Quadripartitum numerorum*,—the *Quadripartitum* is the most noteworthy. It is partly in verse and contains a certain amount of algebra. Among the algebraic equations solved are $x^2 + 12 = 8x$, with the roots 2 and 6; $3x + 18 = x^2$; and one already given by al-Khowârizmî and by Fibonacci, $2\frac{7}{9}x^2 = 100$. It also contains a close approach to a decimal fraction.[2]

The greatest of the French writers of this period was Nicole Oresme,[3] a native of Normandy, sometime professor and "magister magnus" (1355) in the Collège de Navarre at Paris, protégé of Charles V, dean of Rouen (1361), and finally (1377) bishop of Lisieux, Normandy. He wrote *Tractatus proportionum, Algorismus proportionum, Tractatus de latitudinibus formarum, Tractatus de uniformitate et difformitate intensionum*, and *Traité de la sphère*. He also translated Aristotle's *De coelo et mundo*. In the *Algorismus proportionum* is the first known use of fractional exponents, $2^{\frac{1}{2}}$ being written[4] $\frac{1}{2}\,2^{p}$, and $9^{\frac{1}{2}}$ appearing as $\frac{1}{2}.9^{p}$. He also wrote

$$\boxed{1^{p}\,\tfrac{1}{2}}\,4 \quad \text{and} \quad \boxed{\frac{p.1}{1.2}}\,4$$

for $4^{1\frac{1}{2}}$, stating the value to be 8.[5]

In the *Tractatus de uniformitate* there is set forth a suggestion of coordinate geometry, by the locating of points by

[1] For first printed edition, see *Rara Arithmetica*, p. 117. See also A. Nagl, *Abhandlungen*, V, 135; p. 139 for a list of his works.

[2] L. C. Karpinski, *Science* (N. Y.), XLV, 663.

[3] Born probably at or near Caen, *c.* 1323; died at Lisieux, July 11, 1382. Also known as Nicolaus Oresmus, Horem, Horin, and Oresmius. See M. Curtze, *Die Mathematischen Schriften des Nicole Oresme*, Berlin, 1870.

[4] From the 15th century MS. in the University of Basel, used by Curtze in the work above cited. Other MSS. have slightly different forms.

[5] Cantor, *Geschichte*, II (2), 121.

means of two coordinates.[1] Oresme also stands out prominently as a remarkably clear-thinking economist for his generation.[2]

Of much less importance as a writer on mathematics, but of greater reputation in his lifetime, is Petrus de Alliaco,[3] rector of the University of Paris, bishop of Cambray, and cardinal. His work on astronomy[4] throws considerable light on the early computi.

Other Writers. The other contributors to mathematical literature in this century were in general possessed of less ability than those of France and England.

Early in the century Hauk Erlendssön,[5] a Norwegian official, wrote on algorism. This is the first trace that we have of the Hindu-Arabic arithmetic in Scandinavia. There was also a certain Swedish scholar, Master Sven, or Sunon, who lectured on the sphere in 1340.

The leading Jewish mathematician of the 14th century was Levi ben Gerson[6] (1288–1344), who was also well known as a theologian. His *Work of the Computer*[7] was written in 1321. He also wrote a treatise on trigonometry[8] which was translated into Latin under the title *De numeris harmonicis*, but neither work showed any noteworthy power.[9]

Isaac ben Joseph Israeli was apparently a contemporary of Levi ben Gerson, but we are uncertain as to his dates. He

[1] See Volume II, Chapter V. For an early edition of the *Tractatus de latitudinibus* see *Rara Arithmetica*, p. 117. See also *Zeitschrift* (Hl. Abt.), XIII; *Bibl. Math.*, XIII (3), 115, and XIV (3), 210.

[2] *Tractatus de origine, natura, jure, et mutationibus monetarum*, edited by Wolowski, Paris, 1864.

[3] Born at Compiègne, 1350; died at Avignon, August 8, *c.* 1420. The name also appears as Pierre d'Ailly, Alyaco, and Heliaco.

[4] *Cōcordātia astronomie cū theologia.* First printed at Augsburg in 1490.

[5] Born *c.* 1264; died 1334.

[6] Also called Levi ben Gerschom and Gersonides, Leo Ebraeus, and Ralbag (R L B G, for Rabbi Levi ben Gerson), but more commonly known as Leo de Balneolis or Master Leon de Bagnolo, having been born at Balnaolis or Bagnolas, in Catalonia. See J. Carlebach, *Lewi ben Gerson als Mathematiker*, Berlin, 1910.

[7] *Maassei Choscheb*, edited by G. Lange, Frankfort a. M., 1909.

[8] *De sinibus, chordis, et arcubus.* See *Bibl. Math.*, I (3), 372; IV (2), 73; XII (2), 97.

[9] *Bibl. Math.*, XI (2), 103.

wrote a work on astronomy[1] which contains a chapter on geometry and also serves as a source of information on the activity of Jewish and Arabic scholars in Spain.

Among the lesser Jewish scholars of the period were Joseph ben Wakkar of Seville (died 1396), who worked out certain astronomical tables for Toledo; Jacob Poël of Perpignan (fl. *c.* 1360), who did the same for Perpignan; Imanuel Bonfils of Tarascon (died *c.* 1377), whose astronomical tables were highly appreciated and who wrote on the astrolabe; Jacob Carsono (al-Carsi), who wrote both at Seville and Barcelona (*c.* 1375), and whose tables were known to Tycho Brahe; Isaac Zaddik (al-Shadib), who wrote on the astrolabe and prepared various tables of use to astronomers; and Kalonymos ben Kalonymos, a native of Arles (born 1286), known as Master Calo, whose various translations include a paraphrase of Nicomachus.

Of the German writers, two or three are deserving of special mention. The first of these was Heinrich von Langenstein, Heinrich von Hessen, or Henricus Hessianus,[2] bishop of Halberstadt, who taught mathematics at Vienna and had some reputation as a mathematician and astronomer.[3] The second was Chunrad von Megenberg (*c.* 1309–*c.* 1374), who wrote (*c.* 1350) a work based on Sacrobosco's *De sphaera.*[4] About this time there was another Chunrad (Conrad) who was interested in mathematics,—Conrad von Jungingen (*c.* 1400). According to one of the manuscripts he seems to have been the author of the *Geometria Culmensis.*[5] This work consists of five parts, the first two relating to the mensuration of the triangle, the third to the quadrilateral, the fourth to the polygon, and the fifth to curvilinear figures.

[1] *Liber Jesod Olam sive Fundamentum Mundi,* first published in Berlin in 1777, but with later editions in 1846 and 1848.

[2] Born at Langenstein, near Marburg, 1325; died at Vienna, February 11, 1397 (sometimes given as 1394).

[3] *Quaestio de cometa.* He also wrote on the circle.

[4] F. Müller, *Zeittafeln zur Geschichte der Mathematik,* p. 80, with references (Leipzig, 1892); hereafter referred to as Müller, *Zeittafeln.* See also O. Matthaei, *Konrads von Megenberg Deutsche Sphaera,* Berlin, 1912.

[5] *Liber magnifici principis Conradi de Jungegen, magistri generalis Prusie, geometrie practice usualis manualis.* See Cantor, *Geschichte,* II, chap. xlviii.

Contemporary with the first two of these German writers there was Albert of Saxony,[1] or Albertus de Saxonia, who was educated at Prag and Paris, taught at Paris[2] and Pavia,[3] was the first rector of the University of Vienna (1365), and became bishop of Halberstadt (1366–1390). He wrote several scientific works, among them a theoretical treatment of proportion[4] after the manner of Boethius, *De latitudinibus formarum*,[5] *De maximo et minimo*, and *De quadratura circuli*.[6] Like most medieval writers, he took $3\frac{1}{7}$ for the value of π, apparently without considering it a mere approximation.

5. Christian Europe from 1400 to 1500

Influences leading to the Renaissance. Of the influences leading to that revival of learning known as the Renaissance, the two most potent were the transfer of Eastern scholarship to Italy and the invention of printing. It is customary to speak of the former as dating from the fall of Constantinople, that is, from its capture by the Turks in 1453.[7] When Mohammed II, standing on the banks of the Bosporus, repeated the Persian distich,

> The spider has woven his web in the Imperial palace,
> And the owl has sung her watch-song on the towers of Afrasiab,

he epitomized the situation of Greek culture in ancient Byzantium. There was nothing left for the remnant of the Hellenic civilization to do but to seek refuge in other lands.

Had Stephen Dusan not died when he did (1356), or had he left a worthy successor to the Serbian throne, Constantinople might have fallen to a western instead of an eastern conqueror. It is interesting to speculate as to what would have been the

[1] Born at Riggensdorf, Saxony, *c.* 1325; died at Halberstadt, 1390.
[2] Du Boulay, *Historia Universitatis Parisiensis*, p. 362. Paris, 1668.
[3] F. Jacoli, Boncompagni's *Bullettino*, IV, 495.
[4] *Tractatus proportionum*, first printed *c.* 1478. See *Rara Arithmetica*, p. 9.
[5] Printed at Padua in 1505.
[6] F. Jacoli, Boncompagni's *Bullettino*, IV, 493; H. Suter, *Zeitschrift* (Hl. Abt.), XXIX, 81.
[7] For a vivid description see Gibbon, *Decline and Fall of the Roman Empire*, Vol. VI, chap. lxviii.

result of such an event upon civilization in general and upon mathematics in particular.[1] We have already seen, however, that this transfer of Greek civilization began a century before the fall of the city, and both Rome and Florence had begun to acquire collections of Greek and Latin manuscripts that were to be available when the printing press should be ready to make their contents known.

About the opening of the 15th century Niccolò de' Niccoli (1363–1437) made a noteworthy collection of manuscripts at Florence, and in 1414 Poggio Bracciolini, a secretary of the Roman curia, began the copying and collecting of classical works. During the century there were such munificent patrons of learning as the Medici in Florence and Nicholas V in Rome, and the results of their labors are still seen in the Laurentian and Vatican libraries. There was also Federigo, Count of Montefeltro (1422–1482), who had been filled with enthusiasm by no less a teacher than Vittorino da Feltre himself, and who received from Sixtus IV the title of Duc d'Urbino. His library at Urbino, filled with manuscripts, was a rendezvous for the scholars of Italy. The humanist Giovanni Aurispa (*c.* 1369–1460) also brought some 238 manuscripts of the ancient Greek writers from Constantinople to Venice.

With all this activity, however, Italy showed no native ability in mathematics in the 15th century. What Symonds said of Tuscan culture in general applies in particular to mathematics: "Florence borrowed her light from Athens, as the moon shines with rays reflected from the sun. The revival was the silver age of that old golden age of Greece."

It is evident to one who studies the arithmetics of this century that no national spirit had yet developed. It was the city, not the state as we know it, to which men gave their allegiance. Just as we have Venetian art, so we find Venetian and Florentine and Roman arithmetics, quite as distinct from one another as were the Pisan works from those of Nürnberg.

[1] See J. B. Bury, in the *Cambridge Modern History*, Vol. I, chap. iii (London, 1902). This important work, containing different essays, is hereafter referred to as *Cambridge Mod. Hist.*

Origin of Printing. When we consider the effect of printing upon the development of mathematics, we must recall the fact that the art existed long before Gutenberg's time, about 1450. Not only had the Chinese printed from engraved blocks many centuries earlier than this, but they had also made some use of movable types. In Europe, too, block printing had assumed considerable prominence before movable types were invented.

It is also well to consider how mathematical knowledge was disseminated before it was possible to send it abroad by means of the printed page. There were three general periods in the transmission of such knowledge in the Middle Ages:

1. From the founding of the monastery at Monte Cassino by St. Benedict in 529, and continuing until *c.* 1200, the period in which scholars went to the teacher in the monastery and heard his lectures. This was the period of such men as the Venerable Bede and Alcuin of York.

2. From *c.* 1200 to *c.* 1400, when universities appeared in considerable numbers and claimed control of the scribes and booksellers.[1]

3. From *c.* 1400 to *c.* 1460, when the manuscript trade became more general, when large numbers of copies were made, and when these were sold as we sell books today.

All this was, however, a crude way of disseminating knowledge compared with the circulation of printed books. We should therefore expect the 15th century to begin to make widely known the mathematical classics of the ancient civilization, to meet the mathematical needs of a commerce to which new worlds were opening, and to prepare popular summaries

[1] The scribes had stands (stations) and the booksellers had shops near the university. The bookshops near the Sorbonne in Paris today are relics of the houses of the old *librarii* who, in the 13th and 14th centuries, rented their manuscripts. The *stationarius* was required to employ skilled copyists, who were enjoined to perform their tasks "fideliter et correcte, tractim et distincte, assignando paragraphos, capitales literas, virgulas et puncta, prout sententia requirat." See G. H. Putnam, *Books and their Makers during the Middle Ages*, I, 200 (New York, 1896, 1897); hereafter referred to as Putnam, *Books*. On the whole subject of reproduction and sale see J. A. Symonds, *Renaissance in Italy: The Age of Despots*, II, 120 (New York, 1883).

of the mathematics already accumulated. It would hardly be expected that this century would do more than receive these accumulated treasures and transmit them, postponing any new advance in science until the century following. It should be added that the sack of Mainz by Adolf of Nassau (1462) scattered the printers of that city all over Europe, an event comparable in some respects to the results of the fall of Constantinople.

The Universities. The universities were now beginning to mention mathematics more commonly, but that was about all. In the fragments of the Oxford statutes of 1408, only a little arithmetic was required for the bachelor's examination,[1] and in 1431 there was hardly any improvement.[2] In Paris the work was reorganized by the Papal legate in 1452, but nothing was done to better the condition of this science. The baccalaureate demanded only the reading of a little mathematics,[3] but nothing definite was prescribed. In general, little was required beyond arithmetic and a few pages of Euclid.[4]

Italian Writers. The great commercial activity in Italy in the 15th century gave rise to a large number of mercantile arithmetics, and these set a standard for the treatment of the subject that still influences, in some degree, the textbooks of today. Among the best known of the commercial writers were Matteo da Firenze[5] (*c.* 1400), his son Luca da Firenze (*c.* 1400), Giovanni, the son of this Luca (*c.* 1422), and Andrea di Giovanni Battista Lanfreducci (*c.* 1490), an officer of the

[1] "Algorismus integrorum" and "computus ecclesiasticus."


[2] For the licentiate, "Arithmeticam per terminum anni, videlicet Boëthii," with a little more science and mathematics. Suter, *Univ. Mittelalt.*, p. 90.

[3] "Aliqui libri mathematici."

[4] A MS. of 1515 in the Wolfenbüttler Bibliothek gives the following as representing the arithmetic: "Arithmetica communis ex divi Severini Boetii Arithmetica per M. Ioannem de Muris compendiose excerpta; Tractatus brevis proportionum, abbreviatus ex libro de proportionibus D. Thomae Braguardini Anglici, . . . Algorithmus M. Georgii Peurbachii de integris; Tractatus de Minutiis phisicis compositus Viennae Austriae per M. Ioannem de Gmunden." *Monatsberichte der K. P. Akad. d. Wissensch. zu Berlin*, 1867 (Berlin, 1868), p. 43.

[5] Matthew of Florence. See *Rara Arithmetica*, p. 468.

Republic of Pisa in 1505. The works of these four writers exist only in manuscript and contain little of value except what they tell of the commercial customs of their time.



FIRENZE'S WORK, 1422

This portion relates to the calendar. The MS. contains the statement: "questo d dottobre 1422" and the further date "ad 28 dottobre 1422"

Among the illustrious citizens of Florence one of the 16th century historians[1] mentions a mathematician named Benedetto, commonly known as Benedetto da Firenze. His arithmetic[2] was written *c.* 1460, and relates to the mercantile needs of his native city. It is one of the most complete works of this kind that appeared in the 15th century, but it has never been printed.

Of the Italian writers of this period who had some knowledge of mathematics beyond the field of commercial arithmetic the earliest was Prosdocimo de' Beldamandi,[3] a native of Padua and a student and professor in the university in that city (1422–1428). He wrote on arithmetic, music, and astronomy,

[1] U. Verino, *De illustratione urbis Florentiae libri tres.* Paris, 1583.

[2] "Inchomincia el trattato darismetricha," as in the MS. in the library of Mr. Plimpton. See *Rara Arithmetica*, p. 464.

[3] Born at Padua, *c.* 1370–1380; died at Padua, 1428. The name also appears as Beldomandi, Boldomondo, Beldemandi, and Prosdocimo Padvano. Bernardino Scardeone (*De Antiqvitate Vrbis Patavii*, Basel, 1560) speaks of him as "è nobili familia Patauina ortus: egregius Musicus, & eximius philosophus, & clarus astrologus."

and also studied medicine.[1] The arithmetic was limited to the ordinary operations and contained no commercial problems.[2]

About 1425 one Leonardo of Cremona, or Leonardo de' Antonii (born *c.* 1380), wrote a brief practical geometry of which three manuscripts are known.[3] A little later (*c.* 1449) a fellow townsman of his, Jacob of Cremona, a teacher at Mantua and Rome, translated some of the works of Archimedes. Towards the close of the century Georgius Valla (1430–1499), a native of Piacenza, lectured on physics and medicine at Pavia and also at Venice, and in his *magnum opus*, which was merely a compendium of knowledge, treated of Boethian arithmetic, Euclidean geometry, optics, and the astrolabe, as well as a variety of other subjects. The work[4] was printed in 1501, but is notable chiefly for its size.

About 1475 an Italian painter, Pietro Franceschi, also called Pietro della Francesca,[5] wrote a work, *De corporibus regularibus*, in which he treated of the mensuration of regular polygons, the sphere, the five regular polyhedrons inscribed in a sphere, and solid figures in general. His problems may be illustrated by the cases of finding the area of a regular octagon circumscribed by a circle of diameter 7,[6] and the finding of the surface of a cube circumscribed about a sphere also of diameter 7.[7]

[1] This on authority of two early MSS.: "Examen medicinae magistri Prosdocimi de padua M° iiij° undecimo," that is, 1411; "Padovani dottori delle arti e Medicina," with the assertion that he "fu esaminato e dottorato nelle arte . . . 1409."

[2] For description, see *Rara Arithmetica*, p. 13; Boncompagni's *Bullettino*, XII, 1.

[3] *Leonardi Cremonensis artis metrice practice compilatio. Primus tractatus.* The date is doubtful. See *Rara Arithmetica*, p. 474; *Bibl. Math.*, IX (3), 280.


[4] *Georgii Vallae Placentini viri clariss. de expetendis, et fvgiendis rebus opus.*

[5] Born *c.* 1410–1420; died 1492. See E. Harzen, "Ueber den Maler Pietro degli Franceschi und seinen vermeintlichen Plagiarius, den Franziskanermönch Luca Pacioli," *Archiv f. d. zeichn. Künste*, II, 231; W. G. Waters, *Piero della Francesca*, London, 1901; and Mancini's monograph mentioned below.

[6] "Diameter circuli qui circumscribit octagonum est 7. Quanta igitur sit octagoni superficies invenire." Tract. I, Problem XL. He uses $\frac{22}{7}$ for π.

[7] Tract. II, Problem XVI. The work is printed in full in G. Mancini, "L'opera 'De corporibus regularibus' di Pietro Franceschi," in the *Atti d. R. Accademia dei Lincei*, XIV (5), 488.

Printing was introduced into Italy in 1464 by Juan Turrecremata,[1] abbot of the monastery of Subiaco, near Rome, the



FIRST PAGE OF A MANUSCRIPT OF LUCA DA FIRENZE'S WORK (*c.* 1400)
This MS. dates from *c.* 1475. Now in the library of Mr. Plimpton

first book appearing in the following year.[2] Fourteen years later (1478) the first printed arithmetic appeared at Treviso,

[1] The Latin form of the family name Torquemada. He was a native of Valladolid, in Spain.

[2] Putnam, *Books*, I, 405.

then an important commercial town a day's travel from Venice. Three presses had already been established there, and it was from the one of Manzolo, or Manzolino,[1] that this work appeared. That it was anonymous was merely in keeping with the custom of a land where the glory of the individual was absorbed in the glory of the state.[2] Boncompagni learned of only eight copies of this arithmetic in various degrees of preservation,[3] so that the book is rare. It is quite commercial and never had any noticeable influence on the arithmetics of Italy.[4] It was, however, the first step in a remarkable movement, for up to the close of the 15th century there were printed in Italy at least 214 mathematical works, the number rising to 1527 in the following century.[5]

Three years after the Treviso book appeared, the first purely commercial arithmetic[6] was published. The author seems to have been one Giorgio Chiarino, of whom nothing further is known. The work is a mere compilation of measures and of such customs of exchange as were needed by Florentine merchants,[7] but there was enough in it to lead Pacioli (1494) to borrow freely from it.

Three years later Piero Borghi,[8] a Venetian, published the most noteworthy Italian commercial arithmetic of the century,[9]

[1] Domenico Maria Federici, *Memorie Trevigiane*, p. 73. Venice, 1805.

[2] "When Byron swept with superficial yet brilliant eyes the rolls of Venetian history, what did he find for the uses of his verse? Nothing but two old men, one condemned for his own fault, the other for his son's,—remarkable chiefly for their misfortunes." Mrs. Oliphant, in *The Makers of Venice*.

[3] *Atti Pontif.*, Vol. XVI. The only copy of this arithmetic that ever reached America is in the library of Mr. Plimpton.

[4] For a description, see *Rara Arithmetica*, p. 3.

[5] P. Riccardi, *Biblioteca Matematica Italiana*, Parte seconda, XI, XV, seq. (Modena, 1880), hereafter referred to as Riccardi, *Bibl. Mat. Ital.*


[6] *Qvesto e ellibro che tracta di Mercatantie et vsanze de paesi* (Florence, 1481). See *Rara Arithmetica*, p. 10.

[7] "Finito ellibro de tvcti ichostvmi: cambi: monete: pesi: misvre: & vsanze di lectere di cambi: & termini di decte lectere che nepaesi si costvma et in diverse terre."

[8] Piero Borgi, as the name appears in the first edition. The name is also given as Pietro Borghi and Pietro Borgo. He seems to have died after 1494.

[9] *Qui comenza la nobel opera de arithmethica*, the earliest printed books generally having no title pages. In later editions this was sometimes called the *Libro de Abacho*. The first edition was Venice, 1484.

a work intended for the mercantile class of Venice. One of the features of the book is that, like certain others of the



FIRST EDITION OF EUCLID, 1482

First page, reduced, from the Venice edition of 1482

period, it begins with multiplication, putting addition and subtraction after division and therefore presupposing some ability in computation. The work with integers is followed by

fractions, the Rule of Three, and the usual applications of the time,—partnership, profit and loss, barter, and alligation.

These works, however, were mere advance couriers. Arithmetics had been published in Germany and Italy, Euclid's *Elements* had appeared from the famous press of Ratdolt, in Venice,[1] and the psychological moment had arrived for a general treatise summarizing the mathematical knowledge of the time. In such a period of unrest as that centering about the year 1494, when France was at war with most of Italy, when Florence was in arms against her sister towns, when Charles VIII was invading the peninsula and Savonarola was proceeding as an ambassador to Pisa to allay his wrath,—in such a period it would be surmised that a book like this could be prepared only in the peaceful atmosphere of a cloister.

Luca Pacioli. Such was the case, and Luca Pacioli,[2] known from his birthplace as Luca di Borgo, was the author. As a boy he may have come under the influence of his townsman, the artist Pietro Franceschi, already mentioned (p. 247), from whose work he freely took considerable material. At about the age of twenty he went to Venice (1464) and became a tutor to the three sons of a wealthy merchant,[3] and some six years

[1] *Preclarissimus liber elementorum Euclidis*, Venice, 1482. See the facsimile.

[2] Born at Borgo San Sepolcro, Tuscany, *c.* 1445; died probably after 1509. The name is spelled in various other ways, such as Paciolo, Paciolus, and Paciuolo. It does not appear at all in his *Sūma*, and in the *De diuina proportione* (Venice, 1509) it is given only in the Latin genitive,—"Lucae pacioli ex Burgo sancti Sepulchri . . . epistola." In his "Supplica . . . al Doge di Venezia" of December 29, 1508, in which he asks for permission to print the *De diuina proportione*, the name appears as "Luca de pacioli dal borgo sā sepulchro." It is from such contemporary evidence that Boncompagni (*Bullettino*, XII, 420) was led to speak of him as Pacioli, although Bernardino Baldi, writing in 1589, says "Fu de la famiglia de Paciuoli ignobile per quanto mi credo e di poco splendore," which led Cantor to adopt this later spelling. For the contemporary documents, in which he is generally spoken of as Luca or Lucas di Borgo, see Boncompagni, *loc. cit.*

The best sketch of his life is that by H. Staigmüller, "Lucas Paciuolo. Eine biographische Skizze," in the *Zeitschrift* (Hl. Abt.), XXXIV, 81, 121. See also the "Elogio di Fra Luca Pacioli" in B. Boncompagni, *Scritti inediti del P. D. Pietro Cossali* (Rome, 1857), p. 63; B. Boncompagni, in his *Bullettino*, XII, 377.

[3] " . . . nostri releuati discipuli ser Bart°. e francesco e paulo fratelli derōpiasi da la çudeca: degni mercatanti in vinegia: figliuoli gia de ser Antonio." *Sūma*, 1494 ed., fol. 67, *v°*, l. 4.

later he wrote an algebra which he never published.[1] In 1471 he went to Rome and, possibly influenced by his two brothers who had already entered the brotherhood of St. Francis, joined the Minorite order. In 1476 we find him teaching in Perugia and writing a little book for his pupils.[2] Five years later, at Zara, he wrote still another work, "more subtile and rigid,"[3] but neither of these works was printed. He traveled extensively in Italy[4] and possibly in the Orient,[5] and even after he became a Franciscan he was a wanderer. We find him back in Perugia in 1487 and working on his *Sūma*, in Naples in 1494, in Milan in 1496, in Florence and Rome in 1500, and in Venice in 1508.[6]

The great work of Pacioli, summing up not only his previous and unpublished works but also the general mathematical knowledge of the time, appeared in Venice in 1494,[7] having been written seven years earlier, when he was at Perugia.[8] It is a remarkable compilation with almost no originality.[9] He borrowed freely from various sources, often without giving the slightest credit, but in this he merely followed

[1] As to his MSS., now in the Vatican, see G. Eneström, *Bibl. Math.*, XIII (2), 53; B. Boncompagni, in his *Bullettino*, XII, 381, 428.

[2] ". . . alo giouani de peroscia . . . nel. 1476," as he says in the *Sūma* of 1494, fol. 67, *v*.

[3] "E anche in quello che a çara nel. 1481. de casi piu sutili e forti componēmo." *Ibid.*

[4] "Ma da poi che labito indegnamente del seraphyco san francesco ex voto pigliamo: p. diuersi paesi ce conuenuto andare peregrinando." *Ibid.*

[5] So Cossali, writing in the 18th century, says that he "per desio di scienza viaggiasse in Oriente, ed in Arabia precipuamente, dove a que' tempi erano le matematiche dottrine in gran fiore"; but there is no contemporary authority for the statement. Certainly the expression "in gran fiore" is without foundation. On the contrary, there is internal evidence from the *Sūma* that he was never in the East.

[6] See also P. Treutlein, *Abhandlungen*, I, 10.


[7] For the title-page, see page 253. The form *Sūma* is a contraction for *Summa*.

[8] "E al presente q̄ ī peroscia . . . correndo glianni del nostro segnore Jesu Christo. 1487." *Sūma*, 1494 ed., fol. 67, *v*.

[9] G. Mancini, "L'opera 'De corporibus regularibus' di Pietro Franceschi," *Atti d. R. Accademia dei Lincei*, XIV (5), 488. See also Chiarino's work, from which he did not hesitate to take what he wished, and Baldi's article in Boncompagni's *Bullettino*, XII, 426.

the custom of the age. His use of material from Euclid, Ptolemy, Boethius, Fibonacci, Jordanus, Sacrobosco, Biagio of Parma, Prosdocimo, and Suiceth shows that he was at any rate rather well-read. He was, however, a careless writer, so much so that Cardan[1] has a chapter devoted to errors in his book.[2] The work includes such algebra as was then known, a large amount of business arithmetic, a poor summary of Euclid, and a treatment of double-entry bookkeeping.[3]

In 1497, while at Milan, he wrote his *De diuina proportione*, publishing it at Venice in 1509.[4] This is a work of more merit from the standpoint of



Sūma de Arithmetica Geometria Proportioni & Proportionalita.

Continentia de tutta lopera.

De numeri e misure in tutti modi occorrenti.
Proportioni e pportiōalita anotitia del. 5° de Eucli de e de tutti li altri soi libri.
Chiaui ouero euidentie numero. 13. p le q̄nta continue pportiōali del. 6°e. 7°. de Euclide extratte.
Tutte le pti delalgorismo: cioe relevare. petir. multiplicar. sūmare. e sotrare cō tutte sue pue i sani e rotti. e radici e progressioni.
De la regola mercantesca ditta del. 3. e soi fōdamenti con casi exemplari per c°m°. & . G . guadagni: perdite: transportationi: e inuestite.
Partir. multiplicar. summar. e sotrar de le proportioni e de tutte sorti radici.
De le. 3. regole del catayn ditta positiōe e sua origie.
Euidentie generali ouer conclusioni n°66. absoluere ogni caso che per regole ordinarie nō si podesse.
Tutte sorte binomii e recisi e altre linee irratiōali del decimo de Euclide.
Tutte regole de algebra ditte de la cosa e loz fabriche e fondamenti.
Compagnie i tutti modi. e loz partire.
Socide de bestiami. e loz partire
Fitti: pesciōi: cottimi: liuelli: logagioni: e godimenti.
Baratti i tutti modi semplici: compositi: e col tempo.
Cambi reali. secchi. fittitii. e di minuti ouer comuni.

PART OF THE TITLE-PAGE OF PACIOLI'S WORK OF 1494

This shows the general nature of the treatise, forming a kind of table of contents

[1] In his *Arithmetica* of 1539.

[2] "De erroribus F. Lucae quos vel transferendo non diligenter examinavit, vel describendo per incuriam praeteriit, vel inveniendo deceptus est." See also Cossali, "Elogio di Fra Luca Pacioli," in the *Scritti inediti*, p. 63 (published by B. Boncompagni, Rome, 1857). On the variations in each of the two editions (1494 and 1523) see E. Narducci, *Intorno a due edizioni della Summa de Arithmetica*, Rome, 1863.

[3] On this phase of the work see V. Gitti, *Gli Scrittori Classici della Partita Doppia*, a reprint from the *Annali del R. Istituto Indust. e Professionale di Torino*, Vol. V. (Turin, 1877). [4] German translation by C. Winterberg, Vienna, 1896.

geometry than the *Sūma*, but one that could not, in the nature of the case, be as popular. His figures of the regular solids, as given in this book, were the best that had as yet appeared in print,[1] and have been attributed to Leonardo da Vinci. He also published an edition of Euclid in 1509, but the work is of little merit.




TITLE-PAGE OF CALANDRI'S ARITHMETIC, 1491

An example of the fanciful portraits of the Greek mathematicians. Such portraits became common during the Renaissance. The anachronisms in this one are evident

Of the Italian astronomers of the 15th century who gave considerable attention to mathematics the most eminent were Giovanni Bianchini (*c.* 1450), professor of astronomy in the University of Padua, Francesco Capuano, Giovanni Batista de Manfredonia (*c.* 1450–1490), and Domenico Maria Novara da Ferrara (1454–1504). Italian science, however, had hardly yet awakened to the need for more advanced mathematics.

[1] Pacioli here takes his material freely from Franceschi's work, already mentioned.

Of the arithmeticians of this time one of the most noteworthy was Filippo Calandri,[1] whose arithmetic appeared in Florence in 1491 and contained the first printed example in long division by our modern method and the first illustrated problems published in Italy. Nothing further is known of his life, but Sfortunati,[2] writing in 1534, speaks of him as a learned man.[3]



FIRST ILLUSTRATED PROBLEMS
From Calandri's work, Florence, 1491

Francesco Pellos or Pellizzati, a native of Nice, published a commercial arithmetic at Turin in 1492, in which, as will be shown in Volume II, use is made of a decimal point to denote the division of a number by a power of ten.

Austria and Germany. Before considering the names of those who advanced the cause of mathematics in the German states in the 15th century it is necessary to say a word concerning the influence and work of the Rechenmeisters. In the 13th century there had been a great revival of trade in Germany. The Hanseatic League, a union of commercial towns in the Teutonic countries, had shown that the demands of trade must be recognized. It employed force against the pirates,

[1] Italian writers of the period give this form of the name. Latin writers use Philippus Calender and Philippus Calandrus. Calandri himself gives only the Latin genitive form, Philippi Calandri. See *Rara Arithmetica*, p. 47.

[2] Giovanni Sfortunati (born at Siena, *c.* 1500), *Nvovo Lvme Libro di Arithmetica*, Venice, 1534. See page 306.

[3] "Filippo Calādri Cittadino Fiorentino, huomo certamēte in tale disciplina erudito." From the 1545 edition, fol. 3, *r.*

purchased settlements in foreign cities, and even made war upon Denmark and England in order to protect the interests of its members. The Church schools having failed to prepare boys for business, the League undertook this work, offering not only reading and writing, in which the Church schools gave instruction, but also business arithmetic, in which they gave none. Out of this kind of teaching of apprentices there arose a type of commercial school known as the Rechenschule, presided over by a Rechenmeister. In due time the Rechenmeisters formed guilds, claimed a monopoly in their vocation, and finally came to be looked upon as regular officials of the town. Not infrequently their duties included the sealing of measures, the city gaging of casks, and occasionally the minting of money, in those times a function of the city instead of the state. Occasionally they were also the writing masters, and they finally came to be looked upon as among the dignitaries of the town.[1] In the larger cities the Rechenmeisters' guilds admitted apprentices who served for six years, then becoming Schreibers (writers) with the privilege of becoming assistant teachers. When a *locus* opened, the eldest of these cadets was subjected to an examination[2] and, if successful, was given the position with the rank of Meister. The guild of Schreib- und Rechenmeisters continued in Lübeck, for example, until 1813.

In the 15th century the power of the Rechenmeisters was first shown to any considerable extent, and for the next three hundred years they clung more or less tenaciously to their privileges, demanding that arithmetic should be taught by them instead of by the common schoolmasters. Their influence extended to the commercial towns of Holland, and the Rechenmeister is

[1] Thus, the Bürgermeister and Council of Rostock, in 1627, sent an official call, couched in dignified language, as follows: "Wir Bürgermeister und Rat zu Rostock urkunden hiermit, dass wir den ehrenfesten und wohlgelahrten Jeremias Bernstertz zu unserm und gemeiner Stadt Schreib- und Rechenmeister bestellt und angenommen haben." For the entire document see F. Unger, *Die Methodik der praktischen Arithmetik in historischer Entwickelung*, p. 26 (Leipzig, 1888); hereafter referred to as Unger, *Die Methodik*.

[2] One of the questions from an old paper reads: "Wie wird besagte Chilioheptacosioheptacontatetragonal-Zahl, deren Latus 6 formirt und aus solcher gefundenen Polygonal-Zahl die Wurzel wieder extrahirt?"

frequently mentioned in the Dutch arithmetics of the 17th century.[1] The excellent commercial arithmetics which began to appear in Germany in the 15th century and which continued there and in Holland until the 19th century were the work of the Rechenmeisters or of those whom they influenced.[2]

Owing possibly to the early advance in printing in Germany, possibly to the trend of the classical influence from Constantinople through the Balkan states to Vienna, or possibly to the influence of the leisure which wealth afforded, the Teutonic countries forged ahead in the 15th century, taking rank with Italy in their men of mathematical ability. Four of these men were scholars whose standing was recognized abroad; the rest were mediocrities.

Late in the century Johann Widman (born *c.* 1460), a native of Eger in Bohemia, wrote on arithmetic and algebra. He was a student at Leipzig in 1480, B.A. in 1482, M.B. in 1485, and M.A. in 1486. He may have received a doctor's degree in medicine, for a medical work by a Johann Widman appeared in 1497, but this was probably another person.[3] That he gave lectures on algebra, possibly the first given in Leipzig, is shown by a contemporary manuscript.[4] He was probably the

[1] Thus, Coutereels (1631) addresses a problem to one Ghileyn Pietersz "Schepen [sheriff] deser Stadt, goet Reken-meester"; Cardinael (1659) speaks of himself as "Reecken-meester tot Amsterdam"; the printer of Vander Schuere's arithmetic (1634 ed.) speaks of the book "die ik door de correctie van een goet Reken-Mr. het verbetert"; Eversdyck, who revised Coutereels's arithmetic in 1658, is described as the "Reken-meester ter Reken-Kamer van Zeelandt"; and even as late as 1792 an edition of Bartjens is described as due to "den Wel-ervaren Reekenmeester Klaas Bosch."

[2] The Italian "master of the abacus" was probably suggested by the German title, or vice versa. The expression is found in Italian manuscripts of this period but does not seem to have been used in England. Indeed, even the word "abacus" never had as extensive use there as on the Continent.

[3] *Tractatus clarissimi medicinaꝙ doctoris Johānis widman.* In other works the spelling Widmann is sometimes used. Boncompagni's *Bullettino*, IX, 210.

[4] See *Rara Arithmetica*, p. 36. J. G. Bajerus (J. W. Bayer), *De Mathematvm . . . introdvctione* (1704) (hereafter referred to as Bajerus, *De Math.*), says (p. 9): "Joh. Widemann, natione Noricus, patria Egrensis, disciplina Lipzensis, vir in Mathematicis abunde eruditus. Qui capessis in Philosophia insigniis, cum multa admodum in mathematica, & potissime in speciebus in studio Lipzensi, non sine auditorum summo applausu, aliquot annis volvisset . . ." On his life and works see Cantor, *Geschichte*, II, chap. lv.

author of an *Algorithmus Linealis* (Leipzig, *post* 1489), the first printed treatise on calculation by the aid of counters. He wrote the first important German textbook on commercial arithmetic,[1] and in this appear for the first time in print the signs + and −, not as symbols of operation but to express excess and deficiency in packages of merchandise.

More capable as a mathematician but less known by the populace, Johann von Gmünden[2] was educated at Vienna, taught there, and was the first Austrian to occupy a chair devoted wholly to mathematics. He wrote a treatise on sexagesimal fractions,[3] one on trigonometry,[4] and one on the computus.[5]

A few years later, Nicholas Cusa,[6] son of a fisherman, gave proof of what industry and genius will do even for those born in humble estate. He rose rapidly in the Church and held various positions of honor, including the bishopric of Brescia. He was made a cardinal and became the governor of Rome in 1448. He wrote several tractates on mathematics, including

[1] *Behēde vnd hubsche Rechnung auf allen kauffmanschafft*, Leipzig, 1480.

[2] Born at Gmunden on the Traunsee, Gemünd in Lower Austria, or Gemünd in Swabia, *c.* 1380; died *c.* 1442. The name also appears in such forms as Johannes de Gmunden, Johann von Gemunden, Johannes de Gamundia. He has been identified with the Johann Schindel or Sczindel, called Johannes de Praga, a native of Königgrätz, and possibly the identification is correct. His name has also been given as Wissbier and Nyden, but on doubtful authority. See *Bibl. Math*, X (2), 4. Bajerus, *De Math*, has this note (p. 6): "Eo vix A. C. MCCCXCVII. fatis functo, Vindobonam ornare coepit Johannes de Gmunden itidem natione Germanus, (Ricciolus Johannem de Egmunda vocat) Theologus & Astronomus celebris, anno MCCCCXLII. humanis exemtus." See also *Bibl. Math.*, III (3), 140.

[3] *Tractatus de Minucijs phisicis*, first printed at Vienna in 1515.

[4] *De sinibus, chordis, et arcubus.*

[5] It exists in manuscript, closing with the words: "Explicit kalendariū mgrī Joh'is gmünd." See *Rara Arithmetica*, p. 440.

[6] Born at Kues (Cues) on the Mosel, 1401; died at Todi, Umbria, 1464. The name appears in such forms as Nicolaus von Cues, Nicholas Cusanus, Nicholas von Cusa, Nicolaus Chrypffs, Nicolaus Cancer, Nicolaus Krebs. His father's name was Johann Chrypffs or Krebs. "Huomo di mostruoso ingegno impatronissi delle tre lingue migliori, e diede opera all' arti liberali, & alle scienze," Baldi, *Cronica*, 95. Rossi, in his *Niccolo di Cusa* (p. 11), describes him as "originale pensatore in molteplici discipline." There is a good biography by Dr. Schanz, Prog., Rottweil, 1872. The best biographies are those of J. M. Düx, *Der deutsche Cardinal Nicolaus von Cusa*, 2 vols., Regensburg, 1847, and E. Vansteenberghe, *Le Cardinal Nicolas de Cues*, Paris, 1921.

in the subjects treated the quadrature of the circle, the reform of the calendar, the improvement of the Alfonsine Tables, the heliocentric theory of the universe (a theory which was looked upon as a paradox rather than a scientific probability), and the theory of numbers. Wallis[1] asserted that he was the first writer known to have worked on the cycloid, but this is not supported by the evidence.[2] His *Opuscula* appeared *c.* 1490, and his *Opera* appeared in Paris in 1511.[3]

Much better known as a mathematician, Georg von Peurbach[4] studied under Nicholas Cusa and other great teachers, learned Greek from Cardinal Bessarion in order to be able to read Ptolemy, lectured at Ferrara, Bologna, and Padua, and became professor of mathematics at Vienna, making this university the mathematical center of his generation. Although interested primarily in astronomy and trigonometry,[5] he wrote an arithmetic,[6] but this was merely for the use of students in these branches of science. Melanchthon considered the work so excellent that he wrote a preface for the edition of 1534. Peurbach compiled a table of sines, which was extended after his death by his pupil Regiomontanus, and he also wrote various works on astronomy.

The most influential and the best known of the German mathematicians of the 15th century was Johann Müller,[7] generally known, from the Latin name of Königsberg, as Regiomontanus.[8] At the age of twelve he was a student at Leipzig. He

[1] *Philosophical Transactions of the Royal Society of London*, p. 561 (1697); hereafter referred to as *Phil. Trans.*

[2] See *Bibl. Math.*, I (2), 8, 13.

[3] *Rara Arithmetica*, p. 42.

[4] Born at Peurbach, Upper Austria, May 30, 1423; died at Vienna, April 8, 1461. The name is also spelled Peuerbach and Purbach.

[5] *Tractatus Georgii Purbachii super Propositiones Ptolemaei de sinubus et chordis*, Nürnberg, 1541; *Theoricae novae planetarvm*, Venice, 1495, with another edition s. l. a. (Nürnberg) and one at Venice, 1490.

[6] *Elementa Arithmetices Algorithmvs de numeris integris.* It went by various other titles. The first printed edition appeared in 1492.

[7] Born at Unfied, near Königsberg, Lower Franconia, June 6, 1436; died July 6, 1476. Cantor, *Geschichte*, II, chap. lv. In the British Museum there is an interesting block-book almanac, not later than 1474, prepared by him, in which his name appears as Magister Johann van Kunsperck.

[8] Also as Joannes de Monteregio.

afterwards studied under Peurbach, lectured at Venice, Rome, Ferrara, and Padua, and lived for a time in Nürnberg.[1] In 1475 he was called to Rome by Pope Sixtus IV on account of one of the frequent attempts to consider a reform of the calendar, and was made titular bishop of Ratisbon. He studied the mathematics of the Greeks in the original, and was "the first who made humanism the handmaid of science." He wrote *De triangulis omnimodis libri V* (*c.* 1464), the first work that may be said to have been devoted solely to trigonometry.[2] He also wrote an *Introductio in Elementa Euclidis* with some supplementary work on stellar polygons, and had certain definite ideas as to the circumnavigation problem.[3]

The earliest known German algorism was compiled in 1445, more than two centuries after Fibonacci prepared a work on the subject, and about seventy years after the first French manuscript relating to it is known to have been written. The earliest example of a German algebra is found in a Munich manuscript of 1461, with text also in Latin. The first printed German arithmetic appeared at Bamberg in 1482.

France. France received the Italian humanism in the spirit of a sister country speaking a kindred language, but for a number of decades after Gregory Tifernas (1458) went to Paris to teach Greek the taste of the learned class was for letters rather than science.[4]

It is perhaps because of this fact that France produced fewer noteworthy mathematicians than Germany or Austria in the 15th century, but it is more probably due to such causes as the constant turmoil in which the University of Paris found herself at this time. With a continual warfare between Church

[1] Ramus speaks of the great glory he brought to Nürnberg: "Noriberga tum Regiomontano fruebatur: mathematici inde & studii & operis gloriam tantam adepta, ut Tarentum Archyta, Syracusae Archimede, Byzantium Proclo, Alexandria Ctesibio non justius quám Noriberga Regiomontano gloriari possit." *Scholarum Mathematicarum libri unus et triginta*, p. 62 (Paris, 1569); hereafter referred to as Ramus, *Schol. Math.*

[2] First printed edition, Nürnberg, 1533.

[3] A. Ziegler, *Regiomontanus . . . Vorläufer des Columbus*, Dresden, 1874.

[4] See also R. C. Jebb, *Cambridge Mod. Hist.*, Vol. I, chap. xvi.

and State over the control of her work, and with continual protests from within her walls, she was in no mood to foster either science or letters.

A type of the best product of the educational system of France at this time is seen in the person of a very mediocre and almost unknown scholar, one Rollandus (*c.* 1424), who, although probably a native of Lisbon, spent his life in Paris. He was a physician and a minor canon of the Royal Chapel[1] and may have been the Rolland who was rector of the university in 1410. He evidently was acquainted with the general field of pure arithmetic and algebra, as is shown by a manuscript of about 1424 still extant.[2] He also wrote on physiognomy and surgery.

The most brilliant of the French mathematicians of the period, however, was Nicolas Chuquet,[3] who, although a native of Paris, lived in Lyons.[4] He wrote (1484) the *Triparty en la Science des Nombres*, a work touching upon the three fields of arithmetic.[5] The first part relates to computation with rational numbers, the second to irrationals, and the third to the theory of equations. Nothing is known about Chuquet except his statement that he was a bachelor of medicine and that he wrote his work at Lyons.[6]

Great Britain. Great Britain produced relatively few writers on mathematics in the 15th century, and none of any special prominence. John Killingworth may be taken as a type of

[1] He speaks of himself as "prebenda capelle palacij regalis parisiensis."

[2] *Scientia de numero ac virtute numeri*, now in the library of Mr. Plimpton. See *Rara Arithmetica*, p. 446.

[3] Born at Paris; died *c.* 1500.

[4] Estienne de la Roche (1520), speaking of the "plusieurs maistres expertz en cest art," mentions "maistre nicolas chuquet parisien."

[5] It was printed in Boncompagni's *Bullettino*, XIII, 555. See Ch. Lambo, "Une algèbre Française de 1484. Nicolas Chuquet," *Revue des Questions Scientifiques* (Brussels, October, 1902).

[6] "Et aussi pour cause quil a este fait par Nicolas chuquet parisien Bachelier en medicine. Je le nomme le triparty de Nicolas en la science des nombres Lequel fut commance medie et finy a lyon sus les rosne de salut. 1484." From the original copy of his MS. made for Boncompagni by A. Marre, and now in the author's library.

the best scholars of the period. We know little about him,[1] but the records show that he became a fellow of Merton College, Oxford, in 1432 and that he died on May 15, 1445. He seems to have been chiefly interested in astronomy and to have prepared a set of tables for the use of students in this science. There is extant in the Cambridge University Library an algorism written by him in 1444.[2] In this he refers to the use of a slate for purposes of computation, but as to the operations themselves there is no evidence of any originality.

Other Countries. The early mathematics of Russia was largely devoted to questions relating to the calendar and to number puzzles. In some of the medieval manuscripts there are the results of very complicated computations, but we have no knowledge of the methods by which these computations were performed.[3] Several such cases are found in the *Russkaya Pravda*.[4] At the end of the 15th century, work on the calendar is known to have been done by the Metropolitan, Zosima, and by Gennadi, bishop of Novgorod, but the results are not extant. In all this work the numerals seem to have been alphabetic, the system being similar to that of the later Greeks. Even as early as the 14th century, the clergy placed geometry and astronomy under the ban, and not until the 17th century was there any opportunity for the study of mathematics.[5] Even when, in the 18th century, the attempt was made to inaugurate scientific work, all the leaders in mathematics were brought in from abroad.

[1] L. C. Karpinski, "The Algorism of John Killingworth," *English Historical Review* (1914), p. 707.

[2] "Incipit prohemium in Algorismum Magistri Iohannis Kyllyng Worth." The work itself begins: "Obliuioni raro traduntur que certo conuertuntur ordine. Regulas igitur et tabulas ad breua computationem operis calculandi vtiles in formam certam secundum ordinem specierum Algorismi curabo redigere." *Ibid.*, p. 713.

[3] A. N. Peepin, *History of Literature* (in Russian) (Petrograd, 1911 ed.), I, 253.

[4] A work of which there are three versions, one of the 11th century, one of the 12th, and one of the 13th.

[5] V. Bobynin, "De l'étude sur l'Histoire des math. en Russie," in *Bibl. Math.*, II (2), 103.

In the 15th century (*c.* 1450) a Cretan writer commonly known as George of Trebizond[1] (1396–1486) made a new translation of the *Almagest* into Latin and also translated some of Theon's commentaries upon it. He was a quarrelsome man, of little honor to science, to letters, or to manners.

In Hungary there was a certain Georgius de Hungaria who wrote an *Arithmeticae summa tripartita* in 1499,[2] and this perhaps shows the high mark of mathematics in that country in the 15th century.

The Jewish activities in this century were very slight, being chiefly manifest in translations from the Latin. Jacob Caphanton (died by 1439), probably a native of Castile, a physician and a teacher, wrote an arithmetic;[3] and Jehuda Verga (*c.* 1450), known for his compilation on the calamities of the Jews, was also the author of a compendium of the same type.[4] There were various writers who left works on astrology and the calendar, contributions unimportant in themselves but requiring the computation of tables and offering some encouragement to the astronomer.

Sporadic efforts were made here and there in the 15th century to advance the study of mathematics in other countries, but with no effect beyond the establishing of an interest in the science. Thus João (John) II, who was with great difficulty placed (1481) upon the throne of Portugal, sought to elevate scholarship, and particularly astronomy and navigation, by establishing at Lisbon his Junta dos Mathematicos,[5] but the result was not noticeable in pure mathematics. A few scholars, such as the bishops Calsadilha and Don Diogo Ortiz, sought to advance the applications of mathematics in such fields as map drawing, for the purpose of aiding the Portuguese navigators, but this was about all that was done. A little later there were such astrologers as Diogo Mendes Vizinho and

[1] Georgios Trapezuntios.

[2] It was reprinted at Budapest in 1804.

[3] *Bar Noten Ta'am le-Chacham* (a Talmudic phrase). See *Bibl. Math.*, XIII (2), 99.

[4] *Bibl. Math.*, II (3), 62.

[5] A. Marre, Boncompagni's *Bullettino*, XIII, 560; Bensaude, *Astron. Portug.*, 104; R. Guimarães, *Les Mathématiques en Portugal*, 2d ed., p. 11 (Coimbra, 1909) (hereafter referred to as Guimarães, *Math. Portug.*).

Thomaz Torres, but the former knew little beyond the elements of cartography, and the latter seems to have been interested only in the drawing of horoscopes. It was not until the 16th century that mathematics commanded any noteworthy attention in the Portuguese universities.

Some idea of the new Italian spirit in mathematics and letters succeeded in reaching Spain at about the same time. Men like Barbosa lectured on Greek at Salamanca, and Lebrixa (Nebrissensis) returned from Italy in 1474 and lectured at Seville, Salamanca, and Alcalá, but the interests of the learned were in medieval theology and little attention was given to the advancement of the sciences.

The nature of the Spanish works of the 15th century may be inferred from the *Visiõ delectable de la philosophia & artes liberales,* a kind of general encyclopedia, written by Alonso Delatore and published at Tolosa in 1489 and again at Seville in 1538. The fourth chapter of this work is entitled *Dela arismethica y de sus inuẽtores,* but it has no merit and is little better than the treatment of arithmetic given by Capella and Isidorus several centuries earlier. Spain was at this time occupied in suppressing the bandit nobles, in recodifying her laws, and in banishing the last of the Moors, and it was not until 1492 that the political unification necessary to future peace was effected. Nevertheless the foundations seemed in process of being laid for a new type of commercial mathematics and for the rapid development of the science of navigation. Such voyages as those of Columbus and, in the following century, of Ponce de Leon, led to an expansion of the colonial empire and to a rapid increase in wealth. Unfortunately, however, the Spanish leaders were unable to grasp their opportunity, and they squandered their newly acquired possessions in ambitious schemes that led to profitless wars. The chance which the Medici so happily improved was cast aside by the rulers of Spain, and neither art nor science was fostered. During most of the century, therefore, the situation was such as to dampen any scientific zeal, and not a single mathematical work of any consequence was printed in Spain in the 15th century.

TOPICS FOR DISCUSSION

1. Influences at work in the 11th century to improve the intellectual status of Europe.

2. The life and works of Gerbert, who became Pope Sylvester II.

3. A study of the number game of Rithmomachia.

4. The 12th century as a period of translations.

5. Oriental civilization in Spain in the 11th and 12th centuries.

6. Jewish activity in mathematics in the 11th and 12th centuries.

7. Influences tending to foster mathematics in the 13th century.

8. The life and works of Leonardo Fibonacci.

9. The translations of Euclid and their influence on mathematics in the Middle Ages.

10. The life and works of Johannes Sacrobosco and his influence on mathematics in general and on algorism in particular.

11. Roger Bacon; his contemporaries and his projected reforms.

12. The life of Jordanus Nemorarius and his influence on medieval mathematics in general and on algebra in particular.

13. Influences at work in the 14th century to improve the intellectual status of Europe, particularly in the field of mathematics.

14. The mathematics of England in the 14th century. In what ways did the science meet the needs of the time?

15. A comparison of the mathematics of England, France, and Italy in the 14th century.

16. The various methods of diffusing knowledge among scholars during the Middle Ages.

17. The effect upon mathematics of the influx of Greek manuscripts into Italy in the 15th century.

18. Influences leading to the Renaissance, and their bearing upon the development of mathematics.

19. The general nature of the mathematics of Italy in the 15th century, and its effect upon modern education.

20. Influence of the universities in the 15th century, particularly on the development of mathematics.

21. The leading printed mathematical works of the 15th century.

22. Some of the probable reasons why mathematics made such a slight advance in the 15th century.

23. Reasons for the prominence of commercial arithmetics in the latter half of the 15th century.

CHAPTER VII

THE ORIENT FROM 1000 TO 1500

1. China

The General Period. The most interesting period in the history of Chinese mathematics, and indeed one of the most interesting periods in the history of the world, is that of the five centuries from the year 1000 to the year 1500. It is a period not unlike the contemporary epoch in Europe, a time of awakening from sleep. Europe had seen the glories of the classical period die out; she had passed through a season of darkness; she was awakened by the crusades; she arose with a great feeling of refreshment in the 13th century, and in the 15th century she opened a new world of thought through the invention of printing from movable types, and a new world of commercial activity through the discovery of the Western Hemisphere.

China passed through a similar experience, making great advances in algebra in the 13th century and learning of a new civilization through the work of the Jesuit missionaries three hundred years later.

The Eleventh Century. The 11th century saw little progress made in mathematics in China. Indeed, the only name of any importance to be found in her annals is that of Ch'ön Huo (1011–1075), president of the Bureau of Astronomy, a minister of state, and the author of a work in which there is found, perhaps for the first time in China, the summation of a series of any difficulty. This summation appears in the solution of a problem on the number of wine kegs in a pile whose form is a truncated pyramid, the upper row containing 2^2 kegs, the lower one 12^2, and there being 11 rows. The author considers

also the question of finding the length of a circular arc in terms of the radius, and speaks of the difficulty of the problem.

In 1083 the government showed its interest in mathematics by printing Liu Hui's (*c.* 250) *Sea Island Classic*,[1] and a year later (1084) it printed the arithmetic of Ch'ang K'iu-kien[2] (*c.* 575).

The Twelfth Century. The 12th century was about as barren as its predecessor. The *Huang-ti K'iu-ch'ang*[3] seems to have been printed (*c.* 1115), and Ts'ai Yüan-ting[4] wrote on the *I-king*, but these events simply emphasize the fact that the century was barren of achievement. What is much more significant is the fact that the East was again coming in contact with the West; commerce was exchanging the problems of business, and the astrologer was doing the same for the higher and more mysterious strata of mathematical knowledge. We know that in this century metals were transported from Arabia to China, and in 1178 a Chinese work[5] was written in which a great export trade is described and a list of the merchandise is given. In 1128 an immense army from China invaded Turkestan and found there many Chinese residents.[6] The stimulus of world intercourse was becoming stronger, and the effect of this stimulus was to appear in the century following.

Exchange of Thought in the Thirteenth Century. The increase of opportunity for the exchange of thought in the 12th century was carried over into the 13th century, quite as the peripatetic scholars in Europe increased in number and in activity in the same period. Metal was imported from Persia (Po-sz') and from the country of the Arabs (Ta-shi).[7] In 1219

[1] The *Hai-tau Suan-king* (see page 142). This was, of course, from blocks, not from movable type.

[2] The *Ch'ang K'iu-kien Suan-king* (see page 150).

[3] "Yellow Emperor's Nine Sections." One of the versions of the *K'iu-ch'ang Suan-shu* (see page 31).

[4] Born 1135; died 1198; student and historian (see Giles, *loc. cit.*, No. 1985).

[5] The *Ling-wai-tai-ta* of Chóu K'ü-feï. Upon this work Chau Ju-kua's great treatise was based. See the edition by Hirth and Rockhill, Petrograd, 1911.

[6] E. Bretschneider, *Mediaeval Researches*, 2 vols., I, 232 (London, 1910); hereafter referred to as Bretschneider, *Mediaeval Res.*

[7] This is related in the *Si Shi ki* (travels) of Ch'ang-ti. Mention is made of *pin t'ie*, probably steel. See Bretschneider, *loc. cit.*, I, 146.

the conquering Chinghiz Khan[1] carried out his great expedition to Western Asia, established roads connecting Eastern Mongolia with Persia and Russia, sent armies even to Eastern Europe, and opened the way to trade between the Orient and all parts of the Occident. He captured Bokhara, Samarkand, Herat, Merv, Nîshâpûr, Kiev, and probably Moscow (Moscoss). He invaded Poland, Galicia, Silesia, and Hungary, and his astrologers must have mingled freely with their guild in the capitals of Eastern Europe. In 1221 the Chinese traveler K'iu Ch'ang ch'un reached Samarkand[2] and, what is especially significant, records the fact that "Chinese workmen are living everywhere" in the city, and that he saw "peacocks and great elephants which had come from Yin-du" (India). He also speaks of meeting and conversing with an astronomer of Samarkand, as we should naturally expect of a man of his learning. In 1236 the Mongols invaded Bulgaria; in 1238 France received a Mussulman ambassador asking aid against the Oriental forces; in 1241 the Mongols invaded Galicia, and in 1259 they burned Cracow for the second time.

The fact is also significant that in 1266 the king of Ceylon had Chinese soldiers in his service, while about this time the Chinese traveler Chau Ju-kua speaks of the Hindus from his personal knowledge as being "good astronomers and calculators of the calendar."[3]

Europeans in the East. Moreover, Europeans were frequently seen in the East, and the Grand Duke Yaroslav, in 1246, met one Fra Plano Carpini, a Franciscan, at the Mongol court. Only a few years later Rubrouck, another of the friars minor, visited the same court, and both he and Carpini have left accounts of their travels. There were also Eastern travelers who went to the West, such as the Taoist monk Ch'ang ch'un, from 1220 to 1224; Ch'ang-ti, who was sent out in 1259 by Mangu Khan, and who visited Bagdad; and Ye-lü Hi-liang, who traveled in Central Asia from 1260 to 1262.

[1] Jenghiz Khan, Genghes Khan, Jinghis Khan; originally Temuchin.

[2] The Semiscant of medieval writers, known to the Chinese as Sie-mi-sz'-kan and Sün-sz'-kan.

[3] Hirth and Rockhill, *loc. cit.*, p. 111.

The second half of the century saw other travelers from the West in the courts of Mongolia and China. Haithon (Hethum), king of Little Armenia, visited Mongolia in 1254, and the account of his journey is well known. Marco Polo, who set out from Venice in 1271, spent seventeen years in China and traveled in a leisurely way through Arabia, Persia, and India. Rashid ed-dîn, vizier of the Persian Empire in 1298, wrote a history of the Mongols showing thorough familiarity with their country. All these events, unimportant in detail, are very significant taken as a whole, because they afford evidence of the free interchange of ideas between the East and the West. Thus they serve to clear up many questions as to whether the algebra of China, for example, could have found its way to Italy in the 12th century. We repeat that it would be a cause for wonder if it had failed to do so. Moreover, the 13th century was a period of wealth, of luxury, and also of opportunity. Chinghiz Khan, in a letter to Ch'ang ch'un the Taoist traveler, asserted that "Heaven had abandoned China owing to its haughtiness and extravagant luxury."[1] This may have been true; a conqueror would be apt to say so; but it is certain that it was a century in which mathematics ought to have flourished in both continents. We shall see that this was the case.

Mathematical Activity in the Thirteenth Century. The 13th century, in China as in Europe, was a period of awakening. Indeed, it may be said that it was the period of the highest development of native mathematics in the East. Whether as a result of an interchange of thought with the West, or of the leisure which wealth brought to the country before the invasion of Chinghiz Khan, or of the growth of idealism which war is said by its advocates to foster, or of various other causes, China made a noteworthy advance in algebra at this time.

Perhaps the foremost scholar in this movement was Ch'in Kiu-shao, a man whose intimate history is quite unknown, but who was a soldier in his early days, was in government service

[1] Bretschneider, *Mediaeval Res.*, I, 37

in 1244, was governor of two provinces, and wrote the *Nine Sections of Mathematics* in 1247.[1] The work relates chiefly to numerical higher equations, in which the author anticipates to some extent Horner's Method (1819); but it also considers indeterminate equations and the application of algebra to trigonometry. The "nine sections" of his work are not the same as those of the one mentioned on page 31. For the value of π the author gives 3, $\frac{22}{7}$, and $\sqrt{10}$. In his work, too, the symbol ○ is used for zero and the place value is used in the writing of all numbers. The author shows little interest in applying his knowledge of algebra to the solution of practical problems, preferring to look upon it as a pure science.

Li Yeh's Work. The next noteworthy step in this direction was taken by Li Yeh (1178–1265). He wrote the *Sea Mirror of the Circle Measurement*[2] in 1249, the *I-ku Yen-tuan* in 1259, and various other works. In his early life he was engaged in public service, and in 1232 was governor of Chün Chou. He was later held in high esteem by Kublai Khan, whose reign began in 1260.[3] Li Yeh directed his son to burn all his works except the *Sea Mirror*. The *I-ku Yen-tuan* was also preserved, however, and each has since been looked upon as among the great works of China. While Ch'in Kiu-shao had given his attention chiefly to the solution of abstract equations, Li Yeh devoted himself to the forming of equations representing various complicated problems, the solution being neglected.

[1] Chang Ch'i-mei (1616) says that the author called the work *Su-shu* or *Su-hsiao*, but in the Chinese bibliographical works the title is *Su-shu Kiu-ch'ang*. Biernatzki (pp. 13, 28) transliterates the title as *Su schu kiu tschang* and the author as Tsin Kiu tschaou. Vanhée gives the date as 1257 and the author as Ts'in K'ieou-Chao. For details as to his methods see Mikami, *loc. cit.*, p. 63.

[2] *Ts'ö-yüan Hai-king*. Vanhée (*T'oung-pao*, XIV, 537) gives the date as 1248 and the title as *Ts'é yuen hai king*. Biernatzki gives the name of the author as Le Yay and Le yay Jin king, and the title as *Tsih yuen hä king*. Li Yeh also used the *nom de plume* Ching-chai.

[3] The Yuen Dynasty, founded by the Mongols, began in 1280, however. In 1260 the Mongols issued paper money, an event bearing somewhat on the history of arithmetic problems.

About this time another scholar, Liu Ju-hsieh, wrote a treatise on algebra, but the work is not extant.[1]

Chinese Astronomers. That astronomy was the mathematical subject of chief interest at this time, as indeed at all times in early Chinese history, is shown by the considerable list of names of scholars devoted to the science. Among the leaders at this particular period was Ye-lü Ch'u-ts'ai,[2] who lived *c.* 1230. He established a great school at Peking (then Yen-king) and accompanied Chinghiz Khan to Persia, occupying himself with the calculation of eclipses and coming into contact with the Persian astronomers.

Yang Hui's Work. In 1261 Yang Hui wrote *The Analysis of the Arithmetic Rules in Nine Sections*,[3] a work in which he explained some parts of the original *Nine Sections* (see page 31). In this work he gives a graphic representation of the summation of an arithmetic series. In another work[4] he gives rules for summing the series

$$1 + (1 + 2) + (1 + 2 + 3) + \cdots + (1 + 2 + 3 + \cdots + n)$$

and

$$1^2 + 2^2 + 3^2 + \cdots + n^2,$$

but offers no explanations. He wrote several other works,[5] and among his problems are such as that of the hare and hound and several involving simple and compound proportion.

Yang Hui's teacher, Liu I, a native of Chung-shan, wrote (*c.* 1250) a work which is known only by name,[6] but it doubtless related to numerical higher equations.

[1] He is mentioned in a work of 1303 as a prominent contemporary. His work is said to have been written before 1300. Since a commentary upon it was written by Yuen Hao-wen, who was a friend of Li Yeh (1178–1265), the date may be taken as *c.* 1260.

[2] Rémusat, in his *Nouveaux Mélanges Asiatiques*, II, 64, does not speak of his mathematical attainments, but in the *Yuen-Shih* (*Historical Records of the Yuen Dynasty*) these are discussed.

[3] *Hsiang-kieh K'iu-ch'ang Suan-fa.* Yang Hui is also known in European works as Yang Hwuy, Yan Hui, Yang Houei, Yang Hwang, and Kien-kouang.

[4] *Suan-fa T'ung-pien Pen-mo.*

[5] His six works on arithmetic, thought to be lost, were discovered in Shanghai in 1842.

[6] *I-ku Kon-yüan.*

Kóu Shóu-king. In 1267 the Mongols are known to have employed various Arab artillery officers,[1] so that contact with Arabia existed through the army. But this contact is the more apparent in the scientific achievements of Kóu Shóu-king[2] (1231–1316), a man well versed in the astronomy of the Arabs. He was a native of Hsing-t'ai[3] and was remarkable for his attainments even in early childhood, seeming to have inherited the scholarly qualities of his grandfather, Kóu Yung, a mathematician of repute. In early manhood Kóu Shóu-king developed into one of the greatest engineers of China. He was appointed by Kublai Khan to reform the calendar,[4] and for this purpose he replaced the armillary sphere which had been made about 1050, and which was calculated for Peenking, the former capital, differing about 4° in latitude from Peking, by the earliest of the great bronze instruments now on the wall of the latter city. The instruments constructed by him included several that were adapted to observations made in the daytime as well as at night,[5] and show that he had considerable knowledge of spherical trigonometry. Only two of his instruments seem to be extant, and these were found and described by Matteo Ricci[6] when he visited Peking early in the 17th century. Ricci also speaks of having seen similar instruments at Nanking.

With Kóu Shóu-king may be said to have begun the study of spherical trigonometry in China, a subject already far advanced in the Arab schools.

Further contact with the West was made at this time through the sojourn of Friar Odoric in Canton from 1286 to 1331.

[1] Two are called by the names I-se-ma-yin (probably Ismael) and La-pu-tan in the Chinese records.

[2] Also transliterated as Ko-cheou-king, Kouo Cheou-kin, and Kou Shou-ching. Since his cognomen is given as Jŏ-sze, it is possible that he is identical with the Liu Ju-hsieh already mentioned.

[3] Hing-tae, with other transliterations, a district in the prefecture of Shun-tĭh.

[4] He devised this calendar, the *Shóu-shĭ-li*, in 1280 and it was adopted in 1281.

[5] For a list and for the general subject of Kóu Shóu-king see A. Wylie, "The Mongol Astronomical Instruments in Peking," *Travaux de la 3ᵉ session du Congrès internat. des Orientalistes*, Vol. II.

[6] See page 303.

Chu Shï-kié. The 13th century in China closed with the remarkable work of Chu Shï-kié,[1] a native of Yen-shan. As to his private life, we know only that for more than twenty years he was a wandering teacher. He wrote two works, the *Introduction to Mathematical Studies*[2] in 1299 and *The Precious Mirror of the Four Elements*[3] in 1303. With him the old abacus algebra, in which the coefficients were represented by sticks placed on a checkered board,[4] reached its highest mark. In the first of his works there appears the algebraic rule of signs and an introduction to algebraic processes in general. In the second treatise, however, he considers a variety of new questions in higher algebra. He begins with what is called at present Pascal's Triangle, giving the values of the binomial coefficients and referring to the scheme as an old one. He considers higher equations with more than one unknown quantity, his treatment showing some knowledge of elimination by a determinant notation. He shows much ingenuity in his solution of numerical higher equations by the method already used by Ch'in Kiu-shao and which resembles Horner's Method.

2. Japan

Close of the Dark Ages. For Japan, as well as for Europe, there was a period which may properly be spoken of as the Dark Ages. For a thousand years after Buddhism was introduced into Japan there were few other events of intellectual significance to record. Japan was awaiting the world's rebirth, and this period of rebirth came to the East at about the same time as to the West.

Only two men stand out as worthy of mention in the history of Japanese mathematics between the year 1000 and the year 1500. The first of these is Fujiwara Michinori, a daimyō or

[1] Tchou Che-kié, Tchou Che-Kié, Tschu Schi kih, Choo Ché-kié, Chu-Shih-Chieh, Tchou Che-kie, and various other transliterations. He is also called Chu Sung ting.

[2] *Suan-hio-ki-mŏng*, or *Suan-hsiao Chi-mêng*.

[3] *Szu-yuen Yü-kien*, or *Szu-yuen Yu-chien*. The introduction was written by Tsu Yi Chi Hsien Fa.

[4] See Volumé II, Chapter VI.

feudal lord in the province Hyūga, who wrote a work on permutations[1] between the years 1156 and 1159. The work is now lost, but it was thought important enough to be considered by the leaders of mathematics in Japan in the 17th century.

The second name worthy of note is that of the Buddhist priest Genshō, who lived in the first part of the 13th century. No trace remains of any of his writings, but tradition says that he was possessed of remarkable arithmetical ability.

3. India

Śrīdhara. The first of the Hindu writers of this period seems to have been Śrīdhara, commonly known as Śrīdharācārya, Śrīdhara the Learned, who was probably born in 991.[2] His work is known as the *Gaṇita-Sara* (*Compendium of calculation*) but is more commonly designated by the subtitle *Triśatika*, a name referring to its three hundred couplets.[3] The subjects considered are numeration, measures, rules, and problems, and the order bears very close resemblance to the one followed about a century later by Bhāskara in his *Līlāvati*. The latter writer was acquainted with Śrīdhara's work, as he himself testifies in his *Bija-Gaṇita*.

Under the general topics above mentioned are included series of natural numbers, multiplication, division, zero, squares, cubes, roots, fractions, Rule of Three, interest, alloys, partnership, mensuration, and shadow reckoning. The statement relating to zero is noteworthy as the clearest one to be found among the Hindus: "If zero is added to a number, the sum is that number itself; if zero is subtracted, the number remains unchanged; if zero is multiplied, the result is zero; and if a number is multiplied by zero, the product is zero only." The question of division by zero is not considered.


[1] The theory known as *Keishizan*.

[2] The date is uncertain, being placed by one writer three centuries earlier. The question is discussed by N. Ramanujacharia and G. R. Kaye in the *Bibl. Math.*, XIII (3), 203.

[3] The Sanskrit text was published in 1899. The English text is given in the article mentioned above. The name may also have come from the fact that it originally had 103 couplets.

For dividing by a fraction Śrīdhara gives the rule of multiplication by the inverted divisor, a rule already known to Mahāvīra (*c.* 850). Like the latter, too, he uses $\sqrt{10}$ for π.

Bhāskara (1114–*c.* 1185). There is only one other writer who stands out prominently in the history of Hindu mathematics from 1000 to 1500, and that is Bhāskara, commonly known



ŚRĪDHARA'S TRIŚATIKA, *c.* 1025

Two pages from the copy used by Colebrooke in his works on Hindu mathematics. From Kaye's *Indian Mathematics*


as Bhāskara the Learned (Bhāskarācārya),[1] a native of Biddur[2] in the Deccan, but working at Ujjain. An ancient temple inscription refers to him in the following terms: "Triumphant is the illustrious Bhāskarācārya whose feet are revered by the wise, eminently learned, . . ., a poet, . . . endowed with good fame and religious merit. . . ."[3]

[1] The name is variously transliterated. Thus, we have Bhascara Acharya, Bhaskaracharya, and other forms. See J. Garrett, *loc. cit.*, p. 92; Taylor, *Lilawati*, 1; T. W. Beale, *The Oriental Biographical Dictionary*, Calcutta, 1881; Colebrooke, *Bhāskara.*

[2] Probably the modern Bidar.

[3] Kaye, *Indian Math.*, 37.

Bhāskara's Lilavati. Bhāskara wrote chiefly on astronomy, arithmetic, mensuration, and algebra. His most celebrated work is the *Lilāvati*, a treatise based upon Śrīdhara's *Triśatika* and relating to arithmetic and mensuration.[1] This work was translated into Persian by Fyzi[2] in 1587 by direction of the emperor Akbar, a great patron of letters. Fyzi states, though it does not appear upon what authority, that Lilāvati was the name of Bhāskara's daughter and that the astrologers predicted that she should never wed. Bhāskara, however, divined



PALM-LEAF MANUSCRIPT OF THE LILĀVATI

Showing the form in which the Hindu manuscripts appeared before paper became a common medium. This manuscript was copied *c.* 1400. From the author's collection


a lucky moment for her marriage and left an hour cup floating on the vessel of water. This cup had a small hole in the bottom and was so arranged that the water would trickle in and sink it at the end of the hour. Lilāvati, however, with a natural curiosity, looked to see the water rising in the cup, when a pearl dropping from her garments chanced to stop the influx.

[1] The first translation into English is that of Taylor (1816), already mentioned.

[2] Also spelled Faizi and Feizi. He was a brother of Akbar's secretary, Abu Fazil. The work was printed at Calcutta in 1827.

So the hour passed without the sinking of the cup, and Lilāvati was thus fated never to marry. To console her, Bhāskara wrote a book in her honor, saying: "I will write a book of your name which shall remain to the latest times; for a good name is a second life and the groundwork of eternal existence."


The work begins, as is the custom in the East, with an address to the Deity: "Salutation to the elephant-headed Being



FROM BHĀSKARA'S LILĀVATI

From a manuscript of *c.* 1600. The original work was written *c.* 1150. The illustration shows the form of Hindu manuscripts just following the use of palm-leaf sheets. This page has the following statement: "Assuming two right triangles [as shown], multiply the upright and side of one by the hypotenuse of the other: the greatest of the products is taken for the base; the least for the summit; and the other two for the flanks. See" [the trapezoid]. Colebrooke's translation, page 82

who infuses joy into the minds of his worshipers, who delivers from every difficulty those who call upon him, and whose feet are reverenced by the gods." The book includes notation, the operations with integers and fractions, the Rule of Three, the most common commercial rules, interest, series, alligation, permutations, mensuration, and a little algebra. The rules relating to zero are also given, to the effect that $a + 0 = a$, powers of 0 are 0, and $a \cdot 0 = 0$. The statement that $a \div 0 = 0$ (corrected by his commentators) was evidently not clear to him, for his statement is "A definite quantity divided by cipher is the



॥ लीलावती ॥ ८९

PAGE FROM THE FIRST PRINTED SANSKRIT EDITION OF BHĀSKARA'S LILĀVATI

Printed at Calcutta, 1832. This is a continuation of the portion shown in manuscript on page 277. The statement, as translated by Colebrooke, is as follows: "Length of the base, 300. Summit, 125. Flanks, 260 and 195. Perpendiculars, 189 and 224." From these the other parts are found

sub-multiple of nought," while his illustrations are that $10 \div 0 = \frac{10}{0}$ and $3 \div 0 = \frac{3}{0}$,[1] the latter being accompanied by the statement that "this fraction, of which the denominator is cipher, is termed an infinite quantity."

The Bija Ganita. Bhāskara also wrote the *Bija Ganita*,[2] a work on algebra.[3] In this he discusses directed numbers, the negatives being designated in Sanskrit as "debt" or "loss"[4] and being indicated by a dot over each number, as in the case of $\dot{3}$ for -3, and the usual rules being stated correctly. The imaginary is dismissed with the statement, "There is no square root of a negative quantity: for it is not a square." Where several unknown quantities are used, they are mentioned as colors: "'so much as' and the


[1] Colebrooke, *loc. cit.*, pp. 19, 20, 137.

[2] Variously transliterated as Vijaganita, Vija-Gañita, and the like. The term literally means "seed counting" or "seed arithmetic." The work was translated by Colebrooke, *Bhāskara*, p. 129, with the subtitle *Avyacta-Gañita*.

[3] Translated into Persian by Ata Allah Rusheedee in 1634. The English version of this translation was published by Edward Strachey, *Bija Ganita*, London, n. d. (*c.* 1812), apparently with the aid of a translation made, with the help of a pundit, by S. Davis, *c.* 1790.

[4] *Rina* or *cshaya*.

colors 'black, blue, yellow, and red,' and others besides these, have been selected by venerable teachers for names of values



FROM A MANUSCRIPT OF FYZI'S TRANSLATION OF THE LILĀVATI

Fyzi, counselor of Akbar, made the translation into Persian in 1587. This manuscript is dated 1143 A.H., or 1731 A.D. The Persian manuscripts were written in our ordinary book form. The above is page 32 of Taylor's translation of 1816. The problem is: "A number is multiplied by 5; from the product is subtracted one-third of itself, and the remainder is divided by 10; to the quotient is added ½ of ⅓ of ¼ of the assumed number, and the result is 68. What is the number?"

of unknown quantities."[1] Surds are treated extensively, as in many medieval works on algebra, the difficulty of handling

[1] The initial Sanskrit syllables of the names of the colors are used.

irrationals of all kinds being particularly great in the era of poor symbolism. As with Aryabhaṭa and Brahmagupta, the "pulveriser" is given extensive treatment. Simple and quadratic equations receive more attention and are more clearly discussed than is the case with other Hindu writers. Besides numerous problems relating to geometric figures there are the usual poetic types, of which the following may serve as an illustration:

The son of Prīt'há,[1] exasperated in combat, shot a quiver of arrows to slay Carńa. With half his arrows he parried those of his antagonist; with four times the square root of the quiverful he killed his horse; with six arrows he slew Salya[2]; with three he demolished the umbrella, standard, and bow; and with one he cut off the head of the foe. How many were the arrows which Arjuna let fly?

Śrīdhara's Rule for the Quadratic. The rule used for solving the quadratic is given as Śrīdhara's.[3] The method of writing an equation has some evident advantages, the equation $18x^2 = 16x^2 + 9x + 18$ being written

ya v 18 ya o ru o
ya v 16 ya 9 ru 18

which is then transformed into $2x^2 - 9x = 18$, thus:


ya v 2 ya ṡ ru o
ya v o ya o ru 18[4]

Bhāskara's Siddhānta Śiromani. A third work of importance written by Bhāskara is the *Siddhānta Śiromani* (*Head jewel of accuracy*)[5] in which, in the book *Goladhia* (*Theory of the*

[1] The son's name was Arjuna, mentioned later in the problem.
[2] The charioteer of Carńa.
[3] See page 274.
[4] *Ya v* is for *yāvat-tāvat*, "as many of" (the unknown), and is used for the highest power. *Ya* is a color, green, and is used here for the first power. *Ru* is for *rūpa*, the known number. See Volume II and Colebrooke, *Bhāskara*, p. 139.
[5] Various notes on this work will be found in H. T. Colebrooke, *Miscellaneous Essays*, 2 vols., 2d ed., Vol. II (Madras, 1872). It should be stated that the *Siddhānta Śiromani* is thought by various scholars to include the other works mentioned, but this is merely a question of division of material.

Sphere), he treats of astronomy and asserts, as various ancient Greek philosophers have done, the sphericity of the earth.

The ancient inscription referred to on page 275 relates that Bhāskara's grandson, Chaṅgadwa, was chief astrologer to King Siṁghana, and that in his time a college was founded to expound the doctrines of Bhāskara.[1]



FROM BHĀSKARA'S GOLADHIA

From a manuscript of Bhāskara's work on astronomy, being the fourth and last chapter of the *Siddhānta Siromani*. This work was written *c.* 1150. The reproduction is greatly reduced

In the forming of Pythagorean triangles Bhāskara follows Brahmagupta in stating the relations

$$\sqrt{m}, \quad \frac{1}{2}\left(\frac{m}{n} - n\right), \quad \frac{1}{2}\left(\frac{m}{n} + n\right),$$

and adds the two further relations

$$m, \quad \frac{2mn}{n^2-1}, \quad \frac{m(n^2+1)}{n^2-1}$$


and

$$\frac{m(n^2-1)}{n^2+1}, \quad \frac{2mn}{n^2+1}, \quad m.$$

Other Parts of South Asia. In Eastern Sumatra (the Chinese San-fo-ts'i) it is related by Chinese travelers of the 13th century that the people were able mathematicians and could calculate future eclipses of the sun and moon.[2] No doubt the

[1] G. R. Kaye, *Indian Mathematics*, p. 37.
[2] Hirth and Rockhill, *loc. cit.*, p. 64.

same could have been said at that time of other parts of Asia whose records have not come down to us, such calculations



॥ सिद्धान्तशिरोमणिः ॥ १४३

उदाहरणं ।

ये यातायिकमासचोनदिवसा येचापि तच्छेषके नेषा मैक्यमवेच्य जिष्णुजकृतान्काम्बाद्यथैवागतं । भूशैलेंदुखखाभ्रषट्करयुगाष्टाब्ध्यङ्क ।६४८४२६०००१७१ । तुल्यं यदा । काले कल्प गतं तदा वदति यः सब्रह्मसिद्धान्तवित् ॥ १२ ॥

अथ गणानयने ये लब्धा अधिमासाः तयाद्याश्च ये च तच्छेषके नेषामैक्यं भूशैलेंदुखखाभ्रषट्करयुगाष्टाब्ध्यङ्क ।६४८४२६०००१७१ । तुल्यं कृतास्तिथिगोब्ध्यादिभि । ८६३ ४७४८१६८४ । गुणितं जातं । ५५६८३४६८२८२३२६४२ २०३७८६४ । व्येकचन्द्रार्ते । १६०२८८८८८८८८८ । भक्तं लब्धं । ३४८२४१८३२३३६ । जानोऽग्रेपमितोगतं दुद्युगणः । १०३००० । अस्मात्प्राग्वदवमानि । १६१ । अव मग्रेषच्च । २६७४२६००००० । सावनाच्चगणश्च । १०१३८ । अथ पृथग्गतेदुद्युगणेऽधिमासै गुणितो युगेंदुदिनै र्भक्तो लब्धं गताधिमासाः । १० । अधिमासशेषच्च । ३८१०००००००० । लब्धाधिमासैः दिनीकृतैं । ३०० । ऊनद् दुद्युगणः । १०३०० । सौराच्चगणो भवति । १०००० । अतः कल्पगतं सप्तविंशति । २७ । वर्षाणि नव । ८ । मासाः दश । १० । दिनानि अस्योपपत्ति र्बीजगणितेन एको चरज्येदुणकौ विभिन्नावित्यादिना कथमस्य विषय इति चेत् उच्यते गतसौरदिनेभ्यो यावन्तोधिमासाः यच्च शेषं गतचन्द्रदिनेभ्योपि तावन्त एव भवन्ति तावदेव चाव

FROM THE FIRST PRINTED EDITION OF BHĀSKARA'S GOLADHIA

This shows in print the same page shown in manuscript on page 281. The work was printed at Calcutta in 1842

having been a part of the general stock in trade of the astrologers in various countries for many centuries preceding this particular period.

4. Persia and Arabia

Decay of Bagdad. For two centuries after the golden age of the first three caliphs Bagdad continued to be a center of scientific activity, in spite of the fact that it began to lose political prestige after the death of al-Mâmûn (833). By the year 1000, however, the spiritual supremacy of the city had passed and the seats of learning of the Western Arabs had begun to take the place of the capital of the caliphs in Mesopotamia. The Seljuk Turks, an intolerant Tartar tribe, overran much of the territory formerly so well governed by the caliphs, captured the holy cities of Palestine, and by their ruthless behavior gave excuse for the crusades. In 1258 the Mongols took Bagdad, and thenceforth it was little more than a name. Brute force had put an end to the idealism that had been so noticeable in the eastern Mohammedan empire.

Al-Karkhî. Among the last of the real contributors to mathematics in the city of the caliphs was al-Karkhî,[1] who died *c.* 1029. His first work of note was an arithmetic, the *Kâfî fîl Hisâb*,[2] probably written between 1010 and 1016, and drawn largely if not exclusively from Hindu sources.[3] It not only contains the elements of arithmetic as set forth by many writers of the time, but gives the rule of quarter squares,

$$\left(\frac{a+b}{2}\right)^2-\left(\frac{a-b}{2}\right)^2=ab,$$

a rule probably due to the Hindus. It also gives such methods of multiplication as are expressed by the formulas

$$(10a+a)(10b+b)=[(10a+a)b+ab]10+ab$$

and

$$(10a+b)(10a+c)=(10a+b+c)a\cdot 10+bc.$$

[1] Mohammed Abû Bekr ibn al-Ḥasan (or al-Ḥosein), al-Karkhî.

[2] *Book of Satisfactions.* A. Hochheim, *Kâfî fîl Hisâb des Abu Bekr Muhammed Ben Alhusein Alkarkhî*, 3 parts, Halle a. S., 1878–1880; hereafter referred to as Hochheim, *Kâfî fîl Hisâb.*

[3] H. Weissenborn, *Gerbert*, p. 196 seq. (Berlin, 1888). But see Cantor's earlier opinion that it came from Greek sources; *Geschichte*, I (1), 655.

In the approximations for roots al-Karkhî gives, among others,

$$\text{for } m = a^2 + r, \quad \sqrt{m} = a + r/(2a + 1);$$
$$\text{for } r \lessgtr a, \quad \sqrt{m} = a + r/2a.$$

He also considers the mensuration of plane figures, particularly as it involves surd numbers, and includes the Heron formula $\sqrt{s(s-a)(s-b)(s-c)}$.

The work closes with a treatment of algebra, including quadratic equations and the usual Arabic explanation of the terms *al-jabr* and *al-muqâbala*, discussed in Volume II.

Al-Karkhî's Fakhrî. Al-Karkhî is best known, however, for his algebra, the *Fakhrî*,[1] which includes the usual operations on algebraic quantities, roots, equations of the first and second degrees, indeterminate analysis, and the solution of problems. The quadratic equations include such forms as $x^4 + 5x^2 = 126$, and the solution of quadratics in general depends upon rules such as that represented by the equation

$$ax^2 + bx = c$$

and the formula $$x = \left[\sqrt{\left(\frac{b}{2}\right)^2 + ac} - \frac{b}{2}\right] : a.$$

The rules are explained geometrically, as in the works of the earlier Arab writers. Various problems given by him are apparently suggested by al-Khowârizmî and Diophantus, and these include such cases as the finding of integral solutions for

$$x^3 + y^3 = z^2,$$
$$x^2 - y^2 = z^3,$$
$$x^2y^3 = z^2,$$

and

$$x^3 + 10x^2 = y^2,$$

and the finding of fractional solutions for

$$x^2 - y^3 = z^2$$

and

$$x^3 + y^2 = z^3.$$

The work ranks as the most scholarly algebra of the Arabs.

[1] F. Woepcke, *Extrait du Fakhrî . . . par . . . Alkarchî* (Paris, 1853). For an explanation of the term *Fakhrî* see Volume II, Chapter VI.

Minor Writers of the Eleventh Century. Of the minor writers of the 11th century the following deserve brief mention:

Mohammed ibn al-Leiṭ[1] lived about 1000, was interested in the trisection problem, and wrote on the construction of regular polygons of seven and nine sides.

Ḥâmid ibn al-Khiḍr[2] wrote on the astrolabe and asserted that the equation $x^3 + y^3 = z^3$ cannot be solved.

Manṣûr ibn ʿAlî[3] wrote on astronomical instruments, trigonometry, spherical sines, and Ptolemy's *Almagest*.

Al-Nasavî wrote on Hindu arithmetic and on the works of Archimedes.[4]

One of the most brilliant writers on the contemporary history of mathematics at the opening of the 11th century was Albêrûnî.[5] He was one of the *munajjimin*, or astrologer-astronomers of the Arabs. He visited India and made a careful study of that country and of its work in mathematics and the other sciences. He summarized the debased state of knowledge of his day in the words, "What we have of sciences is nothing but the scanty remains of bygone better times." In later life he wrote his work on India, and to this we are indebted for the best summary of Hindu mathematics that the Middle Ages produced.

Avicenna. Among the contemporaries of Albêrûnî there was the famous physician and philosopher known in Christian Europe as Avicen'na[6] (980–1037). He was born in Safar, near

[1] Mohammed ibn al-Leiṭ, Abû'l Jûd.

[2] Ḥamid ibn al-Khiḍr, Abû Mahmûd, al-Khojendî, died *c.* 1000.

[3] Manṣûr ibn ʿAlî ibn ʿIrâq, Abû Nasr, *c.* 1000.

[4] ʿAlî ibn Aḥmed, Abû'l-Ḥasan, al-Nasavî, *c.* 1025. Woepcke in the *Journal Asiatique*, 1863, p. 496. He was born at Nasa, in Khorâsân. On his arithmetic see Suter in *Bibl. Math.*, VII (3), 113.

[5] Mohammed ibn Ahmed, Abû'l Rîhân (or Raiḥân), al-Bêrûnî. Born probably in Khwarezm, 973; died 1048. He may have been born at Byrun in the valley of the Indus, his name also appearing as al-Bîrûnî. On his life and works, see his *Athâr-el-Bâkiya*, *Chronology of Ancient Nations* (London, 1879), and his *India*; Boncompagni's *Bullettino*, II, 153; S. Günther, *Zeitschrift* (Hl. Abt.), XXI, 57; E. C. Sachau, *Zeitschrift d. deutsch. Morgenl. Gesellschaft*, XXIX.

[6] Al-Ḥosein ibn ʿAbdallâh ibn al-Ḥosein (or Ḥasan) ibn ʿAlî, Abû ʿAlî, al-Sheich al-Ra'îs, ibn Sînâ. See K. Lokotsch, *Avicenna als Mathematiker* (Erfurt, 1912).

Kharmîtan, not far from Bokhârâ. He wrote on Aristotle, Euclid, astronomy, music, medicine, and arithmetic, his treatment of numbers being based upon Greek models.

Ibn al-Ṣalâḥ,[1] who died in 1153/54, was one of the later generation of the Persian scholars who made Bagdad so famous. Like so many mathematicians of the East he was also learned in philosophy and medicine,—the latter partly on account of the supposed connection between the healing art and astrology. He was born in Bagdad and finally went to Damascus and died there. He wrote on geometry, and manuscripts of his works, apparently fragmentary, are still extant.

Omar Khayyam. The 12th century saw less attention to mathematics in the ancient Arab seats of learning, and more attention to the science in Persia. Of those whose names added luster to Persian mathematics and letters the most prominent was the poet who is generally known to English writers as Omar Khayyam[2] (*c.* 1100). While he is known to the Western world chiefly as the author of the *Rubaiyat*,[3] he wrote on Euclid and on astronomy,[4] and contributed a noteworthy treatise on algebra.[5]

Minor Writers. A little later than Omar Khayyam another Persian, a native of Khorâsân, made for himself a great name. This writer was al-Râzî,[6] known as one of the leading philosophers, physicians, and mathematicians of the Persians. His contributions to mathematics were chiefly in the domain of geometry.

[1] Aḥmed ibn Mohammed ibn al-Surâ Nejm ed-dîn, Abû'l-Futûḥ.

[2] 'Omar ibn Ibrâhîm al-Khayyâmî, Ġiyâṯ ed-dîn, Abû'l-Fatḥ. He was born at Nîshâpûr (Nishapur, Nishabur) *c.* 1044 and died there in 1123/24.

[3] Largely through the remarkable but not very exact translation of Edward Fitzgerald, London, 1859.

[4] "Ah but my computations people say
Reduced the year to better reckoning."

[5] *L'Algèbre d'Omar Alkhayyâmî*, Arabic and French texts, by F. Woepcke, Paris, 1851. On his life, see various editions of the Rubaiyat, and also J. K. M. Shirazi, *Life of Omar al-Khayyâm*, Edinburgh, 1905.

[6] Mohammed ibn 'Omar ibn al-Ḥosein, Abû 'Abdallâh, Faḫr ed-dîn al-Râzî, ibn al-Khatîb, 1149/50–1210.

Of the Arab scholars of the 12th century one of the best known was Kemâl ed-dîn ibn Yûnis, or ibn Man'a,[1] who was born at Moṣul on the Tigris river. His works on the theory of numbers and conic sections were highly esteemed by his Arab contemporaries.

Contemporary with the last-named writer was Ta'âsîf,[2] a native of upper Egypt, a jurist, an engineer, and a mathematician. He showed his interest in the foundations of mathematics by writing upon Euclid's postulates.

The Mongol Scourge. If the Seljuk Turks had been intolerant in the 11th and 12th centuries, rendering difficult the leading of an intellectual life, the great Mongol scourge, led by Chinghiz Khan between 1206 and 1227, rendered such a life well-nigh impossible. His conquests and his son's included a considerable part of the civilized world from Northern China through Turkestan and Persia, and down to the banks of the Indus. The son, Oktai (died 1241), quite as brutal as his father, ravaged nearly half of Europe, and the result of the total conquest was the impoverishment of all the intellectual centers that were once the glory of central and western Asia. To be sure, two of the successors of these tyrants, Kublai Khan (1216–1294) and Timur, or Tamerlane (1336–1405), contributed to the better things of life, but in general the record of the Mongol invasions for two centuries is one of the blackest in all history.

Decay of Learning. In the 13th century only one Persian writer deserves special mention, and even he spent his closing years in Bagdad. This writer was Naṣîr ed-dîn,[3] a native of Ṭûs, in Khorâsân. He was an all-round scholar, writing upon trigonometry, astronomy, computation, geometry, and the construction and use of the astrolabe.

[1] Mûsâ ibn Yûnis ibn Mohammed ibn Man'a, Abû'l-Fatḥ, Kemâl ed-dîn. Born at Moṣul, 1156; died at Moṣul, 1242.

[2] Qaiṣar ibn Abî'l-Qâsim ibn 'Abdelġanî ibn Musâfir, 'Alam ed-dîn, known under the name of Ta'âsîf. Born 1179 (possibly 1169); died 1251.

[3] Mohammed ibn Mohammed ibn al-Ḥasan, Abû Ja'far, Naṣîr ed-dîn al-Ṭûsî, 1201–1274.

Of the Arab writers, Ibn al-Yâsimîn,[1] who lived in Morocco, is known chiefly for the influence of a poem which he wrote on algebra, the *Arjûza.* Several manuscripts still exist, and it seems to have had some such influence in popularizing algebra as the *Carmen de Algorismo* (p. 226) had with respect to algorism. Ibn al-Lubûdî,[2] a native of Ḥaleb (Aleppo), was known in his century for works on arithmetic, algebra, and Euclid. The Arab interest in learning was rapidly waning, however, and only one other name deserves mention in the record of the 13th century,—that of al-Ṭûsî,[3] another prominent native of Ṭûs. He wrote on geometry and algebra and invented one form of astrolabe known as "Ṭûsî's staff." Islam had lost its hold upon mathematics; the mathematical world was becoming a hyperbola having one focus in China and the other in Christian Europe, with nothing between the branches.

The most notable of the Christian writers in the Near East at this time was Bar Hebræus,[4] whose father, a Jew named Aaron, had entered the Christian church. When the son was twenty years old (1246) he was made Jacobean bishop of Gubos, near Malatia, and later he occupied other positions of ecclesiastical importance. He wrote on astronomy and lectured on Euclid and Ptolemy.[5]

In the 14th century only three Mohammedans of any considerable prominence appear among the world's mathematicians, and no one of these was a genius. Two lived at least part of the time in Egypt, Ibn al-Hâ'im,[6] a writer on

[1] 'Abdallâh ibn Mohammed ibn Ḥajjâj, Abû Mohammed. He died *c.* 1203–1205.

[2] Yaḥyâ ibn Mohammed ibn 'Abdân ibn 'Abdelvâḥid, Abû Zakarîyâ Nejm ed-dîn, 1210/11–1267/68.

[3] Al-Moẓaffar ibn Mohammed ibn al-Moẓaffar Sharaf ed-dîn al-Ṭûsî. He died *c.* 1213.

[4] That is, Son of the Jew. His Arabic name was Jûḥannâ Abû'l-Faraj Bar-Hebræus. Born at Malatia, in Eastern Asia Minor, 1226; died at Moṣul, July, 1286.

[5] F. Nau, in No. 121 of the *Bibliothèque de l'École des Hautes Études.* Paris, 1899.

[6] Aḥmed ibn Mohammed ibn 'Imâd, Abû'l-'Abbâs Shihâb ed-dîn. Born at Cairo, 1352 or 1355; died at Jerusalem, 1412.

arithmetic, and Ibn al-Mejdî,[1] who wrote on astronomy, trigonometry, arithmetic, the calendar, and mathematical tables. A third Arab writer of the period, commonly known as Ibn al-Shâṭir,[2] left works on trigonometry, the astrolabe, and astronomy, and prepared a few mathematical tables.

Ulugh Beg, the Royal Astronomer. Of the representatives of the Arab-Persian interest in mathematics who lived in the 15th century the only one who seems to have possessed any genius was Ulugh Beg[3] (1393–1449). and even this genius was rather perseverance than any unusual endowment of intellect. He was a Persian prince, born at Sultanieh, and his interest in astronomy and astronomical tables was shown in the observatory which he founded at Samarkand. The tables which were worked out under his direction were highly esteemed in Europe as well as in the East.[4] His assistant, al-Kashî,[5] wrote a short treatise in Persian on arithmetic and geometry.[6]

Summary of Arab Achievements. With these names the achievements of the Mohammedan writers practically close. As we sum up these achievements we are struck by the interest of the Arabs in science but by their lack of originality. They received their astronomy first from the Hindus and then from the Greeks, their geometry solely and their algebra chiefly from the Greeks, and their trigonometry largely from the Hindus in connection with astronomy. As already stated (p. 177), they originated nothing of importance either in arithmetic or in geometry, they systematized algebra to some extent,

[1] Aḥmed ibn Rajeb ibn Tîboġâ, Shihâb ed-dîn Abû'l-'Abbâs. Born 1359; died in Egypt, 1447.

[2] 'Alî ibn Ibrâhîm ibn Mohammed al-Moṭ'im al-Anṣârî, Abû'l-Haṣan. Born 1304; died 1375/76 or 1379/80.

[3] Ulûġ Beg.

[4] See L. P. E. A. Sédillot, *Prolégomènes des Tables Astronomiques d'Ouloug Beg*, Paris, 1847; T. Hyde, *Tabulae Longitudinis et Latitudinis Stellarum Fixarum ex Observatione Ulugbeighi*, Oxford, 1665; E. B. Knobel, *Ulugh Beg's Catalogue of Stars*, Washington, 1017.

[5] Jemshîd ibn Mes'ûd ibn Maḥmud, Ġiyât ed-dîn al-Kashî. He is also known as Kazi Zadeh al Rumi and Ali Kushi. Died *c.* 1436.

[6] H. Hankel, *Geschichte der Mathematik*, 289 (Leipzig, 1874) (hereafter referred to as Hankel, *Geschichte*); Taylor, *Lilawati*, Introd., p. 14. The introduction to his *Miftâh al-ḥisâb* (*Key of arithmetic*) was translated by Woepcke.

they improved upon the astronomy of their predecessors, and they made some real contributions to trigonometry. All these matters will be discussed in the appropriate chapters. But on the whole the Arabs of this period were still transmitters of learning rather than creators, and to them Europe is chiefly indebted for preserving in their translations many of the important works of the Greeks. To this rather sweeping assertion, however, one noteworthy exception may justly be taken, for it seems quite certain that it is to an Arab, or rather to a Turkish, scholar that we owe the first actual use of a decimal fraction. This step was taken independently at a later period by European arithmeticians, but the decimal fraction seems certainly to have been in use in Samarkand early in the 15th century. In a work by al-Kashî, or Jemshîd (p. 289), the ratio of the circumference to the radius of a circle is given, in part, as follows:

$$\underset{\text{Integer}}{6}\ 28318\cdots,$$

the full result being correct to sixteen decimal places.

Justice also requires that the Arabs of the four centuries beginning with the year 800 should be judged not with respect to the great achievements of the golden age of Greece but rather in comparison with the very meager results secured by their contemporaries in Europe and the Far East. If we consider Europe during the same period, we shall find the names of few original scholars in the domain of mathematics. The number of Arab, Persian, and Turkish scholars in this period exceeds, so far as we yet know, that of their European contemporaries, and their achievements were more significant. It was only during the period from 1200 to 1400 that European mathematics forged ahead, and even then the Arab influence was one of the prominent moving causes.

Justice further requires the admission that for lucidity of statement the scholars of Bagdad surpassed their contemporaries both in the East and in the West during a period of about six centuries, and were quite their equals in originality.

TOPICS FOR DISCUSSION

1. A comparison of the general nature of Chinese mathematics in the Middle Ages with that of the mathematics of Europe.

2. Evidences of the interchange of thought between the East and the West in the Middle Ages, and the possible effect of this interchange upon the mathematics of Europe and Asia.

3. Influence of astrology upon astronomy and upon pure mathematics, both in the East and in the West.

4. Nature of astronomy in the Middle Ages in the East and the border line between this science and astrology.

5. Nature of the mathematics of China in the period of its greatest development, the 13th century

6. The reliability of Chinese texts in the Middle Ages, based upon a general study of Chinese literature.

7. The justice of the claims of China in the field of numerical higher equations.

8. The nature of the problems that interested Chinese scholars in the Middle Ages.

9. Early history of mathematics in Japan. Reasons for the failure of the science to advance.

10. The nature of the mathematics of India in the Middle Ages. A comparison of this mathematics with that of China.

11. The nature of the problems that interested Hindu scholars in the Middle Ages.

12. The works of Bhāskara; their nature and influence.

13. Causes of the decay of mathematics in the Mohammedan countries, beginning with the 11th century.

14. Nature of algebra as developed by Mohammedan writers after the Golden Age of Bagdad.

15. The contributions to mathematics made by the Persian poet Omar Khayyam.

16. The life and works of the prince-astronomer Ulugh Beg and the reasons why his influence was not more powerful in Mohammedan lands.

17. A summary of the contributions of the Arabs to the science of mathematics.

18. General position of Hebrew mathematics in the Near East in the Middle Ages.

CHAPTER VIII

THE SIXTEENTH CENTURY

1. General Conditions

The Sixteenth Century in General. Until about the year 1500 the mathematics of the world was, so far as any records tell us, limited to a small number of individuals in each century. Printing having only just been invented, there was no simple way for comparatively obscure workers, even in the 15th century, to make their contributions or their interest known, and so they left no record of their achievements. In the 16th century, however, the printed page began to perpetuate names, and it now becomes impossible to do more than select a few out of the many for such comment as space may allow.[1]

Science and Letters. Furthermore, historical events now began to be recorded more freely, the world moved more rapidly, and the influences that bear upon the development of mathematics become more difficult to trace. That the opening of a new world would greatly increase the interest in commercial mathematics is evident, and the fact is abundantly proved by the printed books of the 16th century; but the influence of the great literary movement illustrated by such writers as Shakespeare, Cervantes, and Camoens, or of such world events as the defeat of the Spanish Armada, is not so apparent in the scientific field. Indeed, we may say that it was the influence of such scientists as Leonardo da Vinci, Copernicus, Palissy, and Tycho Brahe that stimulated the literary renaissance of the period, rather than the reverse.

[1] The reader who has access to the seventh edition of the *Encyclopædia Britannica* will find that the "Dissertation Third," by John Playfair, on "The Progress of Mathematical and Physical Science since the Revival of Letters in Europe" forms a very good background for his studies of this period.

Mathematical Conditions. The conditions in the field of mathematics were such as to mark out the course of progress. Euclid's *Elements* had appeared in print in 1482 and the *Conics* of Apollonius was known in manuscript,[1] and hence the most promising field in pure mathematics was in the domain of analysis. The quadratic equation had been fully solved, and so the next step was to attempt the solution of the cubic equation, with which the Greeks and Arabs had been successful only in special cases and by having recourse to the intersection of conics. It was here, then, that mathematics would naturally be expected to advance. Along with this there would be expected to develop a better symbolism, and one that was also suited to the needs of typography. After the cubic was solved, the mathematical world would be expected to try the equations of the fourth and higher degrees. We should anticipate, for geographic reasons, that the lead would be taken in Italy, and we should also expect that the demands of astronomy and navigation would require a more rapid development of trigonometry. All these suggested expectations, as we shall see, were fulfilled. Indeed, in the space of a single century mathematics made more advance than had been achieved since the days when the Alexandrian School dominated the scholastic world.



MATHEMATICS IN THE 16TH CENTURY

Concept of the range of the science. From the title-page of Coignet's *Arithmetica*, Antwerp, 1580

[1] It was first printed in Venice in 1537. The Commandinus edition appeared at Bologna in 1566, and the Halley edition at Oxford in 1710.

2. Italy

Leonardo da Vinci. No list of the Italian mathematicians of the 16th century would be complete without some mention of Leonardo da Vinci[1] (1452–1519), and such were the remarkable attainments of this gifted man that his name may properly be given in its chronological order, thus standing first on the record. He was born at Vinci, near Florence, resided in Florence, Milan, and Rome, went to France at the invitation of the king in the year 1516, and died near Amboise in 1519. Famous as a painter, sculptor, goldsmith, investigator of the circulation of the blood, general scientist, architect, and writer on mechanics, optics, and perspective, he would have ranked as a worthy mathematician had not his talents in this direction been obscured by his unusual gifts in these other lines. In applied mathematics he may be looked upon as one of the founders of the modern theory of optics. In geometry he distinguished between curves of single and double curvature, gave much attention to the subject of stellar polygons, was interested in constructions with a single opening of the compasses, and gave various correct or approximate constructions of regular polygons. In physics he knew the theory of the inclined plane, found the center of gravity of a pyramid, worked in the field of capillarity and diffraction, knew the camera obscura without a lens, and studied the resistance of the air and the effect of friction. The world has rarely produced such an all-round genius.

Early Workers in the Field of Equations. Since the solution of the equations of the third and fourth degrees was the chief mathematical achievement in Italian mathematics of the 16th century, it is proper to group together those scholars who were most prominent in this work. Although the details of their chief contributions will be reserved for Volume II, a brief statement of their achievements will now be given.

[1] P. Duhem, *Études sur Léonard de Vinci*, 2 vols., Paris, 1906–1913. For a popular sketch see D. Merejkowski, *The Romance of Leonardo da Vinci*, New York [1902].

Scipione del Ferro,[1] a native of Bologna, whom Cardan calls by his Latin name of Scipio Ferreus, was professor of mathematics in the city of his birth. In geometry he was interested in constructions depending on a single opening of the compasses. In algebra he found a method of solving the cubic equation for the special case of $x^3 + ax = b$.

In 1506 he revealed this method to his pupil Antonio Maria Fior,[2] a Venetian, who proceeded to turn the information to account in the popular mathematical contests that were then in vogue. Of the life of Fior little is known, but he is said by Tartaglia[3] to have been living in 1536.

Zuanne de Tonini da Coi[4] (*c.* 1530) was a teacher in Brescia and was interested in mathematics from the standpoint of problem solving.[5] In 1530 he sent as a kind of challenge to Tartaglia the two equations

$$x^3 + 3x^2 = 5$$

and

$$x^3 + 6x^2 + 8x = 1000.$$

For some time Tartaglia was unable to solve them, but, as we shall see in Volume II, he finally succeeded in doing so, this being an important step in the general problem of the cubic equation.

Cardan. The first of the two prime movers in the solution of the cubic was Giro'lamo Carda'no.[6] He was the illegitimate

[1] Born *c.* 1465; died at Bologna between October 20 and November 16, 1526. L. Frati, *Bollett. di bibliogr. d. sci. matem.*, XII, 1. The name also appears as Ferri and as Ferreo.

[2] This on the statement of Tartaglia (1546). Cardan (1545) says it was about 30 years earlier than the time of his writing, which would make it *c.* 1514 or 1515. The name appears with various spellings, particularly in the Latin form of Antonius Maria Floridus and in the form of Antoniomaria Fior.

[3] Pronounced tar tä'lya. See page 297.

[4] Also Zuane, Giovanni, Giovanno, John, with his last name sometimes given as Colle or in the Latin form of Colla.

[5] In a letter written in 1540, Tartaglia speaks of "that devil" having returned: "Eglie ritornato qui quel diauolo de Messer Zuanne Colle."

[6] Born at Pavia, 1501; died at Rome, September 21, 1576. The name appears as Hieronymus Cardanus and Jerome Cardan. He is commonly called Cardan by writers of English. H. Morley, *The Life of Girolamo Cardano*, 2 vols., London, 1854 (hereafter referred to as Morley, *Cardan*); Cantor, *Geschichte*, II, chaps. 64, 65, 66; V. Mantovani, *Vita di Girolamo Cardano*, Milan, 1821; Gherardi, *Facoltà Mat. di Bologna*, 47 seq.

son of a jurist, Facio Cardano (1444–1524), who was professor of jurisprudence and medicine in Milan and who edited Peckham's *Perspectiva communis*. Girolamo was a man of remarkable contrasts. He was an astrologer and yet a serious student of philosophy, a gambler and yet a first-class algebraist, a physicist of accurate habits of observation and yet a man whose statements were extremely unreliable, a physician and yet the father and defender of a murderer, at one time a professor in the University of Bologna and at another time an inmate of an almshouse, a victim of blind superstition and yet the rector of the College of Physicians at Milan, a heretic who ventured to publish the horoscope of Christ and yet a recipient of a pension from the Pope, always a man of extremes, always a man of genius, always a man devoid of principle. A certain bitter rival said of Voltaire, "Ce coquin-là has one vice worse than all the rest; he sometimes has virtues." So it was with Cardan.



PORTRAIT OF CARDAN
From the title-page of the first edition (1539) of his arithmetic

His *Ars Magna*,[1] the first great Latin treatise devoted solely to algebra, appeared at Nürnberg in 1545 and set forth the theory of algebraic equations so far as it was then known, including the solution of the cubic, which he seems to have secured from Tartaglia under pledge of secrecy and then dishonorably to have published, and the solution of the biquadratic which had been discovered by his pupil Ferrari. He also wrote on arithmetic,[2] astronomy,[3] physics,[4] and various other branches of knowledge,[5] proving himself a man of remarkable versatility and learning.

Tartaglia. Nicolo Tartaglia,[6] one of the greatest mathematicians of Italy in the 16th century, was born at Brescia. Although known as Tartaglia, we learn from his will[7] that his brother's name was Fontana. It is said that he was present as a child at the taking of Brescia by Gaston de Foix (1512) and at that time received a saber cut in the face which caused an imperfection in his speech. This gave him the nickname of Tartaglia ("the stammerer"), which name he formally used in his published works. He was self-educated but acquired such proficiency in mathematics that he earned a livelihood by teaching the science in Verona, Vicenza, Brescia, and Venice (1535).

Tartaglia seems to have substantially completed the solution of the cubic equation and, as already stated, to have imparted the secret to Cardan, who, in violation of his oath, published it in 1545.[8]

[1] *Artis Magnae, sive de regvlis algebraicis, liber vnvs*, Nürnberg, 1545; Basel, 1570. [2] *Practica arithmetice*, Milan, 1539.

[3] *De revolutione annorum, mensium et dierum . . . liber*, Nürnberg, 1547; *De temporum et motuum erraticarum restitutione*, Nürnberg, 1547; *Aphorismorum astronomicorum segmenta septem*, Nürnberg, 1547.

[4] *De subtilitate*, Nürnberg, 1550, with a Paris reprint in 1551.

[5] His *Opera* appeared in ten volumes, Lyons, 1663.

[6] Born *c.* 1506; died at Venice, December 13/14, 1557. The name is also spelled Tartalea. It is spelled in the text as it appears on the title-page of his work of 1556.

[7] Published by Boncompagni in 1881.

[8] On his possible indebtedness to Ferro see G. Enestrom, in *Bibl. Math.*, VII (3), 38.

Tartaglia was the first to apply mathematics to artillery science,[1] a subject just being perfected by the great French masters Galiot de Genouillac and Jean d'Estrées. He also wrote the best treatise on arithmetic[2] that appeared in Italy in his century, containing a very full discussion of the numerical operations and the commercial rules of the Italian arithmeticians. The life of the people, the customs of merchants, and the efforts at improving arithmetic in the 16th century are all set forth in this remarkable work. Tartaglia also published (1543) editions of Euclid and Archimedes.



From the title-page of *La Prima Parte del General Trattato*, Venice, 1556


On the question of the publication by Cardan of the Tartaglia solution of the cubic after a pledge of secrecy, a biographer of the former, but one who could not feel the influences of the 16th century, has this to say:

The attempt to assert exclusive right to the secret possession of a piece of information, which was the next step in the advancement of a liberal science, the refusal to add it, inscribed with his own name,

[1] *Nuova scienza, cioè Invenzione nuovamente trovata, utile per ciascuno speculativo matematico bombardiero . . .*, Venice, 1537; *Qvesiti ed invenzioni diverse*, Venice, 1546.

[2] *General Trattato di nvmeri, et misvre*, 2 parts (volumes), Venice, 1556–1560; *Tutte l' opere d' arithmetica del famosissimo Nicolò Tartaglia*, Venice, 1592, being substantially Volume I of the *General Trattato*.

to the common heap, until he had hoarded it, in hope of some day, when he was at leisure, of turning it more largely to his own advantage, could be excused in him only by the fact that he was rudely bred and self-taught, and that he was not likely to know better.



TARTAGLIA APPLIES MATHEMATICS TO ARTILLERY SCIENCE

From *Il Primo Libro delli Qvesiti, et Inventioni diverse*, Venice, 1546; 1562 ed.

Any member of a liberal profession who is miserly of knowledge, forfeits the respect of his fraternity. The promise of secrecy which Cardan had no right to make, Tartalea had no right to demand.[1]

As already remarked, however, it is difficult to imagine conditions in the year 1545, and it is hardly just to apply modern ethics to a situation so different from our own.

[1] Morley, *Cardan*, I, 270; Rixner and Siber, *Leben und Lehrmeinungen berühmter Physiker am Ende des XVI. und am Anfange des XVII. Jahrhunderts*, Sulbach, 1820; Firmiani, *Girolamo Cardano*, Naples, 1904. There is a brief but good biography of Tartaglia in D. Martines, *Origine aritmet.*, p. 61 n. See also A. Favaro, "Per la biografia di Niccolò Tartaglia," *Archivio storico Italiano*, 1913, and "Di Niccolò Tartaglia," in *Isis*, I, 329.

Ferrari. Lodovico Ferrari,[1] born in humble circumstances, was taken into Cardan's household in Milan at the age of fifteen. Cardan soon recognized his remarkable ability and made him his secretary. In spite of his ungovernable temper and his blasphemous habits he was later accepted by Cardan as his pupil and friend. Mathematical Italy would have given much to be in his place in Cardan's household; but, such were their quarrels, it would have given more to be out again. At the age of eighteen even Ferrari was glad to sever all relations with his patron and to begin teaching by himself in Milan. He was so successful there and in the mathematical contests of the day as to attract the attention of the court and of the Cardinal of Mantua. Through the favor of the latter he secured a position that brought him abundant means. He then became professor of mathematics at Bologna, but died there in the first year of his service, at the age of thirty-eight, probably poisoned by his only sister.

Zuanne de Tonini da Coi had proposed a problem which involved the equation

$$x^4 + 6x^2 + 36 = 60x.$$

This problem Cardan attempted to solve, and having failed he gave it to Ferrari. The latter succeeded in finding a method, and thus the solution of the equation of the fourth degree was discovered. Ferrari left no written works on mathematics, but Cardan published in the *Ars Magna* (1545) this noteworthy contribution to the theory of equations.

Bombelli. Rafael Bombelli (born *c.* 1530), a native of Bologna, was the last of those Italian mathematicians of the 16th century who contributed in any noteworthy way to the solution of the cubic and biquadratic equations. He wrote *L'Algebra parte maggiore dell' arimetica divisa in tre libri* and

[1] Born at Bologna, 1522; died *c.* 1560. The date of his death is also given as 1562 and 1565. Morley gives 1560. The Christian name is also written Luigi. Cardan left an unpublished *Vita Ludovici Ferrarii Bononiensis*. See also G. de' Sallusti, *Storia dell' Origine e de' Progressi delle Matematiche*, I, 38 (Rome, 1846).

published the work in Bologna in 1572.[1] In this work Bombelli set forth the reality of the three roots of a cubic equation in the case in which the cube root of an imaginary expression is involved in the result secured by the Tartaglia-Cardan rule. The book contained the most teachable and the most systematic treatment of algebra that had appeared in Italy up to that time.

Of Bombelli's life almost nothing is known. He is thought, from the introduction to his algebra, to have been an engineer in the service of the patron to whom he dedicates his book, Alessandro Rufini, bishop of Melfi.



FRANCESCO MAUROLICO
Engraved after a portrait from life

Francesco Maurolico. Of the mathematicians of this period who were interested in the Greek writers the most prominent was Frances'co Mauroli'co,[2] a native of Messina, Sicily, but of Greek parentage. He was a priest, at one time an abbot, and for some years professor of mathematics at Messina. He

[1] A second edition appeared at Bologna in 1579 with another title, *L'Algebra Opera*, with the dedicatory letter reset, but otherwise using the same sheets as the 1572 edition. The name is spelled as above in both editions of this work.

[2] Born at Messina, Sicily, September 16, 1494; died at Messina, July 21, 1575. Latin, Franciscus Maurolycus or Maurolykus; also known as Marullo. D. Scina, *Elogio di Francesco Maurolico* (Palermo, 1808); Martines, *Origine aritmet.*, 65. A life of Maurolico, written by his nephew, was published at Messina in 1613.

translated into Latin the works of Theodosius and Menelaus, the treatise of Autolycus on the sphere,[1] and the *Phaenomena* of Euclid, and published works on Apollonius[2] and Archimedes.[3] He also wrote various general works on mathematics and arithmetic,[4] wrote on mathematical induction,[5] and was a man of some creative power. A few of his more prominent contemporaries are listed below, with a brief statement of their contributions.

Italian Geometers. Federigo Commandino of Urbino (1509–1575) is known as one of the leading translators and editors of the Greek classics in mathematics. His editions of Euclid, Archimedes, Apollonius, Aristarchus, Heron, Ptolemy, and Pappus are highly esteemed.

Frances'co Baroz'zi[6] (*c.* 1538–*post* 1587), a Venetian nobleman, edited the commentary of Proclus on the first book of Euclid.[7] He also wrote on cosmography and geometry[8] and translated Heron's works.

Giambattista Benedetti[9] (1530–1590), a Venetian by birth, wrote on the geometry of a single opening of the compasses,[10] on the gnomon, on optics, and on the theory of numbers, and gave excellent graphic treatments of various problems.[11]

[1] Published at Messina, 1558. [2] Messina, 1654.

[3] Palermo, 1670 (the edition being lost in a shipwreck) and 1685.

[4] *Opuscula Mathematica*, 2 vols., Venice, 1575, although written in 1553. The *Arithmeticorum libri duo* (Venice, 1575, but written in 1557) was the second volume of the *Opuscula* and was republished in 1580. See also the *Elogia* above mentioned, p. 114.

[5] G. Vacca, "Maurolycus, the first discoverer of the principle of mathematical induction," *Bulletin of the Am. Math. Soc.*, XVI, 70; W. H. Bussey, "Origin of Mathematical Induction," *Amer. Math. Month.*, XXIV, 199.

[6] Franciscus Barocius, Francesco Barocci.

[7] *Procli Diadochi . . . in primum Euclidis . . . librum comment.*, Padua, 1560.

[8] *Geometricum problema tredecim modis demonstratum*, Venice, 1586.

[9] Giovanni Battista Benedetti, Joannes Baptista Benedictus.

[10] *De resolutione omnium Euclidis problematum aliorumque . . .*, Venice, 1553. On the general history of this important subject see W. M. Kutta, "Zur Geschichte der Geometrie mit constanter Zirkelöffnung," *Nova Acta . . . Abh. der K. Leop.-Carol. Deutschen Akad. der Naturforscher*, Halle, LXXI, 71, and *Bibl. Math.*, X (2), 16.

[11] *Diversarvm Specvlationvm Mathematicarum, & Physicarum Liber*, Turin, 1580. See *Rara Arithmetica*, p. 364.

Cosimo Bar'toli (1503–1572), a Florentine geometer, translated into Italian the works of the French mathematician Oronce Fine[1] and wrote a popular work on mensuration.[2] Among the features of the book is a table of squares to 662^2.

Pietro Antonio Catal'di[3] (1548–1626) was a native of Bologna and spent the closing years of his life there. He was professor of mathematics and astronomy at Florence (1563), Perugia (1572), and Bologna (1584). He wrote several mathematical works and to him are due the first steps in the theory of continued fractions, although not the first idea of these forms. His *Prima Parte della Pratica Aritmetica* (Bologna, 1602) and *Trattato dei numeri perfetti* (Bologna, 1603) were printed under the pseudonym of Perito Annotio, formed by transposing the letters in his given names. The second part of the *Pratica* appeared under his own name in 1606, and similarly for the third and fourth parts (1617, 1616). He also edited the first six books of Euclid's *Elements*,[4] wrote a brief treatise on algebra,[5] and contributed to the theory of roots (1613), the quadrature of the circle (1612), and various other subjects.

Matteo Ricci. Among the contemporaries of Cataldi the one who did most for the spread of mathematics in remote lands was Matteo Ricci, a man of remarkable energy and of great influence through his work in China.[6] He entered the Jesuit order in 1571, left Rome for China in 1577, and reached

[1] *Opere di Orontio Fineo del Delfinato; Diuise in cinque Parti: Arimetica Geometria, Cosmografia, & Orinoli, Tradotte Da Cosimo Bartoli, Gentilhuome & Academico Fiorentino*, Venice, 1587 (posthumous). See *infra*, page 308.

[2] *Del Modo di Misurare le distantie, le superficie, i corpi, le piante, . .* Venice, 1564, with possibly an earlier edition.

[3] Although the spelling Cattaldi is given on the title-page of his *Dve Lettion di Pietr' Antonio Cattaldi* (Bologna, 1577), the name is generally given in his other works as Cataldi. The above-named book is curious because the printer not having fraction forms, was obliged to insert all fractions by hand. The name also appears as Cataldo.

[4] Bologna, 1620.

[5] *Regola della Quantità, o Cosa di Cosa*, Bologna, 1618.

[6] Born at Macerata, Ancona, October 6, 1552; died at Peking, May 8 (or 11) 1610. His Chinese name was Li-ma-to, derived from Ri (Chinese Li, for Ricci) and Matteo. H. Bosmans, *Revue des Quest. scient.*, January, 1921.

Canton in 1578.[1] Here he did more than any of his predecessors to make known in that country the mathematics and astronomy of the West. With the help of native scholars, the most prominent of whom were two learned mandarins, Hsü Kuang-ching (1562–1634) and Li Chi Ts'ao (died 1631), he translated (1603–1607) into Chinese the first six books of Euclid's *Elements*.[2] He also wrote an arithmetic,[3] which he dedicated to his assistant, Li Chi Ts'ao, and compiled various astronomical works.[4]

Minor Writers. In the second half of the century Silvio Belli (died 1575) wrote on practical geometry[5] and on the theory of proportion.[6] His geometry was very popular, six editions appearing in the 16th century.

Another writer whose work attracted considerable attention in the 16th and 17th centuries was Petrus Bongus,[7] to take the Latin form of his name as it appears in the first edition of his work. He was a native of Bergamo and became canon of the cathedral in that city. His work of nearly 500 pages on the mystery of numbers[8] went through several editions. It contains a mass of information upon such subjects as the religious significance of three, seven, and other numbers.

The Italian Arithmeticians. In all the leading countries many arithmetics were printed in the 16th century,[9] but for our present purposes it suffices to mention only a few of the more prominent Italian writers.

Girolamo and Giannantonio Tagliente,[10] Venetian arithmeticians of *c.* 1500, wrote a work on commercial arithmetic[11] which

[1] Pietro Tacchi Venturi, S. J., *L'Apostolato del P. Matteo Ricci*, 2d ed., Rome, 1010 [2] *Ki-ho-yüan-pen*. [3] *Tung-wen-suan-ki*

[4] P. F. S. Vella, "Del P. Matteo Ricci," *Memorie, Pontificia Accad. dei Nuovi Lincei*, Rome, XXVIII, 51.

[5] *Libro del misurar con la vista*, Venice, 1565, with later editions in 1566, 1560, 1570, 1573, and 1595.

[6] *Della proportione, et proportionalita*, Venice, 1573.


[7] Died at Bergamo, September 24, 1601.

[8] *Mysticae Nvmerorvm significationis liber in dvas divisvs partes*, Bergamo, 1583–1584.


[9] For the list, consult *Rara Arithmetica*. [10] Pronounced tä lyen'tä.

[11] *Opera che insegna A fare ogni Ragione de Mercãtia*.

以乙為心。則乙甲線。與乙丙乙丁線亦等。何者。凡為圜。自心至界各線俱等故。界說十五。既乙丙等于乙甲。而甲丙亦等于甲乙。即甲丙亦等于乙丙。公論一。三遍等如所求。凡論有二種。此以是為論者。正論也。下倣此。



其用法不必作兩圜。但以甲為心。乙為界。作近丙一短界線。乙為心。甲為界。亦如之。兩短界線交處即得丙。



諸三角形。俱推前用法作之。詳本篇二十

第二題

一直線線或內或外有一點。求以點為界作直線與元線

FROM RICCI'S TRANSLATION OF EUCLID, 1603–1607

From a manuscript copy of this translation, made in the 17th century. This page shows the first proposition of Book I

appeared in Venice in 1515 and was so popular that it went through more than thirty editions in the 16th century.

Francesco Ghaligai,[1] a Florentine arithmetician, published in 1521 a mercantile work entitled *Summa De Arithmetica*,[2] of which two other editions appeared later.

Francesco Feliciano da Lazesio[3] published an elementary work on arithmetic, algebra, and practical geometry at Venice in 1517/18.[4] This work went through at least fourteen editions (including the revision of 1526) in the 16th century and several in the century following.

Giovanni Sfortunati[5] published at Venice in 1534 a work on commercial arithmetic[6] which was well received and went through several editions.

The first mercantile tables that had great popularity were published in 1535 by a Venetian arithmetician, Giovanni Mariani,[7] under the title *Tariffa perpetva*. The work was often reprinted.

3. France

Centers of Activity. The centers of mathematical activity in France in the 16th century were Paris and Lyons, the former because of its ancient importance in all matters intellectual and the latter because of its commercial supremacy and the desire to cultivate some of the idealism of the northern capital. The theoretical books appeared more frequently from the Paris presses, while the output of what is known as practical mathematics was fairly large among those whom a recent writer has called "the morose and inhospitable Lyonese, . . . in whose

[1] Pronounced gä lē gä'ē. He died February 10, 1536.

[2] The edition of 1548 had the title *Practica d'Arithmetica*.

[3] Born at Lazisa, near Verona, *c.* 1490; was living in 1536. The family name is pronounced fa lē'chē ä no. Lazesio also appears as Lazisio.

[4] *Libro de Abaco*. There was a second work (Venice, 1526), called the *Libro di Arithmetica & Geometria . . . Intitulato Scala grimaldelli*, but this was only a revision of the work above mentioned.

[5] Pronounced sfor tōō nä'tē. He is also known by the Latin form Johannes Infortunatus. Born at Siena, *c.* 1500.

[6] *Nvovo Lvme Libro di Arithmetica*.

[7] Born at Venice, *c.* 1500. The name also appears as Zuane Mariani.

esteem the pick of humanity is the prosperous silk merchant." Paris was not yet a metropolitan city in the ecclesiastical sense, the ancient Roman political divisions having been retained by the Church, and the capital city being then and until 1622 subordinate to the metropolitan city of Sens. All this had no effect, however, upon her intellectual and political supremacy over all France.

Theoretical Works. The first of the French writers who sought to maintain the standing of the Greek mathematics in the intellectual atmosphere of France was Jacques le Fèvre d'Estaples,[1] known in his Latin works as Jacobus Faber Stapulensis. He was a "Doctor Sorbonnicus," a priest, vicar of the bishop of Meaux, lecturer on philosophy at the Collège Lemoine in Paris, and tutor to the son of François I. He wrote an introduction to the arithmetic of Boethius[2] and a work on geometry, edited (1499) Sacrobosco's *Sphere* and a description of the number game of Rithmomachia, and published various other works. His own writings were heavy and theoretical, and expressed the dying body of medieval mathematics.[3]

Charles de Bouelles,[4] canon and professor of theology at Noyon, wrote on geometry[5] and the theory of numbers.[6] The latter work includes a book on perfect numbers. He is particularly worthy of attention, however, because of his work

[1] Born at Estaples, near Amiens, *c.* 1455; died at Nérac, *c.* 1536. The French forms also appear as Febvre and Étaples.

[2] *Introductio Jacobi fabri Stapulēsis, in Arithmecam Diui Seuerini Boetij*, published in a volume with other works, Paris, 1503, the above title being from the edition of *c.* 1507. In these two editions his geometry and perspective also appear. The first edition of his compendium appeared in 1488. For details, see *Rara Arithmetica*, pp. 27, 30, 80, 82.

[3] Baldi's estimate of his merits is exaggerated: "D'ingegno felicissimo attese con gran frutto ad ogni sorte di dottrina, e giunse all' eccellenza di maniera che fu giudicato meraviglia del suo secolo." *Cronica*, 107.

[4] Born at Saucourt, Picardy, *c.* 1470; died at Noyon, *c.* 1553. The name appears also as Charles Bouvelles, Boüelles, Bouilles, and in the Latin form of Carolus Bovillus.

[5] *Geometricae introductionis libri VI*, Paris, 1503; *Livre singulier et utile, touchant l'art et pratique de Géométrie*, Paris, 1511, with several later editions.

[6] *Liber de duodecim numeris*, part of his general work published at Paris in 1509/10.

on the cycloid, he being one of the first to consider this figure from the scientific standpoint. He also wrote on regular convex and stellar polygons.

Minor Writers. One of the most pretentious of the mathematicians of his time, and one of the least worthy, was Oronce Fine,[1] more commonly known by the Latin form of his name, Orontius Fineus.[2] In his young manhood (1518–1524) he was imprisoned on account of his opposition to the Concordat, an agreement between France and the Pope. Upon being released he devoted himself to teaching, and about 1532 became professor of mathematics in the newly founded institution which was later known as the Collège de France.[3] He wrote extensively on astronomy and produced several works on arithmetic and geometry,[4] including one on the quadrature of the circle.[5] Some of his works were translated into Italian by Cosimo Bartoli. While he enjoyed some reputation, he died in poverty and his works were soon forgotten.

Among several French physicians of the 16th century who devoted much attention to mathematics the only one of great distinction is Jean Fernel[6] (1497–1558). He received his degree in medicine at Paris in 1530, and four years later he had so risen in his profession as to be called to a chair in the faculty. His admirers spoke of him as the modern Galen, and his *Universa Medicina* went through more than thirty editions. In the field of mathematics he published (1528) a work of the

[1] Born at Briançon, 1494; died in Paris, October 6, 1555.

[2] As spelled in the first edition of his *Protomathesis*, Paris, 1530–1532. The name is also spelled Finaeus. There is no warrant for the spelling Finé.

[3] Ramus speaks of him as the one "qui primus regia professione in Galliam mathematicas artes retulit." Introduction to his *Libri dvo*, p. 3 (Basel, 1569).

[4] Both of these subjects and some parts of astronomy are considered in his *Protomathesis*. Among his other works are *In sex priores libros geometricorum elementorum Euclidis demonstrationes*, Paris, 1536; *De re et praxi Geometrica libri III*, Paris, 1555; and *De rebus mathematicis, hactenus desideratis, Libri IIII*, with a biography, Paris, 1556.

[5] *De quadratura circuli*, Paris, 1544. This was severely attacked by Buteo. See *Bibl. Math.*, XII (3), 250.

[6] Joannes Fernelius, as the Latin form appears in his *De Proportionibus Libri duo*, Paris, 1558.

Boethian type on proportion, and his computation of the length of a degree of the meridian was so satisfactory[1] as to entitle him to a worthy place in the history of geodesy.

Among those who may properly be called the dilettanti mathematicians of the time was Claude de Boissière,[2] who wrote on poetry and music as well as on astronomy and arithmetic. His arithmetic,[3] a combination of the mèdieval theory and the contemporary practice in calculation, is one of the many books of the time that, as a result of the Hundred Years' War, related mathematics to the science of warfare.

At about the same time François de Foix, Comte de Candale (*c.* 1502–1594), another of the dilettanti and a bishop in southern France, was interested in a better translation of Euclid's *Elements*,[4] but he contributed nothing to the general theory of geometry.

Ramus. Pierre de la Ramée,[5] better known by the Latin form of his name, Petrus or Peter Ramus, descended from a noble but impoverished family. His grandfather had been driven from his estates in Burgundy and had been forced to become a charcoal burner, and his father was a humble peasant. Pierre early showed unusual intellectual powers, and after many struggles obtained employment as a servant to a rich student in the Collège de Navarre at Paris. By working in this capacity during the day and studying at night he made his way to the master's degree. It was on his examination in 1536, when he was only twenty-one years old, that he attracted

[1] He made it 56,746 French toises instead of 57,024.

[2] Claudius Buxerius. Born in the province of Grenoble, *c.* 1500.

[3] *L'art d'Arythmetiqve contenant tovte dimention, tres-singvlier et commode, tant pour l'art militaire que autres calculations*, Paris, 1554.

[4] *Euclidis . . . Elementa geometrica, Lib. XV . . . restituta. His accessit decimus sextus liber . . .*, Paris, 1566. There was another edition, "Novissime collati sunt XVII^{us} et XVIII^{us} priori editione . . .," Paris, 1578. His name appears in the Latin phrase "Auctore Francisco Flussate Candalla." For various other editors and translators of Euclid in this period see P. Riccardi, *Saggio di una Bibliografia Euclidea*, Bologna, 1887; hereafter referred to as Riccardi, *Saggio Euclid.*

[5] Born at Cust (Cultia, Cusia, Cus, Cuz, Cuth, Cut), Picardy, 1515; died August 26, 1572.

the attention of intellectual Europe by his audacious attack upon one of its idols, his thesis being "All that Aristotle has said is false."[1] Soon after this he began his career of teaching and was not long in attaining a high position in his profession. For many years (from 1546) he was principal of the Collège de Presles and held a professorship (from 1551) in the Collège de France. He was an orator of great power and a skillful debater, but his brilliant career was closed at the massacre of St. Bartholomew's Day, August 26, 1572. Although his work was chiefly in philosophy and the humanities, he devoted much attention to mathematics,[2] editing the *Elements* of Euclid[3] and writing on theoretical arithmetic,[4] geometry,[5] and optics.

Vieta. The greatest of all the French mathematicians of the 16th century was François Viète,[6] Seigneur de la Bigotière, better known by the semi-Latin name of Vieta. As a young man he practiced law in his native town, afterward taking up a political career and becoming a member of the Bretagne parliament. His first work on mathematics appeared in Paris in 1579. In 1580 he became master of requests at Paris, and later was a member of the king's privy council. Under these circumstances he was able to devote much leisure time to the

[1] *Quaequmque ab Aristotele dicta essent, commentia esse.*

[2] There is a good summary of his life in a work by F. P. Graves, *Peter Ramus*, New York, 1912, with a bibliography (p. 219) of the publications of Ramus and of secondary sources of information.

[3] Paris, 1545 and 1549. See Boncompagni's *Bullettino*, II, 389.

[4] *P. Rami, eloquentiae et philosophiae professoris regii, arithmeticae libri tres*, Paris, 1555; Paris, 1557; Basel, 1567; Paris, 1584. Also a *Libri Duo*, Paris, 1569, 1577, and 1581; Basel, 1580; Frankfort, 1586, 1591, 1592, 1596, and 1599; Lemgo, 1599; English translation, London, 1593. See *Rara Arithmetica*, pp. 263, 330, 335.

[5] Paris, 1577; English translation, London, 1636. See also his *Scholarvm Mathematicarvm, Libri vnvs et triginta*, Basel, 1569, in which he criticizes Euclid's arrangement of the *Elements* from the point of view of logic.

[6] Born at Fontenay-le-Comte (Fontenay-Vendée), 1540; died in Paris, December 13, 1603. The French form also appears as Viet, Viette, and de Viette, and the Latin as Vietaeus. There is a good sketch of his life in the *Penny Cyclopaedia*, London, 1843, by De Morgan, with a summary of all his works. See also J. L. F. Bertrand, "La vie d'un savant au XVI. siècle, François Viète," in his *Éloges académiques*, p. 143 (Paris, 1902); F. Ritter, *François Viète*, Paris, 1905; G. Gambier, *Le mathématicien François Viète*, La Rochelle, 1911.

study of mathematics, and the results were such that he ranks as one of several notable instances of a man attaining high standing in this science although not devoting himself chiefly to it until rather late in life. Vieta, indeed, remarked in a letter to Adriaen van Roomen that he did not profess to be a mathematician, but was merely one to whom mathematical studies were delightful in his hours of leisure.



FRANÇOIS VIÈTE (VIETA)

From an old lithograph of a portrait from life

Vieta wrote chiefly on algebra,[1] but he was also interested in geometry,[2] the calendar, and mathematics in general. In connection with the Gregorian reform of the calendar he acquired much unfortunate notoriety through his bitter antagonism to Clavius and through his wholly unscientific attitude. He was an expert in deciphering, for the government, the cryptic writing of diplomatic correspondence.

[1] *Isagoge in artem analyticam, De aequationum recognitione et emendatione libri duo, De numerosa potestatum purarum atque adfectarum ad exegesin resolutione tractatus,* all published privately by Vieta, and republished by Frans van Schooten, Leyden, 1646, in the *Opera mathematica* of Vieta. See also Boncompagni's *Bullettino*, I, 223, 245. There was a French translation of his treatment of equations made by one J. L. de Vaulezard, *Les Cinq Livres des Zetetiques*, published at Paris in 1630. There was an *Algebre de Viete d'vne methode novvelle, claire, et Facile*, by one James Hume (Iac Hvmivs), a Scotchman, Paris, 1636, but it is only after the style of Vieta.

[2] *Effectionum geometricarum canonica recensio*, and *Supplementum Geometriae*, Paris, 1593.

Vieta's Work in Algebra. It will be shown in Volume II that Vieta contributed extensively to the development of algebra and trigonometry, but a brief reference to his work is appropriate in this connection. He was among the first to employ letters to represent numbers in algebra, often using vowels for the unknowns and consonants for the knowns. He found the formula for $\sin n\phi$ in terms of $\sin\phi$; made an advance towards proving that an equation of the nth degree is made up of n linear factors; showed how to increase, decrease, multiply, or divide the roots of the equation $f(x) = 0$ by k; gave one of the earliest methods of evaluating π by infinite products; applied algebra to geometry in such way as to lay a foundation for analytic trigonometry; indicated powers more simply than his predecessors had done, using Aq for the square of the unknown, Ac for its cube, Aqq for its fourth power, and so on; and showed clearly the relation between the problems of the trisection of an angle and the solution of a cubic equation. His interesting combination of infinite products and series is seen in his statement that

$$\frac{2}{\pi} = \sqrt{\tfrac{1}{2}} \cdot \sqrt{\tfrac{1}{2} + \tfrac{1}{2}\sqrt{\tfrac{1}{2}}} \cdot \sqrt{\tfrac{1}{2} + \tfrac{1}{2}\sqrt{\tfrac{1}{2} + \tfrac{1}{2}\sqrt{\tfrac{1}{2}}}} \cdots.$$

Minor Writers. Of the writers on the theory of mathematics at this period the only one of any note who published his works at Lyons was Joannes Buteo,[1] a brother and afterward general of the order of St. Anthony. He wrote chiefly on geometry[2] and arithmetic.[3] His geometry refuted various pretensions of Oronce Fine as to the quadrature of the circle.

[1] Born at Charpey, *c.* 1485–1492; died at Caam, *c.* 1560–1572. These two places are in Dauphiné, France. The dates given by various writers differ greatly. See also Boncompagni's *Bullettino*, XIII, 258, 265 n. This is the form in which the family name appears in his *Logistica*, Lyons, 1559, that is, *Ioan. Bvteonis Logistica*. It is also given as Boteo, Jean Butéon, Bateon, Borrel, and Borell.

[2] *Opera Geometrica*, Lyons, 1554; *De Quadratura Circuli, Libri II*, Lyons, 1559.

[3] *Logistica qvae & Arithmetica vulgò dicitur in libros quinque digesta*, Lyons, 1559; the most original of his works. See *Rara Arithmetica*, p. 292; G. Wertheim, "Die Logistik des Johannes Buteo," *Bibl. Math.*, II (3), 213.

Another instance of a French mathematician of some ability publishing outside of Paris, and in this case outside of France, is that of Francesco dal Sole (born *c.* 1490). He wrote in Ferrara, Italy, and published an arithmetic in Venice.[1] The only feature of the book worth mentioning is the combination which it makes of the concepts of number and space.[2]

Perhaps the most elaborate arithmetics published in France in the 16th century, and among the least practical, were those of Pierre Forcadel.[3] Little is known of this writer except that he lived for a long time in Italy and finally, through the efforts of Ramus, was called to Paris as professor of mathematics in the Collège Royal. He translated Euclid I–VI and parts of the works of Proclus, Archimedes, and other writers.

Practical Mathematics. The earliest of the Lyons school of arithmeticians, and one of the most brilliant as well as most unscrupulous, was Estienne de la Roche,[4] known as Villefranche, although a native of Lyons. He was a pupil of Chuquet, and in his arithmetic[5] he appropriated a large amount of material from a manuscript of the latter which has since been published. Perhaps no other French arithmetic of the 16th century gives a better view of the methods of computation and of the commercial applications of the subject.[6]

The second noteworthy writer of the southern school was Jacques Peletier,[7] a native of Le Mans, who is also known by the Latin name of Peletarius. He was head of a college at Bayeux (1547), secretary to the bishop of Le Mans, a physician at Bordeaux (1550), Poitiers, Lyons, and Paris, and finally head of a college at Le Mans. He contributed to general

[1] *Libretti nvovi con le regole Di Francesco Dal Sole Gallo*, Ferrara, 1546. The first edition was Venice, 1526. There was a third edition in 1564.

[2] Thus he has "Regola delle additione in generalita, tanto geometrica, quanto arithmetica."

[3] Born at Béziers; died in Paris, 1574. For description of the various works see *Rara Arithmetica*, p. 284.

[4] Born at Lyons, *c.* 1480.

[5] *Larismethique nouellement composee par maistre Estienne de la roche dict Villefrãche natif de Lyõ*, Lyons, 1520. *Rara Arithmetica*, 128; Cantor, *Geschichte*, II, chap. 59.

[6] A. Marre, Boncompagni's *Bullettino*, XIII, 573.

[7] Born at Le Mans, 1517; died in Paris, July, 1582.

literature and to elementary mathematics. His arithmetic[1] was published both at Poitiers (1549) and at Lyons (1554). He also wrote on algebra,[2] Euclid's *Elements*, the geometry of lines and angles, and the circle. He equated the terms of an equation to zero and stated that, when all roots are integral, any root is a factor of the last term.

Another Lyons arithmetician of considerable note appeared in the person of Ian Trenchant (born *c.* 1525). His arithmetic[3] includes the usual commercial applications and the operations with counters as well as with common numerals.

Among the practical tables published at Lyons are a set prepared by Monte Regal Piedmontois, professor of mathematics in the University of Paris. These tables[4] are beautifully printed on vellum and copies are very rare. The work contains the products of numbers to 100 × 1000, and the author speaks of having published part of the tables in Venice in 1575.

4. ENGLAND

English Writers. England was later than Italy or France in her appreciation of mathematics, or at least in her publication of works on this subject.[5] Although there is some mention of arithmetic in the early works,[6] it was not until 1522 that a book devoted wholly to mathematics was printed in Great Britain,—the erudite but dull arithmetic[7] of Tonstall.

[1] The 1607 edition has the title *L'Arithmetiqve de Iacqves Peletier dv Mans, Departie en quatre liures.* There were other editions. *Rara Arithmetica*, 245.

[2] *L'algèbre départie en deux livres*, Lyons, 1554.

[3] *L'Arithmetiqve de Ian Trenchant, Departie en trois liures*, Lyons, 1566. This title is from the 1578 edition. There were several editions. On an edition of 1558 see *Bibl. Math.*, II (3), 356.

[4] *Invention novvelle et admirable, pour faire toute sorte de cõpte*, Lyons, 1585.

[5] See also R. C. Jebb, in the *Cambridge Mod. Hist.*, Vol. I, chap. xvi, and J. Gairdner, *ibid.*, chap. xiv.

[6] For example, in Caxton's *Mirrour of the World or Thymage of the same*, London, 1480, translated from the French, there is a chapter (10) beginning, "And after of Arsmetrike and whereof it proceedeth," but this cannot be called a treatise on the subject. There is a recent edition by O. H. Prior, London, 1913 (for 1912).

[7] *De Arte Svppvtandi libri qvattvor Cvtheberti Tonstalli*, London, 1522.



TITLE-PAGE OF THE TONSTALL WORK OF 1522


Tonstall. Cuthbert Tonstall[1] was recognized as a man of great learning and influence. He was born in Hackforth, Yorkshire, went to Oxford in 1491, left on account of the plague and entered Cambridge, and afterward went to Padua, where he took the degree of doctor of laws. In 1511 he became vicar-general to the Archbishop of Canterbury, was presented at court, and soon thereafter received various ecclesiastical and diplomatic appointments. Erasmus speaks in the highest terms of his remarkable attainments at this period of his career. In 1522 he was promoted to the bishopric of London, but continued for some time in his diplomatic career. In 1530 he was made bishop of Durham. In the troublous times of Edward VI, however, no man of prominence was safe, and Tonstall was accused of conspiracy in 1552, was deprived of his bishopric, and was imprisoned in the Tower. On the accession of Mary he was restored to his bishopric of Durham, but under Elizabeth he was again deprived of the honor (1559) and died a few months later.

Tonstall relates in his work that in his dealing with certain goldsmiths he suspected that their accounts were incorrect and therefore renewed his study of arithmetic so as to check their figures. Having been educated in part at Padua, in a country that was still the leader in commercial arithmetic, he was familiar with business methods of computing and was quite prepared to write a book on the subject. On his appointment to the See of London he bade farewell to the sciences by publishing this work. He dedicated the book to one of the greatest scholars and one of the noblest men of his generation, Sir Thomas More, who, in the *Utopia* (1516), "the only work of genius that she [England] can boast in this age," as Hallam characterized it, had spoken of him as his "colleague and companion . . . that incomparable man Cuthbert Tonstal," whose "learning and virtues are too great for me to do them justice."

[1]Born 1474; died at Lambeth Palace, November 18, 1559. The name is also spelled Tonstal and Tunstall. The first name is commonly spelled Cuthbert, but the spelling in the first edition of his work is Cuthebert. The border of the title-page of this edition, shown on page 315, was engraved by Holbein, whose initials appear at the left. The border had been used in an earlier book.

Some further idea of the intellectual group in which Tonstall moved may be obtained from the fact that it was his friend Margaret Roper, More's daughter, whom Erasmus addressed as the "ornament of thine England."[1]

The arithmetic of Tonstall was not original, the material being confessedly drawn from such Italian writers as Pacioli, but the arrangement was good and the presentation was clear even if unnecessarily extended. As for an arithmetic written in Latin, however, the time had passed for such a book to be popular in England.[2]



ROBERT RECORDE

From a recently discovered oil portrait on wood, apparently made from life, and now in the possession of W. F. Bushell, Fleetwood, Lancashire. The painting bears the inscription "Rob^{t} Record. M.D. 1556," but this is now so darkened by age as not to show in the photograph


Recorde. The most influential English mathematician whose works were published in the 16th century was Robert Recorde.[3] He entered Oxford *c.* 1525 and became a fellow of All Souls College in 1531. In 1545 he received the degree of M. D. at Cambridge. He taught mathematics in private classes both at Oxford and at Cambridge, but after receiving his degree in medicine he went to London and became physician to Edward VI and Queen Mary, perhaps absorbing

[1] "Margareta Ropera Britanniae tuae decus."

[2] John Gill, *Systems of Education*, p. 1. London, 1876.

[3] Born at Tenby, Pembrokeshire, *c.* 1510; died in London, 1558. In some editions of his works the name is spelled Record.

some of his educational ideas from Roger Ascham (1515–1568), who was then the Latin secretary to each of these rulers. He was also versed in the law, but this did not keep him from dying in prison. In his will he describes himself as "Robert Recorde, doctor of physicke, though sicke in body yet whole of mynde." The cause of his imprisonment is not known. Although a doubtful tradition says that it was for debt, it is more probable that it was for some misdemeanor in connection with the mines in Ireland, where he was for a time "Comptroller of Mints and Monies." Recorde may be said to have been the founder of the English school of mathematics, inasmuch as he wrote in the English language and showed originality in the treatment of his subjects and in his method of presentation.



FROM THE PROBATE OF RECORDE'S WILL

It is often stated that the will bears the date June 28, 1558. The will is not in existence, so far as known, but the official copy is preserved, and the record shows that this date of probate should be read as xviii die mensis Junii


Recorde's Mathematical Works. Recorde published four books on mathematics and one on medicine, all of which are now extant, and he seems to have written others which have been lost and which were probably never printed. The four mathematical works were written in dialogue, a custom not uncommon at that time,[1] and bore various fanciful names, as shown in the following list:

1. *The Grovnd of Artes*, printed in London between 1540 and 1542, of which no copy of the first edition is known to be extant. This was one of the most popular arithmetics printed in the 16th century. It went through at least eighteen editions

[1] The custom appears in the Middle Ages as well, for example, in a 10th century MS. at Munich (S. Günther, *Geschichte des mathematischen Unterrichts im deutschen Mittelalter*, p. 26 n. (Berlin, 1887), hereafter referred to as Günther, *Math. Unterrichts*), but of course goes back to Plato. It appears in the various editions of the anonymous *Rithmimachia* (*Rara Arithmetica*, p. 63), in the arithmetic of Thierfelder (Nürnberg, 1587), and in numerous other works.

before 1601 and at least eleven more in the next century. Well did a writer of 1662 remark that this book was "entail'd upon the People, ratified and sign'd by the approbation of Time." The work includes computation by counters as well as by written figures, and contains the usual commercial topics which European countries north of the Alps had derived from Italy. It is commonly but incorrectly referred to as the first arithmetic printed in the English language, and it is in fact the earliest one of much lasting influence; but Recorde mentions the existence of other works of this kind, saying:

I doubt not but some will like this my booke aboue any other English Arithmetike hitherto written, & namely such as shal lacke instructers, for whose sake I haue plain-ly set forth the exãples, as no book (that I haue seene) hath hitherto.


The whetstone of witte,

whiche is the seconde parte of Arithmetike: containyng thextraction of Rootes: The Coßike practise, with the rule of Equation: and the woorkes of Surde Nombers.

Though many stones doe beare greate price,
The whetstone is for exercise
As needefull, and in woorke as straunge:
Dulle thinges and harde it will so chaunge,
And make them sharpe, to right good vse:
All artesmen knowe, thei can not chuse,
But vse hir helpe: yet as men see,
Noe sharpenesse semeth in it to bee.
The grounde of artes did brede this stone:
His vse is greate, and moare then one.
Here if you list your wittes to whette,
Moche sharpenesse therby shall you gette.
Dulle wittes hereby doe greately mende,
Sharpe wittes are fined to their fulle ende.
Now proue, and praise, as you doe finde,
And to your self be not vnkinde.

¶ These Bookes are to bee solde, at the Weste doore of Poules, by Iohn Kyngstone.

TITLE-PAGE OF RECORDE'S ALGEBRA, 1557

The first work devoted chiefly to algebra and the theory of numbers to be printed in England

2. *The Castle of Knowledge*, printed in London in 1551, a work on astronomy, and one of the first to bring the Copernican system to the attention of English readers.

3. *The pathewaie to knowledge*, printed in London in 1551, containing an abridgment of Euclid's *Elements*.

4. *The whetstone of witte*, printed in London in 1557, "containyng the extraction of Rootes: The *Cossike* practise, with the rule of *Equation*: and the woorkes of *Surde Nombers*." The "cossic art" was another name for algebra, or, as Recorde says, the subject that begins with "The rule of equation, commonly called Algebers Rule." In this work the modern sign of equality first appears in print.

Such fanciful names as he uses for his titles were then the fashion, as is seen in numerous other works of the time.[1]

As to his arithmetic, England was just beginning to feel the need for such a book. In Elizabeth's reign, extending from 1558 to 1603, mercantile England came to the front, the native powers of the country were developed, new manufactures were introduced, and artisans from all over Europe were encouraged to enter the employ of her nobles and her merchant princes.[2] Never was there a better opportunity for a commercial arithmetic, and never was the opportunity more successfully met.

As already stated, the *Grovnd of Artes* was not the first popular arithmetic in the English language. In 1537 there appeared at St. Albans an anonymous work entitled *An Introduction for to lerne to reken with the Pen and with the Counters, after the true cast of arismetyke or awgrym in hole numbers, and also in broken*, and this was reprinted in 1539, 1546, 1574, 1581, and 1595, although it never ranked with Recorde's work either in scholarship or in popularity.

Recorde's other works were, naturally enough, not so successful, but they filled the needs for which they were written. It was no slight honor to have it said that "he was the first that ever writ of astronomy in the English tongue,"[3] even though the statement is exaggerated.


[1] *E.g.*, *Mirror for Magistrates*, *A Gorgeous Gallery of Gallant Inventions*, *Groat's Worth of Wit*, *Pap with a Hatchet*. See G. Saintsbury, *History of Elizabethan Literature*, p. 11 (London, 1887).

[2] W. Cunningham, in the *Cambridge Mod. Hist.*, Vol. I, chap. xv. London, 1902.

[3] Aubrey, *Brief Lives*, II, 200.

The first rival to Recorde's *Grovnd of Artes*, and the only serious one that appeared in Great Britain for a hundred years, was *The Well spring of Sciences*,[1] written by one Humphrey Baker[2] in 1562 and published in 1568. It went through five editions before 1601 and several after that date. The work is a commercial arithmetic, is evidently under many obligations to Recorde, and was written to meet the criticism of continental scholars on the backward state of the subject in England.[3]

Leonard and Thomas Digges. For about a hundred years, beginning in the second half of the 16th century, the effect of



MILITARY MATHEMATICS IN 1572

From the *Arithmeticall Militare Treatise, named Stratioticos*, by Leonard and Thomas Digges, London, 1572. This cut is from the 1579 edition

the continental wars showed itself in the textbooks on arithmetic and practical geometry, particularly in England and France. Problems relating to military affairs became more numerous, and even the titles of mathematical works bore evidence of this tendency. The most notable example in England is seen in a work by Leonard Digges (died *c.* 1571) and

[1] The form in which the title appears in the 1580 edition.

[2] He was a native of London and died after 1587. In some editions the name is spelled Humfrey.

[3] See the quotations in *Rara Arithmetica*, p. 327.

his son Thomas Digges (died 1595), entitled *An Arithmeticall Militare Treatise, named Stratioticos* (London, 1572). Leonard Digges came of an ancient Kentish family. He studied at University College, Oxford, but took no degree. He was a mathematician of ability, his chief interest lying in the application of the science to surveying, military engineering, and architecture. He published in 1556 a work on mensuration, the *Tectonicon*. He also wrote another work on geometry which his son Thomas published in 1571 and again in 1591 under the title *A Geometricall Practise, named Pantometria*.[1] The *Stratioticos* was begun by him and completed by his son. Thomas matriculated at Cambridge as pensioner of Queens' College in 1546, taking his B. A. in 1551 and his M. A. in 1557, and attained first rank among the English mathematicians of his time. His own works related chiefly to astronomy and navigation, and, like Recorde, he was probably a believer in the Copernican theory, although he did not openly advocate the doctrine.




JOHN DEE
After a portrait from life

Of the 16th century writers on applied mathematics one of the earliest was Richard de Benese, whose *Boke of Measuring of Lande* was published between 1562 and 1575, probably the earliest book on surveying printed in England.

[1] "At the end he discourses of regular solids, and I have heard the learned Dr. John Pell say it is donne admirably well." Aubrey, *Brief Lives*, I, 233.

John Dee. Toward the close of the century there lived one of the most curious characters to be found among the scientific men of his time. This man was John Dee (1527–1608), a native of London and a student of St. John's College, Cambridge. He took his B.A. in 1546 and was selected as one of the original fellows of Trinity College. His studious nature may be inferred from his own record of his rules,—"only to sleep four houres every night; to allow to meate and drink (and some refreshing after) two hours every day; and the other eighteen hours (except the tyme of going to and being at divine service) was spent in my studies and learning." He afterwards traveled on the continent, forming friendships with Gerard Mercator, Gemma Frisius, Jean Fernel, and other scholars, taking courses for two years at Louvain, and publicly lecturing on Euclid before large audiences in Paris. He became interested in alchemy and astrology and his relations to the occult made his life a romantic one.[1] He wrote the preface to the first English translation of Euclid's *Elements*,[2]



The first Booke

Eparte vnder the line B D namely, that the whole shalbe equall to the part which is also impossible Wherefore C D is one right line which was required to be proued.

The 8. Theoreme. The 15. Proposition.

If two right lines cut the one the other: the bed angles shalbe equall the one to the other.

Demonstration

Vppose that these two right lines A B and C D, do cut the one the other in the point E Then I say, that the angle A E C, is equall to the angle D E B For forasmuch as the right line A E, standeth vpon the right line C D, making these angles C E A, and A E D the sum of the 13. proposition the angles C E A, and A E D, are equall to two right angles Againe forasmuch as the right line D E, standeth vpon the right line A B, making these angles A E D, and D E B therfore (by the same proposition) the angles A E D, and D E B, are equall to two right angles and it is proued, that the angles C E A, and A E D, are also equall to two right angles Wherfore the angles C E A, and A E D, are equall to the angles A E D, and D E B Take away the angle A E D, which is common to them both Wherefore the angle remayning C E A, is equall to the angle remayning D E B And in like sort may it be proued, that the angles C E B, and D E A, are equall the one to the other. If therefore two right lines cut the one the other, the bed angles shalbe equall the one to the other which was required to be demonstrated.

Thales Milesius the Philosopher was the first inuenter of this Proposition as witnesseth Eudemus, but yet it was first demonstrated by Euclid And in it there is no construction at all For the exposition of the thing geue, is sufficient enough for the demonstration.

Bed Angles, are appolite angles, caused of the intersection of two right lines and are so called, because the heddes of the two angles are ioyned together in one pointe.

The conuerse of this proposition after Peletarius.

If foure right lines being distant from one point, do make foure angles, of which the two opposite angles are equall the two opposite lines shalbe drawen directly and make one right line.

Suppose that there be foure right lines A B, A C, A D, and A E, drawen from the poynt A, making foure angles at the point A of which let the angle B A C be equall to the angle D A E, and the angle B A D to the angle C A E Then I say, that B E and C D are onely two right lines that is, the two right lines B A and A E are drawen directly

FROM THE FIRST ENGLISH EDITION OF EUCLID

The Billingsley or Dee translation, London, 1570

[1] Charlotte Fell Smith, *John Dee*, London, 1909.

[2] *The Elements of Geometrie of the most auncient Philosopher Evclide of Megara. Faithfully (now first) translated into the Englishe toung, by H. Billingsley, Citizen of London. . . . With a very fruitfull Praeface made by M. I. Dee*, London, 1570.

and may, indeed, have made the translation in whole or in part himself, although it is attributed to Sir Henry Billingsley, who was later (1596) Sheriff and Lord Mayor of London.[1] Dee was a man of erudition and of remarkable powers of exposition, and his influence on mathematics in his day must have been considerable.

Minor Writers. Among the books containing algebra and printed in England in the 16th century there was an arithmetic written by Thomas Masterson and published at London in three books in 1592–1595. Concerning Masterson himself nothing is known. He planned a treatise in six books but completed only three, the third consisting of that part of algebra that has to do with the powers and roots of numbers.

The century closes with the name of Thomas Blundeville, who published (1594) a work entitled *Exercises, containing sixe Treatises*,[2] one of these parts being on arithmetic and the rest being on cosmography. The arithmetic contains "a briefe description of the tables of the three speciall right lines belonging to a circle, called sines, lines tangent, and lines secant," the first fairly complete treatment of trigonometry in England.[3] He also published several other books, partly on cosmography.

5. Germany

Nature of the German Mathematics. When we compare the mathematics of Germany with that of France in the 16th century, we are struck by the same difference that existed in the art of the two countries. The mathematics of Germany was Gothic, unpolished, but virile; the mathematics of France was Renaissance, polished, but generally weak. Germany produced a notable group of arithmeticians; France produced hardly more than one. Germany produced two strong algebraists; France produced one,—a dilettante but a brilliant one. Germany made a definite advance in geometry, in the study of


[1] On this question see A. De Morgan, *British Almanac and Companion* for 1837, p. 38 of the *Companion*.

[2] The seventh edition appeared in 1636.

[3] A. De Morgan, *British Almanac and Companion* for 1837, p. 42 of the *Companion*; *Arithmetical Books*, p. 30.

higher plane curves; France was content to contemplate the past and possibly to dream of the century just ahead.

Of course the greatest influence for advance in the 16th century was printing; but there was also Erasmus (1467–1536),



FROM THE MARGARITA PHYLOSOPHICA (1503)

Showing Arithmetica between the ancient counter reckoning and the modern algorism. This is from the edition of 1504

who, although born in Rotterdam, lived in Germany, England, France, and Italy, and was the world's scholar of the first third of the 16th century; and there was also Martin Luther (1483–1546), who set Germany thinking, not always for the best. It was a century of intellectual awakening and of breaking away from traditions, and all this showed itself in the mathematical

activity of the time. As in other countries, the number of names connected with this activity now becomes so great as to allow for only a limited selection.

The Margarita Phylosophica. The first modern encyclopedia of any note, based upon the late Latin models, was the *Margarita Phylosophica*,[1] first published at Freiburg in 1503. The author was Gregorius Reisch (died 1523), who studied at Freiburg in 1487 and took his bachelor's and master's degrees there. He became a Carthusian and was made prior at Freiburg and confessor to Maximilian I. The work consists of twelve books, and includes considerable material upon arithmetic, geometry, and astronomy. Its popularity is shown by the fact that there were sixteen editions in the course of a century.

Albrecht Dürer. It is not often that the artist of today is confessedly a mathematician; the mathematician is more frequently an artist. But the 16th century has upon its roster the names of several great artists who did something in mathematics, generally in architecture or in perspective. One man, however, stands out with special prominence as a great artist and at the same time as a mathematician with distinctly new interests, and this is Albrecht Dürer.[2] His work as a painter and engraver is well known, but it was in his treatises on geometry,[3] fortification,[4] and human proportion[5] that he showed his

[1] The title of the second (1504) edition, printed at Strasburg, reads, *Aepitoma omnis phylosophiae. alias Margarita phylosophica tractans de omni genere scibili: Cum additionibus: Quę in alijs non habentur.*

[2] Born at Nürnberg, May 21, 1471; died at Nürnberg, April 6, 1528. Cantor, *Geschichte*, II, chap. 63; S. Günther, *Die geometrischen Näherungskonstruktionen Albrecht Dürers*, Ansbach, 1886; S. Günther, "Albrecht Dürer, einer der Begründer der neueren Kurvenlehre," *Bibl. Math.* (1886), p. 137; H. Staigmüller, *Dürer als Mathematiker*, Prog., Stuttgart, 1891; F. Amodeo, "Albrecht Dürer precursore di Monge," *Atti della R. Accademia delle Soc. Fis. e Mat.*, XIII (2), No. 16; W. B. Scott, *Albert Dürer, his Life and Works*, London, 1869.

[3] *Underweysung der messung mit dem zirckel und richtscheyt in Linien, ebnen, vnnd gantzen corporen*, Nürnberg, 1525, with Latin editions, Paris, 1532 and 1555, and Arnheim, 1605; *Institutiones Geometricae*, Paris, 1532.

[4] *Etliche vnderricht zu befestigung der Stett, Schloss und Flecken*, Nürnberg, 1527, 1530, and 1538, with a Latin edition, Paris, 1535.

[5] *Hierin sind begriffen vier Bücher von menschlicher Proportion*, Nürnberg, 1528, with a Latin edition, Nürnberg, 1532.

mathematical powers. The geometry was the first printed work to consider the subject of higher plane curves, and the first to discuss scientifically the question of such approximate constructions as that of the regular heptagon.

Johann Stöffler,[1] although professor of mathematics at the University of Tübingen, can be ranked as a mathematician only because of his computation of astronomical tables. He was one of the first to show how the Julian calendar could be brought into harmony with astronomical events. His calculations led him to the absurd prediction, however, that the Deluge would be repeated in the year 1524. The announcement stirred all Europe, and the number of schemes to protect the race was legion. The people of Toulouse even went so far as to build an ark. Stöffler, however, seems to have survived the storm of protest that ensued upon the failure of his prediction, for he published in the year of his death a new ephemeris, and left a commentary on the *Sphere* of Proclus, which was published posthumously in 1534.[2]

Stifel. The first German writer of the century to devote his life to mathematics and to acquire an enviable reputation in this field was Michael Stifel.[3] A lover of mathematics from his childhood; brought up in Esslingen, a veritable bulwark of the ancient faith; trained in the local Augustine convent and taking holy orders; giving promise of success in the Church; he was finally captured by the eloquence of Luther, and thought himself a reformer when he was really a fanatic. Starting for heaven with a group of peasants on the day which he had prophesied would see the blotting out of this world, he ended ignominiously behind the bars of a jail. Since Luther had launched him on a career that landed him in prison, it was proper that he should get him out, which he did. The state of

[1] Born at Justingen, Swabia, December 10, 1452; died at Blaubeuern, February 16, 1531.

[2] The curious case is discussed in Bayle's *Dictionaire*, under *Stofler*.

[3] Stiefel, Styfel, Stiffelius. Born at Esslingen, April 19, 1487 (some say 1486); died at Jena, April 19, 1567. See Th. Müller, *Der Esslinger Mathematiker Michael Stifel*, Prog., Esslingen, 1897; Cantor, *Geschichte*, II, chap. 62.

Stifel's mind may further be seen by the following line of reasoning in which he indulged:

1. The Latin for Leo Tenth is Leo Decimus.
2. This may be written Leo DeCIMVs.
3. These capitals may be arranged thus: MDCLVI.
4. We may take away M for Mystery, and add X because it is Leo X, and we then have DCLXVI.
5. But this is 666, the "Number of the Beast" in the Book of Revelations; and hence Leo X is the Beast.

Yet this is the man who, in the next few years, produced some of the most original and vigorous mathematical works to be found in the 16th century.

Stifel wrote five works on mathematics, these works treating chiefly of the mysticism of numbers,[1] arithmetic,[2] and algebra.[3] His arithmetic, which is largely on algebra, is a more scholarly work of the kind than any that had yet appeared in Germany, doing for that country what Cardan's and Tartaglia's treatises were doing for Italy.

Christoff Rudolff. Stifel's chief work on algebra was his edition (1553–1554) of a book known as the *Coss,* which appeared in 1525 and was the first algebra of any moment to be published in Germany. It was written by Christoff Rudolff,[4] concerning whose life very little is known. Rudolff published three books, the *Coss* (1525), the *Kunstliche rechnung*[5] (1526),

[1] *Ein Rechen Büchlein,* Wittenberg, 1532.

[2] *Arithmetica Integra,* Nürnberg, 1544, to which Melanchthon wrote the preface; *Deutsche Arithmetica,* Nürnberg, 1545; and *Rechenbuch von der Welschen vnd Deutschen Practick,* Nürnberg, 1546. Before the World War of 1914–1918 there was preserved in the University of Louvain a copy of the *Arithmetica Integra* with marginal notes by Gemma Frisius. Fortunately these notes were copied by H. Bosmans, S. J., and several of the most important ones were published before the destruction of the library.

[3] *Die Coss Christoffs Rūdolffs,* Königsberg, 1553–1554, 1571, and 1615. While, as stated above, this is Rudolff's work, it contains Stifel's commentary. The *Arithmetica Integra* also contains the treatment of radicals that is now considered part of algebra. The word *Coss* is from *cosa* (*causa*), which, as explained in Volume II, refers to the first power of the unknown quantity.

[4] Born at Jauer, *c.* 1500.

[5] The title of the third edition (Nürnberg, 1534) is *Kunstliche rechnung mit der zifer vnnd mit den zal pfenningē.* There were eleven editions in the 16th century.

and a collection of problems[1] (1530). Of these the *Coss* was the most important, doing for algebra in Germany what Pacioli's *Sūma* had done for the subject in Italy. All such works, however, were for the scientific élite of the universities, not for the elementary Latin schools. Of forty-six Schulordnungen of the 16th century only twenty-four gave mathematics any place, and most of these were issued only in the second half of the century.

Johann Scheubel. In marked contrast to Stifel was his contemporary, but slightly his junior, Johann Scheubel.[2] Stifel was brilliant, Scheubel was scholarly; Stifel was popular, Scheubel was heavy; Stifel was eccentric, Scheubel was balanced; and Stifel was effusive, while Scheubel was a man of dignity and poise. The University of Tübingen called Scheubel to a professorship of mathematics at about the same time that Stifel was sent to prison, and at Tübingen he wrote his arithmetics,[3] Latin works that were too heavy for commercial purposes and too light for his own students. Here, too, he wrote his algebras, one of which he published,[4] leaving the other in manuscript,[5] and here he edited the seventh, eighth, and ninth books of Euclid's *Elements* (1558). He also left in manuscript a copy of Robert of Chester's translation of al-Khowârizmî's algebra.[6] He gave the so-called Pascal Triangle a century before Pascal wrote upon it, and extracted roots as high as the 24th by a process similar to the one which employs the Binomial Theorem.

Grammateus. Of about the age of Scheubel and with similar interests, but not so learned, Heinrich Schreyber[7] made for

[1] *Exempel-Büchlin*, Augsburg, 1530, a commercial arithmetic.

[2] Scheybl, Scheubelius, Scheybel. Born at Kirchheim, Württemberg, August 18, 1494; died February 20, 1570.

[3] *De Nvmeris et Diversis Rationibvs seu reȝuiis computationum opusculum, a Ioanne Scheubelio compositum*, Leipzig, 1545; *Compendivm Arithmeticae Artis*, Basel, 1549 (this title from the 1560 edition).

[4] *Algebrae compendiosa facilisque descriptio*, Paris, 1551.

[5] This is now in the library of Columbia University, New York City. His other manuscripts were left to the University of Tübingen.

[6] This is bound with the Columbia MS. above mentioned, and the translation by Professor Karpinski was published in 1915.

[7] Born at Erfurt as early as 1496. The name appears as Schreyber in one of his works of 1523, but is given by modern writers as Schreiber.

himself a worthy place in German history. He was better known in maturity by his Latinized Greek name of Henricus Grammateus, and in his young manhood by the Latin name of Henricus Scriptor. He studied at Cracow and was later (1507)[1] enrolled in the University of Vienna, where he was afterwards an instructor. He took his bachelor's degree in 1511 and his master's degree in 1518, thereafter teaching for a time in the University and privately, and having Rudolff for one of his pupils.[2] Driven from Vienna by the plague (1521), he went to Nürnberg and Erfurt, but returned a little later (1525) and devoted himself to writing. His best-known work was an arithmetic in the German language.[3] It includes arithmetic computations with counters and figures, a little work in the theory of numbers, a chapter on bookkeeping, a few of the simplest rules of algebra, and a brief treatment of the gaging of casks. He is the first German writer to make free use of the signs + and − in the treatment of algebraic expressions, although the symbols had long been used for other purposes. He also published other works on arithmetic,[4] and wrote on the theory of proportion[5] and on mensuration.[6]

Ludolf van Ceulen. One German writer, Ludolf van Ceulen,[7] may quite as well be classified among the Dutch mathematicians, since he spent most of his life in Holland. He seems to

[1] For a record of the time reads: "Anno domini millesimo quingentesimo septimo . . . Henricus Scriptoris de Erfordia."

[2] Rudolff writes: "Ich hab von meister Heinrichen so Grammateus genennt der Coss anfengklichen bericht emphangen. Sag im darumb danck."

[3] *EYn new künstlich behend vnd gewiss Rechenbüchlin vff alle Kauffmanschafft . . . ¶Büchhalten durch das Zornal . . . ¶Visier rûten. . . . M. Henricus Grammateus*, Vienna, 1518, with later editions. This title is from the 1535 edition. C. F. Müller, *Henricus Grammateus*, pamphlet, Zwickau, 1896.

[4] *Behend unnd khunstlich Rechnung nach der Regel und welhisch practic*, Nürnberg, 1521; *Algorismus de integris Regula de tri cum exemplis*, Erfurt, 1523; *Eynn kurtz newe Rechenn unnd Visyrbuechleynn*, Erfurt, 1523, a Visierbuch being a work on the gaging of casks.

[5] *Algorithmus proportionum*, Cracow, 1514.

[6] *Libellus de compositione regularum pro vasorum mensuratione*, Vienna, 1518.

[7] Born at Hildesheim, January 18, 1540; died at Leyden, December 31, 1610. The name also appears as Ludolph van Collen, Cuelen, and Keulen. See D. Bierens de Haan in Boncompagni's *Bullettino*, XIV, 571.

have left Germany in his childhood and to have been educated under Dutch influences. He taught mathematics in Breda, Amsterdam, and Leyden, and became professor of military engineering in the University of Leyden in 1600. He is known chiefly for his value of π, at first given[1] to 20 and then to 35 decimal places.[2] He also published (1615) a work on arithmetic and geometry.

Mention should also be made of Johann Werner,[3] a priest, who was interested chiefly in astronomy but who wrote the first original work on conics to appear in the 16th century.[4]

Pitiscus. The last of the German writers on mathematics whose work falls chiefly in the 16th century and whose contributions entitle him to special mention is Bartholomäus Pitiscus.[5] He was a clergyman by profession but a mathematician by preference. His trigonometry[6] was the first satisfactory textbook published on the subject and the first book to bear this title. He also edited and perfected the table of sines of Rhæticus.[7]

The Classical Group. The name of Philip Melanchthon,[8] the friend and colleague of Luther and professor of Greek at Wittenberg, does not ordinarily suggest the science of mathematics. His wide range of human interest, however, led him not only

[1] *Van den Circkel*, Delft, 1596, with editions in 1615 and (Latin) 1619.

[2] *De arithmetische en geometrische fondamenten*, Leyden, 1615, with a Latin edition by Snell, *Fundamemta (sic) Arithmetica et Geometrica*, Leyden, the same year.

[3] Born at Nürnberg, February 14, 1468; died at Nürnberg, 1528.

[4] *Libellus super viginti duobus elementis conicis*. Nürnberg, 1522. He also wrote *De Triangulis Libri IV*; see *Abhandlungen*, XXIV.

[5] Born at Schlaun, near Grünberg, Silesia, August 24, 1561; died at Heidelberg, July 3, 1613. The name also appears as Petiscus.

[6] *Trigonometriae sive de dimensione triangulorum libri quinque*, Frankfort, 1595, as an appendix to the astronomy of Abraham Scultetus, or Abraham Schultz (1566–1625). Complete editions by Pitiscus were published at Frankfort in 1599, 1608, and 1612, and at Augsburg in 1600. An English translation appeared in London in 1630. In the edition of 1612 Pitiscus makes use of a decimal point, sin 10″ being given as 4.85 for $r = 100{,}000$.

[7] *Thesaurus mathematicus sive canon sinuum ad radium 1,0000 0000 0000 . . .* Frankfort, 1593. On Rhæticus, see page 333.

[8] Greek form for Schwartzerd, his family name. He was born at Bretten, Baden, February 16, 1497; died at Wittenberg, April 19, 1560.

into the fields of philosophy and religion, but also to the study of astronomy and mathematics. He edited (1521–1560) several works on astronomy by Aratus, Peurbach, Schoner, al-Farġânî, Sacrobosco, and Ptolemy. His activity in pure mathematics is shown in a preface to Stifel's *Arithmetica Integra* (1544) and in a work on the mathematical disciplines.[1] He is to be valued, however, for his influence rather than for his contributions in the field of the exact science.

With the name of Melanchthon is naturally connected that of his friend and associate, Joachim Camerarius,[2] a distinguished classicist and a professor at Tübingen and Leipzig. His edition of Nicomachus[3] appeared at Augsburg in 1554, and a work on computation, *De logistica*, was published there at the same time. He also wrote some astronomical verses, which were published with Melanchthon's work of 1540.

Another one of the classical group who did a certain amount of work in mathematics was Jacobus Micyllus.[4] His arithmetic[5] was written for the Latin schools, and it contains a considerable amount of ancient material. His treatment of sexagesimal fractions is unusually extensive.

A little later than Micyllus there lived the well-known scholar Michael Neander,[6] who became professor of mathematics and Greek (1551) in the University of Jena, and later (1560) professor of medicine in the same institution.[7] He wrote an

[1] *Mathematicarvm disciplinarvm, tvm etiam astrologiae encomia*, Leyden, 1540.

[2] Born at Bamberg, April 12, 1500; died at Leipzig, April 17, 1574. The family name was Liebhard, but the office of chamberlain to the Prince-Bishop of Bamberg being hereditary in this family, he took the Latin name of Camerarius (chamberlain).

[3] The title of the Daventer (1667) edition is *Explicatio Ioachimi Camerarii Papebergensis in dvos libros Nicomachi Geraseni Pythagorei Deductionis Ad Scientiam Numerorum.*

[4] Born at Strasburg, April 6, 1503; died at Heidelberg (?), January 28, 1558. The name also appears as Moltzer, Molshem, Molsehm, and Molshehm.

[5] *Arithmeticae logisticae libri duo*, Basel, 1555.

[6] Born in the Joachimsthal, April 3, 1529; died at Jena, October 23, 1581. The family name was Neumann, whence the classical form Neander.

[7] On the relation of medicine to mathematics at this time, see D. E. Smith, "Medicine and Mathematics in the Sixteenth Century," *Annals of Medical History*, New York, 1917, p. 125.

excellent work on metrology (Basel, 1555), one of the first to treat of the subject historically and in a scholarly manner. He also wrote on spherics.[1]

Still another of the classical group was Guilielmus Xylander,[2] professor of Greek at Heidelberg. He translated the first six books of Euclid's *Elements* into German (Basel, 1562), and various works from Greek into Latin, including the *Arithmetica* of Diophantus (Basel, 1575) and the work of Psellus (Basel, 1556). His *Opuscula Mathematica* appeared at Heidelberg in 1577 and contains a certain amount of work on astronomy, arithmetic, algebra,[3] and geometry.

Mathematical Astronomers. Although mathematics and astronomy were no longer synonymous terms, the 16th century produced one or two scholars whose interests in the two sciences were apparently about equal. The first of these was Petrus Apianus.[4] He wrote chiefly on astronomy, but his arithmetic[5] is interesting because it contains the first triangular arrangement of the binomial coefficients (the Pascal Triangle) to appear in print. This appeared some years before Stifel mentioned the subject. Apianus was professor of astronomy at Ingolstadt, was interested in the teaching of trigonometry, and was one of the few university professors of his time to give instruction in the German language.

The leading mathematical astronomer in the Teutonic countries in the middle of the 16th century was Georg Joachim Rhæticus.[6] He studied at Zürich and Wittenberg and was

[1] There was another Michael Neander of the same period (1525–1595), who wrote on physics, theology, and philology.

[2] Wilhelm Holzmann. Born at Augsburg, December 26, 1532; died at Heidelberg, February 10, 1576.

[3] "De svrdis, qvos vocant, nvmeris iis, qvi a qvadratis primò nascuntur, Institutio docendo explicanda."

[4] Born at Leisnig, 1495; died at Ingolstadt, April 21, 1552. Known also by his German name, Peter Bienewitz or Bennewitz.

[5] *Eyn Newe Vnnd wolgegründte vnderweysung aller Kauffmanss Rechnung*, Ingolstadt, 1527. His *Cosmographia* appeared in 1524.

[6] Georg Joachim von Lauchen, his last name being derived from his home region, the ancient Rhætia. Born at Feldkirch, in Vorarlberg, February 16, 1514; died at Kaschau, Hungary, December 4, 1576.

professor of mathematics in the latter university from 1537 to 1542. He published an arithmetic at Strasburg in 1541, but most of his work was on astronomy and trigonometry.[1] He visited Copernicus in 1539, studied with him, and did much to make his theories known.



CHRISTOPHER CLAVIUS
Engraved after a portrait from life

Clavius. Probably the man who did the most of all the German scholars of the 16th century to extend the knowledge of mathematics, although doing little to extend its boundaries, was Christopher Clavius,[2] a Jesuit, who passed the later years of his life in Rome. He was an excellent teacher of mathematics, and his textbooks were highly esteemed because of their arrangement, particularly in the Latin schools. Sixtus V testified to his standing, saying: "Had the Jesuit order produced nothing else than this Clavius, on this account alone should it be praised." His arithmetic[3] was published in Rome in 1583, was translated into Italian in 1586, and went through several editions. His

[1] *Opus Palatinum de Triangulis*, Neustadt a. Hardt, 1596; *Thesaurus Mathematicus*, Frankfort a. M., 1613, both posthumous.

[2] Christoph Klau. Born at Bamberg, 1537; died at Rome, February 6, 1612.

[3] *Epitome Arithmeticae Practicae.* Rome, 1583.

algebra[1] appeared in 1608 and was one of the best textbooks on the subject that had been written up to that time. He published an edition of Euclid in 1574.[2] While not precisely a translation, the book proved very valuable because of the great erudition shown in the extensive scholia which it contains.

Clavius was one of the mathematicians engaged in the reform of the calendar (1582) under the direction of Pope Gregory XIII. His collected works[3] contain, in addition to his arithmetic and algebra, his commentaries on Euclid, Theodosius, and Sacrobosco, his contributions to trigonometry and astronomy, and his work on the calendar.

Johann Schoner. Among the minor German writers on astronomy and mathematics in this period Johann Schoner[4] and his son Andreas Schoner (1528–1590) are perhaps the best known. Johann was for a time a preacher at Bamberg, and was later a teacher in the Gymnasium at Nürnberg. He edited a well-known medieval arithmetic[5] and wrote on geometry, astrology, and astronomy. Andreas wrote on astronomy and dialing and edited his father's works.[6]

German Arithmetic. The German arithmetics in the 16th century were very practical.[7] Commerce was active, and Frankfort was one of the great trading centers of Europe, her agents pushing out through the Hansa towns to England, France, and the northern countries, as well as through Austria and Italy to the Orient. The great house of Fugger multiplied its capital tenfold between 1511 and 1527, possessing five times the wealth which made the Medici so powerful in the preceding

[1] *Algebra Christophori Clavii Bambergensis*, Rome, 1608.

[2] *Euclidis Elementorum Libri XV*, as the title appears in the Frankfort edition of 1654. The earlier editions were 1574, 1589, 1591, 1603, 1607, 1612, which show the popularity of the work.

[3] *Opera Mathematica*, 5 vols., Mainz, 1611 and 1612.

[4] Born at Karlstadt, near Würzburg, January 16, 1477; died at Nürnberg, January 16, 1547. The name also appears as Johannes Schonerus or Schöner.

[5] *Algorithmvs demonstratvs*, Nürnberg, 1534.

[6] *Opera Mathematica*, Nürnberg, 1561.

[7] Hugo Grosse, *Historische Rechenbücher des 16. und 17. Jahrhunderts*, Leipzig, 1901.

century. But whereas the Medici had been the patrons of art and of letters, the Fuggers were little beyond accumulators of wealth. As a result of this national spirit, the German mathematics of this period was more largely commercial than it might have been under more favorable circumstances.

While the classical group was carrying mathematics to the intellectual aristocracy a group containing a few skillful writers was carrying it to the intellectual democracy. One of them, Johann Böschensteyn,[1] like Stifel a native of Esslingen, taught both Luther and Melanchthon,—a sufficient honor for any man. He was a professor of Hebrew at Ingolstadt, Heidelberg, Nürnberg, and Antwerp, but his mathematical contribution was of a humble nature, consisting merely of an arithmetic of a commercial kind, printed at Augsburg in 1514.

Contemporary with Böschensteyn, and indeed but two years his senior, was one whose influence on the people was far greater,—Jakob Köbel,[2] a native of Heidelberg. He and Copernicus were fellow students at Cracow, and each had varied lines of interest. Köbel began as a teacher of arithmetic, a printer, a woodcarver, a poet, and a student of law, and ended as a petty officeholder; while Copernicus began as a priest, physician, and astronomer, and ended as a giant among the thinkers of the world. Köbel wrote three arithmetics, although the editions varied so much as to give an impression of a larger number. These arithmetics were the *Rechenbiechlin*[3] (1514), *Mit der Kryden*[4] (1520), and the *Vysierbuch*[5] (s.a. but 1515). Of these the first was the most important, passing through no less than twenty-two editions in the 16th century. It is purely commercial but shows more vigor and originality than any arithmetic that had yet appeared in Germany. Among other

[1] Among the variants of the name are Beschenstein, Boeschenstain, Bossenstein, Boechsenstein, Buchsenstein, Poschenstein, and Besentinus. Born at Esslingen, Swabia, 1472; died 1540.

[2] Kobel, Kobelius, Kobelinus. Born at Heidelberg, 1470; died at Oppenheim, January 31, 1533.

[3] As spelled on the title-page; *Rechenbüchlein* as spelled in the colophon.

[4] *Mit der Krydē od' Schreibfedern durch die zeiferzal zu rechē Ein neüw Rechēpüchlein*, Oppenheim, 1520.

[5] On gaging. In the 1537 edition, *Eyn new Visir Büchlin*.

features are the crude illustrations, although similar ones had already been used by Widman as early as 1489.

Adam Riese. The greatest of all the Rechenmeisters of this century, however, was Adam Riese.[1] He was the most influential of the German writers in the movement to replace the old computation by means of counters ("auff der Linien") by the more modern written computation ("auff Federn"). He wrote four arithmetics,[2] all of a commercial nature and the second ranking as one of the most popular schoolbooks of the century. It was to Germany what Borghi's book was to Italy, Recorde's to England, and Gemma's to the Latin schools of the Continent. So famous



ADAM RIESE

Germany's best-known Rechenmeister. From the title page of his *Rechenung nach der lenge/auff den Linihen vnd Feder*, Leipzig, 1550

[1] Born possibly at Staffelstein, near Bamberg, *c.* 1489; died at Annaberg, March 30, 1559. The name also appears as Ryse, Ris, and Reis. See B. Berlet, *Adam Riese, sein Leben, sein Rechenbücher und seine Art zu rechnen*, new ed., Leipzig, 1892.

[2] *Rechnung auff der linihen*, printed in 1518 and again in 1525 and 1527. Title from the 1525 edition.

Rechnung auff der Lynihen vn Federn in zal/mass/vnd gewicht auff allerley handierung gemacht, Erfurt, 1522, with at least thirty-seven editions before 1600.

Ein Gerechent Büchlein, Leipzig, 1536.

Rechenung nach der lenge/auff den Linihen vnd Feder, Leipzig, 1550.

was the author that the phrase "nach Adam Riese" is still used in Germany to signify arithmetical accuracy or skill.

It was due in no small degree to the encouragement given by Luther that books like Riese's met with their great success. At the very time that the great Rechenmeister was preparing the second edition of his most successful work[1] Luther was laying down his famous doctrine that all children should study mathematics,[2] a thing unheard of before.

The only other Rechenmeister of the century to deserve special mention is Simon Jacob,[3] who wrote two commercial arithmetics. Bürgi mentions Jacob's treatment of series, and apparently the former's table of antilogarithms, the *Progress Tabulen*,[4] was suggested by the nature of exponents as laid down in these and similar books of the 16th century.

6. The Netherlands

Geographical Limits. When we speak of the Low Countries, the Netherlands, it should be borne in mind that boundaries and governments were constantly shifting at this time, and that when Charles V (died 1558) inherited this territory from his grandmother, Mary of Burgundy, it included seventeen provinces, each with its own government. When Philip II (died 1598) inherited it from Charles, the new ruler was accepted with a pronounced expression of discontent, and finally (1568) the provinces rebelled under William of Orange. The period was therefore one of uncertain geographical limits, and in speaking of the Netherlands it must be understood that we speak of territory of which a part was under Spanish rule during the 16th century. In general, therefore, we shall include for our present purposes what is now Holland and Belgium.

[1] That is, in 1524, the edition appearing in 1525.

[2] This is in his *Schrift an die Ratsherren aller Städte Deutschlands*, wherein he says: "Wenn ich Kinder hätte, und vermöchts, sie müssten mir nicht allein die Sprachen und Historien hören, sondern auch singen und die Musika mit der ganzen Mathematika lernen."

[3] Born at Coburg; died at Frankfort a. M., June 24, 1564. His arithmetics appeared at Frankfort a. M. in 1557 and 1565.

[4] Discussed in Volume II, Chapter VI.

General Conditions. The general conditions in the country, aside from those imposed by dynastic ambitions, were favorable to commercial and intellectual advance. The Netherlands had shown considerable commercial activity in the 15th century. The economic decline in the latter part of that period, owing to British rivalry, had been somewhat overcome by their success in navigation and in the fishing industry.[1] Their arithmetics reflect this commercial activity,[2] showing an extensive trade with Nürnberg, Frankfort, Augsburg, Danzig, and other German towns; with Cracow, Venice, and Lyons; and with Spain.

In matters of scholarship the work of that great humanist of the North, Erasmus of Rotterdam (1467–1536), had told in Holland as it had told in every other intellectual center of Europe. Leyden was becoming one of the great forces of the world. The Netherlands were ready to do their part.

Mathematics in the Netherlands. The first of the writers of consequence in this territory in the 16th century was Joachim Fortius Ringelbergius,[3] a student at Louvain and a teacher of philosophy and mathematics in various places in France and Germany. He wrote on astronomy, optics, and arithmetic, and his collected works,[4] encyclopedic in character, were published at Leyden in 1531. The book was filled with the usual erudition of the time but contained nothing that advanced the bounds of science.

Late in the 16th century Adriaen van Roomen,[5] a student of both medicine and mathematics in Italy and Germany, became

[1] A. W. Ward, in the *Cambridge Mod. Hist.*, I, chap. xiii; W. Cunningham, *ibid.*, chap. xv.

[2] For a list of these books, not complete, see D. Bierens de Haan, "Bibliographie néerlandaise historico-scientifique," in Boncompagni's *Bullettino*, XIV, 519; XV, 355; hereafter referred to as Bierens de Haan, *Bibliog.* See also the *Rara Arithmetica* for a list of books on arithmetic. There is also a list of Dutch arithmetics in the preface to the 1690 edition of Coutereel's *Cyffer-Boeck.*

[3] Born at Antwerp, *c.* 1490; died *c.* 1536. In the vernacular the name appears as Joachim Sterck van Ringelbergh.

[4] *Ioachimi Fortii Ringelbergij Andouerpiani opera*, Leyden, 1531.

[5] Born at Louvain, September 29, 1561; died at Mainz, May 4, 1615. The Latin form Adrianus Romanus and the French form Adrien Romain are often

professor of these two sciences in Louvain. He then became professor of mathematics at Würzburg, and finally was appointed royal mathematician (astrologer) in Poland. While at Louvain he published the first part of a general work on mathematics,[1] and in this he gave the value of π to seventeen decimal places, an unusual achievement at the time. His other works include one on the treatment of the circle by Archimedes (1597) and one on spherical triangles (1609).

There lived in Holland in the middle of the century an engineer who also had the name of Adriaen. His father's name was Anthonis, and so he was called Adriaen Anthoniszoon. From some connection with Metz,[2] he was known as Metius, a name also used by his sons. He suggested $\frac{3}{1}\frac{5}{1}\frac{5}{3}$ as a convenient value of π, probably oblivious of the fact that it had already been given (1573) by a minor writer in Germany,[3] and certainly ignorant of its use in China several centuries earlier. This Adriaen had a son, also named Adriaen, who was called, after the custom of the time, Adriaen Adriaenszoon, but was also called by his father's geographical name of Metius. This younger Adriaen Metius[4] studied both law and medicine and became (1598) professor of mathematics and medicine in the University of Franeker. Although he wrote on mathematics,[5] his chief contributions were to astronomy. He published his father's value of π, and hence it is commonly attributed to the son, although not due in the final analysis to either.

used. The first name is also spelled Adriaan. See H. Bosmans, *Annales* of the *Société Scientifique*, XXVIII and XXIX (Brussels), and *Bibl. Math.*, V (3), 342.

[1] *Ideae mathematicae pars prima, seu Methodus polygonorum*, Antwerp, 1593.

[2] He himself was probably born at Alkmaar, *c.* 1543. He died at Alkmaar, November 20, 1620. This reason, commonly given for the name Metius, is doubtful. The family name was Van Schelvan (haycock; Latin, *meta*).

[3] Valentinus Otto, or Valentin Otho, also called Parthenopolitanus, a native of Magdeburg. See *Bibl. Math.*, XIII (3), 264.

[4] Born at Alkmaar, December 9, 1571; died at Franeker, September 6 (16, or 18), 1635. The Latin form, Adrianus, is also used.

[5] *Praxis Nova Geometrica per usum circini*, Franeker, 1623; *Arithmetica et Geometrica nova*, Leyden, 1625; *Maet-Constigh Liniael*, Franeker, 1626, describing a kind of slide rule; *Doctrinae Sphericae Libri V*, Franeker, 1591; and other works.

A more humble writer but a more progressive teacher appeared early in the century in the person of Giel Vander Hoecke. Although of no marked scholarship, his arithmetic[1] is worthy of study because of its early use of the plus and minus signs as symbols of operation. Widman had already used them as signs of excess and deficiency, and Grammateus had used them with their modern significance, but they now appear for the first time in the Low Countries.



GEMMA FRISIUS

From a contemporary engraving

Gemma Frisius. The most influential of the various Dutch mathematicians of this century was Gemma Regnier.[2] Having been born in Friesland (Frisia), he was called the Frisian, and was known as Gemma Frisius. He was thirty-two years old when his arithmetic[3] was published, and so favorably did this work strike the popular taste, combining as it did the commercial with the theoretical, that it went through at least fifty-nine editions in the 16th century, besides several thereafter. He also wrote on geography and astronomy, suggesting the present method of obtaining longitude by means of the

[1] *Een sonderlinghe boeck in dye edel conste Arithmetica*, Antwerp, 1514; 2d ed., 1537.

[2] Born at Dockum, East Friesland, December 8, 1508; died at Louvain, May 25, 1555. The name is variously spelled, as Rainer, Renier, Reinerus.

[3] *Arithmeticae Practicae Methodvs Facilis*, Antwerp, 1540. From certain internal evidence (1575 ed., fol. B, 5, *v.*) it is probable that the book was written *c.* 1536.

difference in time, and taking one of the first steps toward the modern methods of triangulation.[1] He became professor of medicine at Louvain in 1541, and his son, Cornelius Gemma Frisius (1535–1577), carried on his work, becoming professor of medicine and astronomy in the same university. Cornelius edited one of his father's works and wrote on astronomy and medicine.

While Gemma Frisius wrote for the Latin schools, a man of the people was needed to write for the common schools, and this man was found in the person of Valentin Menher, a native of Kempten. He wrote in the French language three or four arithmetics[2] that occupied the same position in the Netherlands that Borghi's did in Italy and Recorde's in England. His arithmetic of 1573 includes a certain amount of work in geometry and trigonometry.

Among the Belgians who brought a high degree of scholarship into their work in the editing of the early classics of mathematics the one who stood highest at the beginning of the 16th century was Jodocus Clichtoveus, a native of Nieuport, in Flanders. He spent most of his time in France, assisted in editing Boethius on arithmetic, and wrote a *Praxis Numerandi* which was merely an edition of the algorism of Sacrobosco. He died at Chartres in 1543.

The last of the Dutch writers of the century was Jacob Van der Schuere of Meenen (*c.* 1550–1620), a teacher in a French school in Haarlem. His arithmetic,[3] commercial in character, was one of the first of a long series of popular textbooks of this type that appeared in Holland from 1600 to about 1750.

Stevin. The most influential of all the mathematicians produced by the Low Countries in the 16th century was Simon Stevin.[4] In his younger days he was connected with the government service in Bruges. He traveled in Prussia, Poland,

[1] *Libellus de locorum describendorum ratione*, Antwerp, 1533.

[2] See *Rara Arithmetica*, pp. 250, 346.

[3] *Arithmetica, Oft Reken-const, . . . Door Iacqves Van Der Schvere van Meenen, Nu ter tijdt Francoysche School-meester tot Haerlem*, Haarlem, 1600.

[4] The name also appears in such forms as Stevinus, Steven, Stephan, Stevens. Born at Bruges, *c.* 1548; died at Leyden or The Hague, *c.* 1620.

and Norway and later became a quartermaster general in the Dutch army and director of certain of the public works.[1] His most influential but not his most popular work was an arithmetic,[2] first published in Flemish at Leyden in 1585, and republished the same year in a French translation. The importance of this work lies in the fact that it was the first one to set forth definitely the theory of decimal fractions, a subject that had been slowly developing for a century. Stevin also made the first translation into a modern language of the work of Diophantus, apparently from the Latin text of Xylander.[3]

What made Stevin best known among his contemporaries, however, was his contribution to the science of statics and hydrostatics,[4] a subject naturally occupying much attention in a country like Holland.

7. Spain

Spanish Writers. Spain furnished several native mathematicians of considerable merit in the 16th century.[5] The intellectual atmosphere was not favorable to the development of mathematics, however, and many Spanish scholars settled in France and Italy or at least published their works abroad. It is no reflection upon the honesty of purpose of the Church to say

[1] F. V. Goethals, *Notice historique sur la vie de S. Stevin*, Brussels, 1842; M. Steichen, *Mémoire sur la Vie et les travaux de Simon Stevin*, Brussels, 1841.

[2] The title of the 1585 French edition, published at Leyden, is *L'Arithmetique de Simon Stevin de Brvges: Contenant les computations des nombres Arithmetiques ou vulgaires: Aussi l'Algebre, auec les equations de cinc quantitez. Ensemble les quatre premiers liures d'Algebre de Diophante d'Alexandrie, maintenant premierement traduicts en François.*

[3] Xylander's edition, Basel, 1575; Stevin's translation, Leyden, 1585.

[4] *De Beghinselen der Weeghconst*, Leyden, 1586; *De Beghinselen des Waterwichts*, Leyden, 1586; *Weeghdaet*, Leyden, 1586. See also his *Wisconstighe Gedachtenissen Inhoudende 't ghene daer hem in gheoeffent heeft . . .*, Leyden, 1605–1608, with a Latin edition by W. Snell (Leyden, 1608) and a French edition by A. Girard (Leyden, 1634). He also wrote *Problematum Geometricorum Libri V* (Antwerp, 1583) and other works.

[5] J. Rey Pastor, *Los matemáticos españoles del siglo XVI*, Oviedo, 1913; F. Picatoste y Rodriguez, *Apuntes para una Biblioteca Cientifica Española del Siglo XVI*, Madrid, 1891 (hereafter referred to as Picatoste, *Apuntes*); Acisclo Fernández Vallin, *Cultura Científica de España en el Siglo XVI*, Madrid, 1893. The first and second of these works are particularly valuable. See also G. Loria, "Le Matematiche in Ispagna," *Scientia*, XXV (May, June, 1919).

that the religious fervor of Spain from the 15th to the 18th century turned the thoughts of the intellectual class from mathematics, although such work as was done was due to the clergy. This was especially the case after the compact made at Bologna in 1530 between Charles V and Clement VII.[1] To this influence there should be added that which came from the expulsion of the Jews, a race which had done so much in the Middle Ages to foster the science of mathematics, at least with respect to astrology and the theory of the calendar.

Ciruelo. The earliest Spanish mathematician of the century was Pedro Sánchez Ciruelo,[2] who was professor of theology and philosophy at Alcalá, and was later canon of the cathedral of Salamanca. He published an arithmetic[3] at Paris in 1495, and a general work on mathematics[4] in 1516. He also edited the theoretical work on arithmetic of Bradwardine in 1495 and the *Sphaera* of Sacrobosco in 1498, and in general was a learned exponent of the old school of mathematicians that was then in favor in Paris.

Ortega. The second of the early Spanish writers was Juan de Ortega, a Dominican from Aragon, concerning whose career we know little except that he was living in 1512 and in 1567. He wrote an arithmetic[5] which was published in 1512, both in Barcelona and in Lyons, being the first book on commercial computation known to have been printed in France. It was a popular work, being reprinted in Rome, Messina, Seville, Paris, and Granada. It is purely commercial and includes the usual treatment of computation and the common applications of the time.

[1] R. C. Jebb, in the *Cambridge Mod. Hist.*, Vol. I, chap. xvi.

[2] Born at Daroca, Aragon, *c.* 1470; died 1560.

[3] The 1505 edition has the title *Tractatus Arithmetice Pratice qui dicitur Algorismus.*

[4] *Cursus quattuor mathematicarvm artiū liberaliū*, Paris, 1516. First Spanish edition, Alcalá, 1516, but wanting the geometry, at least in the copy examined by the author.

[5] The title of the Rome edition of 1515 is *Svma de Arithmetica: Geometria Pratica vtilissima.* The geometry consists simply of a little mensuration. In the privilege of Leo X (1515), in this edition, the author is addressed as "DIlecto filio Iohāni de Ortega Hispano Clerico Palētino."

Joannes Martinus Blasius,[1] a Spanish astrologer and arithmetician, published in Paris in 1513 a work on computation. It was popular enough to warrant four editions. The author was one of the earliest writers whose works appeared in print with the spelling *substractio* for "subtraction," a custom followed quite generally by the Dutch and English arithmeticians for several generations. He showed a good knowledge of the classical writers in the domain of mathematics.

Another Spanish scholar who found the scientific work more stimulating in Paris than in his native country was Gaspar Lax.[2] He took a course in theology at Saragossa, taught in the University of Paris, and finally returned to Saragossa as a teacher. His principal work was a prolix treatment of theoretical arithmetic[3] based on Boethius. He also wrote on the Greek and medieval theory of proportion.[4]

One of the most noteworthy treatises on mathematics produced in Spain in the 16th century is the work[5] of Juan Perez de Moya (1562). This writer was born in San Stefano, in the Sierra Morena, studied at Alcalá and Salamanca, and became a canon at Granada. His *Arithmetica* includes calculation, applied arithmetic, algebra, and practical geometry, and contains a considerable amount of interesting historical material.

The last of the Spanish writers of any note, in the 16th century, Jerónimo Muñoz,[6] received his bachelor's degree at Valencia in 1537. He traveled in Italy, taught at Ancona, returned to Spain, and taught for ten years in the University of Valencia. He wrote on arithmetic (1566) and Euclid, but his chief work was on astrology.

[1] This is the name as it appears in the 1513 edition of his arithmetic. In the 1519 edition the name appears as "Ioannes Martinus, Scilicevs." The usual Spanish form is Juan Martínez Silíceo. The original name was Juan Martínez Guijeno, the word *guijeno* meaning *silex*, whence Sileceus, Sileceo, or Siliceo. See V. Reyes y Prósper, "Juan Martinez Silíceo," *Revista* of the *Soc. matem. española*, I, 153.

[2] Born at Sariñena, *c.* 1487; died at Saragossa, 1560.

[3] *Arithmetica speculatiua magistri Gasparis Lax Aragonensis de sarinyena duodecim libris demonstrata*, Paris, 1515.

[4] *Proportiones magistri Gasparis lax*, Paris, 1515.

[5] *Arithmetica practica, y specvlatiua del Bachiller Iuan Perez de Moya*, Salamanca, 1562.

[6] Hieronymus Munyos.

Loss of the Jews. As for the Jews who had once added to the brilliancy of Spain, hardly one of their descendants remained in that country at the opening of the 16th century. The edict of banishment of 1492 had driven out hundreds of thousands of this race, some to slavery, some to death at the hands of pirates, some to the plague-stricken towns of Italy, and some to starvation. Persecution after persecution had accomplished the purpose of those in power, but the result had sapped the strength of Spain, and some of the best thought that would have made for the advance of mathematics was turned to the solving of the problem of self-preservation.[1]

8. Other European Countries

Poland. In the 16th century Poland was one of the most progressive countries of Europe in the field of arithmetic, producing several works by native writers and reprinting a number by foreign scholars.[2] The first of her own arithmetics was the *Algoritmus* of Tomas Klos, which appeared at Cracow in 1538.[3] Later in the century (1561) Benedictus Herbestus (1531–1593), a Jesuit priest, published an arithmetic in Latin.[4] One in the Polish language, chiefly commercial in nature, was written by Girjka Gorla z Gorlssteyna and published at Czerny in 1577. It was not through works like these, however, that Poland contributed to human knowledge, but through those of her greatest astronomer, or perhaps we should say the world's greatest astronomer.

Copernicus. Not all of those who have aided in the progress of mathematics have been primarily mathematicians. As we

[1] For a description of one phase of this movement see J. A. Symonds, *Renaissance in Italy: The Age of Despots*, p. 399 (New York, 1883). On the unpublished material relating to the Jewish contributions to mathematics and astronomy, beginning in the 11th century and closing in the 16th century, see B. Cohen, "Ueber unveröffentlichte Schriften jüdischer Astronomen des Mittelalters," *Jahrbuch der Jüdisch-Literarischen Gesellschaft*, XII (1918), 1.

[2] *Rara Arithmetica*, pp. 32, 97, 123, 190, 260, 303, and 353.

[3] A reprint, edited by M. A. Baraniecki, was published at Cracow in 1889.

[4] *Arithmetica Linearis, eiq3 adiuncta Figvrata, cum quibusdam ex compvto necessarijs*, Cracow, 1561. This title is from the edition of 1577.

have already seen, the science of astronomy has always contributed to her sister science, not merely in those centuries in which the name "mathematician" meant an astronomer or an astrologer, but in more recent times, when mathematics and astronomy each outgrew the possibility of mastery by the disciples of the other. Among those whose interest was primarily in astronomy but who stimulated the mathematician to seek for new applications of his science none stands higher than Nicholas Copernicus.[1] He was educated at the University of Cracow (1491–1495), spent some time in the study of law, medicine, and astronomy in the universities of Padua and Bologna, and went to Rome for the purpose of continuing his work in astronomy under the patronage of the pope, Alexander VI. He returned to Poland in 1505, took holy orders, and obtained a canonry at Frauenburg. By 1530 he had completed his theory of the universe, the most significant step ever taken in the science of astronomy, but it was not until 1543 that he published his doctrines.[2] Gutenberg made the free spread of



COPERNICUS

From an early engraving

[1] Born at Thorn, on the Vistula, February 19, 1473; died at Frauenburg, May 24, 1543. He was named after his father, Niklas Koppernigk (died 1483), a native of Cracow. The English form of the name, Copernicus, is so familiar that it is used throughout this work, although the spelling Coppernicus is nearer the original. On his life consult L. Prowe, *Nicolaus Coppernicus*, 3 vols., Berlin, 1883–1884. The literature relating to him is extensive.

[2] *De revolutionibus orbium coelestium*, Nürnberg, 1543.

thought possible, but Copernicus gave the thought; Columbus opened a new world, but Copernicus opened millions of worlds.

The work of Copernicus necessitated the improvement of trigonometry, and for this reason he wrote a treatise on the subject,[1] his single contribution to the literature of pure mathematics.

A generation after Copernicus a Danish mathematician, Thomas Fincke (1561–1656), whose name also appears as Finck, Fink, and Finchius, published a work called *Geometria Rotundi* (Basel, 1583), in which he made a number of contributions to trigonometry.

Switzerland. The best known of the Swiss mathematicians of this century was Henricus Loritus Glareanus,[2] whose last name probably comes from the name of his native canton of Glarus. He was a professor in Basel (1515–1521), at the Collège de France (1521–1524), and later in both Freiburg and Basel. He wrote on arithmetic, metrology, and music.

The only other Swiss writer of the period who need be mentioned is Cunradus Dasypodius.[3] He was professor of mathematics at Strasburg and canon of St. Thomas's Church in that city. He had in mind the editing of all the Greek mathematical works, and made a beginning in that direction. His edition of Euclid's *Elements* appeared at Strasburg in 1564, and he wrote a mathematical dictionary.[4]

Portugal. Only a single Portuguese mathematician acquired any considerable reputation in the 16th century. This man was Pedro Nunes (better known by his Latin name of Nonius but

[1] *De lateribus et angulis triangulorum libellus*, Wittenberg, 1542.

[2] Heinrich Loriti Glarean. Born at Mollis, in Glarus, June, 1488; died at Freiburg, Breisgau, May 28, 1563. H. Schreiber, *Heinrich Loriti Glareanus*, Freiburg, 1837. The date of his death may have been March 27, 1563.

[3] Name as given in his *Lexicon*. This is the Latin-Greek form for his family name of Rauchfuss (rough-foot), Greek δασυπόδειος (of a hare), a δασύπους (rough foot) being a hare. Born at Frauenfeld, *c.* 1530; died at Strasburg, April 26, 1600.

[4] Λεξικον *seu Dictionarium Mathematicum, M. Cunrado Dasypodio*, Strasburg, 1573. He also wrote *Institutionum Mathematicarum voluminis primi Erotemata Logisticae Geometriae Spherae Geographiae*, 2 vols., Strasburg, 1593, 1596; *Volvmen primum: mathematicum disciplinarum principia*, 2 vols., Strasburg, 1567, 1570; *Brevis Doctrina de Cometis* (on astrology), Strasburg, 1578.

often called by the Spanish name of Núñez,[1] a scholar of Jewish origin. He studied at the University of Lisbon and later (1530) became professor of moral philosophy in that institution. He was (1544–1562) also professor of mathematics in the University of Coimbra and held the posts of cosmographer to the king, Don João III, and tutor to the royal princes.[2] His only mathematical work was devoted to algebra, arithmetic, and geometry,[3] but he is best known for his works on navigation and astronomy,[4] and for the instrument for the reading of small angles, often called the "nonius," which was the forerunner of the vernier that is seen on transits and calipers.[5] He left several manuscripts on geometry and navigation.

In the early part of the 16th century the interest in the great voyages of the Portuguese led to the publication of various treatises on the sphere and the use of the astrolabe for nautical purposes. Among the earliest of these works was an anonymous one[6] on the astrolabe and the quadrant, the instruments then chiefly used in navigation.

[1] Born at Alcacer do Sal, 1502; died at Coimbra, August 11, 1578. The year of his birth is often given as 1492 and that of his death as 1577. Nunes himself records, however, ". . . sit anno Domini 1502 quo ego natus . . .," and a 16th century MS. note in a book in the National Library at Lisbon reads: "Natus est hic Doctor año Dñi 1502. Obiit vèro tertio idus Augusti año Dñi 1578." Nothing more authoritative is known. See Guimarães, *Les Math. Portug.*, p. 16, with authorities given. The name also appears as Pedro Núñez Salaciense, and in French as Pierre Nugne. See also Bensaude, *Astron. Portug.*, p. 59; Picatoste, *Apuntes*, p. 218.

[2] His dedicatory epistle of 1564 speaks of himself as "Cosmographo Mayor del Rey de Portugal, y Cathedratico Jubilado en la Cathedra de Mathematicas en la Vniversidad de Coymbra."

[3] *Livro de algebra em arithmetica y geometria*, written *c.* 1532, published at Antwerp in 1564, and reprinted there in 1567 with variations in the spelling of words in the title. The above title is substantially as in Guimarães, *Les Math. Portug.*, pp. 21, 105, 396; but see H. Bosmans, "Sur le 'Libro de algebra' de Pedro Nuñez," *Bibl. Math.*, VIII (3), 154; Cantor, *Geschichte*, II, chap. 59; Picatoste, *Apuntes*, p. 221.

[4] *Tratado da Esphera com a Theorica do Sol e da Lua*, Lisbon, 1537; *Tratado sobre certas duvidas da Navegação*, an appendix to the preceding work.

[5] This is described in his *De Crepusculis Liber unus*, Lisbon, 1542.

[6] *Regimento do estrolabio & do quadrante*, published *c.* 1509. This was reproduced in facsimile in Volume I of J. Bensaude, *Histoire de la Science Nautique Portugaise*, Munich, 1914. See this publication for other works of similar nature in that period.

Jewish Writers in the East. With the expulsion of the Greek Christians at the fall of Constantinople (1453) there returned to the city many Jews who had been subject to persecution under their régime. This movement continued as occasion offered, and notably when the Jews were driven from Spain in 1492 and from Portugal in 1496. The sultan Mohammed II made Moses Kapsali chief rabbi over all the Turkish Jews, and on the latter's death he was succeeded in office by Elia Misrachi.[1] This learned rabbi wrote an arithmetic[2] based to a considerable degree on a work by Rabbi ben Ezra. Knowing both Greek and Arabic, he also drew from each of these sources, particularly from the latter. He was interested in making arithmetic practical and in having the processes thoroughly understood.[3] He seems to have been the first Hebrew writer to treat of finding the sum of the cubes of the first n natural numbers. He also wrote commentaries on the works of Euclid and Ptolemy.

Russia. The only mathematical works known to have been written in Russia in the 16th century were a geometry and an arithmetic, both of which date from about 1587 to 1594.[4] Translations of monographs on western mathematics also appeared in this century, but they only show the low state of science at that time in Russia. This is seen, for example, in the fact that the *Origines* of Isidorus of Seville (570–636) was thought worthy of translation.

9. The Orient

Close of the Dark Ages. We have seen that the East had its dark periods in history just as the West had them, and with each the darkness was greatest just before the dawn. In Asia the gloom was particularly oppressive in the 16th century.

[1] Born probably at Constantinople, *c.* 1455; died at Constantinople, 1526.

[2] *Sefer ha-Mispar (Book of Numbers)*.

[3] G. Wertheim, *Die Arithmetik des Elia Misrachi*, p. 6. Frankfort, 1893.

[4] N. M. Karamzin, *History of the Russian Empire* (in Russian), X, 259, 436 (Petrograd, 1824); Lavrovsky, *Ancient Russian Schools* (in Russian), p. 180 (Kharkov, 1854); Russian *Encyclopedia* (in Russian).

India was intellectually dead. China was just becoming aware of the extent of Western learning, and seemed discouraged in the effort to advance independently. Japan was not yet awake. For these reasons the 16th century was a dark one for Oriental mathematics.

Islam. Of the heirs to the glory of the scholars of Islam in the field of mathematics the name of only one of any note is found in the records of the century. Behâ Eddîn,[1] as he is generally called, was probably a Persian. He wrote on a variety of subjects and among his works was an elementary textbook on arithmetic[2] and the first part of an exhaustive treatise on the subject.[3]

A single other writer may properly be mentioned with Behâ Eddîn, namely, Mohammed ibn Ma'rûf ibn Aḥmed, Taqî eddîn (1525/1526–1585), who seems to have lived in Constantinople. He wrote on algebra, arithmetic, and astronomy.

India. After the death of Bhāskara (*c.* 1175) there was no great interest shown in the advance of mathematics in India except so far as it related to astronomy. From the 16th century only two names of any note have come down to us. Suryadaśa, who flourished *c.* 1535,[4] and Ganeśa,[5] who lived about the same time, were both commentators on the works of Bhāskara. Suryadaśa refers to Śrīdhara's method of finding the area of a cyclic quadrilateral, and Ganeśa quotes one of Śrīdhara's rules for the area of a segment of a circle.

China. China at this time was experiencing a kind of calm before that influx of European mathematics which was heralded by the great Jesuit leader, Matteo Ricci. K'u Ying-hsiang (*c.* 1550), governor of Yünnan, wrote on algebra and geometry; T'ang Shun-ki (1507–1560) wrote on the mensuration of the

[1] Behâ ed-dîn al-'Âmilî, Mohammed ibn Ḥosein. Born probably at Amul (Amol), near the Caspian Sea, 1547; died at Ispahan, 1622. Nazâm ed-dîn Aḥmed, in his biography, says that he was born at Baalbek. See A. Maare, *Beha Eddin*, 2d ed., Rome, 1864.

[2] The *Kholâsat al-Ḥisâb* (*Essence of Arithmetic*). The work has no particular merit. There was an Arabic-Persian edition, Calcutta, 1812; an Arabic-German edition, Berlin, 1843; and a French translation by Marre, Rome, 1864.

[3] The *Bâhr al-Ḥisâb* (*Ocean of Arithmetic*), left unfinished.

[4] *Bibl. Math.*, XIII (3), 205.

[5] *Ibid.*, 205. Also written Ganeça.

circle; Hsin Yun-lu (*c.* 1590) contributed to the subject of the calendar, more in quantity than in quality; and Ch'êng Tai-wei (1593) wrote a work on arithmetic[1] in which we find the earliest description of the *suan-pan* computation. These writers lacked the genius of some of their immediate predecessors; and they contributed nothing that was commensurate in importance with the productions of the century following.

Japan. The 16th century did not see the awakening of the intellectual Japan as it saw the awakening of Europe. The East was at this time about a century behind the West in this great world movement. Perhaps it is more nearly accurate to say that the 16th century in Japan corresponds more closely to the 13th century in the West; it was a century of preparation.

Probably the chief cause which contributed to this preparation in the field of mathematics was the journey to China made by one Mōri Kambei Shigeyoshi, a scholar in the service of two of the powerful lords of Japan. The story goes that the great hero Toyotomi Hideyoshi, better known as Taikō, having subdued all of the country, decided that he would make his court a great intellectual center. In pursuance of this purpose he sent Mōri to China to acquire and bring back that mathematical knowledge which was so lacking in Japan. Mōri, being a man of humble birth, was not well received in China, and for this reason Taikō made him Lord of Dewa,[2] hoping thus to give him high standing among Chinese scholars. Owing to political and military difficulties, chiefly Taikō's invasion of Korea (1592), Mōri's mission was not successful, but he brought back with him a considerable amount of material and is said by some to have made the Chinese abacus[3] known in Japan. His last years were spent in Kyōto in teaching the use of this instrument.

Although this is the story as often told, there is a question as to whether Mōri really visited China or went only to Korea.

[1] *Suan-fa Tong-tsung* (*A systematized treatise on arithmetic*). His name also appears as Tch'eng Ta-wei. [2] *Dewa no Kami.*

[3] The *suan-pan*. It developed later into the Japanese *soroban*. Both of these instruments are discussed in Volume II.

It seems certain, however, that he knew something of Chinese mathematics, that he was an expert with the abacus, that he advertised himself as "the leading instructor in division in the world," and that he was a very successful teacher. Among his pupils were three men,[1] known to their contemporaries as "The Three Arithmeticians," who will be mentioned later. That Mōri was the first who took the abacus from China to Japan is very doubtful, but he seems to have made it popular.

10. The New World

General Conditions. One would not expect to find a treatise on mathematics printed in the New World within sixty-four years of its discovery by Columbus, and still less would he think that only forty-five years after this great discovery there was set up a press for the disseminating of knowledge among the inhabitants of the western hemisphere. Each of these events stands out, however, as a historic fact, a testimony to the zeal and foresight of the early conquerors.

Among the adventurous band organized by Cortés for his first expedition to Yucatan in 1518 was a young chaplain whose name appears in his work of 1556 as Juan Diez. He was of a literary turn of mind, as is shown by three or four books which he published. One of these works was on mathematics, and this appeared in Mexico under the following title:[2]

El Sumario cõpẽdioso delas quẽtas de plata y oro q̃ en los reynos del Piru son neceſſarias a los mercaderes: y todo genero de tratantes. Cõ algunas reglas tocantes al Aritbmetica.

Fecho por Juan Diez freyle.

It should be stated, however, that there were several writers at this time by the name of Juan Diez (Diaz), two of them

[1] Yoshida Shichibei Kōyū, Imamura Chishō, and Takahara Kisshu.

[2] D. E. Smith, *The Sumario Compendioso of Brother Juan Diez*, Boston, 1921.


apparently being in Mexico, and there is much uncertainty among the best Spanish biographers as to which was the author of the *Sumario*.

Printing Established in Mexico. The first viceroy of New Spain, which included the present Mexico, was a man of remarkable genius and of prophetic vision,—Don Antonio de Mendoza. He assumed his office in 1535, and for fifteen years administered the affairs of the colony with such success as to win for himself the name of "the good viceroy." He founded schools, established a mint, ameliorated the condition of the natives, and encouraged the development of the arts. In his efforts at improving the condition of the people he was ably assisted by Juan de Zumárraga, the first bishop of Mexico. Among the various activities of these leaders was the arrangement made with the printing establishment of Juan Cromberger of Seville whereby a branch should be set up in the capital of New Spain.

The idea of setting up a press in Mexico seems to have been considered as early as 1534, even before Mendoza became viceroy, doubtless at the suggestion of Juan de Zumárraga; but it was not until 1536 that the plan was carried out. Juan Cromberger then sent over as his representative one Juan Pablos, a Lombard printer, and so the "casa de Juan Cromberger" was established, prepared to spread the doctrines of the Church to the salvation of the souls of the unbelievers. Cromberger himself never went to Mexico, but his name appears either on the *portadas* or in the colophons of all the early books. From and after 1545, however, the name is no longer seen, Cromberger having died shortly before this time.

Nature of the Sumario Compendioso. The *Sumario Compendioso* consists of one hundred and three folios, generally numbered. After the dedication there is an elaborate set of tables, including those relating to the purchase price of various grades of silver, to per cents, to the purchase price of gold, to assays, and to monetary affairs of various kinds.

The mathematical text consists of twenty-four pages besides the colophon. Of these pages, eighteen relate chiefly to arithmetic and six to algebra. The arithmetic includes problems in the reduction of maravedis to pesos, of ducats to crowns, and the like; but it also considers questions relating to the preceding tables and to simple commercial transactions in general. There are also problems relating to the theory of numbers, some of them involving rules similar to those found in the works of Fibonacci and Diophantus. Of the latter type the following is an example:



PART OF THE TABLE OF THE *SUMARIO*

The entire table covers nearly one hundred and eighty pages, and the above facsimile fills half of one page

Give me a number which, increased by 15, is a square number; and decreased by 4 is also a square number. *Rule*: Add 15 and 4, making 19; then add 1 to this result, making 20. Now take the half of this number 20, which is 10; square this result thus: 10 times 10 is 100. From this subtract 15, and we have 85, and this is the number required, that is, the one from which if you subtract 4 you have 81, the root of

.25.Re.ɣ da.	24.
100.Re ɣ da.	96.
169.Re.ɣ da.	120.
225.Re.ɣ da.	216.
289.Re.ɣ da.	240.
400.Re.ɣ da.	384.
625.Re.ɣ da.	336\ɣ,600.
676.Re.ɣ da.	480.
841.Re.ɣ da.	840.
900.Re.ɣ da.	864.
1156.Re.ɣ da.	960.
1225.Re.ɣ da.	1176.
1212.Re.ɣ da.	1080.
1681.Re.ɣ da.	720.
2025.Re.ɣ da.	1944.
2500.Re.ɣ da.	1344 2400.
2602.Re.ɣ da.	2160.
2704.Re.ɣ da.	1920.
2809.Re.ɣ da.	2520.
3025.Re.ɣ da.	2905.
3364.Re.ɣ da.	3360.
3600.Re.ɣ da.	3456.
3221.Re.ɣ da.	1320.
4225.Re.ɣ da.	2026.

CONGRUENT AND CONGRUOUS NUMBERS FROM THE *SUMARIO* OF JUAN DIEZ

He defines the terms thus: "A congruous number is such a square number that, subtracting from or adding to it another number, called a congruent number, it will still be a square." There are five errors in the table here given

which is 9. The same thing happens if you add the 15, the result being a hundred, the root of which is 10; for 10 times 10 is 100, which checks.

Some idea of the work in the theory of numbers may be obtained from the table of congruous and congruent numbers here shown in facsimile, and of the nature of the algebraic problems from the following example:

A man has mares and cows in quintuple proportion, in such a way that if you square the number of mares and square the number of cows, the products added will be 1664. Required the number of mares and the number of cows.

Apparently the author had some taste for mathematics and was fairly well versed in algebra as it was then known in the best schools of Spain and Italy. When we reflect that only two important treatises on algebra had at that time been issued from the European presses, and that the *Sumario* was the first mathematical work to be published on another continent, the credit due to Diez is the more apparent.

TOPICS FOR DISCUSSION

1. Reasons for the great advance made in mathematics in the 16th century.

2. Causes of the prominence of Italy in mathematical research in the 16th century.

3. Influences leading to the predominance of algebra in the field of mathematics in the 16th century.

4. The leading features in the progress of algebra in the 16th century, with a consideration of the countries and individuals most closely connected with this progress.

5. A comparison of the mathematics of England, France, Italy, and the Teutonic countries in the 16th century.

6. Influences leading to the advance in mathematics in the Netherlands at this time.

7. The leading treatises on algebra in this century, with a consideration of the important features of each.

8. The life and works of Robert Recorde. His influence upon British mathematics.

9. The way in which the arithmetics of the various countries met the commercial needs of the time.

10. The life and works of Michael Stifel.

11. Influences leading to the development of trigonometry in Europe at this time.

12. Effect of the Renaissance upon the nature of mathematics in the 16th century.

13. The nature of Oriental mathematics in the 16th century.

14. A consideration of the causes of the backward state of mathematics in Spain and Portugal in the 16th century.

15. The life and works of Copernicus and his influence upon mathematics in general.

16. The field of mathematical activity developed by the Jewish scholars in the 16th century.

17. The general nature of the literary activity in Mexico in the 16th century, with special reference to the need which produced the work of Juan Diez.

18. The revival of the study of the works of the classical writers in the 16th century, with special reference to its influence upon mathematics.

*

CHAPTER IX


THE SEVENTEENTH CENTURY

1. General Conditions

Political Situation. In order to comprehend the causes of the remarkable advance of mathematics in the 17th century it is necessary to consider briefly the political and social influences that made this century conspicuous in science, in letters, and in the development of human rights. This was the century that saw broken forever in the Anglo-Saxon civilization the doctrine of the divine right of kings; that saw the beginning of the end of this same doctrine in France in the brilliant reign of Louis XIV; and that saw Russia amalgamated into a powerful nation by a powerful leader, Peter the Great. In this century, too, we may possibly find one of the early steps toward the World War of 1914–1918, namely, the founding of the military machine of Prussia at the hands of the Great Elector of Brandenburg. This was also the period in which Europe saw the turning back of the Turks by the Hapsburgs in Austria; in which the New World was definitely opened to colonization and trade; and in which the Thirty Years' War (1618–1648) disturbed the political and religious life of a considerable part of Europe. Great world activities like these could not but affect natural science as well as political, and abstract science as well as natural.

The Trend to the North. There was also the general reason why mathematics, like all intellectual pursuits, should trend to the north,—the reason that had slowly led it away from the warmer countries from time immemorial,—the ability to conquer the cold and the darkness of the long winter nights. Heat and light have always joined with the soil and the

distribution of moisture to make conditions favorable for intellectual work. While coal was known in England as early as the 9th century, the 16th and 17th centuries saw a great advance in the comforts of living north of the Alps, and hence in the ability to utilize the long winter nights in intellectual pursuits. Other influences were evidently more powerful, but this one must be recognized as being somewhat significant.



MATHEMATICS IN MILITARY AFFAIRS

Measuring the distance to a castle by the aid of a form of quadrant. From Mario Bettino's *Apiaria*, Vol. I, Bologna, 1645

Mathematical Situation. Mathematics does not develop by centuries any more than by political units or by religious faiths. Vieta and Harriot were quite as much of the 16th as of the 17th century, just as Erasmus was quite as much of England and Germany and France as of the Low Countries, and as Stifel the Lutheran was quite as good an algebraist as Stifel the monk. Nevertheless it is convenient to mark off blocks of time, and centuries serve the purpose fairly well in mathematics as in art and in politics.

It is impossible to say with truth that this century or that is the greatest in the development of any human interest, but it is entirely within the range of truth to assert that few if any centuries did so much for mathematics as that one which

saw Fermat begin the modern theory of numbers and, with Descartes and Harriot, invent the analytic geometry; Cavalieri pave the way for Newton and Leibniz, who, in their turn, established the calculus; Pascal and Desargues open new fields for pure geometry; Napier reveal to the world a new method of computation; and a large number of brilliant scholars apply the theories thus developed to the study of curves, to difficult problems of mensuration, and to the science of celestial mechanics.


Moreover, the 17th century was characterized by a new spirit,—that of intellectual internationalism, of a free exchange of ideas among members of the learned class, and of the calling of scholars from one country to another, sometimes by universities, sometimes by academies, sometimes by royal command. This spirit was even more manifest in the 18th century, when men like Euler, the Bernoullis, and Lagrange were looked upon no longer as national assets, but as Europeans in the largest sense of the term.

It seems unreasonable to separate these intellectual activities from the general Spirit of the Times. Printing had begun to show its power; not merely the intellectual aristocracy but the people in general were beginning to think; the scholar could now make his discoveries known, and his audience was no longer composed literally of those who heard his voice; and with the breaking of religious and political canons came also the breaking of the canons of science.

As an incident in the military activity of this period the mathematics of warfare became even more prominent than it was in the 16th century,—a symptom of the terrible part that it was to play some three hundred years later in the great World War.

Effect of Skepticism in General. Skepticism (not religious skepticism in particular, but skepticism with respect to tradition in general,—skepticism which Buckle has called "hardness of belief") has been the *sine qua non* of progress. This is as true in mathematics as it was in the combat with the physics of Aristotle, with the music of Boethius, with the canons of

Giotto, or with the divine right of kings. Just so long as Euclid was sacrosanct in elementary geometry, or Apollonius in conics, or Ptolemy in astronomy, or Boethius in arithmetic, the world could not progress in mathematics. It was only when men began to doubt the infallibility of these ancient leaders that they developed a new conception of geometry, a new way of handling conics, a new system of the universe, and a new



MATHEMATICS IN MILITARY AFFAIRS
From Leonhard Zubler's work on geometric instruments, Zürich, 1607

view of arithmetic. The world began to get skeptical of authority in the 16th century, and the leaven of new ideas worked so rapidly that the opening of the 17th century saw the time ripe for an entire recasting of mathematical theories.

2. Italy

Shifting the Center. The 17th century saw a shifting of the center of mathematical activity from Italy northward. One cause has already been mentioned, and other causes are not difficult to see, being largely political. In general, mathematics flourishes where the environment is favorable, and in

the 17th century the political environment was more favorable in France and England than in Italy. The glory of Venice was rapidly fading out; Pisa was no longer a seaport; Florence had ceased to be the source of business customs; Rome had lost her hold upon the most vigorous parts of Europe; and the rivalry between the Italian states was not of that healthy nature which makes for intellectual progress. Italy was still a mathematical power, but it was no longer the world's intellectual center.

Cavalieri. From the standpoint of mathematics alone the Italian writer who influenced the science most in the 17th century was probably Bonaventura Cavalieri,[1] a Jesuit, a pupil of Galileo's, and professor of mathematics at the University of Bologna from 1629 to the time of his death. He wrote on conics,[2] trigonometry,[3] optics, astronomy, and astrology, and was one of the first to recognize the great value of logarithms. His greatest contribution, however, was his principle of indivisibles,—a principle announced by him in 1629 but not set forth in printed form until six years later.[4] The theory is based upon the assertion that a line is made up of an infinite number of points, a plane of an infinite number of lines, and a solid of an infinite number of planes. The theory thus forms the basis

[1] Born at Milan, 1598; died at Bologna, November 30, 1647. The year of his birth may, however, be 1591. He may have died December 1, 1647, but it was the night of November 30–December 1. See P. Frisi, *Elogio del Cavalieri*, Milan, 1778; G. Piola, *Elogio di Bonaventura Cavalieri*, Milan, 1844; F. Predari, *Della Vita e delle Opere di Bonaventura Cavalieri*, Milan, 1843; A. Favaro, *Bonaventura Cavalieri nello studio di Bologna*, Bologna, 1888. The name appears also as Cavallieri, Cavaglieri, Cavalerius, and de Cavalleriis. His place of birth is given by Piola as Milan, although others give it as Bologna. The tomb records the date of his death and the fact that he was a native of Milan.

[2] *Lo specchio ustorio, overo Trattato delle settioni coniche*, Bologna, 1632.

[3] *Directorium generale uranometricum in quo Trigonometriae logarithmicae fundamenta ac regulae demonstrantur, astronomicaeque supputationes ad solam fere vulgarem eruditionem reducuntur*, Bologna, 1632; *Compendio delle regole dei triangoli colle loro dimostrazioni*, Bologna, 1638; *Trigonometria plana et spherica*, Bologna, 1643.

[4] *Geometria Indivisibilibus continuorum nova quadam ratione promota*, Bologna, 1635; 2d ed., Bologna, 1653; *Exercitationes Geometricae sex*, Bologna, 1647.

of a crude kind of calculus, and by its aid Cavalieri found it possible to solve many problems in mensuration that would now be solved by the more scientific methods of integration. The term "indivisible" is ancient, and Cavalieri made no claim to originality in its use.



BONAVENTURA CAVALIERI


From an engraving by G. A. Labus after a drawing by A. Alfieri

Galileo. Much more widely known than Cavalieri,—more widely known, indeed, than most men of his time,—Galileo Galilei[1] was destined to bring great glory to Italy in general and to Florence in particular. Born on the day of Michelangelo's death, and dying in the year of Newton's birth, he seemed to fill the gap between the lives of these two great leaders, a stirring period in the history of art, letters, politics, science, and religious thought; and in the numerous controversies relating to all these lines he played a major part.

He was the son of a certain Florentine nobleman, a dilettante in music and mathematics, whose estate had become so greatly

[1] Born at Pisa, February 18, 1564; died at Florence, January 8, 1642. The literature relating to Galileo is extensive. A good bibliography, particularly as to his life and the controversy over the inquisition question, is given by Karl von Gebler, *Galileo Galilei and the Roman Curia*, English ed., London,

reduced as to indicate for Galileo a life devoted to the restoration of the family fortune,—the life of a cloth merchant. By



GALILEO GALILEI, 1564–1642

From the painting in the Pitti Gallery in Florence, school of Sustermans (Suttermans, 1597–1681). Robinson's *Medieval and Modern Times*

some good chance, however, he was sent to the convent of Vallombrosa, and here he displayed such unusual powers that

1879, p. xxix. See also P. Frisi, *Elogio del Galileo*, Milan, 1774; G. P. C. de' Nelli, *Vita e commercio letterario di Galileo Galilei*, 2 vols., Lausanne, 1793; A. Favaro, *Galileo Galilei e Suor Maria Celeste*, Florence, 1891, with valuable material relating to his contemporaries. Viviani's sketch of the life of his great teacher may be found in the Milan edition of the *Opere di Galileo Galilei* of 1808–1811, Vol. I. The best edition is the one published under the editorship of A. Favaro, Florence, 20 vols., 1890–1909, with a biographical index in Volume XX.

his father changed his mind and decided that he should study medicine. He entered the University of Pisa in 1581, in a period of great intellectual upheaval, but his medical studies were soon put aside by an incident that was to help change the scientific thought of the world. The beautiful lamp that still hangs in the cathedral at Pisa had been moved from its vertical position in order the more readily to light it, and Galileo noticed that the oscillations were at first considerable but gradually became less and less. They seemed, however, to be made in equal periods of time, and this inference he confirmed by comparing them with his pulse. Thus he was able to establish the approximate isochronism of the vibrations (a fact which had been asserted by the Arabs) and to make a beginning in medical diagnosis by accurately timing the arterial beats. Galileo was also led at this time, it seemed by chance, and certainly against his father's wishes, to the study of geometry. His success was such that at last he secured parental consent to give up medicine and devote himself to science. He soon became well known throughout Italy, and in 1589 was made professor of mathematics at Pisa. It is indicative of the low esteem in which mathematics was then held that, whereas the professor of medicine received the equivalent of about $2150 a year, Galileo was rewarded by a salary equivalent to only $65. It was here that he began his experimental work in physics, but owing to local controversies he resigned his chair in 1591. The next year he was offered the professorship of mathematics at Padua, and here he was enabled to carry on some of his most important scientific work.

Of his controversies in astronomy, of his construction of the first satisfactory telescope,[1] of his invention of the modern type of microscope, and of his work in physics this is not the place to speak. It should be said, however, that his interest in mathematics was maintained throughout his stormy life. While at Padua he invented the proportional

[1] On the invention of the telescope the first and one of the best of the standard works is P. Borel, *De Vero Telescopii Inventore*, The Hague, 1655–1656.

compasses,[1] an instrument which was in great favor for a century or more but which of late has been generally discarded.

Torricelli. Of those physicists who sat at the feet of Galileo and who also showed an interest in pure mathematics no one was more celebrated than Evangelista Torricelli.[2] Although more than forty years the junior of Galileo, he survived him by only five years, dying at the age of thirty-nine. He had studied under a pupil of this great master, but was also privileged to receive instruction from the latter himself, then blind and enfeebled by age, in the last year of his life. Not only did Torricelli write on physical questions and give to the world the barometer, but he contributed to geometry as well,[3] anticipating the work of Roberval on the method of tangents. Père Mersenne had announced to Galileo in 1638 that Roberval had squared the cycloid, a curve to which Galileo had first called attention. Galileo thereupon sent the letter to various friends, and Torricelli responded by squaring the cycloid, and Viviani by determining the tangent.



EVANGELISTA TORRICELLI
After a contemporary engraving

[1] *Le Operazioni del Compasso Geometrico et Militare*, Padua, 1606. See also A. Favaro, "Per la storia del compasso di proporzione," in the *Atti* of the *Istituto Veneto*, LXVII, 2, 723.

[2] Born in or near Faenza, possibly in the village of Modigliana, October 15, 1608; died at Florence, October 25, 1647. See the *Opere* of Torricelli, ed. Loria and Vassura, *Introduzione*, Faenza, 1919; G. Loria, *Atti d. R. Accad. dei Lincei*, XXVIII (5), 409.

[3] *Opera geometrica*, Florence, 1644; F. Jacoli, "Evangelista Torricelli ed il metodo delle tangenti detto Metodo del Roberval," Boncompagni's *Bullettino*, VIII, 265.

Vincenzo Viviani. This Vincenzo Viviani[1] was also a disciple of Galileo's. He was interested in physics and in the applications of mathematics, but his tastes led him even more strongly to geometry.[2] His first work (1659) established his reputation, and he was honored by the Medici, made mathematician to Ferdinand II, the grand duke of Tuscany, elected to membership in various learned societies, and invited to France by Louis XIV and to Poland by King Casimir,—invitations which he felt compelled to decline. In 1692 he proposed to scholars a problem which attracted wide attention, and which may be stated briefly as follows: There is among the ancient monuments of Greece a temple dedicated to Geometry. The plan is circular and the temple is surmounted by a hemispherical dome which has four equal windows of such size that the rest of the surface can be exactly squared. Required to find how this is possible.

The problem appeared in the *Acta Eruditorum* under a designation which is an anagram of the words "A postremo Galilei Discipulo," a title which Viviani was always proud to bear.[3] Of this problem there were submitted correct solutions by Leibniz, Jacques Bernoulli, l'Hospital, Wallis, and David Gregory, but Viviani himself gave the simplest one of all. He also solved the trisection problem by the aid of the equilateral hyperbola.

Minor Writers. Among the minor Italian writers of the 17th century Giovanni Antonio Magi'ni,[4] a friend of Kepler, was widely known as the maker of the most perfect maps of Italy up to that time, and as professor of astrology, astronomy, and mathematics in the University of Bologna for nearly thirty

[1] Born at Florence, April 5, 1622; died at Florence, September 22, 1703.

[2] *De Maximis, et Minimis Geometrica Divinatio in Qvintvm Conicorum Apollonii Pergaei adhvc desideratvm*, Florence, 1650; *Quinto libro degli Elementi d' Evclide*, Florence, 1674; *De Locis Solidis Secunda Divinatio Geometrica*, Florence, 1673 and 1701; and other works.

[3] Viviani's solution appeared in his *Formazione e misure di tutti i cieli, con la struttura e quadratura esatta dell' intero, e . . . uno degli antichi delle volte regolari degli architetti*, Florence, 1692.

[4] Born at Padua, June 13, 1555; died at Bologna, February 11, 1617 or 1615.

years (from 1588). While his interest was chiefly in astronomy, on which he wrote extensively, he also contributed to the theory of numbers[1] and to trigonometry.[2]

Marino Ghetaldi,[3] another of the minor writers of the period, was descended from a patrician family and divided his time between scientific and diplomatic pursuits, working in the field of mathematics and acting as ambassador for Venice to Rome and to Constantinople. In his chosen field of science he still further divided his interests, this time between pure mathematics and its physical applications. He wrote upon geometry[4] and algebra[5] but contributed little that was original.

Giovanni Alfonso Borelli,[6] a third in the list of lesser mathematicians, studied at Pisa and then taught philosophy and mathematics at Messina (1649). In 1656 he was recalled to Pisa to take the chair of mathematics. He was also a physician, and his posthumous work *De motu animalium* (1680–1685) was highly esteemed. He edited some of the Greek classics on mathematics.[7]

The Cassini Family. Although classified among the minor writers from the standpoint of their contributions to pure mathematics, several of the members of the remarkable Cassini family would stand among the leaders if we considered

[1] *Tabula tetragonica seu quadratorum numerorum, cum suis radicibus*, Venice, 1592.

[2] *De Planis Triangvlis Liber Vnicus*, and *De Dimetiendi ratione per Quadrantem, & Geometricum Quadratum, Libri Qvinqve*, Venice, 1592; *Tabulae et canones primi mobilis; item calculus triangulorum sphaericorum*, Venice, 1604.

[3] Born at Ragusa, 1566; died probably at Constantinople, but possibly at Ragusa, 1626 or 1627.


[4] *Nonnullae propositiones de parabola*, Rome, 1603; *Apollonius redivivus seu restituta Apollonii pergaei de inclinationibus Geometria*, Venice, 1607; and two other works. See also H. Wieleitner, *Bibl. Math.*, XIII (3), 242; E. Gelcich, *Abhandlungen*, IV, 191.

[5] *De resolutione et compositione mathematica, libri quinque, Opus posthumum*, Rome, 1630,—a work on algebra applied to geometry.

[6] Born at Castelnuovo, near Naples, January 28, 1608; died at Rome, December 31, 1679.

[7] *Euclides restitutus*, Pisa, 1658; *Apollonii Pergaei conicorum libri V, VI, et VII*, Florence, 1661; *Elementa conica Apollonii et Archimedis opera*, Rome, 1679.

their work in the field of astronomy. At any rate they deserve to be mentioned in this connection because of their skillful application of mathematics in their chosen field of science. The founder of the astronomical line was Giovanni Domenico Cassini,[1] professor of astronomy at Bologna (1650). Louis XIV asked that he be sent to Paris (1669), and soon after



AUTOGRAPH OF GIOVANNI DOMENICO CASSINI

From a receipt written on parchment, February 27, 1706. The French form of the name was adopted after he went to France in 1669

he became (1671) the first astronomer royal of France. Since he became naturalized in France and his son Jacques Cassini[2] was born there, the line now ceases to be Italian. Jacques, the second son, succeeded his father (1712) as astronomer royal, and, like him, wrote numerous monographs on astronomy. César-François Cassini de Thury,[3] the son of Jacques, succeeded his father (1756) as astronomer royal, and was in turn

[1] Born at Perinaldo, June 8, 1625; died in Paris, September 14, 1712.
[2] Born in Paris, February 18, 1677; died at Thury, April 16, 1756.
[3] Born in Paris. June 17, 1714; died in Paris, September 4, 1784.

succeeded by a son, Jacques Dominique Cassini de Thury.[1] Each of these members kept up the traditions of the remarkable family with respect to its contributions to science.

3. France

France and England. It is of little moment whether we say that the center of mathematical activity in the 17th century rested in France or in England; perhaps it would be more just to speak of each as one of the foci of an ellipse, as was the case with Bagdad and Cordova, so strong are the claims that each may fairly adduce. When we try to balance such names as Harriot, Napier, Oughtred, Wallis, Barrow, and Newton against so remarkable a group as Fermat, Desargues, Descartes, Pascal, Mersenne, and l'Hospital, it is like comparing two infinities. Perhaps England's mathematics was more usable in the natural sciences, while that of France was more of the nature of *l'art pour l'art*; but any such distinction is easily attacked. If we speak of France before England, it means only that we may consider at random either focus of the ellipse with equal justice.

As to France, Paris had risen in political, intellectual, and artistic splendor, while Lyons had fallen. The Lyonese still had their four great fairs annually, and were content with their lot. The two cities still represented, however, the mathematics of the country.

Early Writers. Among the early French writers of this century, Denis Henrion[2] published the first logarithmic table to appear in France.[3] Contemporary with him was the learned scholar Claude Richard,[4] who entered the Jesuit order in 1606, taught mathematics in Lyons for a number of years, was called to Spain in 1624, and became professor of mathematics at

[1] Born in Paris, June 30, 1748; died at Thury, October 18, 1845.

[2] Born *c.* 1590; died *c.* 1640.

[3] *Traicté des logarithmes*, Paris, 1626. He also published a *Collection . . . de divers Traictez Mathematiques*, Paris, 1621; the *Logocanon, ou Regle Proportionelle*, Paris, 1626; and other works.

[4] Born at Ornans, Burgundy, 1589; died at Madrid, October 20, 1664.

Madrid. He wrote commentaries on Euclid's *Elements* (1645), the *Conics* of Apollonius (1655), and other Greek works.

Another writer of this period, Pierre Hérigone, whose life was one of comparative obscurity, published a work on general mathematics[1] which stands out as a good summary for the time, and which displayed considerable originality in the field of algebra.

Descartes. If one were asked to name the man who was most influential in the revolutionizing of mathematics in the 17th century, he would naturally find it difficult to answer. Probably the name of Newton would lead in any ballot among scholars. Newton's modest assertion that he had seen farther only by standing on the shoulders of giants[2] is capable of easy proof, as is the case with most men of eminence. It is the genius, however, who can pick out his giants. Certainly among those selected by Newton or by Fate was René Descartes,[3] a man whose varied genius led philosophers to rank him primarily as one of themselves, physicists to claim him for their guild, and mathematicians to look upon him as one of the greatest geniuses in their domain, each group being fully justified in its own opinion.

Descartes was fortunate in his birth, his father, Joachim (died 1640), being a counselor in the parliament[4] of Bretagne (from 1586) and possessed of sufficient means to give the son

[1] *Cursus mathematicus*, 5 vols., Paris, 1634–1637, with a *Supplementum*, 1642–1644.


[2] D. Brewster, *Life of Newton*, 1, 142. Edinburgh, 1855.

[3] Born at La Haye, Touraine, March 31, 1596; died at Stockholm, February 11, 1650. Latin, Renatus Cartesius, whence we speak of the Cartesian geometry and the Cartesian philosophy. The earlier English and French writers frequently used the form Des Cartes. Other branches of the family used the older spelling Des Quartes and Des Quartis. René was also called, against his wish, M. du Perron, from a small *seigneurie* belonging to the family and situated near Poitou. This name was used by the family in his younger days to distinguish him from his brother.

The most recent biographies are those of G. Milhaud, *Descartes, Savant*, Paris, 1921, containing various essays theretofore published; Elizabeth S. Haldane, *Descartes, His Life and Times*, London, 1905; and C. Adam, *Descartes*, Paris, 1910. See also J. Millet, *Descartes, sa vie, ses Travaux, ses découvertes, avant 1637*, Paris, 1867.

[4] *Parlement*, little more than a local court.

an opportunity for early advance. When only a child René was placed in a recently founded Jesuit school at La Flèche,



RENÉ DESCARTES

After the painting by Franz Hals, now in the Louvre

in Maine, where he remained for eight years (1604–1612) and where he came to know Mersenne (p. 380), who was seven or eight years his senior. The two established there a friendship

Monsieur

ay receu le contenu de la lettre de change qui
vous a plû m'envoyer et vous en remercie.
ie voudroy garder un peu plus long temps pour
tascher de vous le remettre avec quelque
profit, mais ie ne doute point qu'il ne profite
davantage estant entre vos mains qu'il ne
pourroit faire entre les miennes; et ie suis sur le
point de partir d'icy. Ie ne say que respondre
a la courtoisie de Monsieur t'Huguens sinon que ie
cheris l'honneur de sa connoissance comme l'une de
mes meilleures fortunes, et que ie ne feray iamais
en lieu ou ie puisse avoir le bien de le voir que
ie n'en recherche les occasions, ainsy que ie feray
toujours celles de vous tesmoigner que ie suis

Monsieur

Amsterdam ce 23 May 1632

Voſtre treshumble et
treſaffectionné ſerviteur
Des Cartes

AUTOGRAPH LETTER OF DESCARTES

Written about the time he was completing the manuscript of *La Géométrie*. The M. Huguens mentioned was the father of Christiaan Huygens, then a child of three years

that endured through life. He was sixteen years old when he finished his course at La Flèche and returned to his home "overwhelmed by the blessings and praises of his teachers." There he remained for a year, his father then deciding that he should complete his education in Paris. Here he renewed his acquaintance with Mersenne, now become a Minimite, but the acquaintance was soon interrupted by his friend's departure for Nevers. He now set about to cultivate the study of geometry, made the acquaintance of Mydorge (p. 378), and came in contact with the best exponents of the mathematical thought of the day. It was not long, however, before he felt the need for a broader knowledge of the world and so decided to enlist in army service. Not sympathizing with the aims of the French nobility, he joined the army of Maurice, Prince of Orange, a man who had already begun to attract to himself a group of scholars, and in this way he became acquainted with Stevin (p. 342). He soon became weary of army life, however, and in 1621 withdrew from the service. He now devoted four years to travel, visiting the German states, Denmark, Holland, Switzerland, and Italy, returning to Paris in 1625. His old friend Mersenne had also returned, Mydorge was still there, Desargues (p. 383) had joined the group, and besides these men of the mathematical coterie he met Balzac and the literary world generally, and was presented to Richelieu. Paris was not the place for his meditations, however, and in 1628 he decided to take up his residence in Holland, staying there until 1649, when, at the invitation of Christina of Sweden, he went to Stockholm, where he remained until his death (1650) a few months later.

His first years in Holland (1629–1633) were given to the preparation of a work on philosophy.[1] On finishing this treatise he learned of the condemnation of Galileo for daring to tell the truth, and so he, quite unlike his Italian contemporary, decided that discretion was the better part of valor. As a result, the work was not published during his lifetime. He then devoted himself to the preparation of his great treatise on

[1] *Le Monde*, Paris, posthumously published in 1664.

method in science,[1] a work which included three appendixes, one of which bore the modest title *La Géométrie.*

Descartes's Geometry. It was in this appendix, a small handbook of only about a hundred pages, that analytic geometry first appeared in print.[2] The fundamental idea in Descartes's mind was not the revolutionizing of geometry so much as it was the elucidating of algebra by means of geometric intuition and concepts; in a word, the graphic treatment of the equation. His imagination extended far beyond this, however, to the establishing of a universal mathematics in which algebra, geometry, and arithmetic should be closely related members. He began by extending the ancient idea of latitude and longitude, showing that any point in a plane is uniquely determined by two coordinates, x and y, the equation $F(x, y) = 0$ expressing a property which is true for every point of the curve. By studying the equation, therefore, he could, through the principle of a one-to-one correspondence, transfer his results at any time to the curve itself. It constituted what John Stuart Mill[3] says was "the greatest single step ever made in the progress of the exact sciences." The *Géométrie* is divided into three books. In the first book he relates the fundamental operations of arithmetic to geometry, his use of a certain unit of length being the only novelty in this feature; in the second book[4] he classifies curves and considers the methods of finding tangents and normals; and in the third book[5] he deals with the nature

[1] *Discours de la méthode pour bien conduire sa raison et chercher la vérité dans les sciences*, Leyden, 1637. The appendixes were *La Dioptrique*, *Les Météores*, and *La Géométrie*. A Latin translation of the geometry, by Frans van Schooten, appeared at Amsterdam in 1649.

[2] The *Opera Omnia* of Descartes appeared at Amsterdam in 9 volumes, 1690–1701, with a second impression in 1713. It also appeared at Paris in 13 volumes, 1724–1726. The V. Cousin edition of *Les Œuvres* appeared at Paris in 11 volumes, 1824–1826, with a new edition by J. Simon in 1844. The latest edition, edited by Charles Adam and Paul Tannery, was printed at Paris, Volume I appearing in 1897. For a bibliography of the most important works relating to Descartes see Haldane, *loc. cit.*, p. 387.

[3] *An Examination of Sir W. Hamilton's Philosophy*, p. 617. London, 1878.

[4] In the Latin edition, "De natura linearum curvarum."

[5] In the Latin edition, "De constructione Problematum Solidorum, & Solida excedentium."

of the roots of equations, considering among other things the rule of signs that has since been known by his name. The idea of analytic geometry had already been worked out by Fermat, or at least was conceived by him and Descartes at about the same time; but Fermat only thought, while Descartes not only thought but wrote. Some conception of the plan was probably also in the mind of Harriot, and Descartes was familiar with Harriot's work;[1] but, after all, the real idea of functionality as shown by the use of coordinates was first clearly and publicly expressed by Descartes.

Some idea of his range of knowledge and interest may also be obtained from his work on anatomy, begun in 1634, and of which a Latin edition appeared at Leyden in 1664.[2]

Descartes the Man. Descartes inherited, apparently from his mother, a feeble constitution. He speaks of his "dry cough and pale complexion," which remained with him until he entered the army, and which led his physician to predict that his life would be a short one. He overcame his early ailments, however, and although he died at fifty-four he was able to say in his later life that for thirty years he had been free from any illness that deserved the name. John Stuart Mill remarked of him: "Descartes is the completest type which history presents of the purely mathematical type of mind—that in which the tendencies produced by mathematical cultivation reign unbalanced and supreme." Although the statement may well be questioned, it is interesting as a striking assertion if for no other reason.

Early Commentators on Descartes. Of the two leading commentators on Descartes in the 17th century the first (1649) was his warm personal friend, Florimond de Beaune,[3] an office-holder at Blois. He also wrote on algebra,[4] being one of the

[1] *Artis Analyticae Praxis*, London, published posthumously in 1631.

[2] *De Homine figuris, et Latinitate Donatus a Florentino Schuyl*, Leyden, 1664. This edition has some engravings of great merit.

[3] Born at Blois, 1601; died at Blois, 1652.

[4] *De aequationum constructione et limitibus*, published posthumously at Amsterdam in 1659. There was also an edition, Amsterdam, 1683.

first to treat scientifically of the superior and inferior limits of the roots of a numerical equation. He endeavored to deduce the nature of curves from the properties of their tangents and was also interested in the improvement of astronomical instruments. The second and more prominent of the commentators was Frans van Schooten[1] (*c.* 1615–1661), who will be considered later.



FERMAT

After an old lithograph

Fermat. The life and works of Pierre de Fermat[2] illustrate the fact that no one can account for a genius. Why should the greatest writer on the theory of numbers, at least after the time of Diophantus, suddenly appear in the person of a modest, retiring, punctilious counselor of the parliament of Toulouse, and in the 17th century? Why should this obscure officeholder succeed in making such a name for himself while apparently giving no serious attention to mathematics until he was over thirty? And why, aware of his powers as he must have been, was he content to make his results known chiefly through letters to men like Mersenne, Roberval, Pascal, and Descartes instead of publishing them for the benefit of scholars in general? One answer to each of these questions is that genius is eccentric.

It may have been Bachet's translation (1621) of Diophantus that directed Fermat's attention to the theory of numbers, for

[1] Known also as Franciscus van Schooten.

[2] Born at Beaumont de Lomagne, near Toulouse, *c.* 1608; died at Castres or Toulouse, January 12, 1665. The date of his birth as given by different writers varies from 1590 to 1608. His tombstone in the church of the Augustines in Toulouse (later in the museum) gives the date of death as above and the age as fifty-seven years.

E. Brassinne, *Précis des œuvres mathématiques de Pierre Fermat*, Paris, 1853; *Œuvres de Fermat*, edited by P. Tannery and C. Henry, Paris, 1891–1912. See also C. Henry in Boncompagni's *Bullettino*, XII, 477 and XIII, 437; P. Tannery in Darboux's *Bulletin*, VII (2); G. Libri, "Fermat," in *Revue des Deux Mondes*, May 15, 1845.

he left a series of notes and letters on this work which were published in the form of a commentary (Toulouse, 1670) after his death. At any rate he showed remarkable ability in this field of mathematics. He asserted that no integral values of x, y, and z can be found to satisfy the equation $x^n+y^n=z^n$ if n is an integer greater than 2, and this is commonly known as Fermat's Theorem. No satisfactory demonstration has ever been published, and it is not known whether Fermat himself demonstrated it, few of his proofs having been preserved. What knowledge we have of these proofs is due in part to marginal notes made by him in his reading, and in part to a certain manuscript of Huygens found at Leyden in 1879. He seems to have claimed that he had proved this particular theorem. Fermat's letters show that he had developed the idea of analytic geometry before Descartes published (1637) his work upon the subject. Descartes proposed to represent a curve by an equation, to study this equation, and in this way to discover the properties of the curve itself; while Fermat did substantially the same thing, designating the equation as the "specific property" of the curve and deriving all other properties from it.

In connection with his study of curves Fermat proceeded to apply the idea of infinitesimals to the questions of quadrature and of maxima and minima as well as to the drawing of tangents. In this he seems to have anticipated the work of Cavalieri, but the date of his discovery is unknown.[1]

Mydorge. Of the influential members of the brilliant group of mathematicians that brought the science prominently to the attention of Paris in the first half of the 17th century Claude Mydorge,[2] a friend of Descartes, must be named among the first. He was a man of means, an official of the government, a

[1] The esteem in which he was held by Descartes is shown in one of the letters which the latter wrote to him: "Je n'ai jamais connu personne, qui m'ait fait paraître qu'il sût tant que vous en géométrie." Cantor, *Geschichte*, II, chap. 79. Fermat's *Varia Opera Mathematica*, 2 vols., edited by his son, Samuel Fermat (Toulouse, 1679), was reprinted in facsimile in Berlin in 1861.

[2] Born in Paris, 1585; died in Paris, July, 1647.

physicist of recognized standing, and a mathematician of fair attainments, writing upon optics, conics,[1] and the recreations of mathematics.[2]

Other Writers on the Theory of Numbers. Fermat was not alone in his interest in the theory of numbers at this time. Indeed, he was not the earliest French scholar of the century to consider the subject, for Diophantus had been made known in France before Fermat showed any interest in the subject. The man responsible for this initial step in the theory of numbers was Claude-Gaspar Bachet, Sieur de Méziriac,[3] mathematician, philosopher, theologian, poet, and one of the ablest writers of his day. He came from an ancient and noble family[4] and passed some part of his youth in Italy. He took the first steps toward entering the Jesuit order, but abandoned the idea and went to Paris, where he became a member of the Académie des Sciences. His work on mathematical



MERSENNE
Engraving by Duflos


[1] *De Sectionibus Conicis Libri IV*, Paris, 1631. See also the "Problèmes de Géométrie Pratique" in Boncompagni's *Bullettino*, XVI, 514.

[2] He edited the popular *Recreations Mathematiqves* (title as in the 1628 ed.) of Leurechon (1st ed., 1624). Mydorge's editions were Paris, 1630, 1634, 1639, and later. See Boncompagni's *Bullettino*, XIV, 271.

[3] Born at Bourg-en-Bresse, October 9, 1581; died at Bourg-en-Bresse, February 25, 1638.

[4] For particulars and for biographical information in general see P. Bayle, *Dictionaire historique et critique*, Paris, 1734, with numerous bibliographical references; hereafter referred to as Bayle, *Dictionaire*.

recreations[1] was the best of all that appeared in the 17th century and is still looked upon as a classic in that field, both in style and in content. His well-known translation of Diophantus from the Greek into Latin was published in 1621.[2]



AUTOGRAPH LETTER OF MERSENNE
Written about 1640

Another contributor to the theory of numbers at this time appeared in the person of Marin Mersenne,[3] a Minimite friar, who taught philosophy and theology at Nevers and Paris and was in constant correspondence with the greatest mathematicians

[1] *Problemes plaisans et delectables, qui se font par les nombres*, Lyons, 1612. There were later editions,—Lyons, 1624; Paris, 1874; Paris, 1879; Paris, 1884.

[2] *Diophanti Alexandrini Arithmeticorum libri sex*, Paris, 1621. The text appears in Greek and Latin. Fermat's edition appeared at Toulouse in 1670. Xylander's translation had already appeared at Basel in 1575.

[3] Born at Oizé, Maine, December 8, 1588; died in Paris, September 1, 1648. F. H. D. C., *La Vie dv R. P. Mersenne*, pp. 2, 6 (Paris, 1649).

of his day. He was a voluminous writer, editing some of the works of Euclid, Apollonius, Archimedes, Theodosius, Menelaus, and various other Greek mathematicians. He also wrote on a variety of other subjects, including physics, mechanics, navigation, geometry, and mathematical and philosophical recreations. It is in the theory of numbers, however, particularly with respect to prime numbers and perfect numbers,[1] that he made contributions of real value.



BLAISE PASCAL

After a contemporary drawing

Pascal. Blaise Pascal,[2] whom Bayle[3] appreciatively calls "one of the most sublime spirits in the world," was blessed in having a father who could and did start him in the right direction. Étienne Pascal was an able mathematician and was so desirous of giving the best advantages to his only son that he relinquished his post of Président à la Cour des Aides of his province and went to Paris in 1631. Educated solely under his care, Blaise showed phenomenal ability in mathematics at an early age, and although his father wished him first to have

[1] His *Cogitata Physico-Mathematica*, Paris, 1644, appeared four years before his death. On the nature of Mersenne's Numbers see W. W. R. Ball, *Messenger of Mathematics*, XXI, 34, 121; but consult also L. E. Dickson, *History of the Theory of Numbers*, I, 12 and 31 (Washington, 1919), hereafter referred to as Dickson, *Hist. Th. Numb.*

[2] Born at Clermont-Ferrand in Auvergne, June 19, 1623; died in Paris, August 19, 1662.

[3] Bayle, *Dictionaire*, IV, 500, with an unusually good biography for Bayle, who is commonly more erudite than helpful. See also A. Maire, *L'œuvre scientifique de Blaise Pascal*, Paris, 1912; A. Desboves, *Étude sur Pascal*, Paris, 1878.

a thorough grounding in the ancient languages, and therefore took from him all books on mathematics, he succeeded in beginning geometry by himself and in making considerable progress before his efforts were discovered. Various anecdotes of his youthful activities in mathematics are told by his sister, Madame Périer, who wrote his biography. She relates that he discovered independently most of the first book of Euclid, that his intuition in mathematics seemed miraculous, and that geometry was simply his recreation. He played with conics as other children play with toys, but with the divine enjoyment of discovering eternal truths. When Descartes was shown a manuscript which Pascal wrote on conics at the age of sixteen, he could hardly be convinced that it was not the work of the father instead of the son. At the age of nineteen he invented a computing machine that served as a starting point in the development of the mechanical calculation that has become so important in our time. That he should have been permitted to present one of these machines to the king and one to the royal chancellor shows the esteem in which he must have been held. At the age of twenty-three he became interested in the work of Torricelli in atmospheric pressure, and soon established for himself a reputation as a physicist. Among his discoveries was the well-known theorem which bears his name, that the three points determined by producing the opposite sides of a hexagon inscribed in a conic are collinear,—a theorem from which he deduced over four hundred corollaries. He also wrote (1653) so extensively on the triangular arrangement of the coefficients of the powers of a binomial, which had already attracted the attention of various writers, that this arrangement has since been known as Pascal's Triangle. In connection with Fermat he laid the foundation for the theory of probability.[1] He also perfected the theory of the cycloid and solved the problem of its general quadrature.

[1] I. Todhunter, *A History of the Mathematical Theory of Probability*, Chap. II (Cambridge, 1865), hereafter referred to as Todhunter, *Hist. Probability*. On Pascal's use of induction in this theory see W. H. Bussey, *Amer. Math. Month.*, XXIV, 203. Pascal's work on the triangular arrangement of coefficients was published posthumously in 1665.

His contributions to science and letters often appeared under the nom de plume of Louis (Lovis) de Montalte (as in his *Lettres provinciales*) and the anagram on the same name, Amos Dettonville (as in various problems which he proposed). It is on this account that Leibniz occasionally speaks of him as Dettonville.

Having already, at the age of twenty-five, made for himself an imperishable reputation in mathematics and physics, he suddenly determined to abandon these fields entirely and to devote his life to a study of philosophy and religion. The life of penance which he lived thereafter seems strange, since all must feel that he had little of which to repent, but at any rate it speaks well for the faith of men that he was sincere in his belief and irreproachable in his conduct.

Desargues. In the line of pure geometry the most original contributor of the 17th century was Gérard Desargues,[1] of whose life not much is known except that he was for a time an officer in the army, that he then lived in Paris (1626), where he gave some public lectures, that he was an engineer, and that his later years were spent upon his estate near Condrieux. He published several works, but is known chiefly for his treatise on conics.[2]

Perhaps because it appeared at about the same time as the great work of Descartes, perhaps because the chief interest in mathematics shown by Desargues had been in its applications to the study of perspective, this masterpiece seems to have attracted little general attention, although appreciated by both Pascal and Descartes. At any rate the work was soon forgotten and remained almost unknown until Chasles happened to find a copy in 1845, since which time it has been looked upon as one of the classics in the early development of modern pure geometry. In this work he introduced the notions of the point at infinity, the line at infinity, the straight

[1] Born at Lyons, 1593; died at Lyons, 1662. N. G. Poudra, *Desargues . . . Œuvres*, 2 vols. Paris, 1864.

[2] *Brouillon projet d'une atteinte aux éuénemens des rencontres d'un cone avec un plan*, Paris, 1639; M. Chasles, *Aperçu*, 74; Poudra, *loc. cit.*, pp. 97, 303.

line as a circle of infinite radius, geometric involution, the tangent as a limiting case of a secant, the asymptote as a tangent at infinity, poles and polars, homology, and perspective, thus laying a substantial basis for the modern theory of projective geometry.



MARQUIS DE L'HOSPITAL

He had much to do with the introduction of Newton's mathematical ideas into France

L'Hospital. Guillaume François Antoine de l'Hospital,[1] Marquis de St.-Mesme, a man of ancient and honorable family, was one of the world's infant prodigies in mathematics. When only fifteen he was one day at the Duc de Roanne's and heard some mathematicians speaking of a difficult problem of Pascal's. To their surprise he said that he thought he could solve it, and in a few days succeeded. A career which he sought in the army proved impossible owing to his defective sight, and the latter part of his life was given to his favorite study. He was a pupil of Jean Bernoulli's and introduced the ideas of the new analysis into France.[2] He also wrote on geometry, algebra, and mechanics, most of his works being published after his death.

[1] Born in Paris, 1661; died in Paris, February 2, 1704. He is also known as the Marquis de l'Hospital. The family also spelled the name Lhospital and, somewhat later, l'Hôpital.

[2] *Analyse des infiniment petits*, Paris, 1696.

Frénicle de Bessy and De la Loubère. Among the correspondents of Fermat, Bernard Frénicle de Bessy,[1] an office-holder and a member of the Académie des Sciences at Paris, was known for his work on Pythagorean numbers, that is, numbers which form the sides of a right-angled triangle,[2] and for his interest in magic squares. At about the same time Antoine de la Loubère,[3] a Jesuit and a lecturer on mathematics, rhetoric, theology, and the humanistic subjects, was showing much interest in the study of curves. This interest is seen in his quadrature problem[4] and in his study of the cycloid.[5] His method of tangents, in which the tangent is taken as the direction of a moving point, was quite forgotten until its value was recognized in its applications in kinematics.[6]

Roberval. Among the contemporaries of De la Loubère, Gilles Persone de Roberval[7] became well known for his discoveries in the field of higher plane curves and for his method of drawing a tangent to a curve (already suggested in substance by Torricelli), which was a definite step in the invention of the calculus. He was professor of philosophy in the Collège Gervais at Paris, and later professor of mathematics in the Collège Royal. His chief interest was in physics, but he also wrote on the cycloid (his "trochoid") and other curves, on algebra and indivisibles,[8] and (1644) on the astronomy of Aristarchus.

[1] Born in Paris, *c.* 1602; died 1675.

[2] *Traité des triangles rectangles en nombres*, Paris, 1676 (posthumous).

[3] Born at Rieux, Languedoc, 1600; died at Toulouse, 1664. The name also appears as Laloubère, Lalouère, Lovera, Lalovera, Lalouvère.

[4] *Elementa tetragonismica seu demonstratio quadraturae circuli et hyperbolae ex datis ipsorum centris gravitatis*, Toulouse, 1651. See also Montucla, *Histoire*, II (2), 77.

[5] *Propositio 36ª excerpta ex quarto libro de cycloide nondum edito*, Toulouse, 1659; *Veterum geometria promota in septem de cycloide libris*, Toulouse, 1660.

[6] See Chasles, *Aperçu*, 58, 96; F. Jacoli, Boncompagni's *Bullettino*, VIII, 265.

[7] Born at Roberval, near Beauvais, August 8, 1602; died in Paris, October 27, 1675. The name Roberval, by which he is commonly known, was merely that of his birthplace. The family name, Persone, appears also in the Latin form of Personerius, whence a derived French form is Personier.

[8] *De geometrica planarum et cubicarum aequationum resolutione, De recognitione aequationum*, and *Traité des indivisibles*. These and other of his memoirs were collected, published in 1693, and republished in the *Mémoires de l'ancienne académie, Vol. VI.*

Other Writers of the Period. At about the same time Claude François Milliet Dechales,[1] for some time a Jesuit missionary in Turkey, taught in the schools of his order in Marseilles, Lyons, and Chambéry. He is chiefly known for his editions of Euclid's *Elements*[2] and for a general work on mathematics,[3] but his original contributions to the subject were slight.

Noted as a voluminous writer on theology, Antoine Arnauld (1612–1694)—"the great Arnauld" as the Jansenists called him, hated by the Jesuits and the Calvinists alike because of his bitter attacks upon their beliefs—deserves at least some slight mention for his encouragement of mathematics. He was interested in the works of his great contemporaries, such as Descartes and Pascal, wrote on geometry (1667) and magic squares, and showed interest in the theory of numbers.[4]

Among those who in the latter part of the century did much to make geometry popular was Philippe de Lahire,[5] a pupil of Desargues's and a man of scattered genius. He was at first a painter and architect, then a *pensionnaire astronome* of the Académie des Sciences at Paris, then professor of mathematics in the Collège Royal and the Académie de l'Architecture, and in his later years (from 1679) was connected with the geodetic survey of France. He wrote several works on conics, algebra,[6] and astronomy, besides contributing a large number of memoirs on mathematics, astronomy, and physics to the Académie des Sciences. He also wrote on epicycloids (1694) and roulettes (1694, 1706), and summarized what was then known on magic squares (1705).

[1] Born at Chambéry, Savoy, 1621; died at Turin, March 28, 1678. The name also appears as Deschales and De Challes.

[2] Latin ed., Lyons, 1660; French ed., Paris, 1677.

[3] *Cursus seu Mundus Mathematicus*, Lyons, 1674, with a later edition in 1690.

[4] K. Bopp, "Antoine Arnauld . . . als Mathematiker," *Abhandlungen*, XIV, 187.

[5] Born in Paris, March 18, 1640; died in Paris, April 21, 1718. The *Biographie Universelle* gives the date of his death as 1719, but most authorities give it as 1718. The name is often written La Hire. See Chasles, *Aperçu*, 118, 550, 553; Curtze, in *Bibl. Math.*, II (2), 65.

[6] *Théorie des coniques*, Paris, 1672; *Nouvelle Méthode de Géométrie*, Paris, 1673; *Nouveaux élémens des sections coniques*; *Les Lieux Géométriques*; *La Construction ou effection des équations*, Paris, 1679, with an English translation, London, 1704; *Sectiones conicae in novem libros distributae*, Paris, 1685.

Among the contemporaries of de Lahire, but a few years his junior and like him a *pensionnaire* of the Académie des Sciences, Michel Rolle[1] made for himself a worthy name. He was connected with the war department and apparently was not concerned with teaching. He wrote on both geometry[2] and algebra, his publications on geometry appearing in the form of numerous memoirs, and those on algebra in memoirs and in two books.[3] To him is due the theorem that $f'(x) = 0$ has at least one real root lying between two successive roots of $f(x) = 0$.[4]

As a representative of the other French mathematicians of this period there may be named Pierre Nicolas,[5] a pupil of De la Loubère's and rector of the Jesuit college at Béziers, who wrote on the logarithmic spiral and on conchoids.[6]

In the field of textbook making the most popular French writer of this time was François Barrême, a native of Lyons, who died in Paris in 1703. His *Arithmétique* (Paris, 1677) went through many editions, and his name is still a synonym for a ready reckoner (*barème*).

4. Great Britain

Great Britain in the Seventeenth Century. As already remarked, the two foci of the ellipse that bounded mathematical Europe in the 17th century were located in France and Great Britain. The British center was Cambridge, although the contributions of Edinburgh in the notable discoveries of Napier

[1] Born at Ambert, Auvergne, April 21, 1652; died in Paris, November 8, 1719. The date of his death is also given as July 5.

[2] A certain curve, $xy^2 = a\,(y - mx)^2$, has recently come to bear his name, but apparently with no justification.

[3] *Traité d'algèbre*, Paris, 1690; *Méthode pour résoudre les questions indéterminées de l'algèbre*, Paris, 1699. F. Cajori, "What is the origin of the name 'Rolle's Curve'?" *Amer. Math. Month.*, XXV, 291, and *Bibl. Math.*, XI (3), 300. See also *Bibl. Math.*, IV (3), 399, and *L'Intermédiaire des Mathématiciens*, V, 76, and XVI, 244 (Paris), hereafter referred to as *L'Intermédiaire*.

[4] *Démonstration d'une Methode pour résoudre les Egalitez de tous les degrez*, Paris, 1691.

[5] Born at Toulouse, *c.* 1663; died *c.* 1720.

[6] *De novis spiralibus exercitationes*, Toulouse, 1693; *De Lineis logarithmicis spiralibus hyperbolicis*, Toulouse, 1696; *De conchoidibus et cissoidibus*, Toulouse, 1697.

and Gregory, and of Oxford in the works of such men as Harriot, Briggs, Halley, Wren, and Wallis were such as to challenge the supremacy of the Cambridge school.

Harriot. It is rather surprising to think that the man who surveyed and mapped Virginia[1] was one of the founders of algebra as we know the science today. Such, however, is the case, for Thomas Harriot[2] was sent by Sir Walter Raleigh to accompany Sir Richard Grenville (1585) to the New World, where he made the survey of that portion of American territory. He returned to England (1587) and published (1588) a report upon the colony, and some years later wrote a work that helped to establish the English school of algebraists. Harriot took his B.A. at Oxford in 1579. After his return from America he was introduced to the Earl of Northumberland and was received with other scholars into his household. The earl allowed him a pension of £300 a year for the rest of his life. He was prominent as an astronomer and corresponded with Kepler. He discovered the solar spots, and his observations of the satellites of Jupiter were independent of those made by Galileo at the same time. His great work on algebra[3] was published ten years after his death. In this work he assisted in setting the standard for a textbook in algebra which has been generally recognized since that time. The work includes the formation of equations with given roots, the law as to the number of roots, the relation of the roots to coefficients, the transforming of equations into equations having roots differing from the original roots according to certain laws, and the solution of numerical equations. He used small consonants for the known quantities and small vowels for the unknown quantities.

[1] Or rather North Carolina, since the present boundaries had not yet been fixed. On his map of Virginia see P. L. Phillips, *Virginia Cartography*, Washington, 1896. His report was published in London in 1588. A second edition appeared in Hakluyt's *The Principall Navigations*, London, 1589. F. V. Morley, *The Scientific Monthly*, XIV, 60.

[2] Born at Oxford, 1560; died near Isleworth, July 2, 1621. The name appears as Hariot in several early works.

[3] *Artis Analyticae Praxis, ad Æquationes Algebraicas nouâ . . . Methodo resoluendas*, London, 1631. It was probably written *c.* 1610.

He also took some steps in the direction of analytic geometry. In the further matter of symbols, he used $\surd 3$ for the cube root,[1] and the characters $>$ and $<$ for "is greater than" and "is less than" respectively.

Napier. When Hume, the historian, wrote his appreciation of Napier as "the person to whom the title of 'great man' is more justly due than to any other whom his country has ever produced," he spoke without exaggeration. Burns can be appreciated only in the vernacular, Scott appeals chiefly to a single period in the development of the individual, the fame of those whose effigies justly have place along Princes Street in Edinburgh is generally only national; but Napier's remarkable invention affects the whole world with constantly increasing power. The artisan who carries in his pocket the slide rule is relatively as much indebted to the genius of the Laird of Merchiston as the astronomer, the engineer, the physicist, and the mathematician.



JOHN NAPIER
Engraved by Stewart after the original painting in Edinburgh

John Napier[2] was descended from a strong ancestry which included men who had held positions of prominence because they deserved to do so. Merchiston Castle, built in the 15th

[1] As in $\underbrace{\surd 3.)26 + \surd 675}$ for $\sqrt[3]{26 + \surd 675}$.

[2] Born at Merchiston Castle, now in the city of Edinburgh, 1550; died there, April 4, 1617. The name also appears as Naper, Naperus, Neper, and Neperius.

century, was one of the two strongholds on the outskirts of Edinburgh, and was enlarged from time to time until it became an imposing structure, symbolic of the Napier house.

Napier was born in the period of greatest strife between Protestantism and Catholicism in Scotland. John Knox began his mission only three years before Napier's birth, and the seeds of the bitter antagonism which the latter felt towards Rome were early planted in his soul. In 1563, when only thirteen years old, Napier was sent to the University of St. Andrews, but left without taking a degree. He probably studied abroad, but in 1571 was back again in Scotland, this time at Gartness, in Stirlingshire, where his father had some property. It was probably here that he wrote his popular theological work, *A Plaine Discouery of the whole Reuelation of Saint Iohn* (Edinburgh, 1593),[1] a bitter attack upon the Church of Rome. The common people accused him of dealing in the black art, while the intellectuals recognized him as a man of remarkable ingenuity; but the end was the same. He planned to use burning mirrors, like those of Archimedes but so potent that they should destroy an enemy's ships "at whatever appointed distance"; a piece of artillery that should, as an early writer described it, "clear a field of four miles circumference of all the living creatures exceeding a foot of height"; a chariot which should be like "a living mouth of mettle and scatter destruction on all sides"; and "devises of sayling under water," all of which is of interest in view of the engines of destruction used first in the World War of 1914–1918. It is of interest to compare the mind of Napier, as seen in his vision of future achievements, with that of Roger Bacon. Each seemed to many of his contemporaries, and perhaps to most of those who knew him, as mentally unbalanced and as a mere visionary in his contemplation of future warfare, and yet each prophesied with remarkable success respecting many inventions of the present time.

[1] Its great popularity is shown by the fact that it was reprinted in London in 1594, 1611, and 1641; in French translation at La Rochelle in 1602, with a fourth edition in 1607; in Dutch translation at Middelburgh in 1600; and in German translation at Frankfort a. M. in 1611.

Napier on Logarithms. Napier wrote two works on logarithms,[1] besides one on computing rods[2] and one on algebra.[3] Of all his works he probably thought the *Plaine Discouery of the whole Reuelation of Saint Iohn* the most important, but the world has long since forgotten it. The popular verdict of his day was that the *Rabdologia* was his greatest work, but it is now looked upon only as one of the curiosities of history. The scientific world looked upon his *Descriptio* as epoch-making, and the scientific world was right.[4]

Briggs. The man who did most to start the invention of Napier on its road to success was Henry Briggs,[5] whom Oughtred rather absurdly called the English Archimedes.[6] He entered St. John's College, Cambridge, in 1577, took the degrees of B.A. in 1581 and M.A. in 1585, and was made a fellow in 1588. He was the first professor of geometry at Gresham College, London (1596–1619), after which he became professor of astronomy at Oxford. He saw at once the great importance of Napier's invention, and in a letter to James Ussher, archbishop of Armagh, dated March 10, 1615, he speaks of himself as being "wholly employed about the noble invention

[1] *Mirifici Logarithmorum Canonis descriptio . . . Authore ac Inventore Ioanne Nepero, Barone Merchistonii, &c. Scoto*, Edinburgh, 1614, with other editions in 1616, 1619, and 1889; Leyden, 1620; London, 1616 and 1618; *Mirifici ipsius canonis constructio*, which appeared posthumously and was added to the edition of 1619 mentioned above. On all these editions and on the subject in general see C. G. Knott, *Napier Tercentenary Memorial Volume*, London, 1915.

[2] *Rabdologiae, seu nvmerationis per virgulas libri dvo*, Edinburgh, 1617, published the year of his death. For description, see Volume II. There was an edition at Leyden, 1626; an Italian translation, Verona, 1623, in which Napier is quoted as ascribing all glory and honor "alla Beatissima Vergine Maria," which is the last thing he would have dreamed of doing; and a German translation, Berlin, 1623. There was a free English translation of part of the work by John Dansie (*A Mathematicall Manuel*, London, 1627), the first English version to appear.

[3] Preserved in the Napier family and published in Edinburgh in 1839 under the title *De Arte Logistica Joannis Naperi Merchistonii Baronis Libri Qui Supersunt.*

[4] The subject is considered at length in Volume II.

[5] Born at Warley Wood, Yorkshire, February, 1560/61 (1561 N.S.); died at Oxford, January 26, 1630/31. The date of birth is often given as 1556, but the parish register shows that it was 1560/61.

[6] Aubrey, *Brief Lives*, I, 124.

of logarithms." In the following year (1616) he made a visit to Edinburgh for the purpose of meeting with Napier, and repeated his visit in 1617. It was on the first of these visits that Briggs suggested the base 10, of which Napier had already thought, this being the base of the common system of logarithms that has been in use ever since that time, and on his return to Oxford he prepared a table accordingly.

Briggs published ten works and left six others unpublished. The published works include treatises on navigation, Euclid's *Elements*,[1] logarithms,[2] and trigonometry.[3]

Gellibrand. Briggs's friend Henry Gellibrand,[4] who edited his trigonometry (1633), entered Trinity College, Oxford, in 1615, was granted the degrees of B. A. in 1619 and M. A. in 1623, took holy orders, and entered upon church work. Having heard one of Sir Henry Savile's lectures, he was so impressed that he gave up his curacy and devoted himself entirely to mathematics. Besides editing the trigonometry left unpublished by Briggs he also wrote on navigation and the variation of the magnetic needle, and composed a trigonometry of his own.[5]

Oughtred. One of the greatest of the writers of the early part of the 17th century in his influence upon English mathematics was William Oughtred.[6] As in the case of Harriot,

[1] *Elementorum Euclidis libri VI priores*, London, 1620.

[2] *Arithmetica logarithmica*, London, 1624. The final (French) edition appeared at Gouda in 1628.

[3] *Trigonometria Britannica, sive de doctrina triangulorum libri duo*, Gouda, 1633, published posthumously by his friend Henry Gellibrand.

[4] Born in the parish of St. Botolph, Aldersgate, London, November 17, 1597; died in London, February 16, 1636/37.

[5] *An Institution Trigonometricall wherein . . . is exhibited the doctrine of the dimension of plain and spherical triangles . . . by tables . . . of sines, tangents, secants, and logarithms*, London, 1638. This title is from the second edition (1652).

[6] Born at Eton, March 5, 1574; died at Albury, June 30, 1660. He also wrote his name Owtred, and John Locke preferred the form Outred, from which spellings we infer the pronunciation. On his life and works see F. Cajori, *William Oughtred*, Chicago, 1916, hereafter referred to as Cajori, *Oughtred*. The date of his birth is from Aubrey, *Brief Lives*,—"Gulielmus Oughtred natus 5 Martii 1574, 5 h. P. M."

he was not a professor of mathematics; but, like Harriot also, he knew more mathematics than most professors of his day. He entered King's College, Cambridge, in 1592, became a fellow in 1595, and received the degrees of B. A. in 1596 and M.A. in 1600. Speaking of his college work he says:

> The time which over and above those usuall studies I employed upon the Mathematicall sciences, I redeemed night by night from my naturall sleep, defrauding my body, and inuring it to watching, cold, and labour, while most others tooke their rest.

He vacated his fellowship in 1603 and in the following year began his ministry. He gave much of his time, however, to mathematics and to correspondence with mathematicians. Aubrey,[1] whose gossiping biographies are always entertaining, gives this description of him:

> He was a little man, had black haire, and blacke eies (with a great deal of spirit). His head was always working. He would drawe lines and diagrams on the dust . . . did use to lye a bed till eleaven or twelve a clock . . . Studyed late at night; went not to bed till 11 a clock; had his tinder box by him; and on the top of his bed-staffe, he had his inke-horne fix't. He slept but little. Sometimes he went not to bed in two or three nights.

He thus seems to have violated many of the usual canons of health, and probably continued to do so until his death at the ripe old age of eighty-six.

The Clavis Mathematicæ. Oughtred's best-known work is his *Clavis mathematicæ*,[2] a brief treatise on arithmetic and algebra, composed (*c.* 1628) for the purpose of instructing the son of the Earl of Arundel. It was published in London in 1631. He had already written (*c.* 1597) a treatise on dialing, but this was not published until 1647, when it appeared in an edition

[1] Aubrey, *Brief Lives*, II, 106.

[2] For the full title and for a study of the various editions see Cajori, *Oughtred*, p. 17; H. Bosmans, "La première édition de la 'Clavis Mathematica' d'Oughtred," *Annales de la Société scientifique de Bruxelles*, XXXV, 2me partie, p. 24.

of the *Clavis*.[1] The influence of the latter work was very great. In it appear contracted multiplication and division, the distinction between the two uses of the signs + and −, the symbol (: :) for proportion, and the symbols × for multiplication (already known) and ∽ for the absolute value of a difference.

The Slide Rule. The invention of the slide rule seems unquestionably due to Oughtred,[2] and there also seems to be good reason for believing that he is the author of the *Appendix to the Logarithmes* printed with the English translation of Napier's work (London, 1618) and containing the first natural logarithms. He also wrote a trigonometry[3] to which reference will be made later, and a work on gaging, and he translated and edited Leurechon's French work on mathematical recreations. Among his many pupils were John Wallis and Sir Christopher Wren, the former of whom wrote of the *Clavis* that it "doth in as little room deliver as much of the fundamental and useful part of geometry (as well as of arithmetic and algebra) as any book I know."

Gunter. Connected with the general movement to simplify calculation, initiated by Napier and Briggs, there stands out prominently the name of Edmund Gunter,[4] who left the ministry to become professor of astronomy (1619) at Gresham College, London. He published the first table of logarithmic sines and tangents to the common base,[5] suggested to Briggs the use of the arithmetic complement, invented the surveyors' table and the chain which until recently was commonly known as

[1] It was translated into Latin by Sir Christopher Wren. Another of his works on dialing (1600) was published in 1632.

[2] *The Circles of Proportion and The Horizontall Instrument*, London, 1632. See F. Cajori, *History of the Logarithmic Slide Rule*, New York, 1909, and "On the History of Gunter's Scale and the Slide Rule," *University of California Publications in Mathematics*, I, p. 187.

[3] *Trigonometrie, or, The manner of calculating the Sides and Angles of Triangles, by the Mathematical Canon, demonstrated*, London, 1657.

[4] Born in Hertfordshire, 1581; died in London, December 10, 1626.

[5] *Canon triangulorum or Table of Artificial Sines and Tangents*, London, 1620.

Gunter's chain, and devised a kind of slide rule known as Gunter's scale. Aubrey relates the following incident:

When he was a student at Christ Church, it fell to his lott to preach the Passion sermon, which some old divines that I knew did heare, but they sayd that 'twas sayd of him then in the University that our Saviour never suffered so much since his passion as in that sermon, it was such a lamentable one—*Non omnia possumus omnes.*[1]

The Savilian and Lucasian Professorships. Mention is so often made of the Savilian professorships of geometry and astronomy at Oxford and of the Lucasian professorship at Cambridge that a brief statement concerning them is desirable.

Sir Henry Savile,[2] the founder of two chairs at Oxford, was warden (1585) of Merton College, Oxford, and in 1596 was appointed provost at Eton. He lectured on Euclid,[3] and although he contributed little to mathematics, he contributed much to the extending of the knowledge of the science through the professorships which he founded in 1619, and which have been held by a distinguished line of scholars for three centuries.

Henry Lucas[4] was for a time a student at St. John's College, Cambridge, but seems not to have matriculated. He was, however, admitted M.A. in 1635–1636 and was elected to represent the university in parliament in 1639–1640. By his will he directed that lands yielding £100[5] a year be purchased to found the professorship that bears his name. It was in the year after the founding of the professorship that Barrow was elected (1664) as the first occupant of the chair, and six years later he was succeeded by Newton (1670). In his inaugural address Barrow speaks of Lucas as "a new and benignant star, shining with a ray both true and propitious, such as has not for many years risen above the academical horizon."[6]

[1] *Brief Lives*, I, 276.

[2] Born at Over-Bradley, near Halifax, Yorkshire, November 30, 1549; died at Eton, February 19, 1622.

[3] *Praelectiones XIII in principium elementorum Euclidis, Oxoniae habitae 1620*, Oxford, 1621.

[4] Died in London, June 22, 1663.

[5] In 1860 it had risen only to £155.

[6] Whewell's translation. For Kirby's translation see the English edition of the *Mathematical Lectures*, p. vii (London, 1734).

Barrow. To Isaac Barrow[1] it was given to rank as one of the best Greek scholars of his day, to attain the highest honors as professor of mathematics, to be looked upon as one of the leading theologians of England, to be recognized as a profound student of physics and astronomy, to acquire fame as a preacher and controversialist, and to be perhaps the first to recognize the genius of Newton and to develop his great talents, but his reputation might have been higher had he worked exclusively in only one of his fields of interest, particularly in optics or geometry. He was prepared for the university at Charterhouse, London, and at Felstead School, in Essex. He entered Trinity College, Cambridge, in 1644, took his degree of B. A. in 1648 (when only eighteen), was elected fellow the year following, and left in 1655. His ready wit is illustrated by an incident that occurred in his examination for holy orders, the dialogue being said to have run as follows:

Chaplain. *Quid est fides?*
Barrow. *Quod non vides.*
Chaplain. *Quid est spes?*
Barrow. *Magna res.*
Chaplain. *Quid est caritas?*
Barrow. *Magna raritas.*

The chaplain is then said to have given up in despair and to have reported the candidate's lack of reverence to the bishop. Fortunately the latter had a sense of humor and Barrow was duly admitted.[2]

Barrow traveled for some time after taking orders and in 1662 was elected professor of geometry at Gresham College. Soon after, as already stated, he was elected (1664) the first Lucasian professor. Six years later (1670)[3] he resigned in favor of Newton and devoted his attention to theology. He

[1] Born in London, October, 1630; died in London, May 4, 1677.

[2] W. W. R. Ball, *Cambridge Papers*, p. 109. London, 1918.

[3] These two dates are often given as 1663 and 1669, on account of the style of calendar. His inaugural lecture was given March 14, 1664. See W. Whewell, *The Mathematical Works of Isaac Barrow*, p. 5 (Cambridge, 1860).



BARROW

Statue in Trinity College Chapel, Cambridge. It stands just north of the statue of Newton

became chaplain to Charles II in 1670, master of Trinity College in 1673,[1] and vice-chancellor of the university in 1675. He edited Euclid's *Elements*[2] and the *Data*,[3] the works of Archimedes,[4] the conics of Apollonius, and the spherics of Theodosius. His own contributions to science appear chiefly in his general lectures[5] and in his works on optics[6] and geometry.[7] He also gave a new method for determining tangents,—one that approached the methods of the calculus.[8] In this work he made use of a "differential triangle," which is still essentially the basis of the initial work in differentiation and which we shall consider at some length in Volume II.

Newton. Isaac Newton[9] was born in the stirring times of the Cromwell rebellion and was the posthumous son of a farmer in Lincolnshire. He was a small and feeble child, and as a boy he gave little promise of success in the battle of life. At the age of twelve he was taken from the local day school and placed in a school at Grantham. According to his own statement he was at first extremely inattentive to his studies and ranked among the lowest in the school. His chief interests seemed to be in carpentering, mechanics, the writing of verses, and drawing. Later, however, he began to show considerable

[1] W. W. R. Ball, *Cambridge Papers*, p. 171.

[2] *Euclidis Elementorum Libri XV*, Cambridge, 1655, with many later editions.

[3] *Euclidis Data*, Cambridge, 1657.

[4] The Archimedes, Apollonius, and Theodosius were published together at London in 1675.

[5] *Lectiones Mathematicae XXIII*, London, 1683 (posthumous).

[6] *Lectiones XVIII, . . . in qvibvs Opticorum Phaenomenωn . . investigantur . . .*, London, 1669.

[7] *Lectiones Geometricae*, London, 1670. There is a translation with commentary by J. M. Child, Chicago, 1916.

[8] For a good summary, see Ball, *Hist. Math.*, 6th ed., p. 311.

[9] Born at Woolsthorpe, near Grantham, Lincolnshire, December 25, 1642; died at Kensington, March 20, 1727. One of the best books with which to begin a study of Newton's life is A. De Morgan, *Essays on the Life and Work of Newton*, 2d ed., Chicago, 1914, not so much because of the essays themselves as for the bibliographical and critical notes of the editor, P. E. B. Jourdain. The best biography of Newton is that of Sir David Brewster, *The Memoirs of Newton*, 2d ed., Edinburgh, 1860, hereafter referred to as Brewster, *Memoirs of Newton*. A new biography of Newton and a definitive edition of his complete works are both sadly needed.

Newton's work was so highly appreciated that about this time (1668) he was invited to revise the lectures of Barrow, his former teacher. In 1669 the latter resigned the Lucasian professorship of mathematics at Cambridge, and Newton was appointed his successor. Thus at the age of twenty-seven he had begun his great work on the calculus and mathematical physics, and held already one of the highest academic honors in the world.

For nearly sixty years after receiving his professorship he was looked upon as one of the greatest leaders in the fields of physics and mathematics, receiving all the honors that could be hoped for by a man in academic life. He was elected fellow of the Royal Society in 1672, was chosen in 1689 to represent the university in parliament, and was appointed warden of the mint in 1696 and master in 1699. He was again elected to parliament in 1701, but he took no interest in politics. He was



ROETTIERS'S MEDAL OF NEWTON

Of the numerous portrait medals of Newton, those by Croker and by Roettiers (first struck in 1730) are the best. This particular medal was struck in 1774. The reverse bears the motto from Vergil's Æneid (Lib. V, 378), "Another is sought for him." Its significance appears from the context: "Another is sought for him, nor does any one from so great a band dare approach the man, and draw the gauntlets on his hands"

knighted by Queen Anne in 1705. The latter part of his life was spent in London and was mathematically unproductive.

Newton always hesitated to publish his discoveries. His greatest work, the *Principia*,[1] was begun in 1685 but was not published until 1687, and then only under pressure from his friend Halley, the astronomer. In this work he sets forth his theory of gravitation, "indisputably and incomparably the greatest scientific discovery ever made,"[2] following the methods of the Greek geometry as being more easily understood by students of his time.[3] It was the third of the three great discoveries in the field of mathematical astronomy,— the heliocentric theory, finally established by Copernicus; the elliptic orbits of the planets and the laws relating thereto, finally established by Kepler; and the law of universal gravitation, with which Newton's name will always be connected.

Referring to this great work, there is humor as well as justice in the remark of Lagrange to the effect that Newton was the greatest genius that ever lived, and the most fortunate, since we can find only once a system of the universe to be established. Whether Einstein's Theory of Relativity invalidates this statement remains to be seen.

Newton's *Arithmetica Universalis*, a work on algebra and the theory of equations, was written in lecture form in the period 1673–1683 but was not published until 1707.[4] He wrote a work on analysis by series[5] in 1669, but it was not published until 1711. His work on the quadrature of curves[6] was written in 1671 but was not published until 1704, and his other

[1] *Philosophiae Naturalis Principia Mathematica*, London, 1687, reissued the same year with a new title-page; 2d ed. by Roger Cotes, Cambridge, 1713; 3d ed. by Pemberton, London, 1726. For a list of editions of all of Newton's works see G. J. Gray, *A bibliography of the Works of Sir Isaac Newton*, Cambridge, 1888; 2d ed., Cambridge, 1907.


[2] W. Whewell, *History of the Inductive Sciences*, Bk. VII, ii, § 5.

[3] For a résumé of the work see W. W. R. Ball, *Essay on the Genesis, Contents, and History of Newton's Principia*, London, 1893.

[4] *Arithmetica universalis sive de compositione et resolutione arithmetica liber*, Cambridge, 1707; English translation, London, 1720.

[5] *Analysis per Æquationes numero terminorum infinitas*, London, 1711.

[6] *Tractatus de quadratura curvarum*, printed in the first edition of his *Opticks*, London, 1704.



Tonnage, 1694, &c.

Record' Vic. 4 Apr 1720.

Numb. 168. £ 50...

The 13 Day of April 1720.

Received by me Sr Isaac Newton Kt.

by Virtue of the Order dated the

28th April 1694

To Orelow

Richard Hampden Esq. One of the Four Chiefs of the Receipt of Exchequer, the Sum of

Fifty pounds £ 50.-

In full of all former Directions of the said Order, and

Months Annuity, due at

Lady last, of One hundred Pounds per Annum, by Vertue of an Act of Parliament, (Entituled, An Act for Granting to Their Majesties certain Rates and Duties upon Tonnage of Ships and Vessels, and upon Beer, Ale, and other Liquors, for securing certain Recompences and Advantages in the said Act mentioned, to such Persons as shall voluntarily advance the Sum of fifteen hundred thousand Pounds towards carrying on a Vigorous War against France;) I say Received by me—

Isaac Newton

AUTOGRAPH OF NEWTON

The document was written after he had retired from his work at Cambridge and is interesting politically

works were similarly held from the reading public until after Newton had given up his lectures at Cambridge. His *Fluxions*,[1] treating of the subject by which he is probably the best known, did not appear until nine years after his death, although the theory had by that time become well known through the publication of such works as that of Charles Hayes (1704).

It was a noble and a generous tribute that Leibniz paid when he said that, taking mathematics from the beginning of the world to the time when Newton lived, what he did was much the better half.

Newton had the eccentricities of genius, always being so absorbed in his work as to be oblivious of life about him. Many stories are related of his absent-mindedness, some of them also told of other mathematicians and very likely most of them apocryphal.

Of these stories one relates that, when giving a dinner to some friends, he left the table to fetch a bottle of wine; but on his way to the cellar he forgot all about the errand, went to his room, put on his surplice, and ended up in chapel. It is also related that once in riding he dismounted and started in his absent-minded way to lead his horse up the hill, but found when he came to remount that the horse had wandered off, leaving only the bridle in Newton's hand.

Newton died in 1727 and was buried in Westminster Abbey, where his tomb is still seen.

Voltaire attended the funeral, and we shall see later that when he was very old he did much to make Newton's philosophy known in France. It is said that "his eye would grow bright and his cheek flush" when he said that he had once lived in a land where "a professor of mathematics, only because he was great in his vocation," had been buried "like a king who had done good to his subjects."[2]

[1] *The Method of Fluxions and Infinite Series . . . from the Author's Latin Original not yet made publick . . .*, London, 1736, translated and edited by John Colson. There was another translation published shortly after this, London, 1737.

[2] S. G. Tallentyre, *Life of Voltaire*, p. 57. New York, n. d.

Halley. In his youth Edmund Halley[1] attended St. Paul's School, London, distinguishing himself in both mathematics and the classics. He entered Queen's College, Oxford, in 1673, and before he was twenty he communicated a paper to the Royal Society. So noteworthy had been his progress that in the very month in which he reached his twentieth birthday (November, 1676) he set out for St. Helena for the purpose of making astronomical observations. On the day before he was twenty-one he made the first complete observation of a transit of Mercury. So remarkable was his work at St. Helena that Flamsteed[2] called him the "southern Tycho" and the Royal Society elected him to a fellowship when he was only twenty-two (1678). Although his supposed materialistic views prevented his election as Savilian professor of astronomy at Oxford in 1691, he followed Wallis as Savilian professor of geometry in 1703. He succeeded Flamsteed as astronomer royal in 1721. He was the first to predict the time of the return of a comet, and the comet of which he announced the period has since then been known by his name.



EDMUND HALLEY
Friend of Newton, illustrious as an astronomer

[1] Born in London, November 8 (October 29, O.S.), 1656; died at Greenwich, January 14, 1742.

[2] John Flamsteed (1646–1719), first astronomer royal of England.

While Halley's tastes were chiefly astronomical, he was deeply interested in geometry,[1] algebra,[2] and the construction of logarithmic tables.[3] He solved the problem of the construction of a conic, given a focus and three points.

It was due to Halley's insistence as well as his financial help that Newton published his *Principia* in 1687. Besides editing the conics of Apollonius (1710) he edited the works of Serenus[4] and Menelaus.[5] He also compiled a set of mortality tables, thus giving a practical basis to the subject of life insurance.

Wallis. Of the contemporaries of Newton one of the most prominent was John Wallis.[6] He studied theology at Emmanuel College, Cambridge, and took the degree of B.A. in 1637 and that of M.A. in 1640, the year in which he was ordained. He became a fellow of Queens' College in 1644. His tastes, however, were in the line of physics and mathematics, and in 1649 he was elected to the Savilian professorship of geometry at Oxford, a position which he held until his death. He was awarded the degree of doctor of divinity in 1653,[7] became chaplain to Charles II in 1660, and was one of the founders of the Royal Society (1663).

Wallis was a voluminous writer, and not only are his writings erudite, but they show a genius in mathematics that would have appeared the more conspicuous had his work not been so overshadowed by that of his great Cambridge

[1] *Apollonii Pergaei Conicorum Libri octo*, Oxford, 1710. His version from the Arabic of the treatise of Apollonius *De Sectione Rationis*, with a restoration of the two lost books *De Sectione Spatii*, appeared at Oxford in 1706.

[2] *A new and exact Method of finding the Roots of any Equation*, London, 1720; "De numero radicum in aequationibus solidis ac biquadraticis," *Phil. Trans.*, 1687; "Methodus nova, accurata et facilis inveniendi radices aequationum," *Phil. Trans.*, 1694. See also the abridgment of the *Phil. Trans.*, 4th ed., I, 63, 68, 81, 137 (London, 1731).

[3] "A most compendious and facile method for constructing the logarithms . . .," *Phil. Trans.*, 1695.

[4] *De Sectione Cylindri et Coni*, Oxford, 1710.

[5] *Menelai Sphaericorum Libri III*, published by Dr. Costard, Oxford, 1758.

[6] Born at Ashford, Kent, November 23, 1616; died at Oxford, October 28, 1703.

[7] According to the *Dict. Nat. Biog.* Older authorities give 1654.



JOHN WALLIS, 1616–1703

From D. Loggan's drawing from life. Published in the *Opera Mathematica*, Oxford, 1695

contemporary. He was one of the first to recognize the significance of the generalization of exponents to include negative and fractional as well as positive and integral numbers. He recognized also the importance of Cavalieri's method of indivisibles, and employed it[1] in the quadrature of such curves as $y = x^n$, $y = x^{\frac{1}{n}}$, and $y = x^0 + x^1 + x^2 + \cdots$. He failed in his efforts at the approximate quadrature of the circle by means of series because he was not in possession of the general form of the binomial theorem. He reached the result, however, by another method. He also obtained the equivalent of $ds = dx\sqrt{1 + \left(\frac{dy}{dx}\right)^2}$ for the length of an element of a curve, thus connecting the problem of rectification with that of quadrature. In 1673 he wrote his great work *De Algebra Tractatus; Historicus & Practicus*, of which an English edition appeared in 1685.[2] In this there is seen the first serious attempt in England to write on the history of mathematics, and the result shows a wide range of reading of the classical literature of the science. This work is also noteworthy because it contains the first record of an effort to represent the imaginary number graphically by the method now used. The effort stopped short of success but was an ingenious beginning. Wallis was in sympathy with the Greek mathematics and astronomy, editing parts of the works of Archimedes, Eutocius, Ptolemy, and Aristarchus; but at the same time he recognized the fact that the analytic method was to replace the synthetic, as when he defined a conic as a curve of the second degree instead of a section of a cone, and treated it by the aid of coordinates. His writings include works on mechanics, sound, astronomy, the tides, the laws of motion, the Torricellian tube, botany, physiology, music, the calendar (in opposition to the Gregorian reform), geology, and the compass,—a range too wide to allow of the greatest success in any of the

[1] *Arithmetica Infinitorum, sive Nova Methodus Inquirendi in Curvilineorum Quadraturam, aliaque difficiliora Matheseos Problemata*, Oxford, 1655.

[2] The Latin edition is in Volume II of his *Opera Mathematica*, Oxford, 1693–1695.

lines of his activity. He was also an ingenious cryptologist and assisted the government in deciphering diplomatic messages.[1]

Among his interesting discoveries was the relation

$$\frac{4}{\pi} = \frac{3 \cdot 3 \cdot 5 \cdot 5 \cdot 7 \cdot 7 \cdots}{2 \cdot 4 \cdot 4 \cdot 6 \cdot 6 \cdot 8 \cdots},$$

one of the early values of π involving infinite products.[2]

Gregory. Scotland seems frequently to attempt to conceal her great men by a kind of protective coloring. Probably it is because she does not wish to seem to boast of her success. She had in the 17th century, and still has, a distinctive school of mathematical teaching, but she seems reluctant to proclaim the fact. Occasionally, however, she fails to keep her scholars and writers under cover, and so in mathematics we have such names as Napier, Gregory, and Kelvin.

James Gregory[3] was one of the first Scotchmen to make for himself a great name both in mathematics and in physics. He lived for some years in Italy, but in 1668 returned to Scotland to assume the professorship of mathematics at St. Andrews. In 1674 he became professor of mathematics at Edinburgh, where he died a year later. Like most of the leading British mathematicians of this period, he was equally interested in mathematics and physics. In 1661 he invented but did not practically construct the reflecting telescope which bears his name.[4] He also originated the photometric method of estimating the distances of stars. In the field of pure mathematics he expanded $\tan^{-1}\theta$, $\tan\theta$, and $\sec^{-1}\theta$ in series (1667), distinguished between convergent and divergent series, calculated the areas of successive polygons as the number of sides

[1] D. E. Smith, in the *Bulletin of the Amer. Math. Soc.*, XXIV (2), p. 82.

[2] See his *Opera Mathematica*, I, 441, where he explains that he uses the symbol $\square$ for the ratio of the square on the diameter to the area of the circle. In our symbolism this is $4r^2 : \pi r^2$, or $4 : \pi$. On page 469 he gives the value stated above. See also his Volume II, cap. lxxxiv.


[3] Born at Drumoak, near Aberdeen, November, 1638; died at Edinburgh, October, 1675.

[4] Described in his *Optica promota, seu abdita radiorum reflexorum et refractorum mysteria geometrice enucleata*, London, 1663.

was doubled, and gave an ingenious but unsatisfactory demonstration of the incommensurability of π.[1] The series

$$\theta = \tan\theta - \tfrac{1}{3}\tan^3\theta + \tfrac{1}{5}\tan^5\theta - \cdots$$

is commonly known by his name. He died in the thirty-seventh year of his age, shortly after being stricken with blindness as a result of the strain upon his eyes in carrying on his astronomical observations.



AUTOGRAPH OF LORD BROUNCKER

Part of a document dated December 31, 1661, just before Brouncker's appointment as Chancellor of Queen Catherine

Other British Writers. First among the other British writers of the period made memorable by the great work of Barrow, Wallis, and Newton, was William, Viscount Brouncker.[2] He was one of the founders and was the first president of the Royal Society, a linguist with the genius of a mathematician,

[1] *Vera circuli et hyperbolae quadratura*, Padua, 1667, and again in 1668. *Geometriae pars universalis*, Venice, 1667; "De circuli quadratura," in the *Phil. Trans.*, London, 1668. See also G. Heinrich, *Bibl. Math.*, II (3), 77.

[2] Born at Castle Lyons, Ireland, *c.* 1620; died at Westminster, April 5, 1684. The name is also given as Brounker. He was the second viscount.

and a man who was justly esteemed by the mathematical world of his day. He was interested in the rectification of the parabola[1] and the cycloid[2] and in the quadrature of the hyperbola[3] (1668) and the circle. The last of these investigations led him to state that

$$\frac{4}{\pi} = \cfrac{1}{1+\cfrac{1^2}{2+\cfrac{3^2}{2+\cfrac{5^2}{2+}}}} \cdots,$$

a form which he derived from the continued product which Wallis had discovered. This was the first use of continued fractions by an English writer. In 1662 he was appointed Chancellor of Queen Catherine, and in 1681 he became Master of St. Catherine's Hospital in London. Aubrey, who knew him, relates that "he was of no university, he told me. He addicted himselfe only to the study of the mathematicks, and was a very great artist in that learning."[4]

John Pell,[5] who was about ten years older than Brouncker, was admitted to Trinity College, Cambridge, at the age of thirteen, and also studied at Oxford. At the age of twenty he had mastered Latin, Greek, Hebrew, Arabic, Italian, French, Dutch, and Spanish. He became a professor of mathematics at Amsterdam (1643–1646) and at Breda (1646–1652), was Cromwell's representative in Switzerland (1654–1658), entered the ministry (1661), was given the degree of D.D. at Lambeth (1663), and in the same year was made a Fellow of the Royal Society. He spent the latter part of his life in parish work and in writing on mathematics.

He wrote on the quadrant (1630), on other matters relating to astronomy, on the magnetic needle, and on the quadrature problem,[6] and computed a table of 10,000 square numbers

[1] "On the Proportion of a curved line of a paraboloid to a straight line," *Phil. Trans.*, III, 645.

[2] "Of the Finding of a straight Line equal to that of a cycloid," *Phil. Trans.*, VIII, 640.

[3] "An algebraical paper on the squaring of the hyperbola," *Phil. Trans.* abridgment, 4th ed., I, 10. London, 1731.

[4] Oxford ed., 1898, I, 128.

[5] Born at Southwick, Sussex, March 1, 1611; died in London, December 12, 1685.

[6] *Refutation of Longomontanus's Pretended Quadrature of the Circle*, 1646; in Latin, 1647.

(1672). He edited (1668) the algebra of Rhonius[1] (1622–1676), which had been translated by Thomas Branker[2] and which contains the first mention of the Anglo-American sign (÷) of division. Through an error on the part of Euler, Pell's name is commonly connected with a certain equation of the form

$$x^2 - Ay^2 = 1,$$

although he had but little to do with it.[3]

One of the most promising pupils of Wallis was William Neile,[4] who gave (1660) the rectification of the semicubical parabola $y^3 = ax^2$. He died too soon, however, to fulfill his early promise.

If it had not been for the great fire of London (1666), Sir Christopher Wren[5] would have been known as a mathematician rather than as the architect of St. Paul's Cathedral. He was educated at Westminster School and in 1649 or 1650 was entered at Wadham College, Oxford. He was graduated B.A. in 1650–1651 and M.A. in 1653. In 1653 he was elected fellow of All Souls College, Oxford, where he resided until 1657. From 1657 to 1660 he was professor of astronomy at Gresham College, London, and from 1661 to 1673 was Savilian professor of astronomy at Oxford. He was a Fellow of the Royal Society and was its president from 1680 to 1682. He wrote on "The law of nature in the collision of bodies,"[6] on the grinding of hyperbolic mirrors, on perspective, and on the rectification

[1] J. H. Rahn, *An Introduction to Algebra,* London, 1668. The book was first published in Zürich in 1659, in German.

[2] Or Brancker. He was a clergyman, born in Devonshire, 1636; died at Macclesfield, 1676. He also taught at Macclesfield. He wrote a work of no merit on astronomy.

[3] E. E. Whitford, *The Pell Equation,* New York, 1912, with a good bibliography; H. Konen, *Geschichte der Gleichung* $t^2 - Du^2 = 1$, Leipzig, 1901; G. Enestrom, "Ueber den Ursprung der Benennung Pell'sche Gleichung," *Bibl. Math*, III (3), 204.

[4] Born at Bishopsthorpe, December 7, 1637; died in Berkshire, August 24, 1670.

[5] Born at East Knoyle, Wiltshire, October 20, 1632; died in London, February 25, 1723.

[6] *Phil. Trans.*, 1669, but originally separately printed in Latin as *Lex naturae de collisione corporum.*

problem, and he discovered (1669) the two systems of generating lines on a hyperboloid of one sheet. After the great fire he took a prominent part in the rebuilding of St. Paul's Cathedral and more than fifty other churches and public buildings in London. His noble epitaph in the cathedral is well known.[1]

Among the other pupils of Wallis who acquired some reputation was John Caswell.[2] He matriculated at Wadham College, Oxford, at the age of sixteen. Six years later he began teaching mathematics in Oxford, and so marked was his success that in 1709 he was elected Savilian professor of astronomy. He published a trigonometry in 1685 and at one time thought of publishing the work of Menelaus on spherics, but gave up the plan.[3]

James Gregory's nephew, David Gregory,[4] was professor of mathematics at Edinburgh from 1684 to 1691, after which he became Savilian professor of astronomy at Oxford. He published a work on geometry[5] when he was only twenty-three years of age, and in a work on optics[6] set forth the possibility of achromatic lenses. He also wrote on the Newtonian theory.[7] In 1703 he brought out at Oxford an elaborate edition of Euclid's works.

Textbook Writers. In the 17th century we reach a time in the development of elementary mathematics when textbooks became so standardized and numerous as to require in this connection the mention of only the most important. Their mission henceforth was to improve the method of presenting theories already largely developed and to adapt the applications of these theories to the needs of the world. From that time on they ceased to be a great factor in the presentation of mathematical discoveries.

[1] "*Lector, si monumentum requiris, circumspice.*"

[2] Born at Crewkherne, Somerset, 1655; died April, 1712.

[3] L. C. Karpinski, *Bibl. Math.* XIII (3), 248.

[4] Born at Kinnairdie, Banffshire, June 24, 1661; died at Maidenhead, Berkshire, October 10, 1708.

[5] *Exercitatio Geometrica de dimensione figurarum*, Edinburgh, 1684.

[6] *Catoptricae et Dioptricae Sphaericae Elementa*, Oxford, 1695.

[7] *Astronomiae Physicae et Geometriae Elementa*, Oxford, 1702.

The most prominent British textbook writer on elementary arithmetic in the 17th century was Edmund Wingate.[1] This is somewhat strange, because he entered the profession of the law after leaving Oxford,[2] went to Paris in 1624, where he taught English to the princess Henriette-Marie, future wife of Charles I, and returned to England in 1650 and became member of parliament for Bedford, none of which activities related to mathematics or even to textbooks. His tastes, however, seem to have led him to look upon mathematics as an avocation, for in the same year that he went to Paris he published a work on Gunter's scale,[3] two years later a work on logarithmic arithmetic,[4] and in 1630 his popular work *Of natural and artificial Arithmetick* (London), a work that went through many editions and was popular for more than a century. He also prepared a set of logarithmic tables[5] and a work[6] on a mathematical instrument which he had invented.

Of the textbook makers of this period in the domain of elementary algebra the best known was John Kersey,[7] a self-made teacher. He was highly esteemed in London as an instructor in mathematics and was a friend of Wingate's. His algebra[8] presents the subject in a logical and teachable manner. One of the most interesting features is the preliminary explanation of the analogies between proportion, which then held a high place, and the modern treatment of equations. The work was, however, altogether too elaborate to meet with great success.

William Leybourn (1626–*c.* 1700) was well known in London in his day, not only as a writer of textbooks but as a

[1] Born at Flamborough, Yorkshire, 1596; died in London, December 13, 1656.

[2] He edited the second edition of Britton's famous *Collections* on English law, London, 1640.

[3] *Construction, description et usage de la règle de proportion*, Paris, 1624.

[4] *Arithmétique logarithmique*, Paris, 1626; English edition, London, 1635.

[5] *Tables of the Logarithms of the sines and tangents*, London, 1633.

[6] *Ludus Mathematicus*, London, 1654.

[7] Born probably at Bodicote, near Banbury, Oxfordshire, 1616 (at any rate he was baptized there November 23, 1616); died May, 1677. See the *Bibl. Math.* XII (3), 263, with evidence for this date.

[8] *The Elements of that Mathematical Art commonly called Algebra*, London, 1673.

teacher of mathematics and a surveyor. He wrote on astronomy,[1] surveying,[2] arithmetic,[3] the logarithmic rule,[4] the Napier computing rods,[5] and mathematical recreations.[6] In 1690 he published a *Cursus Mathematicus* containing the substance of his other works. Three years later he published the best known of his works, a ready reckoner,[7] the first elaborate book of the kind to appear in English.

Leybourn was not, however, as popular as Edward Cocker,[8] who is described in 1657 as living "on the south side of St. Paul's Churchyard, over against St. Paul's Chain . . . where he taught the art of writing and arithmetick in an extraordinary manner." He was also a bibliophile, and in his "public school for writing and arithmetic" where he "takes in boarders" he had a large library of manuscripts and printed books in various languages. He wrote three or four books on arithmetic and penmanship, but it was his *Arithmetick, being a Plain and Easy Method*, edited by John Hawkins in 1678, that fixed the expression "according to Cocker" in the common speech of England. This remarkable book went through upwards of a hundred editions and had great influence upon British textbooks for more than a century. He also wrote on algebra,[9] but there has always been a question as to how much of this work

[1] *Urania Practica*, London, 1648, written with Vincent Wing. One of the first books on astronomy written in English, but not the first, Recorde's *Castle of Knowledge* having appeared in 1551.

[2] *Planometria, or the Whole Art of Surveying of Land . . . by Oliver Wallinby*, London, 1650, the pseudonym being formed by transposing the letters of his name, with minor changes.

[3] *Arithmetick, Vulgar, Decimal, and Instrumental*, London, 1657.

[4] *The Line of Proportion or Numbers, commonly called Gunter's Line, made easie*, London, 1667.

[5] *The Art of Numbering by Speaking-Rods: Vulgarly termed Nepeirs Bones*, London, 1667 and 1685. Leybourn, as will be shown in Volume II, mistakes the etymology of *Rabdologia*, and translates the word as "Speaking-Rods."

[6] *Pleasure with Profit; consisting of Recreations of divers kinds, Numerical, Geometrical, Mechanical, Statical, Astronomical*, London, 1694.

[7] *Panarithmologia, being a Mirror Breviate, Treasure Mate for Merchants, Bankers, Tradesmen, Mechaniks, . . .* London, 1693.

[8] 1631–1675, probably a descendant of the Northamptonshire family of Cokers.

[9] *Algebraical Arithmetic, or Equations*, London, 1684.

as well as the arithmetic was due to Cocker and how much to Hawkins. The world is as well off either way, the important thing being that in them we have two books that represent the popular view of the elementary science at the period in which they were written.

5. GERMANY

Germany in the Seventeenth Century. In the 16th century Germany moved toward the front in the mathematical progress of Europe, but with Italy always in the lead. In the 17th century she fell behind, producing only a single mathematician of the first class. The first part of the century was the period of the Thirty Years' War (1618–1648), and indeed the whole period was one of unrest in the Teutonic countries,—a period quite unsuited to intellectual progress.

Kepler. Although known chiefly for his work in the domain of astronomy, Johann Keppler,[1] or Kepler, as the name more often appears in English, ranked high as a mathematician. He studied in the cloister school at Maulbronn and at the University of Tübingen. At the age of twenty-two he began teaching mathematics and moral philosophy in the Gymnasium at Gratz in Steyermark.[2] For two years (1599–1601) he was an assistant of Tycho Brahe, and in 1601 became court astronomer to Kaiser Rudolph II and later to his successors Matthias (1612) and Ferdinand III (1615). His life was, however, made almost unendurable through domestic infelicity, troubles at court, and financial difficulties. In his work in mathematics he shows himself an excellent geometer and was also much interested in algebra. He set forth (1604) the idea of continuity in elementary geometry,[3] made (1615) some advance in the use of the infinitesimal,[4] and did much to further the cause of logarithms.[5]

[1] Born at Weil der Stadt, Württemberg, December 27, 1571; died at Regensburg, November 15, 1630. From the family of von Kappel.

[2] Or Graz, in the former Austrian crownland of Styria.

[3] In his *Ad Vitellionem Paralipomena.* See the Frisch edition of his *Opera Omnia,* II, 119, 187 (Frankfort, 1858–1871).

[4] In his *Stereometria*; e. g., see the Frisch edition of his *Opera Omnia*, IV, 583.

[5] His *Chilias Logarithmorum* appeared at Marburg in 1624, with a *Supplementum* in 1625.

Tschirnhausen. Ehrenfried Walther, Graf von Tschirnhausen,[1] often known as Tschirnhaus, is one of the few minor German mathematicians of the time to require special mention. He served for a year or two (1672–1673) with the Dutch army, traveled extensively, and then settled down on his estates, devoting his time largely to mathematics and physics. He is known chiefly for his study of curves,[2] including caustics and catacaustics,[3] and for his work in maxima and minima[4] and the theory of equations.[5]

Leibniz. Gottfried Wilhelm, Freiherr von Leibniz,[6] was the only pure mathematician of the first class produced by Germany during the 17th century. He early showed great proficiency in mathematics, having read the most important treatises on the subject before he was twenty. He studied law at Nürnberg, entered upon a diplomatic career, traveled extensively, and made the acquaintance of leading mathematicians in Holland, France, and England. He finally went to Hannover and became librarian to the duke. He lived in good style, and the visitor to Hannover today may see his palatial house, now used as a museum.

The leisure which his office allowed him gave Leibniz the opportunity to develop the differential and integral calculus. He seems to have begun to think about the subject in 1673,[7]

[1] Born at Kiesslingswalde, near Görlitz, April 10, 1651; died at Dresden, October 11, 1708. H. Weissenborn, *Lebensbeschreibung des Ehrenfried Walther von Tschirnhaus*, Eisenach, 1866.

[2] "Nova methodus tangentes curvarum expedite determinandi," *Acta Eruditorum*, I (1682), and various other memoirs in the same publication.

[3] All published in the *Acta Eruditorum*. The terms are due to Jacques and Jean Bernoulli. See Jacques Bernoulli, *Opera*, I, 466, *et passim*.

[4] "Nova methodus determinandi maxima et minima," *Acta Eruditorum*, II (1683).

[5] "Methodus auferendi omnes terminos intermedios ex data aequatione," *Acta Eruditorum*, II, 204.

[6] Born at Leipzig, June 21, 1646 (O.S.); died at Hannover, November 14, 1716 (N.S.). His father wrote his name Leibnütz. It is often written Leibnitz. The Latin form used by our Leibniz was at first Leibnüzius or Leibnuzius, but later the name appears as Leibnitius. The preferred spelling in modern German scientific works is Leibniz.

[7] At least, this is his claim. See J. M. Child, *The Early Mathematical Manuscripts of Leibniz*, p. 37 (Chicago, 1920).

some years after Newton had explained the fluxional calculus to his pupils. Two years later he had his theory well developed, but it was not until 1684 that he published, in the *Acta Eruditorum*, a description of the method and its possibilities. There is no longer any doubt that Leibniz developed his calculus quite independently, and that he and Newton are each entitled to credit for their respective discoveries. The two lines of approach were radically different, although the respective theories accomplished results that were practically identical. Leibniz knew or could easily have known what Newton was doing, and this may have suggested the line of work; and he knew the contribution already made by Barrow[1] in the form of the "differential triangle," but at any rate he was original in much that he accomplished. In a word, it may be said that he made Cavalieri scientific. He also laid the foundation for the theory of envelopes and defined the osculating circle and showed its importance in the study of curves.

Leibniz was a diplomat in the days when Machiavelli was a model, and this is not a flattering way in which to characterize him. Of no other man does the visitor to the portrait gallery of the mathematical world carry away such varied impressions. Some of the engravings show him as a man of great refinement and dignity, while others show the mean, dishonest, disappointed face of a man whose word would never be accepted. In the old controversy as to the invention of the calculus, the word of Leibniz would not have the weight usually given to the statement of a scholar. Nevertheless it must be repeated that he showed great originality in his theory and in the symbolism of the calculus, and is entitled to a high degree of credit for this work.

Minor Writers. Among the minor writers of the first half of the 17th century Johann Faulhaber[2] is known as a successful teacher of mathematics in Ulm. He published various works on algebra and on the curious phases of elementary mathematics.

[1] J. M Child. *loc. cit.*, p. 19 n. *et passim*, and his *Geometrical Lectures of Isaac Barrow*, Chicago, 1918.


[2] Born at Ulm, May 5, 1580; died at Ulm, 1635.



LEIBNIZ

After an engraving by Ficquet

His contemporary, Benjamin Ursinus,[1] professor of mathematics at the University of Frankfort a. d. O., was one of



AUTOGRAPH LETTER OF LEIBNIZ

First page of a letter to Christian Wolf, dated February 10, 1712


the first to introduce logarithms into Germany.[2] Still another contemporary, Daniel Schwenter,[3] was professor of

[1]Born at Sprottau, July 5, 1587; died at Frankfort a. d. O., September 27, 1633 or 1634. Ursinus is the Latin form for Behr, his family name.

[2]*Trigonometria logarithmica, . . . cum magno logarithmorum canone*, Frankfort a. d. O., 1618; 2d ed. 1635; *Magnus canon triangulorum logarithmicus*, Köln a. d. Spree, 1624.

[3]Born at Nürnberg, January 31, 1585; died at Altdorf, January 19, 1636.

Hebrew (1608), oriental languages (1625), and mathematics (1628) at the University of Altdorf. He wrote a well-known work on mathematical recreations.[1] A fourth of the minor



AUTOGRAPH LETTER OF LEIBNIZ
Second page of the letter shown on page 420

writers of this period, Peter Roth,[2] was a Rechenmeister at Nürnberg. He wrote an unimportant algebra[3] in which he treats of equations of the third and fourth degrees.

[1] *Deliciae physico-mathematicae oder Mathematische und philosophische Erquickstunden*, Nürnberg, 1636 (posthumous).

[2] Born at Ingolstadt, *c.* 1580; died at Nürnberg, 1617.

[3] *Arithmetica philosophica, oder schöne, neue, wohlgegründete, überaus künstliche Rechnung der Coss oder Algebra*, Nürnberg, 1608.

For his general learning and his great gift to the world through the museum at Rome which bears his name, if not for any important contribution to mathematics, Athanasius Kircher[1] should be mentioned, although he might quite as appropriately be recorded in the list of Italian scholars, since he spent the better years of his life in Rome. He was a Jesuit, professor of mathematics and philosophy, and later of Hebrew and Syriac, at the University of Würzburg, after which he went (1635) to Avignon and then to Rome, where he taught mathematics and Hebrew. He was a voluminous writer on a great variety of subjects and a zealous collector of curios. His study of optics may have led to the invention of the stereopticon, or to its improvement, but the claims of Kircher's friends are questionable. His mathematical works relate to instruments[2] and to the occult in number,[3] and are not to be taken seriously.

6. The Netherlands

Geographical Limits. In speaking of the Netherlands we must again mention the fact that reference is made to geographical rather than political boundaries. The 17th century was a strenuous one for those countries which we now designate as Holland and Belgium. Roughly speaking, the latter was then under the sovereignty of Spain until Louis XIV began his conquest in 1667, and the former was nearly lost to the same invader in 1672. In the midst of these strenuous times, but generally before the wars, the Netherlands produced several mathematicians who stood high in their respective lines.

Snell. There lived at Leyden at the opening of the 17th century a professor of mathematics (1581) and Hebrew by the name of Rudolph Snell.[4] He wrote a few unimportant works on mathematics, all of which were published in the 16th

[1] Born at Geisa, near Fulda, May 2, 1602; died at Rome, November 28, 1680.

[2] *Pantometrum Kircherianum, h. e. Instrumentum geometricum novum*, Würzburg, 1660.

[3] *Arithmologia sive De occultis numerorum mysteriis*, Rome, 1665.

[4] The Latin form, Snellius, is often used. Born at Oudewater, October 8, 1546; died at Leyden, April 2, 1613.

century. He had a son, Willebrord Snell van Roijen,[1] who succeeded him as professor of mathematics at Leyden (1613), devoting himself chiefly to astronomy, physics, and trigonometry. He wrote on the mensuration of the circle,[2] and set forth the properties of the polar triangle in spherical trigonometry.[3] To him is due the name "loxodrome" for the rhumb line in navigation, the latter name being due to the Portuguese navigator and mathematician Nunes (Nonius).

Girard. Little is known of the life of the mathematician Albert Girard.[4] He wrote on trigonometry,[5] fortifications, practical geometry,[6] and algebra,[7] and edited the works of Simon Stevin.[8] He was one of the first to appreciate the significance of the negative sign in geometry, and was successful in his use of imaginary quantities in the theory of equations. He inferred by induction, as others had done, that an equation of the nth degree has n roots, expressed the sum of the first four powers of the roots of an equation as functions of the coefficients, and discussed general polygons, both cross and simply convex.[9]

Huygens. Although he was known chiefly as one of the world's greatest physicists, particularly in relation to the study of the pendulum, the invention of pendulum clocks, and the laws of falling bodies, Christiaan Huygens[10] should be ranked

[1] Born at Leyden, 1581; died at Leyden, October 31, 1626.

[2] *Cyclometricus, de Circuli Dimensione secundum Logistarum Abacos*, Leyden, 1621.

[3] *Doctrinae Triangulorum Canonicae . . . Libri IV*, Leyden, 1627 (posthumously published). Vieta had already given them.

[4] Born at St. Mihiel, Lorraine, 1595; died at The Hague, December 8/9, 1632. He seems to have lived chiefly in Holland.

[5] *Tables de sinus, tangentes et secantes*, The Hague, 1626.

[6] *Géométrie contenant la théorie et la pratique d'icelle, escrite par Sam. Marolois, revue, augmentée et corrigée*, Amsterdam, 1627. Marolois was a writer on fortifications.

[7] *Invention nouvelle en l'algèbre*, Amsterdam, 1629. Reprinted by Bierens de Haan, Leyden, 1884.

[8] This edition appeared posthumously, Leyden, 1634.

[9] Thus he classified quadrilaterals as "la simple, la croisée et l'autre ayant l'angle renversée."

[10] Born at The Hague, April 14, 1629; died at The Hague, June 8, 1695. The name often appears as Huyghens or Hugenius, with several other variants. There is a biography in his *Opera Varia*, Leyden, 1724.

high among those who improved the new geometry and made known the power of the calculus. He introduced the notion of evolutes,[1] rectified the cissoid, determined the envelope of a moving line, investigated the form and the properties of the catenary,[2] wrote on the logarithmic curve, gave in modern form the rule for finding maxima and minima of integral functions, wrote on the curve of descent (1687), published a work on probability, proved that the cycloid is a tautochronous curve, and contributed extensively to the application of mathematics to physics.



CHRISTIAAN HUYGENS

From an engraving by Edelinck, after a drawing by Drevet

Minor Writers. Among the minor writers the name of Grégoire de Saint-Vincent[3] is one of the best known. He was a Jesuit, taught mathematics in Rome and Prag (1629–1631), and was afterwards called to Spain by Philip IV as tutor to his son Don Juan of Austria. He wrote two works on

[1] *Horologium oscillatorium*, Paris, 1673. It is in this treatise that his work appears at its best.

[2] "Solutio problematis de linea catenaria," *Acta Eruditorum*, 1691.

[3] Born at Bruges, 1584; died at Ghent, January 27, 1667. Among the common variants of his name are Gregorius a Sancto Vincentio and Gregorius von Sanct Vincentius. For a biographical sketch by Quetelet, see the *Annales Belgiques*, VII (1821), 253, and one by Bosmans in the *Annales* of the Société Scientifique of Brussels, XXXIV, 1, 174.

geometry,[1] giving in one of them the quadrature of the hyperbola referred to its asymptotes, and showing that as the area increased in arithmetic series the abscissas increased in geometric series.

The Van Schooten family produced three generations of professors of mathematics at Leyden, all sympathetic with the science but no one of them of first rank. The first was Frans van Schooten,[2] with a trigonometric table to his credit (1627). The second, his son, was also called Frans van Schooten.[3] He wrote on mathematics[4] and was a professor in the engineering school at Leyden and the teacher of Huygens. He edited Vieta (1646), wrote on perspective (1660), and is well known for his Latin edition of the geometry of Descartes (1649). His half brother, Petrus van Schooten,[5] occupied the chair of mathematics at Leyden, being later (1669) transferred to the chair of Latin, but he contributed nothing of permanent value in either of his fields of interest.

René François Walter, Baron de Sluze[6] was a man of standing in the Church, with a taste for mathematics. He contributed to the geometry of spirals and the finding of geometric means, and also invented a general method for determining points of inflection of a curve. One of his contemporaries, but a few years his junior, Johann Hudde[7] was an officeholder in Amsterdam. He was interested in the theory of maxima and minima

[1] *Opus geometricum quadraturae circuli et sectionum coni*, 2 vols., Antwerp, 1647; *Opus geometricum posthumum ad mesolabium per rationem*, . . . Ghent, 1668. See also K. Bopp, *Abhandlungen*, XX, 87.

[2] Born at Leyden, 1581; died at Leyden, December 11, 1646. The name often appears in the Latin form Franciscus. There are a number of interesting MSS. of works by the various Van Schootens now in the library at Groningen. For a list of these, see H. Brugmans, *Catalogus codicum*, No. 108 seq. (Groningen, 1898).

[3] Born at Leyden, *c.* 1615; died at Leyden, May 29, 1660.

[4] *Principia Matheseos Universalis*, Leyden, 1651; *Exercitationum Mathematicarum Libri quinque*, Leyden, 1657. See *Bibl. Math.*, XII (3), 156.

[5] Born at Leyden, February 22, 1634; died at Leyden, November 30, 1670.

[6] Born at Visé on the Maas, July 7, 1622; died at Liége, March 19, 1685. The name also appears as Sluse or Slusius.

[7] Latin, Huddensis. Born at Amsterdam, probably in 1628 or 1629; died at Amsterdam, April 16, 1704.

(1658) and the theory of equations (1657), and in his work on the latter subject he separated into factors the polynomial which he equated to zero.

About this time Cornelis van Beugham wrote a *Bibliographia Mathematica* which appeared at Amsterdam in 1688. This seems to have been the first printed book devoted solely to mathematical bibliography.

7. Other European Countries

Countries Considered. Of the other European countries that exerted substantial influence on mathematics in the 17th century Switzerland stands easily at the head. This seems due not to any particular intellectual influences but to the efforts of one of the most interesting families known in the history of science, and to the labors of a man who came near being the inventor of logarithms.[1]

The other countries demanding attention are Spain and Denmark.

The Bernoullis. Students of heredity have called attention to the extraordinary number of distinguished scholars who descended from the protestant population expelled from the catholic countries in the 16th and 17th centuries.[2] Presumably the same result would be found among the descendants of Catholics, Jews, political refugees, or others who maintained their faith in any manner that tries the souls of men.[3] Of those who descended from Belgian stock that was rooted up during the reign of terror of the Duke of Alba were the members of the Bernoulli family, a family that furnishes one of the most remarkable evidences of the power of heredity or of early home influence in all the history of mathematics. No

[1] L. Isely, *Histoire des Sciences Mathématiques dans la Suisse Française*, Neuchâtel, 1901.

[2] A. de Candolle, *Histoire des sciences et des savants*, 2d ed., p. 338. Paris, 1885.

[3] Among similar cases of descendants of religious refugees are those of Jean Trembley, Simon Lhuilier, Georges-Louis Le Sage, Louis Bertrand, and Élie Bertrand.

less than nine of its members attained eminence in mathematics and physics,[1] and four of them were honored by election as foreign associates of the Académie des Sciences of Paris.

Jacques Bernoulli. The first of the family to attain any reputation in mathematics was Jacques Bernoulli.[2] He first studied theology, but his taste was in the direction of astronomy, mathematics, and physics, and he traveled in France, Holland, Belgium, and England for the purpose of devoting his time to these studies and to meeting learned men. He returned to Switzerland in 1682, took up the study of the new calculus as set forth by Leibniz, and in 1687 became professor of mathematics in the University



JACQUES BERNOULLI

The elder of the two brothers who founded the famous Bernoulli family of mathematicians

[1] For the relationships, see P. H. von Fuss, *Correspondance mathématique . . . de quelques célèbres géomètres du XVIII. siècle*, Vol. I, p. xviii (Petrograd, 1843) (hereafter referred to as Fuss, *Correspondance*); also a note in Terquem's *Nouvelles Annales de Math., Suppl. Bulletin*, Vol. 17, p. 85 (Paris, 1858); P. Merian, *Die Mathematiker Bernoulli*, Basel, 1860.

[2] Born at Basel, December 27 (o.s.), 1654; died at Basel, August 16, 1705. Jacobus Bernoulli, often called Jacques (I) Bernoulli to distinguish him from another Jacques (1759–1789) in the 18th century. English writers often call him James and the German writers Jakob. Since, however, he wrote in French or Latin, the preferable form of his given name is Jacques or Jacobus.

of Basel. He wrote (1683–1701) a large number of memoirs for the *Acta Eruditorum.* These memoirs include such lines of research as series (1686), the quadrisection of a general triangle by two normals (1687), conics (1689), lines of descent (1690), mensuration (1691), cycloids (1692, 1698, 1699), transcendent curves (1696), and isoperimetry (1700). He wrote the second book devoted to the theory of probability,[1] although the subject had been studied in Italy and France much earlier than this.

He solved by infinitesimal analysis the problem of the isochronous curve which had already attracted the attention of various writers and had been solved by Huygens, Leibniz, L'Hospital, and Newton. He determined the length of the catenary curve. Because of his study of the logarithmic spiral, $r = a^{\theta}$, he directed that this curve should be engraved upon his tombstone, with the words *Eadem mutata resurgo*,[2] and the visitor to the cloisters at Basel may still see the rude attempt of the stonecutter to carry out his wish.[3]

The fact that his father was emphatically opposed to his study of astronomy and mathematics, placing all possible obstacles in his way, led him to choose for his device Phaethon driving the chariot of the sun, with the legend, *Invito patre sidera verso.*[4]

His brother Jean Bernoulli[5] was thirteen years his junior,[6] which may account for an attitude of superiority on the part of Jacques and one of resentment on the part of Jean which caused the ill feeling that long existed between them. His father learned no wisdom by his failure to have his brother

[1] *Ars conjectandi,* Basel, 1713 (posthumous). With this was published his *Tractatus de Seriebus Infinitis, Earumque summa Finita, et Usu in Quadraturis Spatiorum & Rectificationibus Curvarum,* and a letter *De Ludo pilae reticularis.* His *Opera* in two volumes appeared at Geneva in 1744.

[2] "I shall arise the same, though changed."

[3] There is an engraving of the design in J. J. Battierius, *Vita . . . Jacobi Bernoulli,* p. 40 (Basel, 1705).

[4] "I study the stars against my father's will"

[5] Born at Basel, July 27 (o.s.), 1667; died at Basel, January 1, 1748. The first name also appears as Johann or John, and often as Jean (1).

[6] Jacques was the fifth child in his father's family, and Jean was the tenth, a fact which may interest those who have faith in the theory of primogeniture.

Jacques become a theologian, and so he determined to make Jean a merchant. The latter thought that he preferred medicine or literature, but soon found that his real taste was for mathematics, and so the world was saved the loss of a genius.

Jean first studied medicine and wrote his doctor's dissertation *De effervescentia et fermentatione* (Basel, 1690). As Jacques found theology uncongenial, so Jean found medicine equally so, and each sought in the study of mathematics the mental activity that he required. Jean became professor of mathematics in the University of Groningen in 1695, and on his brother's death (1705) was elected to fill the place thus vacated at Basel. Each was made a foreign member of the Paris Académie des Sciences in 1699.




JEAN BERNOULLI

The younger brother of Jacques (I) Bernoulli

Jean was even more prolific than his brother in his contributions to mathematics, writing on a wide range of topics, including caustic curves (1692), differential equations (1694), the rectification and quadrature of curves by series (1694), the cycloid (1695), catoptrics and dioptrics (1701), the multisection of angles and arcs (1701), isochronous curves and curves of quickest descent (1718), and various related subjects. He

wrote on the calculus and was one of the most influential scholars on the Continent in making its power appreciated. His collected works appeared six years before his death.[1]



AUTOGRAPH OF JEAN (I) BERNOULLI

The "second fils à Petersbourg" mentioned in the letter was Nicolas (II) Bernoulli, who died at Petrograd only a little over a month after this letter was written

To him is due the use of the term "integral" in its technical sense in the calculus, the first attempt to construct an integral calculus, and the invention of the exponential calculus.[2] He was the first, except in such obvious cases as those known to Cardan, to obtain real results by the use of $\sqrt{-1}$; for example, in finding $\tan n\phi$ in terms of $\tan \phi$.

[1] *Opera omnia*, 4 vols., Lausanne, 1742.
[2] "Principia calculi exponentialium," *Acta Eruditorum*, 1697.

The Later Bernoullis. The work of the later Bernoullis falls in the 18th century, but it is appropriate to mention it briefly at this time. The leading one of the descendants of the two brothers was Daniel Bernoulli.[1] He was the son of Jean (I) Bernoulli, who, curiously enough, made the same mistake as his own father in trying to force his son into trade. Daniel spent some years (1725–1733) in Petrograd as professor of mathematics in the Academy of Petrograd, but in 1733 returned to Basel, where he became a professor in the university. He was a prolific writer, most of his work appearing in the memoirs of the Academy of Petrograd, but he published one volume on mathematics before going to Russia.[2] Most of the memoirs were upon physical questions, but a few related to pure mathematics, including the computation of trigonometric functions (1772, 1773), continued fractions (1775), and the Riccati problem.



DANIEL (I) BERNOULLI
After a portrait from life

Dr. Hutton[3] relates these incidents concerning him:

He used to tell two little adventures, which he said had given him more pleasure than all the other honours he had received. Travelling with a learned stranger, who, being pleased with his conversation, asked his name; "I am Daniel Bernoulli," answered he with great modesty; "And I," said the stranger (who thought he meant to

[1] Usually designated as Daniel (I) Bernoulli. Born at Groningen, February 9 (January 29, o.s.), 1700; died at Basel, March 17, 1782.

[2] *Exercitationes quaedam mathematicae*, Venice, 1724.

[3] *Philosophical and Mathematical Dictionary*, I, 205. London, 1796.

laugh at him), "am Isaac Newton." Another time having to dinner with him the celebrated Koenig the mathematician, who boasted, with some degree of self-complacency, of a difficult problem he had resolved with much trouble, Bernoulli went on doing the honours of his table, and when they went to drink coffee he presented Koenig with a solution of the problem more elegant than his own.

Nicolas (I) Bernoulli[1] was a nephew of Jacques (I) and Jean (I). In his younger days he was professor of mathematics at Padua (1716–1719), but returned to Basel, where he became a professor in the university. He was trained in the law, and his first mathematical treatise was upon the use of the theory of probability in legal matters.[2] He wrote extensively on differential equations and geometry.

Nicolas (II) Bernoulli[3] was the son of Jean (I). He studied law, traveled extensively, became professor of law at Bern (1723–1725), and was finally called to Petrograd as professor of mathematics. He wrote on the geometry of curves, but his death at the age of thirty-one closed a promising career.

Jean (II) Bernoulli[4] was the youngest son of Jean (I). He studied law but spent his later years as professor of mathematics in his native city. His work was chiefly on physics.

Jean (III) Bernoulli[5] was the son of Jean II. Like his father, he studied law but soon turned to mathematics, becoming director of the mathematics class at the Academy of Sciences at Berlin. He was much interested in the history of astronomy but also wrote on the doctrine of chance (1768), recurring decimals (1771), factoring (1771), and indeterminate equations (1772).

The other Bernoullis who were interested to a greater or less degree in mathematics were Daniel (II) (1751–1834), son

[1] Born at Basel, October 10, 1687; died at Basel, November 29, 1759. A note in Terquem's *Nouvelles Annales, Suppl. Bulletin*, 1858, p. 86, says November 25, 1749, but this is an error.

[2] *De Usu Artis Conjectandi in Jure*, Basel, 1709. He also edited the *Ars Conjectandi* of his uncle Jacques (1713).

[3] Born at Basel, January 27, 1695; died at Petrograd, July 26, 1726.

[4] Born at Basel, May 18, 1710; died at Basel, July 17, 1790.

[5] Born at Basel, November 4, 1744; died at Köpnick, near Berlin, July 13, 1807.

of Jean (II); Jacques (II) (1759–1789), son of Jean (II); Christoph (1782–1863), son of Daniel (II); and Jean Gustave (1811–1863), son of Christoph; but none of these attained great fame. The Bernoulli blood had lost its strength.

Other Swiss Writers. Two other Swiss mathematicians of the 17th century deserve mention,—one a genius, the other a plagiarist. The genius was Jobst Bürgi,[1] from 1579 to 1603 court watchmaker to Landgraf Wilhelm IV of Hesse, and later (until 1622) to Kaiser Rudolph II. He wrote on the proportional compasses[2] and on astronomy, but is best known for his invention of logarithms independently of Napier. He was led to the idea by an entirely different route from that taken by the latter, approaching it through the theory of exponents. He did not publish anything upon the subject until after Napier had made known his discovery, and when he finally concluded to print his work it was in the form of a small table of antilogarithms, issued anonymously at Prag in 1620.[3] The book never attracted any attention and remained practically unknown except to historians of mathematics.

The other Swiss writer was of a different character. He was a professor while Bürgi was a watchmaker; his name has been known for three centuries, while Bürgi's has been almost forgotten; but he was a plagiarist, while Bürgi was a genius. Paul Guldin[4] began his work as a goldsmith. He later entered the Jesuit order, lived for a long time in Rome, and became professor of mathematics at the University of Vienna and

[1] Born at Lichtensteig, February 28, 1552; died at Cassel, January 31, 1632. The first name also appears as Joost and Justus; the last as Burgi, Byrgi, Borgen, and Byrgius. See R. Wolf, "Zwei Kleine Notizen zur Geschichte der Mathematik," *Bibl. Math.*, III (2), 33.

[2] Invented by Galileo. They were also described (1607) by Levinus Hulsius in his *Dritter Tractat der mechanischen Instrumente*, Frankfort a. M., 29 pp., being the third of four tractates, 1603–1615. In the fourth of these tractates Hulsius describes a pedometer ("Instrument Viatorii oder Wegzählers"), the earliest mention of this instrument.

[3] *Arithmetische und Geometrische Progress Tabulen*, Prag, 1620.

[4] Born at St. Gall, June 12, 1577; died at Gratz, November 3, 1643. His name was originally Habakuk Guldin, but he changed it when he went from Protestantism to Catholicism.

later at Gratz. He wrote on physics and mathematics,[1] but is chiefly known for the fact that his name attaches to a theorem of Pappus on the volume of a solid generated by the revolution of a plane about an axis,[2]—a theorem which he included in his works without credit, fully aware that it was in the works of Pappus, to which he is known to have had access.

Denmark. In the 17th century Holstein, then a part of Denmark, produced one and only one mathematician of note,—Nicolaus Mercator.[3] He was one of the leading writers of the time on cosmography,[4] and also wrote on trigonometry,[5] the method of computing logarithms,[6] and astronomy, besides editing Euclid's *Elements*.[7] He lived for some time in London and was one of the first members of the Royal Society. In his *Logarithmotechnia* he gives the series that bears his name,

$$\log(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \cdots.$$

Spain. Whatever may be said for mathematics in the 16th century in Spain, less can be said for the century following. Philip II was a bigot, but he was a great bigot. He commanded the respect and won the loyalty of his people, and his reign was one of great works in art and in letters. After his death (1598), however, there occupied the throne a sorry line of kings, and the ruin of the country began. The population of Madrid fell one half in a century, and Seville's sixteen thousand looms were reduced to less than three hundred. From being a prosperous world power the country became a wreck among nations, and in the general destruction mathematics suffered with the other sciences and with letters and art.

[1] *Problema arithmeticum de rerum combinationibus*, Vienna, 1622.

[2] *Centrobaryca seu de centro gravitatis . . . liber I*, Vienna, 1635; *Centrobarycarum pars altera*, Vienna, 1641.

[3] Born near Cismar, in Holstein, *c.* 1620; died in Paris, February, 1687.

[4] *Cosmographia sive Descriptio coeli et terrae in circulos . . .*, Danzig, 1651.

[5] *Trigonometria sphaericorum logarithmica . . .*, Danzig, 1651.

[6] *Logarithmotechnia, sive Methodus construendi logarithmos nova accurata et facilis . . .*, London, 1668.

[7] *Euclidis Elementa Geometrica Libri VI novo ordine ac methodo fere demonstrata*. London. 1678

Russia. As for Russia, she had not yet awakened. The first Russian arithmetic to appear with Hindu-Arabic numerals was written by a teacher named Magnitzky and was printed in 1703. Under Peter the Great there was some development of vocational mathematics, but it was only after the founding of the Academy of Sciences at Petrograd in 1725 that pure mathematics had any standing.[1]

8. The Orient

Effect of Western Civilization. The introduction of Western civilization into India, China, and Japan is interesting because of its diverse effects. As to India, mathematics was already stagnant, and the European influence gave it no stimulus. India has always been content to take her time. Not since Bhāskara (12th century) has she produced a single native genius in this field. In the 17th century only one name, that of Raganātha (*c.* 1621), attracted any general attention even in India, and he contributed nothing that was original. China, which had once done so much in algebra, was content in the 17th century to adopt the European astronomy while allowing her own undoubted abilities to lie dormant. Japan alone of all the Orient developed her native mathematics, although with more or less suggestion from France, Belgium, Italy, and Germany, through the Jesuit missionaries in China; from Holland, through the Dutch traders at Nagasaki; and possibly through scholars who secretly visited the universities of the Low Countries. As to Persia and Arabia, mathematics was dead and forgotten.

China. The mathematical feature of importance at the opening of the 17th century in China was the work of the Italian Jesuit Matteo Ricci already mentioned in Chapter VIII. After his death (1610) other missionaries carried on this work, not only in a religious line but also in the introduction of Western

[1] V. Bobynin, "De l'étude sur l'Histoire des mathématiques en Russie," *Bibl. Math.*, II (2), 103; A. N. Peepin, *History of Literature* (in Russian), I, 253 (Petrograd, 1911).

science. Among these were the Jesuit Nicolò Longobardi[1] (who went to China in 1596), Giacomo Rho,[2] Johann Adam Schall von Bell,[3] Smogolenski[4] (1611–1656), and Ferdinand Verbiest.[5] Of these Schall and Verbiest were particularly prominent in astronomical work. Smogolenski made known the use of logarithms,[6] and his pupil, Sié Fong-tsu, published the first Chinese work on the subject (*c.* 1650). It is interesting to observe that Vlacq's tables (1628) were reprinted in Peking in 1713.[7]

In this century there were several Chinese scholars who wrote important works, but they were all inspired by the Jesuits, and their works are based on European models.[8] Mention should also be made of Mei Wen-ting (1633–1721), a profound scholar, well versed in European as well as native science, who wrote on a variety of subjects and to whom we are indebted for much information concerning the history of Chinese mathematics.

The reason for the activity of the Jesuits in teaching the Western astronomy to Chinese scholars is apparent. It is only by establishing an intellectual superiority that a foreign religion

[1] Born at Calatagirone, Sicily, 1565; died at Peking, December 11, 1655. The Chinese name was Lung Hua-ming.

[2] Born at Milan, 1593; died at Peking, April, 1638. He reached China in 1618. The name also appears as Jacomo Russ, whence the Chinese, Lo Ya-ku.

[3] Born at Cologne, 1591; died at Peking, August 15, 1666. His Chinese name was T'ang Jo-wang. He arrived in China in 1622, and in 1630 was called upon by the government to reform the calendar. He and his colleagues wrote a large number of works.

[4] Chinese name, Mu Ni-ko.

[5] Born at Pitthem, near Courtrai, Belgium, October 9, 1623; died at Peking, January 28, 1688. Chinese name, Nan Huai-jên. He arrived in China in 1659. See H. Bosmans, "Ferdinand Verbiest," in the *Revue des Quest. scientifiques*, pp. 195, 375 (Brussels, 1912), and "Le problème des relations de Verbiest avec la Cour de Russie," in *Annales de la Société d'Émulation pour l'étude de l'hist. . . . de la Flandre*, p. 103 (Bruges, 1913). It was under his direction that most of the large astronomical instruments were made (1674) for the emperor.

[6] In his *T'ien-pu Chên-yüan*.

[7] In the *Lü-li Yüan-yüan*. This work also contained a treatment of algebra on European lines.

[8] For example, Tu Chih-ching, who wrote a geometry, *Chi-ho Lun-yüeh*, based on Euclid, and the *Su-hsiao Tao*, based on European mathematics; Huang Tsung-i, who wrote on the calendar; and Ch'ên Chin-mo (*c.* 1650), who gave 3.15025 as the value of π.

ever makes permanent progress. The Jesuits were not long in seeing that the two sciences in which Europe far surpassed the East were geometry and astronomy, and on these they concentrated their attention.[1]

The Intellectual Awakening of Japan. When Japan, in the 17th century, finally awoke to her intellectual possibilities, it was in a blaze of glory not unlike that which characterized her awakening to her national possibilities in the 19th century. Her progress in mathematics was strangely comparable to the remarkable progress that was going on at the same time in Europe, for in this century she developed a native calculus at almost the same time that Newton and Leibniz were working out their epoch-making theories.

Of the pupils of Mōri Kambei Shigeyoshi (p. 352), one of those to achieve renown was Yoshida Shichibei Kōyū, or Mitsuyoshi (1598-1672), whose *Jinkō-ki*[2] was the first great work on arithmetic to appear in Japan. In this the value of π is given as 3.16. So familiar was the name of this work that it was often used subsequently as a synonym for arithmetic.[3]

The second of Mōri's pupils to contribute to mathematics in a noteworthy manner was Imamura Chishō, whose *Jugai-roku*, devoted to stereometry as well as to arithmetic, appeared in 1639. In this work the value of π is given as 3.162, the area of a circle as $\frac{1}{4}\,cd$, and the volume of a sphere of radius $\frac{1}{2}$ as 0.51.

The third of Mōri's celebrated pupils was Takahara Kisshu, or Yoshitane, but he published nothing on mathematics.

The middle of the century saw also a number of minor writers whose works show considerable ability in mathematics. Among these was Isomura or Iwamura Kittoku, known also as Yoshinori. In his *Ketsugi-shō* (1660) there are a number

[1] "Ce fut alors que des Jésuites pénétrèrent dans la Chine pour y prêcher l'évangile. Ils ne tardèrent pas à s'appercevoir qu'un des moyens les plus efficaces pour s'y maintenir . . . étoit d'étaler des connoissances astronomiques." Montucla, *Histoire*, I (2), p. 468.


[2] The full title means "Small number, large number, treatise," that is, a treatise on numbers from the smallest to the largest.

[3] Compare the word *algorismus* as synonymous with arithmetic, and the name "Euclid" as synonymous with geometry.

of interesting problems proposed by Yoshida Kōyū, of which the following, referring to measurements, are typical:

There is a log of precious wood 18 feet[1] long, whose bases are 5 feet and 2½ feet in circumference. . . . Into what lengths should it be cut to trisect the volume?

A circular piece of land 100 measures in diameter is to be divided among three persons so that they shall receive 2900, 2500,



ONE OF ISOMURA KITTOKU'S PROBLEMS, 1660

His *Ketsugi-shō* appeared in 1660. This is from the 1684 edition. It represents the early state of the advanced native mathematics of Japan and shows the rise of a crude integration

and 2500 measures respectively.[2] Required the lengths of the chords and the altitudes of the segments.

There is also in this work a rough approach to an integral calculus.

In his later years Isomura devoted much attention to magic squares, magic circles, and magic wheels.[3] By a very ingenious method he showed that the surface of a sphere, which he at first thought was equal to $\pi^2 r^2$, has the value πd^2.

[1] In the original, "3 measures." For this and other problems see Smith-Mikami, p. 66.

[2] That is, square measures, by drawing parallel chords.

[3] For particulars, see Smith-Mikami, p. 69.

In 1663 Muramatsu Kudayū Mosei began the publication of a work on arithmetic and mensuration which contributed to the knowledge of the circle and the regular polygons, but only in respect to measurement.

In 1664 Nozawa Teichō published a work called *Dōkai-shō*, in which some ingenious problems in mensuration appear and a step is taken in advance of Isomura in the integral calculus.

In 1666 Satō Seikō wrote the *Kongenki*, a work in which the custom is continued of proposing and solving ingenious problems. This is the first Japanese work in which the ancient Chinese method of solving numerical higher equations appears.

In 1670 Sawaguchi Kazuyuki wrote a work entitled *Old and New Methods in Mathematics*.[1] In this there again appears an approach to an integral calculus, somewhat after the method of Cavalieri, and also a treatment of numerical equations.

Seki Kōwa. The most distinguished Japanese mathematician of the 17th century, and in some respects the most distinguished of all Japanese mathematicians, Seki Shinsuke Kōwa, or Takakazu (1642–1708), was born of a samurai[2] family and showed his great mathematical ability at an early age. He acquired his knowledge to a large extent without the aid of teachers, showing great ingenuity in the affairs of life, in mechanics, in mathematics in general, and in problem-solving in particular. He improved upon the Chinese methods of solving higher equations, systematized the early Chinese use of determinants, possibly invented the circle principle (*yenri* method) which was later developed into a kind of calculus, and proposed numerous problems of an intricate nature. Two problems proposed by Sawaguchi Kazuyuki and solved by Seki Kōwa are substantially as follows:

In a circle three circles are inscribed, each tangent to the other two and to the original circle. They cover all but 120 square units of the circumscribing circle. The diameters of the two smaller circles are equal and each is 5 units less than the diameter of the next larger one. Find the diameters of the three inscribed circles.

[1] *Kokon Sampō-ki.* [2] Feudal lords.

In a certain triangle ABC there is a point P such that $PA = 4$, $PB = 6$, and $PC = 1.447$; such that the sum of the cubes of the longest and shortest sides is 637; and such that the sum of the cubes of the other side and the longest side is 855. Find the lengths of the sides.[1]


The Chinese had some idea of the determinant, as we have seen, but it is to Seki that the honor must be given of expanding a determinant in solving simultaneous equations,—a discovery which anticipated the one made by Leibniz.

Seki's reputation was such as to attract to him a large number of pupils, and his influence upon them was so great as to make itself felt up to the time when the native mathematics became absorbed in the Western science which was so completely adopted in the 19th century.

On the other hand, Seki made no great discovery in mathematics, with the exception of his anticipation of determinants. He was a great teacher, he did pioneer work in the awakening of a scientific spirit in Japan, he showed ingenuity in improving upon the work of his predecessors, but he was not the author of any new method that is now recognized as valuable, and he wrote no great treatise that stands forth today as anything more than a historical document. Because of his efforts to give to his people a knowledge of the mathematical sciences, however, His Majesty the Emperor of Japan very justly paid honor to his memory in 1907 by bestowing upon him the highest posthumous honor ever awarded to such a scholar.

Space does not permit of the further mention of Japanese scholars, with the exception of Nakane Genkei (1661–1733), a contemporary and disciple of Seki, whose works on astronomy (see the illustration on page 441) were influenced by European treatises which had begun to find their way into Japan through the Dutch traders at Nagasaki. He is also known to have been familiar with certain works of the Jesuit missionaries in China and to have recognized their superiority in the astronomical field.

[1] For the problems and suggestions as to solutions see Smith-Mikami, pp. 96, 100.



JAPANESE ASTRONOMY INFLUENCED BY THE WEST

From a work by Nakane Genkei, printed in 1696, showing the explanation of the phases of the moon. Influenced by the Dutch astronomy which had begun to be known in Japan

Contact with Europe. There were certain periods in the history of Japan when contact with the outer world was very difficult. Even when the Dutch traders had a monopoly of bartering with the country through the port of Nagasaki, it was practically impossible for students to leave the island. We find, however, mention of two Japanese students on the records of Dutch universities in Seki's time.[1] Nothing further is known of these men, but the important fact is that they represent contact with intellectual Europe. The problem which this suggests is to ascertain whether through these or similar channels any suggestion of the status of European mathematics reached Japan in Seki's time. There is a tradition, too, that Hatono Sōha, a physician, went to "the Spanish lands"[2] at this time, and that he later returned to Japan. If this is true, European learning of some kind entered the country in the second half of the 17th century. Whether or not it suggested the calculus, which reached its highest native development in Japan in the 18th century, is unknown.

There is a tradition that Seki made a pilgrimage to the ancient shrines at Nara, having learned of certain treatises which were carefully preserved in the Buddhist temples there, and which no one was able to understand. These proved to be Chinese works on mathematics, and Seki is said to have spent three years in mastering their contents. But again, we do not know whether they contained fragments of Western learning that the Jesuits had brought from Europe, or the ancient algebraic science of the Chinese, or possibly traces of Hindu astrology. No doubt, however, the Japanese scholars will in due time search out the sources of the mathematics of Seki's school.

[1] In the *Album Studiosorum Academiae Lugduno Batavae* (The Hague, 1875) it appears that one "Petrus Hartsingius Japonensis," aged 31, was studying philosophy at Leyden in 1654. He is also mentioned by van Schooten in his *Tractatus de concinnandis demonstrationibus geometricis ex calculo algebraico*, in Descartes's *La Géométrie*, 1661 and 1683 editions, p. 413. He also appears on the roll in 1660, as a student of medicine, and again in 1669. In the *Album* there is also the entry, under the date September 4, 1654, "Franciscus Carron Japonensis." Of course the names are not Japanese, and Franciscus Carron is that of a Christian missionary of a century earlier.

[2] Which were then interpreted by the Japanese as including Holland.

TOPICS FOR DISCUSSION

1. Conditions particularly favorable to the development of mathematics in the 17th century.

2. Causes of the decline of Italy's position in mathematics in the 17th century.

3. Forerunners of Newton and Leibniz in the development of the calculus.

4. Cases of relatively late development of mathematical ability.

5. Cases of the influence of inheritance or of early environment upon the development of mathematical power.

6. The relative standing of mathematics in France and in England in the 17th century.

7. The rise of analytic geometry.

8. The greatest mathematical discovery in the 17th century, with statements showing why it was the greatest.

9. The four greatest mathematical books of the century.

10. From the standpoint of the individual and his life, the most interesting mathematical personage of the century.

11. The steps taken in the 17th century to improve geometry.

12. The five most interesting mathematicians of France in the 17th century.

13. The five most interesting British mathematicians in the 17th century.

14. The influence of Newton's English predecessors upon him and upon his work.

15. Certain early steps in the use of continued fractions, infinite series, and infinite products in the 17th century.

16. The rise of books on mathematical recreations.

17. The influence of astronomy and mathematics upon each other in the 17th century.

18. Men who were prominent in both mathematics and physics in the 17th century.

19. The six leading mathematical countries in the 17th century, arranged according to their importance, with reasons justifying the arrangement.

20. The introduction of Western mathematics into the East.

21. The work, the general standing, and the influence of the Japanese scholars in the 17th century.

CHAPTER X

THE EIGHTEENTH CENTURY AND AFTER

1. General Conditions

Status of Elementary Mathematics. Since this work is concerned primarily with the history of elementary mathematics, it would be quite justifiable to set its limit at the close of the 17th century. By that time arithmetic as we ordinarily speak of it, referring to the operations with numbers for commercial and industrial purposes, was practically what it is today. We have changed the way of teaching it, and we have added new applications from time to time as the requirements of business dictated; but the mathematical part of the subject has been very nearly static. We even preserve certain traditional topics and methods that might profitably have been discarded long ago, rarely recognizing that logarithms are more easily handled than roots, and that the algebraic equation is superior to the method of proportion, which we still retain for certain purposes where it might better be discarded.

The algebra that is taught in the secondary schools and in the freshman course in college was practically all in use before 1700. The symbolism has changed but little, and although the elementary textbook is more extensive, it contains no mathematics that was not generally known before that date. The changes that have been made relate chiefly to methods of teaching and to the applications of the subject.

Elementary geometry as ordinarily taught to beginners has made no advance, although, scientifically speaking, the foundations have been explored with far-reaching results. The pupil who studies geometry in a secondary school today is not getting as good mathematics as the one who studied it in the 17th century, simply because it was the selected boy who took the work

at that time. Euclidean geometry is what it was then; it has been rearranged for educational purposes, but the modern textbook of the popular type is not mathematically as scientific as its predecessor. Geometry has made giant strides, but not in the field that teachers generally cultivate in the secondary schools.

Elementary trigonometry and analytic geometry were well known to the mathematical world at the close of the 17th century, and even our modern geometry had made some progress in the work of Desargues.

The calculus has been greatly improved with respect to its foundation principles and the method of presentation, but the elementary calculus that is taught in our colleges, both differential and integral, with its most important applications, was familiar by the year 1700.[1]

The elementary theory of equations, the solution of numerical equations, the symmetric functions of the roots, such forms as continued fractions, the actual handling of complex numbers, the use of infinite series, and even the elementary use of determinants,—all these and various similar topics were well understood before the 18th century. From the standpoint of elementary mathematics, therefore, a large part of the history of the subject closes with the year 1700.

It would be a mistake, however, to suppose that the further progress of the science has an interest only to the student of higher mathematics. Great achievements in any line of work are always stimulating, and some knowledge of these achievements and of the men who made them is necessary to the well-informed teacher or student of the science.[2]

Limitation of the Study. It is the purpose, in this chapter, to limit the study chiefly to a consideration of those mathematicians whose achievements were so noteworthy that everyone who is interested in mathematics should be informed concerning

[1] On the mathematics of the 18th century, particularly in Great Britain, consult J. Leslie, "Dissertation Fourth . . . The Progress of Mathematical and Physical Science, chiefly during the eighteenth century," in the seventh edition of the *Encyclopædia Britannica*.

[2] For a summary of the history of mathematics in the 19th century see J. Pierpont, *Bulletin of the Amer. Math. Soc.*, XI, 136.

them. In the mathematical world names like De Moivre, d'Alembert, Euler, Laplace, Lagrange, Legendre, Gauss, Monge, Galois, Poncelet, von Staudt, and Steiner are so often seen that all teachers of mathematics should know something of the achievements of the men. The student of the higher branches will know this in connection with his researches, but for the teacher in the elementary field of the science a brief résumé will be helpful. It should be understood, however, that no mention will ordinarily be made of certain names which, had they been met in the formative period of elementary mathematics, would have found place. Various men who might properly be mentioned in this connection will be referred to in Volume II of this work.

Royal Patronage. With the 18th century the king of France no longer stands out as the sole royal patron of science.[1] Queen Anne bestows knighthood upon Newton; George I shows an interest in scientific laboratories; Peter the Great is at pains to meet with learned men and founds (1724) an academy at Petrograd to which there later come such mathematicians as the Bernoullis and Euler; George III, in spite of his parsimony, endows the observatory of Herschel; and Frederick II calls to the Berlin Academy Maupertuis, d'Alembert, a Bernoulli, and Lagrange, not to speak of Voltaire, who, as we shall see, had some claim to the title of mathematician. In spite of all this, France held high place in the fostering of all the sciences,—perhaps the highest; and no other nation could boast, at the close of the 18th century, such a galaxy of stars in the mathematical firmament.

2. Great Britain

Nature of the Work. As would naturally be expected, the influence of Newton determined to a large extent the nature of the work in Cambridge, Oxford, London, and Edinburgh in the 18th century. The improvement of the calculus and the widening of the range of applications of the subject were the

[1] A. Rambaud, *Histoire de la Civilisation Française*, 12th ed., II, 473. Paris, 1911.

characteristic features. A few of the leading names connected with this movement will be mentioned, together with a brief statement of the contributions of each.

It should be observed, also, that the 18th century saw mathematics made popular for the first time in Great Britain. Schools were established for the poor to attend on Sunday, the only day that they could attend them at all, and this was done much against the opposition of many in the Church; circulating libraries were established; printing ceased to be largely a London monopoly and spread throughout the country; elementary handbooks appeared; and such popularizers of mathematics as the *Ladies' Diary* (1704–1840) and the *Gentleman's Diary* (1741–1840) had a wide circulation. Even such classical works as Newton's *Principia* were printed in the vernacular. Mathematics had ceased to be aristocratic; democracy had begun to assert its rights in intellectual as well as political matters.

Cotes. Newton is credited with the statement, "If Cotes had lived, we had known something," a remark that might, it would seem, be made with greater force with respect to certain others who, like Pascal, Galois, and Clifford, died at a relatively early age. Nevertheless it is true that few scholars showed such powers of analysis before the age of thirty-four, the age at which Roger Cotes[1] died. He had shown a taste for mathematics when only about twelve years of age. He was educated at St. Paul's School, London, where he also developed a taste for metaphysics, philosophy, and divinity, and thence proceeded to the mathematical Mecca of England, Trinity College, Cambridge. When only twenty-four years of age he was appointed (1706) to the Plumian professorship of astronomy, the first to fill the chair which had just been established (1704) by Dr. Plume, archdeacon of Rochester. In 1713 he published at Cambridge the second edition of Newton's *Principia*. Only two of his memoirs appeared during his lifetime, but most of his writings were collected and published.

[1] Born at Burbage, Leicestershire, July 10, 1682; died at Cambridge, June 5, 1716.

shortly after his death, by his cousin, Dr. Robert Smith (1689–1768),[1] who succeeded him in the Plumian professorship.

He discovered an important theorem on the *n*th roots of unity, partly anticipated the method of least squares, and discovered a method of integrating rational fractions with binomial denominators. His theorem on the harmonic mean between the segments of a secant to a curve of the *n*th order, reckoned from a fixed point, is well known.[2]

The Calculus in English. The first work on the Newtonian calculus to appear in the English language was published in London in 1704.[3] It was written by Charles Hayes,[4] a member of Gray's Inn, London. The purpose of the work is set forth in the preface:

> The Author has been well assur'd that there are in England as many Lovers of the Mathematicks as in any part of the World; . . . that in other Nations the best pieces of Learning are written in their own mother Tongues, for the good of their Country which we seem purposely to slight, seeking a little empty applause by writing in a Language not easily attain'd.

The work is clearly written, but was overshadowed later by such treatises as those of James Hodgson[5] and John Rowe.[6] Hayes also wrote on the finding of longitude (1710) and began but did not live to complete a *Chronographia Asiatica et Aegyptica.*

[1] *Harmonia Mensurarum, sive Analysis et Synthesis . . .*, Cambridge, 1722. The second part of the volume comprised his *Opuscula Mathematica.*

[2] W. W. R. Ball, *A History of the Study of Mathematics at Cambridge*, p. 88 (Cambridge, 1889), hereafter referred to as Ball, *Hist. Math. Cambridge*; Chasles, *Aperçu*, p. 147.

[3] *A Treatise of Fluxions: or, an Introduction to Mathematical Philosophy*, London, 1704.

[4] Born 1678; died in London, December 18, 1760.

[5] Master of the mathematical school in Christ's Hospital, London. He was born in 1672 and died in London, June 25, 1755. Among his several works was *The Doctrine of Fluxions*, London, 1736, with an edition in 1758.

[6] *Introduction to the Doctrine of Fluxions*, London, MXCCLI (*sic* for MDCCLI, 1751), with editions in 1757, 1767, and 1809. This was the first really popular presentation of the subject in English.

In connection with these works Joseph Raphson's[1] history of the calculus[2] should be mentioned. The purpose of the book is thus stated:

To assert the Principal Inventions of this Method, to their First and Genuine Authors; and especially those of Sir Isaac Newton, who has vastly the Advantage of all others as well in respect of Priority of Time, as the Great and Noble Nature of his Discovery.

Such a book, written in the heat of the controversy as to the priority of the works of Newton and Leibniz, would naturally be open to the charge of partisanship.

James Stirling. A brief list of some of the other important 18th century writers will tend to show the nature of the work being done in Great Britain.

James Stirling[3] was educated at Glasgow and at Balliol College, Oxford. He left Oxford (1715), partly on account of his relations to the Jacobites, and went to Venice to accept a professorship. In Italy he formed the acquaintance of Nicolas Bernoulli, who was then at Padua, and was probably encouraged by him to write his well-known work on lines of the third order.[4] Of these lines he added four to those already discussed by Newton, and he also wrote a paper on the differential method in the treatment of infinite series.[5] He returned to London in 1725, devoting himself to mathematics, meeting with Newton in the latter's closing years, writing several important memoirs, and corresponding with many noted mathematicians of the day. His sojourn in Venice gave him the nickname "the Venetian," and by this he was commonly

[1] Or Ralphson. He wrote an *Analysis aequationum universalis*, London, 1690, and died before 1715.

[2] *The History of Fluxions, Shewing in a compendious manner The first Rise of, and various Improvements made in that Incomparable Method*, London, 1715 (posthumous).

[3] Born at Garden, Stirlingshire, 1692; died at Edinburgh, December 5, 1770.

[4] *Lineae tertii ordinis Newtonianae sive illustratio tractatus Newtoni de enumeratione linearum tertii ordinis*, Oxford, 1717.

[5] *Methodus differentialis sive Tractatus de summatione et interpolatione serierum infinitarum*, London, 1730. An English translation by Francis Holliday appeared in 1749.

known to his friends. He made an important survey of the Clyde, and in later life (1735) became the manager of a mining company in Lanarkshire; and, strange to say of such a scholar, he made a great success of this venture.

De Moivre. Although born in France, Abraham de Moivre[1] spent his life from the age of eighteen in London, and may properly be ranked with the English school of mathematicians. Compelled by narrow circumstances to forego the life of a student, he supported himself by private teaching, by lecturing, and by giving answers to mathematical puzzles. It is said that he passed most of his time in a London coffee house, where his genius in solving problems brought to him sufficient return for his humble needs. Having come by chance upon a copy of Newton's *Principia*, he discovered his weakness in the higher range of mathematics and by assiduous application soon became recognized as a man of genuine ability in research. He was admitted to membership in the Royal Society and into the academies of Paris and Berlin. His work was chiefly on trigonometry,[2] probability,[3] and annuities.[4] In his discussion of trigonometry he gave the theorem which bears his name, $(\cos x + i \sin x)^n = \cos nx + i \sin nx$, a relationship already stated in substance by Cotes,[5] and one which leads to numerous interesting identities in connection with complex numbers. Indeed, it stands as one of the basic propositions in the theory of such numbers. He is also known for having given the various quadratic factors of $x^{2n} - 2kx^n + 1$, for having stated the rule for finding the probability of a compound event, for his work on recurring series, and for his extension of the quadrature of the lunes of Hippocrates.[6]

[1] Born at Vitry, Champagne, May 26, 1667; died in London, November 27, 1754.

[2] *Miscellanea Analytica, de seriebus & quadraturis* . . ., London, 1730.

[3] *Doctrine of Chances*, London, 1718, with later editions in 1738 and 1756. It was dedicated to Newton. There was an Italian edition, Milan, 1776.

[4] *Annuities upon Lives*, London, 1725, with later editions.

[5] See Volume II, Chapter IV.

[6] See the summaries of his papers in the *Phil. Trans.* abridgment, 4th ed., I, 1, 29, 81, 90, *et passim*; IV, 3, 25, 77 (London, 1731).

There is often told a story of his death, to the effect that he had declared it to be necessary to sleep a quarter of an hour longer each day than on the preceding one. If he was sleeping six hours a day when he began this series, it is evident that the first day thereafter he would sleep $6\frac{1}{4}$ hours, and on the 73d day he would reach the limit.

Whiston. Born in the same year as De Moivre, but under seemingly more favorable stars, and dying only two years before him, William Whiston[1] lived a life as ideal as De Moivre's was discouraging. He received many honors, wrote numerous scientific and theological works, and held the Lucasian chair of mathematics in Cambridge (1703–1710); and yet he left a name that is today by no means so well known in the history of mathematics as that of his humbler contemporary. His chief interests were in astronomy.

Brook Taylor. The name of Brook Taylor[2] is familiar to every student who knows the rudiments of the calculus, Taylor's Theorem being one of the first instruments that he uses. The discoverer of this theorem was educated at St. John's College, Cambridge. He early gave great promise of success in mathematics, wrote various papers for the *Philosophical Transactions*, was admitted to the Royal Society, and became its secretary. When only thirty-four, however, he gave up his secretaryship and devoted himself to writing. In 1715 he published a work[3] in which is contained his well-known proposition,

$$f(x+h)=f(x)+hf'(x)+\frac{h^2}{2!}f''(x)+\cdots,$$

and some treatment of the calculus of finite differences, of interpolation, and of the change of the independent variable. He published two works on perspective,[4] giving the first

[1] Born at Norton, Leicestershire, December 9, 1667; died in London, August 22, 1752.

[2] Born at Edmonton, August 18, 1685; died in London, December 29, 1731.

[3] *Methodus Incrementorum Directa et Inversa*, London, 1715. He had announced the discovery in 1712.

[4] *Linear Perspective*, London, 1715; *Principles of Linear Perspective*, London, 1719.

general enunciation of the principle of vanishing points. He was also the author of various memoirs on physics, logarithms, and series.[1]



From an engraving by J. Dudley

Maclaurin. Associated with Taylor's name, on account of the theorem above mentioned, is that of Colin Maclaurin,[2] a Scotch mathematician, who entered the University of Glasgow (1709) at the age of eleven. He soon showed a taste for

[1] For a biography see the preface to his posthumous work, *Contemplatio Philosophica*, London, 1793.

[2] Born at Kilmodan, Argyllshire, February, 1698; died at York, June 14, 1746. See C. Tweedie, *Math. Gazette*, IX, 303.

mathematics, and at the age of twelve, having accidentally run across a copy of the work, he mastered the first six books of Euclid in only a few days. At the age of fifteen he took the degree of M.A., publicly defending with much success a thesis on the power of gravity. At the age of nineteen he was elected to the chair of mathematics in the Marischal College, Aberdeen, and at the age of twenty-one (1719) he took to his London printer his first important work.[1] After traveling for some time as tutor to the son of Lord Polwarth he became (1725) an assistant at the University of Edinburgh, finally being elected to a professorship.



COLIN MACLAURIN

Known chiefly for the formula which bears his name

To him is due a method of generating conics which bears his name. His treatise on fluxions[2] contains the well-known identity,

$$f(x)=f(0)+xf'(0)+\frac{x^2}{2!}f''(0)+\cdots,$$

a relationship easily deduced from Taylor's Theorem, and one which had been announced by James Stirling twelve years earlier. Maclaurin greatly generalized the theory of the mystic hexagram and was the first to publish a work on the subject, for Pascal's essay, although written more than a century earlier, did not appear in print until 1779. He wrote an algebra

[1] *Geometria Organica: sive Descripto Linearum Curvarum Universalis*, London, 1720. Part of the propositions were worked out in his 16th year.

[2] *Treatise of Fluxions*, 2 vols., Edinburgh, 1742.

that was published posthumously,[1] and various memoirs on geometry and physics. Ball[2] has very well summed up his influence in these words:

Maclaurin was one of the most able mathematicians of the 18th century, but his influence on the progress of British mathematics was on the whole unfortunate. By himself abandoning the use both of analysis and of the infinitesimal calculus, he induced Newton's countrymen to confine themselves to Newton's methods, and it was not until about 1820, when the differential calculus was introduced into the Cambridge curriculum, that English mathematicians made any general use of the more powerful methods of modern analysis.

Saunderson. Nicholas Saunderson[3] deserves mention as one of the mathematicians of this period, not so much because of his great achievements in advancing the science as on account of the inspiration that his history offers to those who labor under difficulties such as discourage most men and lead them early to abandon hope.

When only one year of age he became blind through an attack of smallpox. He was a pupil of Whiston's, who was then Lucasian professor of mathematics at Cambridge, and succeeded him in 1711. He was created doctor of laws in 1728 by command of George II, and became a Fellow of the Royal Society in 1736. He was very successful as a teacher, and is especially known for his *Algebra*, published posthumously in 1740–1741 and translated into French by É. de Joncourt (Amsterdam, 1756). His *Method of Fluxions* also appeared after his death (1751). Saunderson counted among his friends such well-known scholars as Newton, Cotes, and De Moivre, and did much to make the philosophy of Newton known to the mathematicians of his time. He could carry on long and complicated mathematical problems mentally, which partly accounts for his success in spite of his misfortune.[4]

[1] *Treatise of Algebra*, London, 1748, with several later editions. On his attempted proof of Taylor's Theorem see *Bibl. Math.*, I (3), 438.

[2] *Hist. Math.*, 6th ed., 388.

[3] Born at Thurlston, Yorkshire, January, 1682; died at Cambridge, April 19, 1739.

[4] For his biography consult the *Algebra*, mentioned above.



NICHOLAS SAUNDERSON
After Vanderbanck's painting

Other British Writers of the Century. Among those who contributed to the advance of the Newtonian philosophy at this time Humphrey Ditton[1] deserves mention. He was a man without university training but interested in Church work. He left this work, however, and devoted the latter part of his life to the study of mathematics and to teaching. He was much esteemed by Newton, on whose recommendation he was elected mathematical master at Christ's Hospital, London. He published a number of memoirs on mathematics and physics, a work on fluxions,[2] a revision of an algebra,[3] and a work on perspective (1712).

All students of the history of geometry will recognize the name of Robert Simson,[4] who, although educated as a physician, became professor of mathematics (1711) at the University of Glasgow. He was a thorough student of Greek mathematics,[5] and most of the English editions of Euclid are based upon his edition of the *Elements*. He was averse to the use of algebraic analysis in geometry, and his methods are those of the Greeks.

Another well-known Scotch writer of this period, Matthew Stewart,[6] entered the University of Glasgow in 1734, coming under the instruction of Simson. He also attended Maclaurin's lectures at Edinburgh, and succeeded him in his professorship in 1747. He was particularly interested in geometry and in the introduction of the simple form of the Greek synthetic demonstration into modern higher mathematics.[7] He also devoted

[1] Born at Salisbury, May 29, 1675; died in London, October 15, 1715

[2] *An Institution of Fluxions*, London, 1706. There was an enlarged edition in 1726.

[3] *Synopsis Algebraica* of John Alexander, London, 1709.

[4] Born at Kirktonhall, Ayrshire, October 14, 1687; died at Glasgow, October 1, 1768. See William Trail, *Account of the Life and Writings of Robert Simson*, Bath, 1812.

[5] *Sectionum Conicarum Libri V*, Edinburgh, 1735, with an enlarged edition in 1750; *Apollonii Pergaei Locorum Planorum libri II, restituti . . .*, Glasgow, 1749; *Euclid's Elements*, Glasgow, 1756, with numerous editions. Some of his works were published posthumously at Glasgow in 1776.

[6] Born at Rothesay, Isle of Bute, 1717; died at Edinburgh, January 23, 1785.

[7] *General Theorems*, Edinburgh, 1746; *Tracts, Physical and Mathematical*, Edinburgh, 1761.

much of his energy to astronomy, particularly to the problem of the sun's distance from the earth.[1] Several propositions of modern geometry bear his name.[2]

Contemporary with these Scotch writers, but living in England, there was that strange mathematical genius, Thomas Simpson.[3] He was brought up by his father to be a weaver, and hence his early education was confined to the reading and writing of English. Since Thomas persisted in reading beyond what his father thought necessary, resulting in a vigorous paternal protest, the boy decided to run away from home. A peddler having given him a copy of Cocker's arithmetic containing an appendix on algebra, he began the study of mathematics. His life was a turbulent one, and it suffices for our purposes to say that he struggled against poverty in London with sufficient success to allow him to publish works on the calculus,[4] probability,[5] algebra,[6] and various other subjects, and to prepare numerous monographs of importance. He was a man of undoubted genius, and his abilities were recognized in his election as professor of mathematics at the Woolwich Military Academy in 1743 and as a Fellow of the Royal Society in 1745. As a teacher he was a failure; equally was he a failure in the home; and, as with many other human failures, drink finally asserted the mastery.

Another self-made mathematician of this period appeared in the person of John Landen,[7] known for his work on residual analysis,[8] a theory by which he solved a number of problems more simply than had been done by fluxions, and for his memoir

[1] See *Essays of the Phil. Soc. of Edinburgh*, 1756.

[2] Chasles, *Aperçu*, 173.

[3] Born at Market Bosworth, Leicestershire, August 20, 1710; died at Market Bosworth, May 14, 1761.

[4] *A new Treatise of Fluxions*, London, 1737; *The Doctrine and Application of Fluxions*, London, 1750.

[5] *A Treatise on the Nature and Laws of Chance*, London, 1740; *The Doctrine of Annuities and Reversions*, London, 1742.

[6] *An Elementary Treatise of Algebra*, London, 1745; 2d ed., 1755.

[7] Born at Peakirk, near Peterborough, January 23, 1719; died at Milton, near Peterborough, January 15, 1790.

[8] *Discourse concerning Residual Analysis*, London, 1758; *The Residual Analysis*, London, 1764.

(1755) on the rectification of the arc of a hyperbola. He also wrote on astronomy, series, elliptic transcendents (1771), and physics, and was one of the early contributors to the theory of the top. He was admitted to the Royal Society in 1766.

Charles Hutton,[1] one of the best-known English writers on mathematics at the close of the 18th century, owed his prominence more to his perseverance than to his scientific ability. He was professor of mathematics at the Military Academy at Woolwich (1772–1807) and is chiefly known for his mathematical tables[2] and his dictionary.[3]

Among the contemporaries of Hutton one of the intellectual leaders, but by no means one of the best-known, was William Wallace.[4] As a young man he was a printer, later becoming interested in bookselling. Meantime he developed a taste for mathematics, gave private lessons, and at the age of twenty-six became a teacher in the Perth Academy. He finally became a professor in the University of Edinburgh. He wrote a number of memoirs on logarithms, trigonometry, the pantograph, and geodesy, but his chief work was in relation to the quadrature of the hyperbola and to hyperbolic functions.[5]

Contemporary with Landen and Hutton there was a writer who did very much less but possibly gained somewhat more in the estimation of many people, the accident of a single discovery having given his name a place in the history of the number theory. John Wilson[6] was one of those men who did one thing well in the field of mathematics and then failed to

[1] Born at Newcastle upon Tyne, August 14, 1737; died in London, January 27, 1823.

[2] *Mathematical Tables, containing Common, Hyperbolic, and Logistic Logarithms, with other Tables, and a large and original History of the Discoveries and Writings relating to those Subjects,* London, 1785.

[3] *Mathematical and Philosophical Dictionary*, London, 2 vols., 1795, 1796, with a second edition in 1815.

[4] Born at Dysart, Fifeshire, September 23, 1768; died at Lauriston, near Edinburgh, April 28, 1843. S. Günther, *William Wallace, ein Vorläufer der Lehre von den Hyperbelfunktionen,* Prog., Ansbach, 1880.

[5] Günther, *loc. cit.*, and the *Trans. of the Royal Soc. of Edinburgh*, VI, 269, 271, 302, etc.

[6] Born at The How, Westmoreland, August 6, 1741; died at Kendal, October 18, 1793. See a note on his life by Cantor, *Bibl. Math.*, III (3), 412.

meet the expectations of his contemporaries. He entered Peterhouse, Cambridge, in 1759, and while still an undergraduate he discovered that if p is a prime number, then $1+(p-1)!$ is a multiple of p,—a fact already known to Leibniz but not published. The statement has generally been known as Wilson's Theorem.[1] He was senior wrangler in 1761 and became a Fellow of the Royal Society in 1782.

The Early Nineteenth Century. Of British writers born in the 18th century but whose work was done in the century following, relatively few stand out as brilliant mathematicians. Among those who contributed in some noteworthy way to the progress of their science, one of the earliest was Robert Woodhouse,[2] who was educated at Caius College, Cambridge, became Plumian professor in the university, and did much to replace the calculus of fluxions by the differential calculus. He sought, moreover, to put the latter subject on a firm scientific foundation, and it is due in no small degree to his efforts that this was done.

Probably the name best known of all this group, in books on elementary mathematics, is that of William George Horner.[3] Although not a man of great ability as a mathematician, he succeeded in making for himself a name that is well known to students of algebra. He was a teacher at Bath when he came independently upon an ancient Chinese method of approximating the roots of a numerical equation. This method, which had been practically forgotten in China, was made known in a paper read before the Royal Society in 1819, and since that time Horner's Method has become familiar in all parts of the English-speaking world.[4]

His contemporary, George Peacock,[5] was a very different type of man. A student at Trinity College, Cambridge, he

[1] On the history of the theorem see Dickson, *Hist. Th. Numb.*, I, 50.

[2] Born at Norwich, April 28, 1773; died at Cambridge, December 23, 1827.

[3] Born in 1786; died at Bath, September 22, 1837.

[4] Republished in the *Ladies' Diary* for 1838 and again in revised form in *The Mathematician* for 1843. His proof (1826) of Euler's Theorem (see Volume II, Chapter I) should be mentioned as showing his ability in number theory. As to the validity of the Chinese claim, see Volume II, Chapter VI.

[5] Born at Denton, April 9, 1791; died at Ely, November 8, 1858.

came to represent the solid, substantial mathematics of England and to do much to improve its status. He was appointed Lowndean professor of astronomy and geometry in 1836, and three years later became dean of Ely cathedral, spending the last twenty years of his life there. He was interested in the movement to introduce the differential notation into the work in the calculus, in the founding of the observatory at Cambridge, and in the preparation of scholarly treatises on elementary mathematics. He was one of the prime movers in all mathematical reforms in England during the first half of the 19th century, although contributing no original work of particular value.

One of his contemporaries, and also a Trinity man, Charles Babbage[1] was a worthy representative of the output of Cambridge in this period. He became Lucasian professor (1828–1839), assisted in founding the Astronomical Society (1820), the British Association for the Advancement of Science (1831), and the Statistical Society of London (1834), and did much to introduce the differential notation into British mathematics. He worked on an elaborate calculating machine, the most noteworthy effort in this direction, from the point of view of originality, since that of Pascal.[2]

Perhaps the British mathematician of this period who showed the greatest genius, or at any rate the greatest perseverance, was Peter Barlow.[3] Born of humble parents, he became one of the leading writers of England on the theory of numbers,[4] professor of mathematics at the Woolwich Military Academy, and a Fellow of the Royal Society (1823). His contributions to the magnetic theory, the strength of materials, and optics were also noteworthy, and his mathematical dictionary[5] is still a valuable source of information. His tables of factors,

[1] Born at Teignmouth, Devonshire, December 26, 1792; died in London, October 18, 1871.

[2] His *Calculating Engines*, a work including much historical information, was edited by his son, General H. P. Babbage, and published at London in 1889.

[3] Born at Norwich, October 13, 1776; died March 1, 1862.

[4] *An Elementary Investigation of the Theory of Numbers*, London, 1811.

[5] *A New Mathematical and Philosophical Dictionary*, London, 1814.

reciprocals, powers, roots, hyperbolic logarithms, and primes should also be mentioned as the best of the earlier publications of the kind, being still looked upon as standards.[1]

Sir William Rowan Hamilton. Of those whose work added to the prestige of Great Britain and Ireland in the mid-Victorian period, only a few need be mentioned in a work devoted chiefly to elementary mathematics. Of these, one of the best known is Sir William Rowan Hamilton.[2] He was one of the great mathematical products of Ireland in the 19th century. Although descended from Scotch stock, he was proud to proclaim himself an Irishman. He was one of the infant prodigies that occasionally arise in the history of mathematics, usually, as we have seen, with disappointing results. At the age of three he read English fluently and was somewhat advanced in arithmetic; at five he could read Latin, Greek, and Hebrew; at eight he could also write Latin and read French and Italian; at ten he was studying Arabic and Sanskrit; at twelve he had a working knowledge of all these languages, together with Syriac, Persian, Hindustani, and Malay, and was contesting with the American prodigy, Zerah Colburn, in long mental calculations; and at thirteen he had written an algebra, which, fortunately no doubt, was not offered for publication. At fourteen he was able to write Persian, and at sixteen he had made known an error in one of the demonstrations in the *Mécanique Céleste* of Laplace. At Trinity College, Dublin, he continued his brilliant record, receiving his appointment to the professorship of astronomy while still an undergraduate. Hamilton was knighted in 1835. In 1843 he made his great discovery of quaternions, but his first work on the subject was not published until 1853. His second work appeared posthumously.[3] If the theory has not led to the results anticipated by Hamilton and his friends,

[1] *New Mathematical Tables*, London, 1814. De Morgan published an edition in 1856.

[2] Born at Dublin, August 3, 1805; died at Dublin, September 2, 1865. For biography, see A. Macfarlane, *Ten British Mathematicians*, p. 34 (New York, 1916); hereafter referred to as Macfarlane, *Ten Brit. Math.*

[3] *Lectures on Quaternions*, Dublin, 1853; *Elements of Quaternions*, London, 1866 (posthumous).

both it and the *Ausdehnungslehre* of Grassmann have greatly extended the vision of both mathematicians and physicists.

Salmon, De Morgan, and Boole. Among the prominent mathematicians produced by Ireland in the 19th century there should also be mentioned George Salmon,[1] whose works on conics, higher plane curves, analytic geometry of three dimensions, and higher algebra are recognized as standard authorities.

Less well known than Hamilton for any single achievement, less well known than Salmon for his thoroughness in science, familiar to a much larger circle of readers because of his wide range of interest and his skill in popularizing the science, was that eccentric but brilliant teacher, Augustus De Morgan.[2] Educated at Trinity College, Cambridge, he became professor of mathematics in London in 1828, displaying unusual gifts as a teacher and scattering his energies recklessly. His *Trigonometry and Double Algebra* (1849) contained certain features of quaternions, but he did not follow this or any other theory to the conclusion that seemed within his reach. He wrote various textbooks, each a mine of information for the teacher and entirely hopeless for the pupil. His contributions to the theory of probability still rank as among the best in English, and the same may be said for his contributions to logic. He devoted considerable attention to the history of mathematics, but his articles are not only eccentric but unreliable. His best work in this line is to be found in Smith's *Dictionary of Greek and Roman Biography* (London, 1862–1864), the *Penny Cyclopædia* (London, 1833–1843), the *Companion to the Almanac* for various years, and his *Arithmetical Books* (London, 1847). His *Budget of Paradoxes*, edited by Mrs. De Morgan after his death,[3] is an interesting satire on circle squarers and their kind. Had he been able to confine himself to one line, he might have been a much greater though a less interesting man.

If poverty, delayed education, and general lack of early advantages were a bar to progress in abstract science, George

[1] Born at Cork, September 25, 1819; died at Dublin, January 22, 1904.
[2] Born at Madura, India, June 27, 1806; died in London, March 18, 1871.
[3] London, 1872; 2d ed., by D. E. Smith, Chicago, 1915.

Boole[1] would never have been professor of mathematics at Queen's College, Dublin, the theory of invariants and covariants would not have been what it is today, and the mathematical theory of logic might not have reached its present position. Boole's circumstances did not permit of his beginning any serious study of mathematics until he was twenty, although he picked up by himself some knowledge of Latin and Greek and was able to do a little teaching to help him on with his scholastic work. When he was twenty years old he decided that he was able to open a school of his own, and this he did, using the small income to assist his aged parents and to buy books for himself. Thus, without any university training, he advanced, and in 1849 was appointed to the chair of mathematics in Queen's College. His *Mathematical Analysis of Thought* (1847), *Laws of Thought* (1854), *Differential Equations* (1859), and *Finite Differences* (1860) are still looked upon as standard authorities.

Sylvester. There were two men, companions for many years but, like many companions, of very different character, who stand out with special prominence at this period. The first of these was James Joseph Sylvester.[2] Educated at St. John's College, Cambridge, and one of the most gifted members of his class, he was not allowed to take a degree because of his Jewish faith, and for the same reason he was barred from a fellowship. For the degree he went to Dublin, but after the abolition of the theological tests in 1872 the University of Cambridge awarded him both the bachelor's and master's degrees. Soon after leaving the university he was appointed (1837) professor of natural philosophy in University College, London, and two years later (1839) was elected a Fellow of the Royal Society. Opportunities for advancement not being promising in England, he accepted an appointment (1841) as

[1] Born at Lincoln, November 2, 1815; died at Cork, December 8, 1864. See Macfarlane, *Ten Brit. Math.*, p. 52.

[2] Born in London, September 3, 1814; died in London, March 15, 1897. See F. Franklin, in *Bulletin of the Amer. Math. Soc.*, III, 299; M. Noether *Math. Annalen*, L, 133; P. A. MacMahon, *Nature*, March 25, 1897.

professor of mathematics in the University of Virginia. His election took place on July 3, 1841, and he began his work in the autumn. He made a failure of his teaching, had a serious personal encounter with a student who is said to have attacked him, and hurriedly left the university in the following March. The official records of March 22, 1842, contain the following resolution:



JAMES JOSEPH SYLVESTER


His influence was strongly exerted in establishing university research in mathematics in the United States

Resolved, That the resignation of Mr. Sylvester be accepted, to take effect from and after the 29th day of the present month or at any earlier period that he may elect;—that a copy of this resolution be forthwith communicated to him by the Secretary, and that he be informed that in accepting his resignation the Board has not deemed it necessary to investigate the merits of the matter in difference between himself and the student Ballard, and does not mean to impute to Mr. Sylvester any blame in the matter.

Evidently, therefore, the university authorities were convinced that Sylvester's further relations with the faculty were not desirable.

He seems to have returned to London about three years after leaving Virginia. Here he took up actuarial work, became a student in the Inner Temple (1846), and was called to the bar (1850). He became professor of mathematics at the Military Academy in Woolwich in 1855 and remained there until

his forced retirement on account of age (1869). In 1877 he was called to Johns Hopkins University and did more than any other man of his time to establish graduate work in mathematics in America. Among his other contributions to the advance of the science in this country was his founding of the *American Journal of Mathematics*. In 1883 he was elected to succeed H. J. S. Smith in the Savilian professorship at Oxford, but his lectures were not popular and in 1892 he gave place to a deputy professor and spent his last years in London.




ARTHUR CAYLEY
From a photograph made in 1870

Sylvester was often looked upon as unsystematic, domineering, impractical, conceited, and unhappy, but those who knew him well have testified to his genial nature and his enthusiasm in his work with students. His contributions show that he was a genius in mathematical investigation, his chief line of interest being in higher algebra, including the study of invariants.

Cayley. The companion of Sylvester already referred to was Arthur Cayley.[1] He was the son of an English merchant who had settled in Petrograd and who looked forward to his son's taking part in the business which he had established. Soon after young Cayley, at the age of fourteen, was sent to King's College School, London, it was found that he showed such ability in mathematics that his father decided that he should proceed to Cambridge. He accordingly entered Trinity College at the age of seventeen, and his progress was such that he graduated with the highest honors, secured a fellowship, and

[1] Born at Richmond, Surrey, August 16, 1821; died at Cambridge, January 26, 1895. For biography, see Macfarlane, *Ten Brit. Math.*, p. 64.

devoted himself to the preparation of a number of important memoirs. Forced to find some remunerative employment, he then took up the law, and for fourteen years made a specialty of conveyancing, devoting his leisure to the preparation of further scientific memoirs. Sylvester was at this time an actuary in London, and the two were close friends and were in frequent consultation. About 1860 the Sadlerian professorship of pure mathematics was established at Cambridge,[1] and Cayley



FROM AN AUTOGRAPH LETTER FROM CAYLEY TO SYLVESTER
The Miller referred to is W. J. C. Miller, for many years the editor of the mathematical columns in the *Educational Times*, London

was the first to occupy the chair (1863). Although he wrote but one extensive work, the *Treatise on Elliptic Functions* (Cambridge, 1876), he contributed a large number of important memoirs to various scientific publications. In 1889 the Cambridge University Press began the publication of his papers, nearly a thousand in number, in collected form. Seven volumes appeared under his own editorship, the remaining six volumes being published under the supervision of Professor Forsyth.

Cayley's papers cover a very wide range, but it may be said that his chief interest was in the fields of elliptic functions, the theory of invariants, and analytic geometry. Of these the theory of invariants was the one which he did most to advance.

[1] Through funds bequeathed originally by Lady Sadler to found certain lectureships (1710).

America is indebted to him for his course of lectures at Johns Hopkins University in 1882 on the Abelian and theta functions, whereby he again coöperated with Sylvester, who was then helping to place mathematics in this country on a university basis. Well might Sylvester say of him that whatever he touched he embellished.[1]

H. J. S. Smith. Less well known because working in a narrower field, but in the same class of genius as Sylvester and Cayley, Henry John Stephen Smith[2] began his education under the care of his mother, a woman of unusual ability. In 1841 he went to Rugby, where he came under the influence of Dr. Arnold. From Rugby he went (1844) to Balliol College, Oxford, later spending some time at the Sorbonne and the Collège de France. In 1860 he became Savilian professor of geometry at Oxford and a year later was made a Fellow of the Royal Society. His time was so taken up in public duties of various kinds that he did not achieve the success in mathematics of which he was capable. His interest in this science was chiefly in the direction of the theory of numbers and the study of binary and ternary quadratic forms. It was probably in relation to a problem in this theory that he is said to have remarked, "It is the peculiar beauty of this method, gentlemen, and one which endears it to the really scientific mind, that under no circumstances can it be of the smallest possible utility." His various writings were collected by Dr. Glaisher and published in 1894.[3]

Clifford. Among the most promising mathematicians produced in England in the 19th century, but one whose early death prevented the maturing of his genius, was William Kingdon Clifford.[4] He was educated at King's College, London

[1] "Cayley, of whom it may so truly be said, whether the matter he takes in hand be great or small, 'nihil tetigit quod non ornavit.'" *Phil. Trans.*, XVII (1864), 605.

[2] Born at Dublin, November 2, 1826; died at Oxford, February 9, 1883. For biography see Macfarlane, *Ten Brit. Math.*, p. 92.

[3] *The Collected Mathematical Papers of H. J. S. Smith*, 2 vols., Oxford, 1894.

[4] Born at Exeter, May 4, 1845; died in Madeira, March 3, 1879.

(1860–1863) and Trinity College, Cambridge. At the university his mathematical genius was at once recognized, and in 1868 he was elected to a fellowship at Trinity. In 1871 he was made professor of applied mathematics at University College, London, and three years later was elected a Fellow of the Royal Society. He was among the first to protest against the analytic bias of the Cambridge mathematicians, and he assisted in introducing into England the graphic methods of Möbius and other German writers. His most important works were in relation to Riemann's surfaces, biquaternions, and the classification of loci. His *Common Sense of the Exact Sciences* is a classic on the foundations of mathematics, and suggests, as other works (including those of Copernicus and Kepler) had already done, the idea of relativity in all physical measurements. His *Mathematical Papers*, edited by R. Tucker, appeared in 1882.[1]



WILLIAM KINGDON CLIFFORD
From a photograph made shortly before his death

Todhunter. Of all the English mathematicians of this period the one most widely known to elementary students and to teachers is Isaac Todhunter.[2] His name is familiar to everyone who has studied the history of textbook making in the 19th century, although he was much more than a textbook writer. As a young man he attended evening classes at the University of London, where he came under De Morgan's influence, receiving the B.A. degree in 1842 and the M.A. degree two years later. He then entered St. John's College, Cambridge, and

[1] These contain (pp. xv, xxxiii) a brief biography.

[2] Born at Rye, Sussex, November 23, 1820; died at Cambridge, March 1, 1884. See Macfarlane, *Ten Brit. Math.*, p. 134.

in 1848 took his second B.A. degree. He remained at Cambridge until 1864, beginning the textbook writing which made him financially independent and resulted in a series of works that exercised great influence on education in all the English-speaking world. He also wrote (1865) a work on the history of probability and one (1873) on the history of the mathematical theories of attraction, each a classic in its line. He was a good mathematician but not a great one, an excellent linguist, and a man who stood for sound scholarship. As is commonly the case with men in his line of work, numerous stories are told of him, one being that he used to remark that he knew two tunes, the first of which was "God save the Queen" and the second wasn't, and that he recognized the former by the fact that people stood up when it was sung.

Other British Mathematicians. Among the other British mathematicians of prominence in this period it is possible at this time to mention only a few of those whose names should be familiar to the general student of mathematics. Others will be found in the second volume of this work. Dr. George Berkeley (1684–1753) is known in the history of the calculus for his work, *The Analyst* (London, 1734), in which he attacked the foundations of the new science; Edward Waring (1734–1798) was interested in the theory of numbers, and a theorem relating to powers is known by his name; Sir James Ivory (1765–1842), with mathematics as an avocation, did much to advance the progress of analytic methods in England and contributed to the theory of attraction; James Booth (1806–1878) wrote on modern geometry; Sir John Frederick William Herschel (1792–1871) contributed to the study of analysis but finally followed in his father's steps and devoted himself to astronomy; James MacCullagh (1809–1846) contributed to the theory of quadric surfaces; Thomas Penyngton Kirkman (1806–1895) made the attempt to extend the theory of quaternions and was interested in the subject of "analysis situs"; George Biddel Airy (1801–1892), Astronomer Royal of England, contributed to the lunar and planetary theory; John Couch Adams (1819–1892), independently of the French

astronomer Leverrier, determined mathematically the position of the planet Neptune; Sir George Howard Darwin (1845-1912), son of Charles Darwin the naturalist, contributed to the theory of three bodies; Sir Robert Stawell Ball (1840–1913), Astronomer Royal of Ireland and later Lowndean Professor of Astronomy and Geometry at Cambridge, wrote on the theory of screws (1876); Peter Guthrie Tait (1831–1901), sometime professor at Belfast and later at Edinburgh, is well known for his work in quaternions and physics; Lord Kelvin (William Thompson, 1824–1907) contributed extensively to the application of mathematics to physical problems; James Clerk Maxwell (1831–1879) is especially known for his application of mathematics to the study of electricity; and Lord Rayleigh (John William Strutt, 1842–1919) is similarly known with respect to mathematics and the study of vibrations.[1]

3. France

Nature of the Work. France took the lead again in the 18th century, as she did in the first half of the century preceding. She may have had no more brilliant intellects than England, but she found in the differential and integral calculus a set of tools that she could use more deftly than the British mathematicians could use the heavy machinery of the calculus of fluxions. It was perhaps as well that the two nations should experiment on different lines of approach, and it is evident that each cultivated its own peculiar power of attack by so doing, but the total results secured in France show for more than those produced in Great Britain.

Early Eighteenth Century Writers. There will first be mentioned a few names of those who, although born in the 17th century, completed their work after its close. Among the first of these writers was Pierre Varignon,[2] professor of mathematics at the Collège Mazarin (1688) and later at the Collège Royal, and

[1] For a list of biographies of mathematicians dying between 1881 and 1900, see G. Enestrom, *Bibl. Math.*, II (3), 326, covering all European countries.

[2] Born at Caen, 1654; died in Paris, December 22, 1722.

a member of the Académie des Sciences at Paris. Although intended for the Church, he accidentally came across a copy of Euclid, and thus was led, as so many others have been, to the study of mathematics. He then read Descartes's *Géométrie*, and thereafter devoted himself to the mathematical sciences, with special emphasis upon physical problems. He was one of the first of the French scholars to recognize the value of the new calculus. His chief contributions were to the science of mechanics, although he wrote upon pure mathematics as well.[1]

A little younger than Varignon, although dying before him, Pierre-Rémond de Montmort[2] rose to a position of some prominence. He was born to fortune and thus had ample means to enable him to follow his tastes in the study of law and philosophy. He was made a canon of Notre-Dame at a time when piety was not the chief qualification, but finally married and devoted the rest of his life to travel, to Paris, and to mathematics. He was chiefly interested in the doctrine of chance,[3] a subject which brought him into cordial relations with De Moivre and with Jean and Nicolas Bernoulli. He also wrote on infinite series[4] and summed to n terms the series

$$S = na + \frac{n(n-1)}{2!}\Delta a + \frac{n(n-1)(n-2)}{3!}\Delta^2 a + \cdots.$$

Antoine Parent,[5] a private teacher of mathematics in Paris and (1699) a member of the Académie des Sciences, was another of the minor writers who helped to advance his subject. His interest was chiefly in mechanics and physics, although he also wrote on arithmetic, the cycloid, geometry, and perspective. He is known in the history of mathematics for his

[1] *Éclaircissements sur l'analyse des infiniment petits, Paris*, 1725 (posthumous); *Éléments de mathématiques*, Paris, 1731 (posthumous); *Manière de trouver une infinité de portions de cercle toutes quarrables moyennant la seule géométrie d'Euclide*, Paris, 1703; and other works.

[2] Or Monmort. Born in Paris, October 27, 1678; died in Paris, October 7, 1719. His family name was Rémond, the "de Montmort" being assumed from his estates.

[3] *Essai d'anlyse sur les jeux de hasard*, Paris, 1708; 2d ed., 1714.

[4] "De seriebus infinitis tractatus," in the *Philosophical Transactions* for 1717.

[5] Born in Paris, September 16, 1666; died in Paris, September 26, 1716.

work in analytic geometry of three dimensions.[1] His most important contributions were published in his collected works in 1705.[2]

Among Parent's contemporaries, Joseph Saurin,[3] a priest, wrote on the determination of tangents at multiple points of an algebraic curve, on the curve of least descent, and on various other geometric questions.

There was also Thomas-Fantel de Lagny,[4] who gave up the law for the purpose of devoting himself to mathematics. He wrote on new methods of extracting roots (1692), the cubature of the sphere (1702), binary arithmetic (1703), and methods of solving problems.[5] The story is told that Maupertuis, called to his deathbed and finding him in a comatose state, asked him suddenly for the square of 12; whereupon De Lagny started up, gave the answer, and at once passed away.

Less of a mathematician but contributing worthily to the science, Amédée François Frézier,[6] a French infantry officer (1702–1707) and later an engineer in South America and San Domingo, by means of his works on stereometry as applied to stone cutting and architecture[7] laid part of the foundation for the theory of descriptive geometry.

Among those who formed a brilliant group in Paris at this time was one of the youthful prodigies that, as we have seen, arise from time to time,—François Nicole.[8] He was a boy of unusual promise, having shown his genius in geometry by rectifying the cycloid at the age of nineteen. His interest in the

[1] In a paper read before the Académie in 1700. See also Chasles, *Aperçu historique*, p. 138.

[2] *Recherches de mathématiques et de physique*, Paris, 1705; rev. ed., 3 vols., Paris, 1713.

[3] Born at Courthézon, Vaucluse, September 1, 1655; died in Paris, December 29, 1737.

[4] Born at Lyons, November 7, 1660; died in Paris, April 12, 1734.

[5] *Analyse Générale, ou Méthodes Nouvelles pour résoudre les Problèmes de tous les Genres et de tous les degrez à l'Infini*, Paris, 1733.

[6] Born at Chambéry, 1682; died at Brest, October 16, 1773.

[7] *La théorie et la pratique de la coupe des pierres et des bois, ou Traité de Stéréotomie*, Strasburg, 1738; 2d ed., 3 vols., Paris, 1754, 1768, 1769; *Éléments de Stéréotomie, à l'usage de l'architecture*, Paris, 1750–1760.

[8] Born in Paris, December 23, 1683; died in Paris, January 18, 1758.

study naturally led to a consideration of roulettes in general,[1] a subject in which he showed great insight. He also wrote on the calculus of finite differences,[2] lines of the third order (1729), probability (1730), conics (1731), cubics (1738, 1741, 1743), and the trisection problem (1740). As in most cases of unusually early development, however, his work was not of the highest order.



PIERRE LOUIS MOREAU DE MAUPERTUIS

For some years a favorite at the court of Frederick the Great, interested chiefly in geodesy


Maupertuis. The work of Pierre Louis Moreau de Maupertuis[3] was of a more stable kind. In his younger days (1718) he was a captain of dragoons in the French army, but he later retired to private life and devoted himself to the study of mathematics. He was made a member of the Académie des Sciences in 1731, directed the measurement of a meridian degree in Lapland in 1736, became president of the physical class in the Berlin Academy (1745–1753), basked in the sunshine of the favor of Frederick the Great for the usual brief period, learned that he could "climb, but heights are cold," and after falling from favor spent the last six years of his life in his native country. His chief work was in astronomy and geodesy,

[1] *Méthode générale pour déterminer la nature des courbes formées par le roulement de toutes sortes de courbes sur une autre courbe quelconque*, Mém., Paris, 1707; *Manière de déterminer la nature des roulettes formées sur la superficie convexe d'une sphère*, Paris, 1708, 1722.

[2] *Traité du calcul des différences finies*, Paris, 1717, 1723, 1724.

[3] Born at St. Malo, July 17, 1698; died at Basel, July 27, 1759.

but he also wrote on maxima and minima (1724), quadrature problems (1727), curves in general (1727–1730), and various physical questions.[1] He taught mathematics to his friend the Marquise du Châtelet and, considering her friendship for Voltaire, deserved something better than the harsh treatment which the latter gave him in his *Diatribe du Dr. Akakia*, written with



AUTOGRAPH LETTER OF MAUPERTUIS

This letter was written to Frederick the Great in 1750, while Maupertuis was still president of the physical class in the Berlin Academy

the desire to defend a learned but indiscreet Swiss mathematician, Samuel Koenig (died 1757), who had accused Maupertuis of plagiarism. One of the biographers of Voltaire speaks of Maupertuis as "the pompous and touchy mathematician," and the phrase is probably appropriate.

Minor Writers. Alexis Fontaine des Bertins[2] was more promising in his youth but less successful in his later years than his contemporary, Maupertuis. He was a man of means, was

[1] *Œuvres de Mr. de Maupertuis*, 4 vols., Paris, 1752; Lyons, 1768.

[2] Born at Bourg-Argental, Loire, *c.* 1705; died at Cuiseaux, August 21, 1771.

therefore able to devote his life uninterruptedly to study, and became a member of the Académie des Sciences in 1733, when only about twenty-eight years of age. His early promise was not fulfilled, however, his efforts not being directed in lines of probable success. He wrote on tautochronous curves (1734 and 1768) and differential equations, and proposed various problems in geometry and astronomy. He suggested the common notation of partial derivatives of a function of several variables.

Among his contemporaries, and a man of considerable influence in the scientific circle of Paris, there should be mentioned Jean Paul de Gua de Malves.[1] He belonged to a family that had been impoverished by John Law's Mississippi scheme, and, seeing no career open to him, he entered the Church and secured a benefice which enabled him to live comfortably and to devote his life to study. He wrote a work on the Cartesian analysis[2] which gave him admission to the Académie des Sciences (1740) and a professorship (1743) of philosophy in the Collège de France. He seems to have suggested the idea of the *Encyclopédie* which was finally carried out by Diderot, d'Alembert, and Voltaire. He perfected the proof of Descartes's Law of Signs (1741) and wrote on geometry and trigonometry.

Clairaut Family. One of several noteworthy instances in the history of mathematics, showing the influence of heredity or early environment, is seen in the case of the Clairaut family. Jean Baptiste Clairaut[3] was a teacher of mathematics in Paris about the middle of the 18th century. He was a correspondent of the Berlin Academy and published three memoirs on geometry in the *Miscellanea* (1734, 1737, 1743). One of his sons was Alexis Claude Clairaut,[4] the most prominent member of the family, an infant prodigy who read l'Hospital's *Analyse des*


[1] Born at Carcassonne, *c.* 1712; died in Paris, June 2, 1786.

[2] *Usage de l'analyse de Descartes pour découvrir, sans le secours du calcul différentiel, les propriétés . . . des lignes géométriques de tous les ordres*, Paris, 1740.

[3] Died soon after 1765. The name is also spelled Clairault.

[4] Born in Paris, May 7, 1713; died in Paris, May 17, 1765.

infiniment petits and his *Traité des sections coniques* at the age of ten, presented a paper on geometry before the Académie des Sciences when he was thirteen, and was admitted to membership in the Académie and published a work on curves of double curvature[1] when he was only eighteen. His solutions of the



AUTOGRAPH OF ALEXIS CLAUDE CLAIRAUT

Written about twenty years before his death and while he was working on his algebra

problem of tangents drawn to such curves and of the quadrature of the curves themselves are still found in current treatises. Clairaut was only twenty-three when he was made a member of the commission which went to Lapland to measure the length of a degree. He now began to devote most of his attention to problems of celestial mechanics, but still found time to write on geometry (1741), algebra (1746), algebraic

[1] *Recherches sur les courbes à double courbure*, Paris, 1731.

curves on a cone (1732), maxima and minima (1733), the calculus (1739), and similar topics, most of his contributions being in the form of memoirs presented to the Académie. He demonstrated Newton's theorem that all curves of the third order are projections of one of five parabolas. Possibly it was his interest in Newton that first brought him under the spell of the Marquise du Châtelet in the months just preceding her death, when she was hastening to complete her translation of the *Principia*.

Arago, in his eulogy of Laplace, remarked:

> Five geometers—Clairaut, Euler, d'Alembert, Lagrange, and Laplace—shared among them the universe of which Newton had disclosed the existence. They explored it in all directions, penetrated into regions which had been thought inaccessible, pointed out there a multitude of phenomena which observation had not yet detected, and finally, and herein lies their imperishable glory, they brought within the domain of a single principle, a single law, all that is most refined and mysterious in the movements of the celestial bodies.

Alexis had a brother[1] who died when he was sixteen, but who at the age of fourteen had read a memoir on geometry before the Académie des Sciences, and who published a work on geometry[2] when he was only fifteen.

Voltaire and the Marquise du Châtelet. The world does not often connect the name of Voltaire with mathematics, and when it connects that of the Marquise du Châtelet with the science, it is largely by courtesy. Each, however, did something to make the Newtonian theory known, and each absorbed enough mathematics to make the labor fairly serious.

François Marie Arouet,[3] known tc the world as Voltaire and as the foremost leader of the 18th century in the contest for human liberty,[4] was interested in mathematics chiefly because he was interested in all things English, was interested in Newton,

[1] Born in Paris, 1716; died in Paris, 1732.

[2] *Traité des quadratures circulaires et hyperboliques*, Paris, 1731.

[3] Born in Paris, November 21, 1694; died in Paris, May 30, 1778.

[4] Among his monographs is the *Essai sur les Probabilités en fait de Justice*.

was interested in getting out a work on Newton's philosophy,[1] and was interested in Émilie, Marquise du Châtelet.[2] Daughter of the Baron de Breteuil, the marquise married at nineteen, and turned her brilliant mind to Euclid, to Newton, to the literary classics of Greece and Rome, to Locke, and to Voltaire. She had studied mathematics under Maupertuis and Koenig, read Newton and understood him,—at least in part,—and in due time translated the *Principia*,[3] completing it a few days before her death. Frederick the Great, who loved an epigram far more than he loved the courtesies of life, suggested this epitaph: "Here lies one who lost her life in giving birth to an unfortunate child and to a treatise on philosophy." Madame du Châtelet also wrote on physics, but at best she was only an amatrice in science.[4]



ÉMILIE, MARQUISE DU CHÂTELET
After a lithograph from a contemporary drawing by N. H. Jacob

[1] *Élémens de la philosophie de Neuton*, Amsterdam, 1738.

[2] Gabrielle Émilie Le Tonnelier de Breteuil; born in Paris, December 17, 1706; died at Commercy, September 10, 1749.

[3] It was published posthumously at Paris in 1759. There is a bibliography of her works in A. Rebière, *Les femmes dans la science*, 2d ed., p. 65 (Paris, 1897).

[4] Voltaire, in one of his many epigrams about her, wrote:

"Son esprit est très philosophe,
Mais son cœur aime les pompons."

In his work on Newton he addresses a poem to her, beginning:

"Tu m'appelles à toi, vaste & puissant Génie,
Minerve de la France, immortelle Emilie,
Disciple de Neuton, & de la Vérité."

D'Alembert. There are certain names in the history of mathematics to which there attaches a special human interest apart from the mere recital of a list of discoveries. One of these is d'Alembert.[1]



JEAN BAPTISTE LE ROND D'ALEMBERT
From an engraving made by P. Maleuvre in 1775 after a drawing made by A. Pujos in 1774

On the night of November 16, 1717, a gendarme, while making his rounds in Paris, found near the church of Saint-Jean le Rond a newly born infant who had been abandoned to the fate of winter, and had him hurriedly christened with the name of his first resting place, Jean Baptiste le Rond. Foster parents were found and Jean grew up, known but unrecognized by his mother, pitied and somewhat helped by his father, and soon showed remarkable intellectual powers that spoke for intellectual parentage. His mother, Madame de Tencin, sister of a cardinal, has been described by one of d'Alembert's biographers as "small, keen, alert, with a little sharp face like a bird's, brilliantly eloquent, bold, subtle, tireless, a great minister of intrigue, and insatiably ambitious." His father, General Destouches, was a man of large heart, and at his death in 1726 left enough to provide for the boy's education. When Jean was eighteen (1735) he took his bachelor's degree and soon, for reasons unknown, adopted the name of d'Alembert. He prepared for the bar, then took up medicine, and

[1] Born in Paris, November 16, 1717; died in Paris, October 29, 1783.

finally devoted his life to mathematics. Friend of Voltaire, collaborator on the *Encyclopédie*, admirer of Madame du Deffand, and lover of her companion Mademoiselle Julie de Lespinasse, he knew those in France who were best worth knowing and experienced all the joys and sorrows that Paris affords. One of his biographers says:

> In himself d'Alembert was always rather a great intelligence than a great character. To the magnificence of the one he owed all that has made him immortal, and to the weakness of the other the sorrows and the failures of his life. For it is by character and not by intellect the world is won.[1]

D'Alembert wrote upon mathematics in general,[2] the calculus and its applications,[3] the theory of differential equations,[4] and dynamics.[5]

Minor Writers. Among the minor writers of the middle of the 18th century one of the best known is Johann Heinrich Lambert.[6] Born in humble surroundings, leading a roving life, acting as bookkeeper, secretary, private tutor, and architect, and living in Germany, Holland, France, Italy, and Switzerland, he was, in spite of such an unsettled life, a voluminous writer, his fields of interest being as varied as his occupations and his places of abode. He wrote on perspective, light, astronomy, logarithms, pyrometry, transcendent quantities, theory of equations, the slide rule, psychology, ballistics, photometry, and a variety of other subjects, most of his efforts displaying respectable mediocrity,—all save one, hyperbolic trigonometry, and this gave him an enduring place in history.[7]

[1] S. G. Tallentyre, *The Friends of Voltaire*, chap. i, London, 1907.

[2] *Opuscules mathématiques*, 8 vols., Paris, 1761–1768.

[3] In the memoirs of the Berlin Academy, 1746 and other dates.

[4] Cantor, *Geschichte*, II, chap. 118. [5] J. Bertrand, *D'Alembert*, Paris, 1889.

[6] Born at Mulhouse, Alsace, August 26, 1728; died at Berlin, September 25, 1777. Mulhouse was then Swiss territory, so that he may also properly be ranked among Swiss scholars.

[7] For bibliography see Engel and Stäckel, *Die Theorie der Parallellinien*, p. 151 (Leipzig, 1895). Consult also F. Rudio, *Archimedes, Huygens, Lambert, Legendre*, Leipzig, 1892; D. Huber, *Johann Heinrich Lambert*, Basel, 1829; J. Lepsius, *Johann Heinrich Lambert*, Munich, 1881; F. Schur, *Johann Heinrich Lambert als Geometer*, Karlsruhe, 1905.

Alexandre Théophile Vandermonde,[1] member (1771) of the Académie des Sciences at Paris and director (1782) of the Conservatoire des Arts et Métiers, was another of the relatively minor writers of this period. He contributed to the theory of equations through two memoirs (1771, 1772), and to the general theory of determinants.

Étienne Bézout[2] was also one of the writers of this period on the theory of equations. He was an examiner for the navy and is known for several memoirs and textbooks. He was among the first to recognize the value of determinants. His method of elimination by the aid of symmetric functions (1764 and 1779) is well known to students in the theory of equations.



ANTOINE-NICOLAS CARITAT
MARQUIS DE CONDORCET
After an engraving from a drawing from life

During the Reign of Terror the revolutionists spared most of those whose mathematical genius is now recognized, but they did so reluctantly in the case of Marie-Jean-Antoine-Nicolas Caritat, Marquis de Condorcet.[3] Brought up under the Jesuits, he admired their learning and hated their doctrines. He was admitted to the Académie des Sciences when only twenty-six (1769) and at thirty became its secretary. At an age when all his family

[1] Born in Paris, February 28, 1735; died in Paris, January 1, 1796. On his given name see *Zeitschrift* (Hl. Abt.), XLI, 83.

[2] Born at Nemours, March 31, 1730; died in Paris, September 27, 1783.

[3] Born at Ribemont, near Saint-Quentin, September 17, 1743; died near Bourg-la-Reine, in the vicinity of Paris, March 29, 1794. See Arago's *Biographie*, read before the Académie des Sciences, December 28, 1841.

was demanding that he should be a captain of cavalry he was making for himself a name by his essay on the integral calculus and by his work on the problem of the three bodies.[1] He then took up the theory of differential equations, wrote extensively on the calculus, applied himself to the study of probability, wrote various eulogies on deceased academicians which are still read as classics in French literature, and lived the life of a scholar and, in Voltaire's words, of "the man of the old chivalry and the old virtue." D'Alembert spoke of him as a volcano covered with snow; in other words, he was an intellectual aristocrat, and that was enough to condemn him in the days of 1794. "If I have one night before me," he had said, "I fear no man; but I will not be taken to Paris." When the jailer to whom the gendarme had taken him for the night opened the door of his cell, Condorcet was dead, with an empty poison ring by his bed. He had kept the faith with himself.

Lagrange. Joseph Louis, Comte Lagrange,[2] was Tourangean by descent, Italian by birth, German by adoption, and Parisian by choice. He began his teaching as professor of mathematics in the artillery school at Turin (1755) when only nineteen years of age, succeeded Euler (1766) as mathematical director in the Berlin Academy, and was called to Paris (1787), where he became a member of the Académie des Sciences and, somewhat later (1795), was professor of mathematics at the newly founded École Normale and (1797) at the École Polytechnique. Under Napoleon he was made a senator and a count, and was awarded other honors appropriate to his genius.

Lagrange was not one of the infant prodigies in mathematics. Indeed, it is said that he showed no interest in the

[1] *Essai sur le calcul intégral*, Paris, 1765; *Analyse de la solution du problème des trois corps*, Paris, 1768, the memoir which first called attention to his powers.

[2] Born at Turin, January 25, 1736; died in Paris, April 10, 1813. As an Italian by birth, the name might be given as Giuseppe Luigi; but the family was originally French, affiliated to that of Descartes, and Lagrange spent the best years of his life in France. His complete works were published in Paris, 14 vols. (1866–1892), with a biography by Delambre in Volume I. See also G. Loria, "G. L. Lagrange nella vita e nelle opere," in the *Annali di Matematica pura ed applicata*, XX (3), p. ix.

subject until he was seventeen; but from that time on he made such marvelous progress that in a few years he became recognized as the greatest living scholar in his science. When he was twenty-three years old he published two memoirs[1] which at once attracted attention. Euler wrote (October 2, 1759) an enthusiastic letter to him about the problem of isoperimetry which is here solved and on which the great Swiss mathematician had long been working, and d'Alembert was equally appreciative of its importance. It is here that we find the beginning of the calculus of variations, and it is here that Lagrange took the first step toward Berlin and Paris, although it was not until 1766 that Frederick the Great wrote that "the greatest king in Europe" wanted "the greatest mathematician of Europe" at his court. As a result of this letter, Lagrange went to Berlin and remained there more than twenty years. At about the time that Frederick was urging him to go to Berlin he solved Fermat's problem relating to the equation $nx^2 + 1 = y^2$, n being integral and not a square,[2] an intellectual feat that added greatly to his reputation. He now began a series of investigations on partial differential equations, numerical equations, the theory of numbers, the calculus of variations, and the application of mathematics to physical problems, and made some progress in the theory of elliptic functions. To one of his memoirs (1773) may be traced the first important step in the theory of invariants, and in another there is evidence that the notion of a group was in his mind. At this time, too, he composed his monumental work on analytic mechanics,[3] although this was not published until a year after he left Berlin.


The death of Frederick (1787) brought many changes to Prussia. Lagrange, whose frail constitution had never found the climate of Berlin salutary, and whose sensitive nature now

[1] *Recherches sur la méthode de maximis et minimis*, Turin, 1759; *Sur l'intégration d'une équation différentielle à différences finies, qui contient la théorie des suites récurrentes*, Turin, 1759. These appeared in the *Miscellanea Taurinensia*.

[2] "Sur la solution des problèmes indéterminés du second degré," published in the *Miscellanea Taurinensia* in 1767.

[3] *Mécanique analytique*, Paris, 1788.

found the intellectual atmosphere far from agreeable, decided to accept the invitation of Louis XVI to take up his residence in Paris. It was about the time of the agitation for the



DOCUMENT SIGNED BY LAGRANGE

This official document was written by Laplace and was signed by Lagrange and himself

metric system, and Lagrange was made president of the commission to carry out the work. The value of such an undertaking could appeal even to a Sans-culotte, and so, although all foreigners were banished from France, the Committee of Public Safety expressly excepted Lagrange from the decree. Nevertheless the fate of Lavoisier and Bailly, both of whom

met their death by the guillotine, led Lagrange to decide on leaving France. He spoke bitterly to Delambre of the death of Lavoisier, saying that the mob had removed in an instant a head that it would take a century to reproduce. Prussia knew his genius and seriously wanted him back; Paris knew his name and vaguely wished him to remain. The Prussia of that day wished to be scientific; the Paris of that day merely wished to be thought so. But just as Lagrange was reaching a decision, new forces were created in France, and these forces were more potent than any fear of the guillotine, than any discouragement at the acts of the revolutionary leaders, or than any call of the successor of Frederick. France had decided to establish a school with the humble name of École Normale, and a little later she established a second one, the École Polytechnique, and to each school Lagrange was called and to each he gave a mathematical impetus that it has never lost. In the first he saw a chance to found the training of teachers on the most thorough scholarship, and no similar institution either before or since that time has so thoroughly recognized the value of this principle, and none has ever stood so high in the esteem of the world. Similarly, in the École Polytechnique he saw the opportunity for basing the technical work on a foundation of the highest type of mathematical skill, and this institution, like the other, has ever since been a constant inspiration to the world of science. It was at the École Normale that he gave those lectures on algebra and arithmetic[1] that he had to temper to the revolutionary demand before he could bring the work up to the standard that permitted him to present the calculus of functions.[2] It was at the École Polytechnique that he lectured on analytic functions,[3] setting forth in new fashion the differential and integral calculus and the calculus of fluxions. Here, too, he expounded his noteworthy work on numerical higher equations.[4]

[1] *Leçons d'arithmétique et d'algèbre*, 1794–1795.

[2] *Leçons sur le calcul des fonctions*, Paris, 1801.

[3] *Théorie des fonctions analytiques contenant les principes du calcul différentiel*, Paris, 1797.

[4] *Traité de la résolution des équations numériques de tous degrés*, Paris, 1798.

It is probable that his work more profoundly influenced later mathematical research than did that of any of his contemporaries, although it was an era of giants in this field.

Laplace. Pierre-Simon, Marquis de Laplace,[1] was born in poverty and owed his early education to the interest which his promise excited in men of intellectual power. Of these days of struggle he never spoke. Almost the first reliable records that we have of his life show him studying and afterwards teaching mathematics in the military school at Beaumont and making such a reputation as to lead to his call to succeed (1784) Bézout as examiner of the artillery corps. He later took part in the organization of the École Polytechnique and the École Normale. Napoleon made him a count and appointed him minister of the interior (1799). After standing his eccentricities for six months, the consul dismissed him with the remark that he carried into his work the spirit of the infinitesimal.[2] Laplace then entered the senate but made no worthy record. After the restoration, Louis XVIII raised him to the peerage and (1817) made him a marquis.



PIERRE-SIMON LAPLACE
From Goutière's engraving after a painting by Naigeon

[1] Born at Beaumont-en-Auge, Calvados, March 23, 1749; died in Paris, March 5, 1827. For Arago's eulogy on Laplace, see the English translation in the Smithsonian Institution *Report* for 1874, p. 129 (Washington, 1875).

[2] "L'esprit des infiniment petits."

Laplace was a political opportunist. At heart he was a royalist, but for his personal interests he became a follower of Napoleon. He was a friend of the people but not a believer in the people's judgment.

His name is chiefly connected with astronomy and celestial mechanics,[1] but he also wrote on probability,[2] the calculus, differential equations, and geodesy.[3] As a master of the theory of celestial mechanics he stands unrivaled.[4] As to his style of exposition, Nathaniel Bowditch (1773–1838), the self-made American astronomer, remarked: "I never come across one of Laplace's 'Thus it plainly appears' without feeling sure that I have hours of hard work before me to fill up the chasm and find out and show how it plainly appears."[5]

Legendre. The third of the great trio, of which the first two were Lagrange and Laplace, appeared in the person of Adrien-Marie Legendre.[6] He was educated at the Collège Mazarin in Paris, where he early showed his taste for mathematics, and with the help of his teacher, the Abbé Marie,[7] and of d'Alembert, he became (1775) professor of mathematics in the École Militaire at Paris, resigning in 1780. Two years later (1782) he won the prize of the Berlin Academy for his essay on the path of a projectile.[8] In elementary mathematics

[1] *Exposition du système du monde*, 2 vols., Paris, 1796; *Traité de mécanique céleste*, 5 vols. and suppl., Paris, 1799–1825.

[2] *Théorie analytique des probabilités*, Paris, 1812; *Essai philosophique sur les probabilités*, Paris, 1814.

[3] His collected works were published in seven volumes in Paris in 1843–1847. A later and better edition was published in fourteen volumes by the Académie des Sciences, 1878–1912.

[4] With that felicity of speech which characterizes the French, Fourier enumerated his great discoveries, and added: "Voilà des titres d'une gloire véritable, que rien ne peut anéantir. Le spectacle du ciel sera changé; mais à ces époques reculées, la gloire de l'inventeur subsistera toujours; les traces de son génie portent le sceau de l'immortalité."

[5] For an appreciation of Laplace and for his influence upon the century following, see R. S. Woodward, *Bulletin of the Amer. Math. Soc.*, V, 133.

[6] Born at Toulouse, September 18, 1752; died in Paris, January 10, 1833.

[7] Joseph François Marie (1738–1801), who edited Lacaille's *Tables de Logarithmes* (1768) and his *Leçons élémentaires de Mathématiques* (1798), which had appeared in 1760 and 1741 respectively.

[8] *Recherches sur la trajectoire des projectiles dans les milieux résistants*

he is known chiefly for his geometry,[1] a work which had a generous reception in various countries and which justly ranks as one of the best textbooks ever written upon the subject. In it he sought to rearrange the propositions of Euclid, separating the theorems from the problems and simplifying the proofs, without lessening the rigor of the ancient methods of treatment. To Legendre is largely due the abandoning of Euclid as a textbook in American schools.



ADRIEN-MARIE LEGENDRE
After a lithograph by Delpech

In higher mathematics Legendre is known for his works on the theory of numbers[2] and on elliptic functions.[3] He is also known for his treatises on the calculus, higher geometry, mechanics, astronomy, and physics. To him is due the first satisfactory treatment of the method of least squares,[4] although Gauss had already discovered the method. In his theory of numbers appears the law of quadratic reciprocity which Gauss called the "gem of arithmetic." The treatise on elliptic functions appeared almost simultaneously with the works by Abel and Jacobi on the same subjects; and although Legendre had spent thirty years on the theory, he

[1] *Éléments de géométrie*, Paris, 1794.

[2] *Essai sur la théorie des nombres*, Paris, 1798; 2d ed., Paris, 1808, with supplements in 1816 and 1825; 3d ed. under the title *Théorie des nombres*, Paris, 2 vols., 1830.

[3] *Mémoire sur les transcendantes elliptiques*, Paris, 1794; *Traité des fonctions elliptiques et des intégrales eulériennes*, Paris, 1827–1832.

[4] On the history of this subject see M. Merriman, *Method of Least Squares*, p. 182 (New York, 1884); *Transactions of Connecticut Academy*, IV, 151 (1877), with complete bibliography to that date; Todhunter, *Hist. Probability*, Cambridge, 1865.



Institut National

Classe des Sciences Physiques et Mathématiques

Paris, le an XIV de la République française

Le ~~Secrétaire perpétuel pour les Sciences~~ Président de la Classe des Sciences Physiques et Mathématiques

à Monsieur le président de l'Institut

Monsieur et cher Confrère

Je m'empresse de vous informer que la Classe des Sciences physiques et Mathématiques a adopté unanimement le projet d'arrêté délibéré dans l'assemblée des Bureaux. Elle a pensé en même temps qu'il suffisait que cet objet fut rapporté à l'assemblée générale prochaine de l'Institut sans convoquer aucune séance extraordinaire à cet effet.

Agréez, Monsieur et cher Confrère, l'assurance de la Considération distinguée avec laquelle j'ai l'honneur de vous salue

Le Gendre

AUTOGRAPH LETTER OF LEGENDRE

In some of his letters the form "Le Gendre" appears, as in this case. In general the name is spelled Legendre

recognized at once the superiority of the treatment given to it by these younger men, and posterity has agreed with his judgment.

Failing to yield to the government in its desire to dictate to the Académie, he was deprived of his pension, and his last days were spent in poverty. His letters of this period are depressing, showing how one of the greatest scientists of France had lost heart at the failure of a nation to recognize his honesty of purpose and his powers of intellect.



GASPARD MONGE, COMTE DE PÉLUSE
After a lithograph of a drawing by Hesse

Beginning of the Nineteenth Century. Of those who made France a great mathematical center in Napoleon's day, Gaspard Monge[1] was, after Lagrange, Laplace, and Legendre, one of the leaders. He was the son of an itinerant tradesman, and was one of many in the history of science who early showed promise of success. At the age of fourteen he constructed a fire engine which was put into service, and at sixteen he was teaching in a secondary school (*collège*) in Lyons. At twenty-two he was professor of mathematics in the military school at Mézières, and from this time on he continued to progress, with the excitement of just escaping the guillotine, until he reached a professorship (1794) in the École Polytechnique. He also became a member of Napoleon's staff in Egypt, a senator

[1] Born at Beaune, May 10, 1746; died in Paris, July 28, 1818. See M. Brisson, *Notice historique sur Gaspard Monge*, Paris, 1818; F. Arago, "Gaspard Monge," in Arago's *Œuvres complètes*, II, 427 (Paris, 1854); Ch. Dupin, *Essai historique sur . . . Monge*, Paris, 1819.

(1799), and, as Comte de Péluse, a member of the nobility. With the restoration, however, all his honors were taken from him, and his last years were a period of disappointment.

He is known chiefly for his elaboration of descriptive geometry, a theory which, as we have seen, was suggested by Frézier in 1738, but which Monge worked out independently while at Mézières. It was some years before anything was published on the subject, the idea being held as a military secret of great value in the designing of fortifications. The opening of the École Polytechnique gave him an opportunity to lecture upon the theory, and finally, in 1799, he published his treatise upon it.[1] He also wrote numerous memoirs on differential equations, curves on various surfaces, and physical problems.

Among the most unfortunate scholars on the roll of the world's mathematicians there will always rank the name of Pierre-François-André Méchain,[2] a man who rose from the position of private tutor to become one of the leading astronomers of France, charged with duties of greatest importance in connection with the metric system, a collaborator with the great Delambre in the field work on which the units were based, and one who was recognized as a scientist of genuine ability. It was his duty to measure that part of the meridian lying between Rodez and Barcelona. After his report was sent to Paris he discovered that he had made an error of 3″ in the latitude of Barcelona. In his endeavor to conceal this error, which he knew would ruin his scientific reputation, he sought to extend the meridian, cutting out Barcelona altogether, but died from yellow fever while carrying out the plan. Instead of being known as a scientist of repute he is thought of as the man who made the chief mistake in the determining of the standard meter. It should be said, however, that the fault was not really his, for the obstacles placed in his way were such as to make accurate observations almost impossible.

[1] *Géométrie descriptive. Leçons données aux Écoles normales, l'an 3 de la République* (17^4–1795), Paris, l'an VII (1798–1799).

[2] Born at Laon, August 16, 1744; died at Castellon de la Plana, near

Sylvestre François Lacroix,[1] whose work falls in this period, was one of those men who succeed by persevering rather than by distinguished scientific ability. In his early years he occupied various positions in the naval and military schools, but finally became connected with the École Normale, the École Polytechnique, and the Collège de France, positions of highest prominence. He was a voluminous writer on higher algebra, geometry, probability, and the calculus, but he is not known for the original development of any great theory. The translation (1816) of his calculus into English by Charles Babbage, Sir John Frederick William Herschel, and George Peacock, however, did much to introduce the Continental methods and notation into the work of the Cambridge school of mathematicians.




JEAN-BAPTISTE-JOSEPH DELAMBRE
After a drawing made two years before Delambre's death

Delambre. Jean-Baptiste-Joseph Delambre[2] furnishes one of the interesting cases of a man who turned late to the study of mathematics and yet rose to be a leader in the science. As a young student in Amiens he was steeped in the classics, and it was not until he was thirty-six years of age that he seems to have even begun the serious study of astronomy, a subject which required him at the same time to begin his mathematical work. He was forty before he published anything on the

[1] Born in Paris, 1765; died in Paris, May 25, 1843. See the eulogies pronounced at the Institut in 1843 by Libri and Despretz, and various biographical articles of the time.

[2] Born at Amiens, September 19, 1749; died in Paris, August 19, 1822.



mais l'on peut sans erreur sensible faire sin MQm ou sinus de la déviation de la lunette du quart de cercle dirigée sur M = Distance du quart de cercle à la lunette / Distance de la lunette à sa mire.

Soit donc $LQ = 4^M = \frac{2}{3}$ d.m.s. $\quad$ log $\frac{2}{3}$ 9.823991

$MQ = 1000^T$ ..... log 1000 : 3.000000

sin.MQm = sin 2'.17" $\quad$ 6.823991

Comp. sin 1' $\quad$ 5.314443

MQm = 2'.17".51 ..... 2.138434

ou log.15 $\quad$ 1.176091

MQm en mes. = 129".6) 0.962 25

Il suffirait donc de donner au cercle un mouvement azimuthal de 2'.17".5 vers m pour le conduire dans la ligne Mm

en général voici le petit calcul

log. comp. sin 5.314443

log. distance du q.d.c. à la lunette 4^m. 6.602060

Comp. arithm. log. distance mire 6000 pieds ... 6.221853

déviation horizontale 2'.17".51 .... 2.138434

ici le calcul est emploié, et il peut paraître inutile de compter aux distances de 4 pieds et 6000 pieds on substituerait les distances véritables si on connoit à peu près la distance de la mire.

Si on ne la connoit pas, il faudra mettre une seconde mire en m' à une distance Mm = LQ : mais la distance Mm doit être prise sur la perpendiculaire à la méridienne : si le bâtiment avait une déviation sensible il faudrait y avoir égard : si par exemple le bâtiment avoit la direction Mm' la distance Mm' seroit plus grande que Mm : en général $Mm' = \frac{Mm}{\cos. mMm'}$

voilà je crois tout ce qu'on peut dire sur ce petit Problème.

agréez mes remercimens et comptez sur ma fidélité à vous les témoigner

Delambre

AUTOGRAPH LETTER OF DELAMBRE

With computations relating to the survey made for establishing the standard meter

subject,[1] and it was some years later that he was awarded a prize by the Académie for his tables of Uranus.[2] From this time on he was known as one of the leading astronomers of France, and his various works on the history of astronomy are still looked upon as authorities. In the history of elementary mathematics he is chiefly known for his work in measuring the arc of the meridian between Dunkirk and Barcelona for the purpose of establishing the basis for the metric system. He was a scholar, a persistent worker, and a man of highest character.[3]

Carnot. Another interesting illustration of the development of mathematical talent rather late in life is seen in the case of Lazare-Nicolas-Marguerite Carnot,[4] a member of an old and respected family of France. After the manner of so many sons of the well-to-do landowners, he studied for the army, and was thus led to the military school at Mézières. Here he came under the influence of Monge, and thus his tastes were turned toward geometry. He developed into one of the great military leaders of France, held various important offices, voted for the execution of Louis XVI, suffered in the general upheavals of the Revolution, was exiled by Napoleon, and spent his later years in Magdeburg.

His scientific work showed itself in various lines, but especially in his contributions to geometry.[5] It was in these contributions that he assisted in laying the foundations for modern synthetic geometry.

[1] *Tables de Jupiter et Saturne*, Paris, 1789.

[2] Published with other tables in 1792.

[3] The great scientist Cuvier, in his address at the burial of Delambre in the cemetery of Père-Lachaise, said of him: "Qu'il me soit permis, au moment où je vous dis ce triste et dernier adieu, de rendre témoignage à cet admirable caractère que, pendant vingt ans de liaison intime et de rapports journaliers, je n'ai pas vu se démentir un instant. Jamais, pendant ce long intervalle, un seul mouvement n'a troublé votre inaltérable douceur, . . . il ne vous est échappé une parole qui ne fût dictée par la justice et la raison."

[4] Born at Nolay, Côte d'Or, May 13, 1753; died at Magdeburg, August 2, 1823. F. Arago, *Biographie de . . . Carnot*, eulogy delivered before the Académie des Sciences, August 21, 1837.

[5] *Géométrie de Position*, Paris, 1803; *Sur la relation qui existe entre les distances respectives de cinq points quelconques pris dans l'espace, suivi d'un Essai sur la théorie des transversales*, Paris, 1806.

Gergonne and his Time. With this work is also connected the name of Joseph-Diez Gergonne.[1] In his younger days he was lieutenant in the artillery, then becoming a teacher of mathematics at Nîmes and later a professor at Montpellier. His great work, however, was as editor (1810–1831) of the mathematical journal[2] which commonly bears his name. In his later years he gave himself up to the life of a retired student. He was a prolific writer, chiefly on questions of geometry, the terms 'polar' (1810) and 'class' of a curve (1827) originating with him.

While not, like Gergonne and many other men of Napoleon's time, himself an army man, Siméon-Denis Baron Poisson[3] was the son of a soldier. He showed unusual abilities in mathematics when very young, and on this account was sent (1798) to the École Polytechnique, where his powers came to the attention of men like Lagrange and Laplace. Soon after finishing his prescribed work he was given a place on the faculty and devoted the rest of his life to teaching there and in the university, and to contributing to the literature of mechanics, mathematical physics in general, and pure mathematics. In the field of mathematics his chief contributions were to the theory of probability, algebraic equations, differential equations, definite integrals, surfaces, and the calculus of variations.

There have been several instances in the history of mathematics where a man's name has become known for a single discovery, not in itself remarkable, but striking in its peculiar interest. Such an instance is seen in the case of Charles-Julien Brianchon,[4] a student in the École Polytechnique (1804) and later (1808) an artillery officer. Brianchon had the ingenuity, when only twenty-three (1806), to take the dual of Pascal's proposition concerning a hexagon inscribed in a conic. The

[1] Born at Nancy, June 19, 1771; died at Montpellier, May 4, 1859. M. A. Lafon, *Gergonne, Sa vie et ses travaux*, reprint (n. d.) from the *Extraits des Mém. de l'Acad. de Stanislas*.

[2] *Annales de Mathématiques pures et appliquées*.

[3] Born at Pithiviers, June 21, 1781; died at Sceaux, April 25, 1840.

[4] Born at Sèvres, December 19, 1783; died at Versailles, April 29, 1864. J. Boyer, "Charles-Julien Brianchon d'après des documents inédits," *Revue scientifique*, I (4), 592.

result is Brianchon's Theorem with respect to the concurrence of lines joining opposite points of a circumscribed hexagon.[1] He became a professor in the artillery school and wrote several memoirs on geometry, particularly on curves of the second degree (1806) and lines of the second order (1817).

Poncelet. The life of Jean-Victor Poncelet[2] illustrates the military activity of many mathematicians of the disturbed Napoleonic period. A pupil of Monge's in the École Polytechnique (1807–1810), he entered the army (1812) as lieutenant of engineers. On the French retreat from Moscow he was captured by the Russians and was taken to Saratoff, on the Volga River. Here he devoted his time to the contemplation of certain possibilities in the domain of mathematics, and on his return to Metz (1814) he began to put the results of his thoughts into form for publication. The result was his great contribution to the theory of projective geometry.[3] He devoted the latter part of his life to military duties, his leisure being given to writing on mechanics, hydraulics, series, and geometry. He was one of the founders of modern geometry, probably the most important one. The Germans were more strongly influenced by his works, however, than were his own countrymen. It was Chasles who awakened France to the importance of his contributions and to a recognition of his genius.

Cauchy. The great technical and military schools founded or encouraged by Napoleon began at this time to enroll the most brilliant scientists of France. Among these was Augustin-Louis Cauchy,[4] who entered the École Polytechnique in Paris at the age of sixteen, proceeding thence to the École des Ponts et Chaussées. After a certain amount of engineering experience he was elected to the chair of mechanics in the École Polytechnique and to membership in the Académie des Sciences. On


[1] Chasles, *Aperçu*, p. 370.

[2] Born at Metz, July 1, 1788; died in Paris, December 23, 1867.

[3] *Traité des propriétés projectives des figures*, Paris, 1822; *Applications d'analyse et de géométrie*, 2 vols.. Paris, 1862, 1864.

[4] Born in Paris, August 21, 1789; died at Sceaux, May 23, 1857. J. Bertrand, "Éloge" in the *Mémoires* of the Académie d. Sci., Paris, Vol. 47, pp. clxxxiii–ccv

account of the political situation he went to Turin in 1830, where he became professor of mathematics in the university. Two years later he went to Prag and in 1838 returned to Paris and taught in certain Church schools. In 1848 he was made professor of mathematical astronomy in the university. His life was one of unrest on account of his own marked eccentricities as well as because of the changing political situation in France; but in spite of this fact he published upwards of seven hundred memoirs on mathematics and showed himself a man of uncommon scientific ability. Although usually displaying an affable manner, he was not a man of good breeding, being possessed of an unfortunate conceit, narrow in his views, and disposed to argue endlessly over trifles. He was an indefatigable worker, and his contributions to mathematics include researches into the theory of residues, the question of convergence, differential equations, the theory of functions, the elucidation of the imaginary, operations with determinants, the theory of equations, the theory of probability, the foundations of the calculus, and the applications of mathematics to physics. He was one of the first to use the imaginary as a fundamental instead of a subsidiary quantity, was the first to use Gauss's word "determinant" in its present sense, did much to establish the modern theory of convergence, and perfected the theory of linear differential equations and the calculus of variations.



AUGUSTIN-LOUIS CAUCHY
One of the foremost mathematicians of France in the 19th century

Chasles. Michel Chasles,[1] one of the leading French geometers of the 19th century, was, like his leading contemporaries in the field of mathematics, a student at the École Polytechnique (1812–1814). He went into business for a time but again returned to scientific work. He began publishing important memoirs on geometry as soon as he left school (1814), but it was his semihistorical work on the development of geometry[2] and his treatise on higher geometry[3] that gave him a world-wide reputation. These works were followed by various important memoirs on the different branches of geometry.[4] Chasles became professor of geometry and mathematics at the École Polytechnique in 1841 and professor of geometry in the faculty of sciences in 1846. In 1867 he prepared a noteworthy report on the progress of geometry in France.[5] He also received (1865) the Copley medal of the Royal Society for his work in conics.

Galois. The mathematician has not always been as conservative or as engrossed in his studies as the world seems to think. As an illustration of this fact one of the most interesting is that of Évariste Galois.[6] Educated at the Lycée Louis-le-Grand and the École Normale, at Paris, a rabid republican, twice imprisoned for his political views, a hot-blooded lover who fought a duel at twenty which cost him his life, he was able in the space of three or four years, even in his boyhood, to make for himself a lasting reputation as a genius. His life was mentally brilliant, but physically, politically, and morally it was a failure.

[1] Born at Épernon, November 15, 1793; died in Paris, December 18, 1880. For an obituary notice see Boncompagni's *Bullettino*, XIII, 815.

[2] *Aperçu historique sur l'origine et développement des méthodes en géométrie*, Paris, 1837; 2d ed., Paris, 1875; 3d ed., Paris, 1889; German ed., Halle, 1839. Some of the editions of his works bear the imprint of both Brussels and Paris.

[3] *Traité de géométrie supérieure*, Brussels, 1852; *Traité des sections coniques*, Paris. 1865.

[4] For example, "Construction de la courbe du troisième ordre déterminée par neuf points" (1853), *Journal de math. pures et appliquées*, XIX (1854).

[5] *Rapport sur les progrès de la géométrie*, Paris, 1871.

[6] Born in Paris, October 26, 1811; died in Paris, May 30, 1832. P. Dupuy, "La vie d'Évariste Galois," *École normale, Annales*, XIII (3), 197; G. Sarton, *The Scientific Monthly*, XIII, 363.

To him is due, however, one of the first important modern advances in the theory of groups, and hence to him we owe much of our modern theory of algebraic equations of higher degree. His most important memoir[1] was written the year before his death but was not published until 1846.[2]

Poincaré. Of the French mathematicians of the close of the 19th century no one ranked so high in the estimation of his contemporaries as Henri Poincaré.[3] There was hardly a branch of mathematics, pure or applied, to which he did not contribute in one way or another. His reputation was first made in his treatment of Abelian functions and in the more general type to which he gave the name of Fuchsian functions. His memoirs on these subjects began to appear in the *Comptes rendus* in 1880 and in the first volume of the *Acta Mathematica*. His contributions to elliptic functions, modular functions, double integrals, and the general theory of analysis are well known. He is equally well known for his important contributions to astronomy and physics and for his profound researches in the field of philosophy.

Other Contributors. Among the many others who added to the reputation of France in the field of mathematics during this period there may be mentioned Jean-Baptiste-Marie-Charles Meusnier de la Place (1754–1793), usually known as Meusnier, who wrote on the theory of surfaces; Jean-Baptiste Biot (1774–1862), who successfully applied mathematics to problems in physics and astronomy; Jean-Nicolas-Pierre Hachette (1769–1834), who wrote on algebra and geometry; Sophie Germain (1776–1831), known for her work on the theory of elastic surfaces; Louis Poinsot (1777–1859), who

[1] "Mémoire sur les conditions de résolubilité des équations par radicaux," Liouville's *Journal*, XI (1846). His *Manuscrits*, edited by J. Tannery, appeared at Paris in 1908.

[2] On the history of the group theory see the bibliography by C. Alasia in the *Rivista di fisica, matematica e scienze naturali*, XVIII–XXII. See also Miller, *Introduction*, p. 97.

[3] Born at Nancy, April 29, 1854; died in Paris, July 17, 1912. E. Lebon, *Henri Poincaré*, Paris, 1909; V. Volterra, "Henri Poincaré," *Rice Institute Pamphlets*, I, 133 (Houston, Texas).

contributed to the theory of numbers, to geometry, and to mechanics; Gabriel Lamé (1795–1870), primarily a physicist but writing on probability and surfaces; Théodore Olivier (1793–1853), especially concerned with descriptive geometry; Louis Arbogast (1759–1803), whose *Calcul des Dérivations* appeared in 1800; Jean Robert Argand (1768–1822), who wrote on the graphic representation of $\sqrt{-1}$; Joseph Fourier (1768–1830), known for his work in series, particularly with respect to Fourier's series, which is used in studying the flow of heat; Charles Dupin (1784–1873), prominent because of his works on mechanics and differential geometry; Georges-Henri Halphen (1844–1889), who contributed to the theory of invariants; Jean Gaston Darboux (1842–1917), contributor to differential geometry, one of the editors of the *Bulletin des sciences mathématiques et astronomiques*, and permanent secretary of the Académie des Sciences; Edmond Laguerre (1834–1886), a contributor to the theory of equations; Charles Hermite (1822–1901), who proved the transcendence of e and who wrote on the theory of functions; Joseph Liouville (1809–1882), long the editor of Liouville's *Journal*; Charles Méray (1835–1911), original in his ideas of the foundations upon which elementary geometry should be built; Joseph Alfred Serret (1819–1885), best known for his *Cours d'algèbre supérieure* (1849) but also a prolific writer on the function theory, groups, and differential equations; Joseph-Louis-François Bertrand (1822–1900), professor of mathematical physics in the Collège de France and secretary of the Académie des Sciences, a writer on the theory of probability, the calculus of variations, and differential equations as applied to dynamics; Pierre Duhem (1861–1916), contributor to mathematical physics and especially, by his study of original sources, to the history of science; and Louis Couturat (1868–1914), writer upon the interrelation of mathematics and logic. Any such list is necessarily fragmentary, and the student who wishes to carry his investigations farther should consult such works as the French or German editions of the encyclopedia of mathematics.

4. Germany

General Survey. Germany began to show her real strength in mathematics at the close of the 18th century. Theretofore she had depended largely on imported men, such as Euler, the Bernoullis, and Lagrange. Now she produced Gauss, and his influence on German mathematics made Göttingen a focus for scholars; it placed Germany among the leading nations in the cultivation of this science, and gave her a position of supremacy during part of the 19th century.



CHRISTIAN VON WOLF
After a mezzotint by Jacob Haid

Of the work accomplished in the 18th century a fair example is that of Freiherr Christian von Wolf,[1] a philosopher of merit and a mathematician of erudition if not of brilliancy. He took his master's degree at Leipzig in 1703 and at once became Dozent in the university. Soon after this (1706) he went to Halle as professor of mathematics and, somewhat later, of physics. Because of his religious views he was banished from the university in 1723, but was immediately invited to accept the professorship of philosophy at Marburg. He was recalled to Halle by Frederick the Great, to whom a religious question was not a matter of much moment, became chancellor of the

[1] Born at Breslau, January 24, 1679; died at Halle, April 9, 1754.

university in 1743, and was raised to the rank of baron (Freiherr) in 1745. He was a member of various learned societies, did much to popularize the theories of his friend Leibniz, and was a voluminous writer but not an original thinker. His most extensive works are his *Elementa* and *Anfangsgründe*,[1] but he also prepared an unimportant set of logarithmic tables (1711) and wrote a mathematical dictionary (1716).[2]

In the field of elementary education, Germany produced a number of important writers, but few whose names can be rated as international. Among the most industrious of the group was Christian Pescheck,[3] who wrote a large number of textbooks and was one of the first of the German writers to consider seriously the methods of teaching the subject.

Gauss. The real founder of modern German mathematics, however, is Carl Friedrich Gauss,[4] one of the many mathematicians who rose to highest eminence from very humble birth.



AUTOGRAPH OF GAUSS

Signed as Director of the Royal Society of Sciences at Göttingen

The son of a day laborer, his abilities showed themselves so early as to attract attention, and he was sent to the Carolineum at Braunschweig (1792–1795) and thence to the University of Göttingen (1795–1798). During his university career he conceived the idea of the theory of least squares,[5] discovered the

[1] *Elementa matheseos universae*, 4 vols., Halle, 1713, with later editions; *Anfangsgründe aller mathematischen Wissenschaften*, 4 vols., Halle, 1710, with several editions.

[2] J. C. Gottsched, *Historische Lobschrift des . . . Christians . . . Freyherrn von Wolf*, Halle, 1755; W. Arnsperger, *Christian Wolff's Verhältnis zu Leibniz*, Weimar, 1897.

[3] Born at Zittau, July 31, 1676; died at Zittau, October 28, 1747.

[4] Born at Braunschweig (Brunswick), April 30, 1777; died at Göttingen, February 23, 1855. In his autographs the first name begins with C. The name was originally Johann Carl Friedrich Gauss.

[5] Legendre (1805) was the first to write upon the subject, introducing it in his work on the orbit of comets. The deduction of the law was effected by an Irish-American writer, Adrain, in 1808.

at the University of Berlin, became Privatdozent in 1825, and two years later was made professor of mathematics at Königsberg. He was a prolific contributor in various lines of mathematics, but his chief work was in the fields of elliptic functions,[1] determinants, the theory of numbers,[2] differential equations, the calculus of variations, and infinite series.

Belonging to about the same period as Jacobi, Peter Gustav Lejeune-Dirichlet[3] was educated at Göttingen, studied under Gauss, and became professor of mathematics at Breslau, Berlin, and Göttingen. He was chiefly interested in algebra, the number theory, and quadratic forms.

That a university professorship is not a *sine qua non* to success in mathematics is a fact again illustrated in the case of Hermann Günther Grassmann,[4] who was the son of a teacher of mathematics in the Gymnasium at Stettin and himself occupied a similar position in the same school. The father wrote some textbooks of no particular moment, and both he and the son gave much attention to physical questions. There was also another son, Robert, with whom Hermann collaborated in writing an arithmetic (1860). The entire output of the family, however, was as nothing compared with Hermann's *Ausdehnungslehre*.[5] In this he set forth a theory that covered much the same ground as the theory of quaternions, then being independently developed by Sir William Rowan Hamilton.

Ernst Eduard Kummer[6] is another instance of a man of genius who spent some years as a Gymnasium teacher before being called to a university chair. Educated for theology as well as mathematics, he began his teaching at Sorau, afterwards going to Liegnitz, where he taught for ten years

[1] *Fundamenta nova theoriae functionum ellipticarum*, Königsberg, 1820.

[2] *Canon arithmeticus*, Berlin, 1839.

[3] Born at Düren, February 13, 1805; died at Göttingen, May 5, 1859.

[4] Born at Stettin, April 15, 1809; died at Stettin, September 26, 1877. See F. Engel, *Jahresbericht* of the *Deutsche Math.-Verein.*, XIX, 1.

[5] *Die Wissenschaft der extensiven Grösse oder die Ausdehnungslehre*, Leipzig, 1844; completed in 1862. A list of his works may be found in the *Mathematische Annalen*, XIV, 43.

[6] Born at Sorau, Nieder-Lausitz, January 29, 1810; died at Berlin, May 14, 1893.

(1832–1842) in the Gymnasium, having Kronecker for one of his pupils. He then became (1842) professor of mathematics in the University of Breslau, later (1855) being transferred to Berlin, where he remained until 1884. In Crelle's *Journal* may be found his valuable contributions to the theory of hypergeometric (Gaussian) series (1836), the Riccati equation (1834), the question of the convergency of series (1835), the theory of complex numbers (1844, 1850), and cubic and biquadratic remainders (1842, 1848). He created the theory of ideal prime factors of complex numbers (1856) and laid down the principles applicable to Kummer surfaces.[1]

His contemporary, Georg Friedrich Bernhard Riemann,[2] also proved himself a genius in the study of surfaces. He studied at Berlin and Göttingen, receiving his doctorate at the latter university in 1851. His dissertation[3] has since been recognized as a genuine contribution to the theory of functions. Three years later (1854) he became a Privatdozent in Göttingen and in 1857 became a professor[4] of mathematics in the university. His introduction of the notion of geometric order into the theory of Abelian functions, and his invention of the surfaces which bear his name, led to a great advance in the function theory. He also set forth (1854) a new system of non-Euclidean geometry,[5] and wrote on partial differential equations,[6] elliptic functions,[7] and physics.[8]

[1] Surfaces of the fourth degree, 16 knot points, 16 singular tangent planes. See "Allgemeine Theorie der gradlinigen Strahlensysteme," Crelle's *Journal*, LVII (1860), 189. For a biographical sketch and a list of his works see the *Jahresbericht* of the *Deutsche Math.-Verein.*, III, 13.

[2] Born at Breselenz, Hannover, September 17, 1826; died at Selasca, Lago Maggiore, July 20, 1866.

[3] *Grundlagen für eine allgemeine Theorie der Functionen einer veränderlichen complexen Grösse*, Göttingen, 1851; 2d ed., Göttingen, 1867.

[4] He succeeded Dirichlet as ordentlicher Professor in 1859.

[5] *Ueber die Hypothesen welche der Geometrie zu Grunde liegen*, Leipzig, 1867. Like several of his works, this appeared posthumously.

[6] *Partielle Differentialgleichungen*, Braunschweig, 1869; 2d ed., Hannover, 1876; 3d ed., Braunschweig, 1882; 4th ed., Braunschweig, 1900–1901.

[7] *Elliptische Functionen, Vorlesungen mit Zusätzen*, Leipzig, 1899.

[8] See his *Gesammelte mathematische Werke und wissenschaftlicher Nachlass*, Leipzig, 1876; 2d ed., Leipzig, 1892; French translation, Paris, 1898; contains a biography

One of the most brilliant and promising mathematicians of Germany in the middle of the 19th century appeared in the person of Ferdinand Gotthold Max Eisenstein.[1] Gauss, in a moment of enthusiasm, and without sufficiently weighing his words, said of him: "There have been but three epoch-making mathematicians,—Archimedes, Newton, and Eisenstein." He was brought up in poverty, showed no particular taste for mathematics until he was nineteen, died at the age of twenty-nine, and yet in ten years developed powers so remarkable as to place him in the first rank of scholars. His most important contributions were to the theory of ternary and quadratic forms, the theory of numbers, and the theory of functions. He has been spoken of by his countrymen as the real founder of the theory of invariants.

Weierstrass. As a type of those great leaders who, towards the close of the 19th century, made Germany a great gathering place for scholars there may be mentioned Karl Weierstrass.[2] He studied law and finance at Bonn (1834), taught in various Gymnasien, went to Berlin in 1856 as a teacher in the Gewerbeinstitut, and became ordentlicher Professor of mathematics in the University of Berlin in 1864. Here he became one of the great leaders in the theory of elliptic and Abelian functions, in the theory of functions in general, and in the development of the theory of irrational numbers. The Berlin Akademie der Wissenschaften began the publication of his collected works in 1894.[3]

Dedekind, Cantor, and Fuchs. Julius Wilhelm Richard Dedekind[4] stands out as one of the most prominent contributors of the 19th century to the theory of algebraic numbers. He studied at Göttingen and in 1854 became a Dozent in the university. In 1858 he went to the polytechnic school at

[1] Born at Berlin, April 16, 1823; died at Berlin, October 11, 1852. F. Rudio, "Eine Autobiographie von Gotthold Eisenstein," *Abhandlungen*, VII, 145.

[2] Born at Ostenfelde, October 31, 1815; died at Berlin, February 19, 1897.

[3] On his life, see *Acta Mathematica*, XXI, 79; XXII, 1.

[4] Born at Braunschweig, October 6, 1831; died at Braunschweig, February 12, 1916.

Zürich, and four years later became a professor in a similar institution at Braunschweig. He wrote various important memoirs on the binomial equation and on the theory of modular and Abelian functions, but is best known for his treatises *Was sind und was sollen die Zahlen?* (1888) and *Stetigkeit und irrationale Zahlen* (1872).[1] In the latter work he set forth his idea of the *Schnitt* (cut) in relation to irrational numbers,—an idea which he had in mind as early as 1858.

Although Georg Cantor[2] was the son of a Danish merchant and was born in Russia, he should properly be ranked among the German mathematicians, having spent the greater part of his life in German universities. He studied at Zürich, Göttingen, and Berlin, and became a Dozent at Halle in 1869 and a professor three years later. He was a man of original ideas, and the theory of assemblages is practically his creation. His researches on this subject were first published in the *Annalen* in 1879.

Emmanuel Lazarus Fuchs[3] became professor of mathematics at Greifswald in 1869 and afterwards occupied similar positions at Göttingen, Heidelberg, and Berlin. His earlier labors were in the fields of higher geometry and the theory of numbers, but he attained his highest reputation in his work on linear differential equations.

Other Writers. Among the other German writers of the 19th century not many were at the same time leaders in advanced research and contributors to elementary mathematics. A few of the best-known names of those who extended the boundaries of mathematics should, however, be mentioned. These are Johann Friedrich Pfaff (1765–1825), professor at Helmstädt (1788) and Halle (1810), known for his work in astronomy, geometry, and analysis; Ludwig Otto Hesse (1811–1874), one of the foremost writers on modern pure geometry, analytic geometry, and determinants; Christoph Gudermann (1798–1852), who wrote on hyperbolic functions;

[1] English translations by Professor W. W. Beman, Chicago, 1901.
[2] Born at Petrograd, March 3, 1845; died at Halle, January 6, 1918.
[3] Born at Moschin, May 5, 1833; died at Berlin, April 26, 1902.

Johann August Grunert (1797–1872), editor of the *Archiv*; Ernst Ferdinand August (1795–1870), known for his work on mathematical physics; Rudolff Friedrich Alfred Clebsch (1833–1872), professor at Carlsruhe, Giessen, and Göttingen, a contributor to modern geometry; Hermann Ludwig Ferdinand von Helmholtz (1821–1894), a contributor to many fields of scientific research including that of non-Euclidean geometry; Leopold Kronecker (1823–1891), a leading writer on the theory of equations and on elliptic functions; Friedrich Wilhelm Bessel (1784–1846), one of the leading astronomers of the century, a physicist, and well known for the functions which bear his name; Paul Du Bois-Reymond (1831–1889), known for his work on Fourier's series, the problem of convergence, the calculus of variations, and integral equations; Siegfried Heinrich Aronhold (1819–1884), professor in the technical high school at Berlin, well known for his work on invariants; and Karl Theodor Reye (1837–1919), whose *Geometrie der Lage* (3d ed. 1886–1892) is one of the best-known textbooks on the subject.

5. Italy

Nature of the Work. In the 18th and 19th centuries Italy produced a worthy line of mathematicians, but until recently she has not made a serious effort to regain her earlier standing. Her scholars seemed to work in isolation during much of this period,—a result due in some measure, no doubt, to the lack of political homogeneity in Italy herself. A few names in the 18th and the early 19th century, however, deserve our attention.

Ceva Brothers. About the middle of the 17th century there were born two brothers, Giovanni Ceva[1] and Tommaso Ceva,[2] each of whom contributed to geometry and physics. The latter was a teacher of mathematics in the Jesuit college at Milan, while the former was in the service of the Duke of Mantua. Tommaso wrote on the cycloid, the mechanical trisection of an

[1] Born at Mantua, December, 1647; died May 13, 1736. These dates are uncertain.

[2] Born at Milan, December 20, 1648; died at Milan, February 3, 1737.

angle,[1] and mathematics in general,[2] but Giovanni was the more original and prolific. To him is due a well-known theorem which asserts that if three lines from the vertices A, B, C of a triangle ABC are concurrent in P and meet the opposite sides in X, Y, Z respectively, then

$$\frac{AZ}{ZB} \cdot \frac{BX}{XC} \cdot \frac{CY}{YA} = 1.$$

This theorem was published by him in 1678.[3] His work extended into the 18th century.[4]

Manfredi Brothers. Another interesting example of a family devoted to mathematics is seen in the case of the Manfredi brothers. Eustachio Manfredi[5] was a jurist who at the age of twenty-five (1699) became professor of mathematics in the University of Bologna. He was one of the founders of the Institute of Bologna, and is particularly known for his elaborate ephemerides[6] and his general works on astronomy. He contributed also to the textbook literature of geometry.[7]

Gabriele Manfredi[8] was professor of mathematics (1720) in the University of Bologna and wrote on differential equations[9] and geometry. Eraclito Manfredi[10] was professor of medicine and later of geometry in the University of Bologna, his major interest being in mechanics.

Riccati Family. Still another family that showed decided genius in mathematics and physics was that of Jacopo Francesco, Conte Riccati,[11] a man of private fortune, who studied

[1] *Acta Eruditorum*, 1695. [2] *Opuscula mathematica*, Milan, 1699.

[3] *De lineis rectis se invicem secantibus statica constructio*, Milan, 1678.

[4] *Tria problemata geometris proposita*, Mantua, 1710; *De re numeraria, quod fieri potuit, geometrice tractata*, Mantua, 1711. Chasles, *Aperçu*, note vii, p. 294; T. Perelli, in A. Fabbroni, *Vitae Italorum doctrina excellentium*, Vol. XVIII (Pisa, 1778–1805).

[5] Born at Bologna, September 20, 1674; died at Bologna, February 15, 1739.

[6] Bologna, 1714, 1715.

[7] *Elementi della geometria piana e solida, e della trigonometria*, Bologna, 1755 (posthumous).

[8] Born at Bologna, March 25, 1681; died at Bologna, October 13, 1761.

[9] *De Constructione Æquationum Differentialium Primi Gradus*, Bologna, 1707, a work highly esteemed by Leibniz.

[10] Born at Bologna, *c.* 1682; died at Bologna, September 15, 1759.

[11] Born at Venice, May 28, 1676; died at Treviso, April 15, 1754.

in Padua and spent the latter part of his life in Venice and Treviso. He wrote on philosophy, physics, differential equations, mensuration, and related subjects, and did much to make the theories of Newton known. His collected works were published after his death.[1] He is known chiefly for his elaborate study of the so-called Riccati Equation $\frac{d\eta}{dx} = A + B\eta + C\eta^2$, where A, B, C are functions of x, of which he gave solutions for certain special cases. The equation had already been studied by Jacques Bernoulli. His second son was Vincenzo Riccati,[2] a Jesuit, professor of mathematics in the college of his order at Bologna.[3] His line of interest was quite like that of his father, his work in mathematics including the study of differential equations (1752), series (1756), quadrature problems (1767), and the hyperbolic functions.[4] Jacopo's third son was Giordano Riccati,[5] who wrote on the Newtonian philosophy (1764, 1777), geometry (1778,



JACOPO FRANCESCO, CONTE RICCATI

After an engraving by Comirato. Jacopo was the first of the well-known mathematical family of Riccati

[1] *Opere del Conte Jacopo Riccati*, 4 vols., Treviso, 1758.

[2] Born at Castelfranco, province of Treviso, January 11, 1707; died at Treviso, January 17, 1775.

[3] Collegio di San Francesco Saverio.

[4] His *Opuscula ad Res Physicas et Mathematicas pertinentia*, 2 vols., appeared at Bologna in 1757–1762.

[5] Born at Castelfranco, February 25, 1709; died at Treviso, July 20, 1790. D. M. Federici, *Commentario sopra la Vita e gli Studj del Conte Giordano Riccati*, Venice, 1790.

1779, 1790), cubic equations (1784), and physical problems (1763, 1764, 1767). The fifth son was Francesco Riccati,[1] who wrote on geometry as applied to architecture.

Other Writers. In order to give some further idea of the general nature of Italian mathematics at this time it will suffice to record the names of a few scholars who did the most to advance their science. Among the earliest of these in the 18th century was Luigi Guido Grandi,[2] a member of the order of the Camaldolites, professor of philosophy (1700) and later (1714) of mathematics at the University of Pisa, and author of a number of works on geometry,[3] in which are considered the analogies of the circle and the equilateral hyperbola, curves of double curvature on a sphere, and the quadrature of certain parts of a spherical surface.

Slightly younger than Guido Grandi, Giulio Carlo Fagnano de Fagnani, Marchese de' Toschi e S. Onorio,[4] commonly known as Fagnano, a man of private means and a savant at sixteen, was an important contributor to mathematics.[5] He was chiefly interested in geometry and in the study of algebraic equations. One of his interesting discoveries (1719) was the relation

$$\frac{c}{4} = 2 \log \left(\frac{1-\sqrt{-1}}{1+\sqrt{-1}} \right)^{\frac{1}{2}\sqrt{-1}}$$

where $c = 2\pi$. He further showed that the integral which expresses the arc of a lemniscate has properties analogous to those of the integral which represents the arc of a circle. Euler, in extending the theory of elliptic functions, gives Fagnano credit for having first directed attention to it.

[1] Born at Castelfranco, November 28, 1718; died July 18, 1791.

[2] Born at Cremona, October 7, 1671; died at Pisa, July 4, 1742.

[3] *Geometrica demonstratio Vivianeorum problematum*, Florence, 1699; *Geometrica demonstratio theorematum Hugenianorum circa logisticam seu logarithmicam*, Florence, 1701; *Quadratura circuli et hyperbolae per infinitas hyperbolas et parabolas geometrice exhibita*, Pisa, 1703; *Istituzioni geometriche*, Florence, 1741.

[4] Born at Senigallia, December 6, 1682; died September 26, 1766. *Opere Matematiche*, ed. Volterra, Loria, and Gambioli, 3 vols., Milan, 1911, 1912.

[5] His *Produzioni Matematiche* appeared at Pesaro, 2 vols., 1750.

Giulio had a son, Giovanni Francesco (1715–1797) or Gianfrancesco, who deduced from his father's formula the relations[1]

$$\frac{c}{4} = \sqrt{-1}\log\left(-\sqrt{-1}\right)$$

and

$$\frac{c^2}{4} = \log(+1)\cdot\log(-1).$$

To students of geometry, one of the most familiar names of this period is that of Giovanni Francesco Giuseppe Malfatti.[2] He was educated in his early youth at a Jesuit college in Verona, but at the age of seventeen (1748) he went to Bologna and studied under Vincenzo Riccati. In 1771 he became professor of mathematics at Ferrara, where he spent the rest of his life. He wrote a variety of works, but his name is chiefly remembered for a problem which he published[3] in 1803: In a triangular prism to inscribe three cylinders of altitude equal to that of the prism and of maximum volume and so that the remaining volume of the prism shall be a minimum. This at once reduces itself to the inscription, in a given triangle, of three circles, each tangent to the others and to two sides of the triangle. Malfatti solved the problem analytically, Steiner solved it geometrically, and it has been the subject of extended study by various other scholars.[4]

Students of higher algebra will possibly be more familiar with the name of Paolo Ruffini,[5] who taught mathematics and medicine at Modena and contributed extensively to the theory of equations. He wrote several works and important

[1] Montucla, *Histoire*, III, 285.

[2] Gianfrancesco Malfatti. Born at Ala di Trento, 1731; died at Ferrara, October 9, 1807.

[3] *Memoria sopra un problema stereotomico*, in the *Memorie di . . . Società Italiana delle Scienze*, Modena, X, parte I[a]., p. 235.

[4] M Baker, "The History of Malfatti's Problem," in the *Bulletin of the Philosophical Society of Washington*, II, 113; Boncompagni's *Bullettino*, IX, pp. 361, 383; A. Wittstein, *Geschichte des Malfatti'schen Problems*, Diss, Munich, 1871.

[5] Born at Valentano, September 23, 1765; died at Modena, May 10, 1822. G. Bianchi, *Elogio*, pronounced at Modena, November 25, 1822.

monographs on the subject.[1] In connection with his work on equations he made a beginning in the theory of groups.[2]

In the second half of the century the study of the geometry of the compasses, which had already attracted attention at various times, was successfully undertaken by Lorenzo Mascheroni.[3] He was one of those who succeeded in making a name for himself in mathematics although beginning the study relatively late. His first interests were in the humanities, and he taught Greek and poetry at the liceo in his native town and at Pavia. He also took holy orders and became an abbot, which did not hinder him, however, from being a good fighter for the Cisalpine republic. After his experience in teaching the humanities he took up the study of geometry and became professor of mathematics at Pavia. He wrote on physics, the calculus, and the proposed metric system, but is known chiefly for his geometry of the compasses.[4] In the last-named work he showed that the ordinary constructions of elementary



LORENZO MASCHERONI

After an engraving by F. Redenti, from a drawing made from life

[1] Among them, *Teoria generale delle equazioni*, 2 vols., Bologna, 1799.

[2] H. Burkhardt, *Abhandlungen*, VI, 119.

[3] Born at Castagneta (Castagnetto), near Bergamo, May 14, 1750; died in Paris, July 30, 1800. See G. Mangili, *Elogio di Lorenzo Mascheroni*, Milan, 1812; G. Savioli, *Memorie . . . Mascheroni*, Milan, 1801; F. Landi, *Elogio di Lorenzo Mascheroni*, Modena, 1804. These writers disagree slightly as to the date of his birth.

[4] *Geometria del Compasso*, Pavia, 1797; new ed., Palermo, 1901. There was a French translation by A. M. Carette, 1798; 2d ed., Paris, 1828. For bibliography see *L'Intermédiaire*, XIX, 92.

geometry can be performed by the use of the compasses alone; that is, that all critical points of an elementary figure can be found with no assistance from the straightedge. The idea seems to have been suggested to him by Benedetti's work of 1553, already mentioned.

There were in this period a number of Italian mathematicians who, following the French school, made notable advance in mathematical astronomy. Among these was Ruggero Giuseppe Boscovich[1] who, as the family name indicates, descended through his father from a Herzegovina family, although his mother was a native of Ragusa, Italy. He entered the Jesuit order and in 1740 became professor of mathematics and philosophy in the Collegio Romano at Rome. After traveling extensively in various countries of Europe he became a professor (1764–1770) at Pavia. He afterwards lived in Paris for some years, returning to Italy in 1783 and taking up his residence in Milan. He wrote extensively on astronomy, giving the first geometric solution of the problem of determining the equator of a planet by three observations of a spot and determining its orbit from three observations of the body.[2] He also worked on the general problem of the determination of the orbits of comets and wrote a general treatise on mathematics.[3]

In the field of pure geometry perhaps the most original of the Italian writers of this period was Giusto Bellavitis,[4] professor of mathematics at Padua and Vicenza, who contributed to the theory of projective geometry and was the first to set forth the method of equipollence.[5]

Luigi Cremona (1830–1903) is even better known in this field, having written an important textbook on projective

[1] Born at Ragusa, May 18, 1711; died at Milan, February 13, 1787. G. Dionisi, *Ruggero G. Boscovich*, Zara, 1887; F. Ricca, *Elogio . . . Ruggiero Giuseppe Boscovich*, Milan, 1789.

[2] *De determinanda Orbita Planetae*, Rome, 1749.

[3] *Elementa universae matheseos*, 3 vols., Rome, 1754.

[4] Born at Bassano, near Padua, November 22, 1803; died at Padua, November 6, 1880.

[5] *Saggio d' applicazioni del calcolo delle equipollenze*, Padua, 1837; *Metodo delle equipollenza*, Padua, 1837. See O. Brentari, *Biografia di Giusto Bellavitis*, Bassano, 1881.

geometry. He was, however, more than a textbook writer, having contributed to the theory of cubic surfaces (1866) and to the theory of transformation of curves.

Eugenio Beltrami (1835–1900) was a prolific writer on the theory of higher geometry and on physical problems, higher algebra, and invariants, most of his memoirs appearing in the *Annali di Matematica.* He was professor of mathematics at Pisa, Bologna, Pavia, and Rome, and was president of the Accademia dei Lincei in Rome. He was also minister of public instruction (1862) and a senator. Giuseppe Battaglini (1826–1894) was well known for his interest in the theory of groups, his work in line geometry, and his editorial work on the *Giornale di Matematiche* (Naples, beginning in 1863). His contemporary, Barnaba Tortolini (1808–1874), founded (1850) the *Annali di Scienze Matematiche e Fisiche* and was a voluminous writer on analysis. Of the later writers, Francesco Brioschi (1824–1897), professor at Pavia, contributed to the study of mathematical physics; Enrico Betti (1823–1892), professor at Pisa, added to the theory of binary forms; Felice Casorati (1835–1890), professor at Pavia, did much to advance the study of analysis in Italy; and Ulisse Dini (1845–1918) wrote upon the theory of the functions of a real variable (1877) and contributed to the theory of series.



MARIA GAETANA AGNESI

Professor of mathematics at Bologna and a writer on analytic geometry

Among the women of Italy who have added to the store of the world's knowledge of mathematics the most erudite one of this period was Maria Gaetana Agnesi,[1] who occupied for a time the chair of mathematics in the University of Bologna. Her work on analytic geometry is well known.[2]

6. Switzerland

Nature of the Work. The influence of the Bernoullis showed itself in Switzerland in the 18th century, but with the exception of two great scholars no one arose to maintain the standard which they had set. There was certainly no Swiss school of mathematicians, hardly even a Basel school. To have produced in one century both Euler and Steiner is, however, quite enough of glory for any country.

Minor Writers. Among the minor writers the earliest of any considerable note was Nicolas Fatio de Duillier.[3] At various times a resident of Geneva, Paris, The Hague, and London, pilloried for his relations with the mystic sect of Camisards, a Fellow of the Royal Society, an extensive traveler, and a friend of Giovanni Domenico Cassini, Huygens, and Leibniz, his life was filled with adventure rather than with study. Nevertheless he found time to write on the curve of least descent,[4] on navigation, and on astronomy.

Jean Pierre Crousaz[5] also belongs to the period between the 17th and 18th centuries. Although born in Switzerland and spending the last fifteen years of his life there, he was also professor of mathematics at Groningen for four years (1724–1728) and spent some time in Germany. He wrote on the

[1] Pronounced än yā'zē. Born at Milan, March 16, 1718; died at Milan, January 9, 1790.

[2] *Istituzioni analitiche ad uso della gioventù Italiana*, Milan, 1748. There is an English translation, 2 vols., London, 1801, with a brief biography.

[3] Facio, Faccio. Born at Basel, February 16, 1664; died at Maddersfield, Worcestershire, May 10, 1753.

[4] *Lineae brevissimi descensus investigatio geometrica duplex . . .*, London, 1699, which helped to precipitate the contest between Newton and Leibniz.

[5] Born at Lausanne, April 13, 1663; died at Lausanne, March 22, 1750. The name also appears as Crouzas.

geometry of lines and surfaces (Amsterdam, 1718), on algebra (Paris, 1726), and on the new calculus (Paris, 1721). He also wrote an essay on arithmetic, one of the latest to give a complete discussion of the ancient finger reckoning.[1]

Of the pupils of the Bernoullis only a few rose even to moderately high rank. One of the first to acquire considerable reputation, although now generally forgotten, was Jacob Hermann.[2] He was a pupil of Jacques (I) Bernoulli and was professor of mathematics in the University of Padua (1707–1713), at Frankfort a. d. O. (1713–1724), and at Petrograd (1724–1731), after which he became professor of moral philosophy at Basel. He was one of the early writers on the differential calculus (1700) and contributed to the *Acta Eruditorum* (from 1702) and other periodicals numerous memoirs on the application of the subject to geometric problems.

Among the Swiss mathematicians of the 18th century Gabriel Cramer[3] was much better known than Hermann. He belonged to a Holstein family which settled for a time at Strasburg but finally went to Geneva. At the age of twenty (1724) he was made professor of mathematics, becoming a colleague of Giovanni Ludovico Calendrini[4] at the University of Geneva. Cramer's work related chiefly to physics, but he also wrote on geometry (1732), the history of mathematics (1739, 1741, 1748, 1750), and algebraic curves,[5] most of these contributions appearing in the form of memoirs. He had a wide personal acquaintance with the mathematicians of England, Holland, and France. The revival of the subject of determinants, already suggested by Leibniz in Europe and by Seki in Japan, is due largely to him (1750).

Euler. Basel had achieved enough glory in the history of mathematics through being the home of the Bernoullis, but she

[1] *Réflexions sur l'utilité des mathématiques . . . avec un nouvel essai d'arithmétique démontrée,* Amsterdam, 1715.

[2] Born at Basel, July 16 (o.s.), 1678; died at Basel, July 11, 1733.

[3] Born at Geneva, July 31, 1704; died at Bagnols, near Nîmes, January 4, 1752.

[4] Born at Geneva, 1703; died at Geneva, 1758. He wrote on Newton's theories and on conics.

[5] *Introduction à l'Analyse des Lignes Courbes Algébriques,* Geneva, 1750.

doubled her glory when she produced Léonard Euler.[1] While the Bernoullis had generally been destined by their parents for commerce or the law, but had finally entered the field of mathematics, Euler had been taught mathematics by his father, who had himself studied under Jacques (I) Bernoulli. Léonard studied theology, but also put himself under the instruction of Jean (I) Bernoulli. To his study of theology and mathematics he added that of medicine, the oriental languages, astronomy, and physics, and became not only the greatest mathematician and astronomer of his generation but one of its most all-round savants. He went to Petrograd in 1727 and taught mathematics and physics there. In 1735, through excessive work, he lost the use of his right eye. "J'aurai moins de distractions" was his comment,—the



LEONARD EULER

From a contemporary engraving

[1] Born at Basel, April 15, 1707; died at Petrograd, September 18, 1783. Since he wrote chiefly in French when not employing, as he usually did, the international Latin, the French spelling of his Christian name has been retained. In German it appears as Leonhard. In his Latin books he uses Leonhardus Eulerus. For a bibliography of recent articles, see *Bibl. Math.*, X (3), 284. See also P. Stäckel, in the *Vierteljahrschrift der Naturf. Gesellsch. in Zürich*, Jahrg. 54, and the recent edition of his works. Condorcet's *Éloge de M. Euler* was pronounced before the Académie des Sciences in 1785 and is well known, as is also N. Fuss, *Lobrede auf Herrn Leonhard Euler*, Basel, 1786.

comment of a practical philosopher with a sense of humor. In 1766 he lost the use of the other eye, but continued his labors without complaint to the day of his death. Few writers ever contributed so extensively or so fruitfully. Indeed, he was in a sense the creator of modern mathematical expression. In his lifetime there appeared about six hundred important memoirs from his hand, besides various treatises. He published three monumental works on analysis,[1] in which he "freed the analytic calculus from all geometric bonds, thus establishing analysis as an independent science."[2] He also wrote on algebra,[3] arithmetic, mechanics, music,[4] and astronomy, and had an extensive knowledge of botany, chemistry, medicine, and *belles-lettres*. It is said that he could repeat the *Æneid* from beginning to end.[5] Arago said of him that he calculated without effort, just as men breathe and as eagles sustain themselves in the air.

Several algebraic expressions bear his name. Euler's constant is the limit of $1+\frac{1}{2}+\frac{1}{3}+\cdots+\frac{1}{n}-\log n$ as $n \to \infty$, the value being 0.577215664901532 8 as found by him, but since carried much farther. Euler's Equation is $\frac{dx}{\sqrt{X}}=\frac{dy}{\sqrt{Y}}$, where X and Y are two quartic functions of x and y, differing only in the variable. From it is derived a formula that is of great importance in the theory of elliptic functions. The equation $e^{iv}=\cos x+i\sin x$ is also called by his name.

De Morgan relates an interesting anecdote concerning Euler's meeting with Diderot at the Russian court. Diderot

[1] *Introductio in analysin infinitorum*, 2 vols., Lausanne, 1748; German edition, Berlin, 1788–1790; *Institutiones calculi differentialis*, Berlin, 1755; German edition, 3 vols., Berlin, 1790–1793; *Institutiones calculi integralis*, Petrograd, 1768–1770, with Vol. IV in 1794 (posthumous), and later editions. Eneström has published a complete list of his works (Leipzig, 1910–1913).

[2] H. Hankel, *Die Entwickelung der Mathematik in den letzten Jahrh.*, p. 13. Tübingen, 2d ed., 1884.


[3] *Vollständige Anleitung zur Algebra*, 2 vols., in Russian, Petrograd, 1768; German edition, Petrograd, 1770; French edition by Lagrange in 1774.

[4] Fuss remarked that his *Tentamen novae theoriae musicae* (Petrograd, 1729) "contained too much geometry for musicians and too much music for geometers."

[5] D. Brewster, *Letters of Euler*, I. 24. New York, 1872.

had somewhat displeased the Czarina by his antireligious views, and so she persuaded Euler to assist her in suppressing him.

Diderot was informed that a learned mathematician was in possession of an algebraical demonstration of the existence of God, and would like to give it him before all the Court, if he desired to hear it. Diderot gladly consented: though the name of the mathematician



AUTOGRAPH LETTER OF EULER

Written during the period of his residence in Petrograd

was not given, it was Euler. He advanced toward Diderot, and said gravely, and in a tone of perfect conviction:

"Monsieur, $\frac{a+b^n}{n} = x$, donc Dieu existe; répondez!"

Diderot, to whom algebra was Hebrew, was embarrassed and discomfited: while peals of laughter rose on all sides. He asked permission to return to France at once, which was granted.[1]

[1] *Budget of Paradoxes*, 2d ed., II, 4. Chicago, 1915.

His death was as he would have wished,—a discussion on mathematics, a dinner, a cup of tea, a pipe, and in an instant, in the words of Condorcet, "il cessa de calculer et de vivre."

Euler's eldest son, Johann Albrecht Euler,[1] was also a scientist of high attainments, particularly in the line of physics.[2]

Steiner. Johann Heinrich Pestalozzi[3] is not generally thought of as a mathematician, although no man of his time did so much as he to improve the teaching of elementary arithmetic. He deserves a place in the history of mathematics, however, for having taken a poor Swiss boy who could not write a single word before reaching the age of fourteen, admitting him at the age of seventeen to his school at Yverdon, and giving him a love for mathematics. The boy was Jacob Steiner,[4] the last one who would have been selected, by those who knew him in childhood, as destined to become one of the greatest geometers of modern times. Not long after leaving Yverdon he went to the University of Heidelberg (1818), where he soon showed his ability in mathematics, and in 1821 he began giving private lessons in Berlin, becoming soon after



JACOB STEINER
After a photograph

[1] Born at Petrograd, November 16 (o.s.), 1734; died September 6 (o.s.), 1800.

[2] For a list of his works see P. Stäckel, in the *Vierteljahrschrift der Naturforschenden Gesellsch. in Zürich*, 1910.

[3] Born at Zürich, January 12, 1746; died at Brugg, Aargau, February 17, 1827.

[4] Born at Utzensdorf, March 18, 1796; died at Bern, April 1, 1863.

(1825) a teacher in the Gewerbeakademie. In 1834 he became a professor in the University of Berlin, and from that time on he was a prolific contributor to geometry. The later years of his life were spent in Switzerland. He wrote several treatises of highest rank[1] and stands out as one of the prominent men in his field of work.[2] He extended Carnot's treatment of the complete quadrilateral to the n-gon in space, discussed the properties of ranges and pencils, and perfected the theory of curves and surfaces of the second degree.[3]

Lhuilier. Of the later Swiss contributors to elementary mathematics one of the best known was Simon-Antoine-Jean Lhuilier.[4] In his younger days he was a private tutor in Warsaw, after which he spent some time in Tübingen, then (1795) becoming professor of mathematics in Geneva. He wrote upon geometry while in Warsaw (1780) but first attracted attention by winning (1786) the prize offered by the Academy of Berlin on the nature of infinity. He wrote several elementary works but is known chiefly for his numerous memoirs on the measure and construction of polygons and polyhedrons and on the analogy between rectilinear and spherical right-angled triangles.[5] In the latter study he showed that the theorem analogous to the Pythagorean is

$$\sin^2 \tfrac{1}{2} a = \sin^2 \tfrac{1}{2} b \cdot \cos^2 \tfrac{1}{2} c + \sin^2 \tfrac{1}{2} c \cdot \cos^2 \tfrac{1}{2} b.$$

His contemporary, Jean Trembley (1749–1811), also a native of Geneva, a member of a family that contributed not a little to the appreciation of mathematics in Switzerland, wrote extensively on the calculus and its applications.

[1] *Systematische Entwickelung der Abhängigkeit geometrischer Gestalten von einander*, Berlin, 1832; *Die geometrischen Constructionen, ausgeführt mittelst der geraden Linie und eines festen Kreises*, Berlin, 1833; *Vorlesungen über synthetische Geometrie*, Leipzig, 2 vols., 1867.

[2] See his *Gesammelte Werke*, 2 vols., Berlin, 1881–1882.

[3] J. H. Graf, *Der Mathematiker Jakob Steiner von Utzensdorf*, Bern, 1897; J. Lange, *Jacob Steiners Lebensjahre in Berlin, 1821–1863*, Berlin, 1899; Cajori, *History of Math.*, 2d ed., p. 290.

[4] Born at Geneva, April 24, 1750; died March 28, 1840.

[5] "Analogie entre les triangles rectangles, rectilignes et sphériques," Gergonne's *Annales Math.*, I (1810–1811).

7. Other European Countries

Countries Considered. The Low Countries, Russia, Hungary, and Norway all added to the roll of eminent mathematicians in the 18th and 19th centuries, although space allows for the mention of only a few names.

Holland. Of the mathematicians of Holland in the 18th century Wilhelm Jacob Storm van s'Gravesande[1] may be taken as a representative. Beginning with the law, he later took up the teaching of mathematics, first (1717) at The Hague and then (1734) at Leyden. His was another case of the early display of mathematical ability, his essay on perspective having attracted attention when he was only nineteen years old. His first important publication[2] was devoted to an exposition of Newton's philosophy. His mathematical work consisted rather in making known the new theories of the calculus together with its applications than in original contributions,[3] but in physics his inventions and discoveries were considerable.

Belgium. As Belgium's representative may be selected Lambert-Adolphe-Jacques Quetelet.[4] He studied in Ghent and was only eighteen when (1814) he began teaching mathematics in the lyceum in his native city, an institution that was converted into a university in 1815. In the same year he received the degree of doctor of science, the first to be conferred by the new faculty. In 1818 he became director of the observatory at Brussels, an empty honor at the time because there was no building. In 1826 the erection of the observatory was decided upon and Quetelet's plans were adopted. Owing to political troubles it was not completed until 1832. In 1836 he became professor of astronomy and geodesy in the military school in

[1] Born at Herzogenbusch (Bois-le-Duc), September 27, 1688; died at Leyden, February 28, 1742.

[2] *Physices elementa mathematica, experimentis confirmata, sive Introductio ad philosophiam Newtonianam*, 2 vols., Leyden, 1720–1721.

[3] *Œuvres philosophiques et mathématiques*, Amsterdam, 1774 (posthumous).

[4] Born at Ghent, February 22, 1796; died at Brussels, February 17, 1874. Ed. Mailly, *Essai sur la vie et les ouvrages de Quetelet*, Brussels, 1875; English abstract, Smithsonian Institution *Report* for 1874, p. 169 (Washington, 1875).

Ghent. He was much interested in various phases of geometry, including spherical polygons, conics, three-dimensional figures, caustics, and stereographic projections; in astronomy and terrestrial magnetism; and, most of all, in the theory of probability,[1] a subject in which he was a recognized leader. He also wrote the leading works on the history of mathematics in Belgium.[2]

Scandinavia. Scandinavia produced a number of prominent mathematicians in this period, but the one of greatest genius was Niels Henrik Abel.[3] Like Eisenstein, Cotes, Clifford, Pascal, and Galois, he died young and yet attained very high rank in the domain of mathematical research. His memoir on elliptic functions did more for the theory than Legendre was able to accomplish in a long life. His name also attaches to certain functions which have since his time been the object of extended study. To him is due the first proof that an algebraic solution of the general equation of the fifth degree is impossible.[4]

Marius Sophus Lie[5] was the most prominent Scandinavian mathematician of his generation. He spent some time in France, but in 1872 became a professor at Christiania and from 1886 to 1898 taught at Leipzig. He is known chiefly for his work in differential equations, the theory of transformation groups (1873), differential geometry, and the theory of infinite continuous groups.

The Bolyais. The rise of non-Euclidean geometry is so closely connected with the name of Bolyai that a few words are appropriate concerning the family. The name is found in

[1] *Instructions populaires sur le calcul des probabilités*, Brussels, 1828; *Théorie des probabilités*, Brussels, 1845.

[2] *Histoire des sciences mathématiques et physiques chez les Belges*, Brussels, 1864; *Sciences mathématiques et physiques chez les Belges au commencement du XIX siècle*, Brussels, 1866.

[3] Born at Findöe, Norway, August 5, 1802; died at Arendal, April 6, 1829. L. Sylow, *Discours* at the centenary of his birth, Christiania, 1902.

[4] C. A. Bjerknes, *Niels-Henrik Abel*, French translation, Paris, 1885. This work is taken chiefly from articles in the *Nordisk Tidsskrift*, Stockholm, 1880.

[5] Born at Christiania, December 17, 1842; died at Christiania, February 18, 1899. G. Darboux, *Bulletin of the Amer. Math. Soc.*, V, 367; F. Engel, *Bibl. Math.*, I (3), 166; M. Noether, *Mathem. Annalen*, LIII, 1.

the Magyar records as early as the 13th century, the family belonging to the landed gentry, its estate lying in Bolya, a small town in Hungary. Farkas[1] Bolyai, a professor in a college[2] at Maros-Vásárhely,[3] had a son Farkas Bolyai,[4] who, after finishing his preparatory work, went to Göttingen (1796), where Kästner was closing his somewhat mediocre career and Gauss was beginning his brilliant one. Here he and Gauss became very intimate, exchanging ideas, taking long walks together, and together indulging in the few social recreations that they allowed themselves. Circumstances compelled Bolyai to return to his home in 1799, much against his personal desires. In 1804 he became professor of mathematics, physics, and chemistry in the college at Maros-Vásárhely, and here he remained until 1851. Here he wrote to Gauss two letters[5] on geometry, and here he published (1830) a little work in the Magyar language on arithmetic[6] and also (1832) his work on elementary mathematics.[7] The letters outline a book on geometry and show that he was interested in the subject of parallels. The book itself includes both algebra and geometry, and raises the question of the validity of Euclid's postulate of parallels. The general ideas of this book were in his mind when he went to Göttingen, 1796, and for a generation he had pondered upon the foundations of geometry. Among them appears the principle of the permanence of equivalent forms,[8] which English writers assign to Peacock (1830) and the Germans to Hermann Hankel (1867).

[1] Latin Wolfgangus; German Wolfgang. The standard authority on the subject is P. Stäckel, *Wolfgang und Johann Bolyai*, 2 vols., Leipzig, 1913, with excellent bibliography. There is also a "Vita di Giovanni Bolyai," by M. Darvai, in the *Atti* of the *Congresso internaz. di sci. storiche*, XII, 45.

[2] Corresponding to the German Gymnasium.

[3] *I.e.*, "market-place on the Maros" River.

[4] Born at Bolya, February 9, 1775; died at Maros-Vásárhely, November, 1856. Referred to in German works as Wolfgang Bolyai, and in other works occasionally in the Magyar form, Bolyai Farkas.

[5] September 16, 1804, and December 27, 1808.

[6] *Az arithmetica eleje* (*Elements of Arithmetic*), Maros-Vásárhely, 1830

[7] *Tentamen Juventutem Studiosam in Elementa Matheseos Purae . . . introducendi*, 2 parts, Maros-Vásárhely, 1832, 1833. The book bears the Imprimatur of 1829.

[8] Stäckel, *loc. cit.*, I, 35.

To the work of Farkas there was an appendix[1] written by his son, János Bolyai,[2] of whom the father had written to Gauss (1816) that this boy of fourteen already had a good knowledge of the differential and integral calculus and could apply it to mechanics, to the tautochronism of the cycloid, and to other lines of work, and that he knew Latin and astronomy. János went to the engineering school in Vienna when he was sixteen, and at twenty-one entered the army. In 1825 or 1826 he worked out the theory of parallels which he set forth in the appendix (1832) above mentioned, and in this is a clear discussion of the validity of Euclid's postulate of parallels and a presentation of a non-Euclidean geometry.[3]

The last years of Farkas Bolyai were unhappy ones, owing to the loss of his wife and the estrangement of his son. He wrote several other works,[4] however, including some on poetry.

Lobachevsky. In connection with the Bolyais it is natural to mention the work of Nicolai Ivanovitch Lobachevsky.[5] Although the son of a Russian peasant, he early showed a remarkable genius. He studied at the University of Kazan and when only twenty-one became professor of mathematics in that institution. In 1826 he made known through his lectures his conception of a geometry which should not depend upon the Euclidean postulate of parallel lines. These ideas were published in 1829, and in various later works.[6]

[1] *Appendix. Scientiam spatii absolute veram exhibens: a veritate aut falsitate Axiomatis XI Euclidei (a priori haud decidenda) independentem. . . .*

[2] Born at Klausenburg (in Magyar, Kolozsvár), December 15, 1802; died at Maros-Vásárhely, January 27, 1860. He also uses the Latin form, Johannes Bolyai.

[3] For a further discussion see Volume II, Chapter V.

[4] For a list of his works see Stäckel, *loc. cit.*, I, 205.

[5] Born at Nijni-Novgorod, November 2 (October 22, O.S.), 1793; died at Kazan, February 24 (February 12), 1856. The name is transliterated from the Russian in various ways, such as Lobatschevskij, Lobatschewsky, and Lobatcheffsky.

[6] *Ueber die Principien der Geometrie*, Kazan, 1829–1830; *Geometrische Untersuchungen zur Theorie der Parallellinien*, Berlin, 1840; French translation by Hoüel, 1866; English by Halsted, 1891; *Pangéométrie ou Précis de géométrie fondée sur une théorie générale et rigoureuse des parallèles*, Kazan, 1855. He also published various memoirs on other subjects. Vassilief's eulogy on Lobachevsky has been translated into French (1896) and English (1894).

Of the independent discovery of the non-Euclidean geometry by Lobachevsky and Bolyai there can be no doubt. The subject was in the general intellectual atmosphere of the time. Gauss, who was considering the question as early as 1792, had doubtless stimulated the elder Bolyai to study the problem, and no doubt had been stimulated in return. Both Lobachevsky and the younger Bolyai had been influenced by the Göttingen school. Each in his own way had attacked the question, and each had worked out his theory at about the same time (1825–1826); Lobachevsky published his theory first (1829), but János Bolyai published his independently (1832).[1]



NICOLAI IVANOVITCH LOBACHEVSKY
After a contemporary drawing

Kovalévsky. Among the Russian mathematicians of the latter part of the 19th century none was better known in western Europe than Sónya Krukovsky, who married Vladímir Kovalévsky and is commonly called by the name of Sophia Kovalévsky.[2] She was a pupil of Weierstrass, took her doctor's degree at Göttingen, and became professor of mathematics at Stockholm, where she was associated with Mittag-Leffler in the study of the function theory. Her own work was largely connected with the theory of differential equations,—a theory to which she made contributions of recognized value.

[1] On Bolyai's appreciation of Lobachevsky's work, see Stäckel, *loc. cit.*, I, 140.

[2] The date of her birth is unknown, but it was probably 1850. She was married in 1868 and died at Stockholm, February 10, 1891. See her autobiography, translated by Isabel F. Hapgood, New York, 1895.

Wronski. In the 19th century Poland produced only one mathematician who succeeded in attracting much attention abroad, and this was Hoëné Wronski.[1] He spent most of his life in France and wrote on the philosophy of mathematics. His *Introduction to a Course in Mathematics* appeared in London in 1821.

In Bohemia the subject of infinite series was studied by Bernhard Bolzano (1781–1848), but there were few other writers of prominence in the field of mathematics in this part of the world during the period now under consideration.



HOËNÉ WRONSKI

After an etching by Mme Frédérique O'Connell and autographed by Wronski

8. United States

Brief History. The United States inherited its first mathematics almost wholly from Great Britain. Early in the 19th century there was an influx of French mathematics.[2] Until Johns Hopkins University brought Professor

[1] Born August 24, 1778; died August 9, 1853. Since he wrote chiefly in French, the French spelling of his name is used. S. Dickstein has various references to and articles upon him in the second series of the *Bibliotheca Mathematica*, particularly VI (2), 48. See also his *Catalogue des œuvres imprimées et manuscrites de Hoëné Wronski*, Cracow, 1896.

[2] F. Cajori, *The Teaching and History of Mathematics in the United States*. Washington, 1890; J. Pierpont, "The History of Mathematics in the Nineteenth Century," *Bulletin of the Amer. Math. Soc.*, XI, 156; R. S. Woodward, "The Century's Progress in Applied Mathematics," *ibid.*, VI, 133; T. S. Fiske, *ibid.*, XI, 238; C. J. Keyser, *Educ. Rev.*, XXIV, 346.

Sylvester to this country (1876) for his second sojourn, and gave him the facilities for graduate work, little effort was made to encourage the study of modern mathematics. Several mathematicians of genuine ability had, however, developed before that time. Among these may be mentioned Benjamin Peirce,[1] who became a professor at Harvard at the age of twenty-four. He was for some time in charge of the U. S. Coast Survey, but is best known for his work on linear associative algebra (1870). Nathaniel Bowditch,[2] a self-made mathematician, was a generation older than Peirce. At the age of seventeen he began the study of Latin for the purpose of reading Newton's *Principia*, and later became proficient in French, Spanish, Italian, and German in order to study mathematics in these languages. His *New American Practical Navigator* (1802) and his translation of Laplace's *Mécanique céleste* (published in 1829–1839) gave him an international reputation.

It is proper at this time to mention Robert Adrain,[3] who contributed in a noteworthy manner to the progress of mathematics in his adopted country. His work on the form of the earth and on the theory of least squares showed that he was possessed of genuine mathematical ability. He founded the *Analyst* (1808) and the *Mathematical Diary* (1825).

In the domain of mathematical astronomy the most prominent of our native scholars in the 19th century was George William Hill.[4] He was for many years connected with the Nautical Almanac, but is known chiefly for his work in the lunar theory. When Poincaré visited this country he met Hill, and his first words were, "You are the one man I came to America to see," and he meant them.

Simon Newcomb (1835–1909) was a native of Nova Scotia but spent most of his life in the United States. He was

[1] Born at Salem, Massachusetts, April 4, 1809; died at Cambridge, October 6, 1880.

[2] Born at Salem, Massachusetts, March 26, 1773; died at Boston, March 16, 1838.

[3] Born at Carrickfergus, Ireland, September 30, 1775; died at New Brunswick, New Jersey, August 10, 1843.

[4] Born at Nyack, New York, 1838; died at Nyack, August 17, 1916.

connected with the Nautical Almanac and was also professor at Johns Hopkins. He was largely a self-made mathematician and was possessed of undoubted ability in the science.

Of the new school of mathematicians the best-known representative among those who have passed away was Maxime Bôcher (1867–1918). His special fields of research were those of linear differential equations, higher algebra, and the function theory. He was educated at Harvard and Göttingen and was called to the mathematical faculty at Harvard in 1891. In 1913–1914 he lectured at Paris on "les méthodes de Sturm dans la théorie des équations différentielles linéaires et leurs développements modernes."[1]

Josiah Willard Gibbs (1839–1903) was one of the most original of the American physicists of his time. He received his doctor's degree at Yale in 1863 and afterwards studied in Paris, Berlin, and Heidelberg. His interest in mathematics was great, particularly in the application of the science to mechanics.[2]


In the theory of probability, and in actuarial science in particular, Emory McClintock (1840–1916) is the best-known American contributor. He was for a time an assistant professor in Columbia College (later Columbia University) and then became a leader in life insurance work. He was the first president of the American Mathematical Society, was one of the founders of its *Transactions*, and was a frequent contributor to various mathematical journals.

9. The Orient

Nature of the Work. Except for the case of Japan, the Orient lost its initiative in mathematics when Western science was made known. India produced nothing that was distinctive in the 18th or 19th century. China, while occasionally protesting against Western mathematics, in reality sacrificed on the altar of the Jesuit missionaries her own originality in the science.

[1] W. F. Osgood, *Bulletin of the Amer. Math. Soc.*, XXV, 337.
[2] P. F. Smith, *Bulletin of the Amer. Math. Soc.*, X, 34.

Japan, owing to her policy of isolation, had little besides herself on which to depend, at least until the close of the 18th century, and so she continued to show her remarkable ingenuity in the application of her native calculus. This policy was unchanged until her doors were opened to Western civilization, when she, like China, lost that power of initiative along the lines of native research which her scholars had shown in the 17th and 18th centuries.



CHINESE TABLES OF SQUARES

This work of the 18th century, by Chang Tsu-nan, Fan Ching-fu, and Chiung Lin-tai, also contains a table of logarithms. It illustrates the early stages of modern tables in China

China. Chinese mathematics in the 18th and 19th centuries was at first characterized by the continued appreciation and study of the science introduced by the Jesuits. With this, however, there went a parallel study and appreciation of the native mathematics, and later in the 18th century there was a revulsion of feeling against the Jesuits, with the result that native mathematics came even more into favor, although with no appreciable progress. Among those who were most influential in the European movement was the Jesuit missionary Pierre Jartoux (1670–1720), who went to China in 1700. He was in correspondence with Leibniz and is known to have

interested the Chinese scholars in certain algebraic series. Among these scholars was Ch'ên Shih-jen (fl. 1715), who was skillful in the use of series of this kind. The century is noteworthy also for the general introduction into China of tables of various kinds, including logarithmic tables. In 1799 Yüan Yüan (1764–1849) published his great work[1] on the biographies of astronomers and mathematicians,—the most valuable book on the history of mathematics that has appeared in China. Supplements were afterwards added which made it more nearly complete.

Japan. In the 18th century the mathematics of Japan was enriched by a native calculus known as the *yenri*, or "circle principle," a name which may have been suggested by the title of the Chinese work of Li Yeh (1248), the *Tsĕ-yüan Hai-ching*.[2] Tradition asserts that Seki, the greatest mathematician of Japan in the 17th century, discovered the method, and there is strong circumstantial evidence to confirm this belief, but no works of his that are now known make any mention of the principle.

One of the pupils of Seki, Takebe Hikojirō Kenkō (1664–1739), may, however, have been the discoverer of the *yenri*, for he was an excellent mathematician and was probably familiar with certain European books. At any rate, he wrote upon the subject, the chief problem being to express an arc of a circle in terms of the versed sine. In Takebe's work $(\sin^{-1}x)^2$ is expressed in terms of versin x in three ways by means of three different infinite series. The question of the further development of the theory in Japan is not as interesting as that of the possible source of these series. Takebe himself does not seem to have understood their development, for his attempted explanations are very obscure. The series resemble those given by Wallis,[3] but they have not as yet been identified with any of the latter. In the form given by Ōyama (or Awayama)

[1] *Ch'ou-jen Ch'uan.*


[2] *Tsĕ-yüan* means "to measure the circle," and *Hai-ching* means "mirror of sea." See Smith-Mikami, 49, 143, and consult this work on the entire topic.

[3] *De Algebra Tractatus*, cap. xcvi (Oxford, 1693).

Shōkei in 1728, a modification of the most general form given by Takebe, the square of the arc may be expressed as

$$a^2 = 4\,dh\left[1 + \sum_{1}^{\infty} \frac{2^{2n+1}(n!)^2}{(2n+2)!} \cdot \left(\frac{h}{d}\right)^n\right],$$

where a is the length of the arc, d the diameter, and h the height of the arc (versed sine). This series is said to have been known to the Jesuit missionary Pierre Jartoux, already mentioned. There seems, therefore, to be good reason for believing that the *yenri* principle was suggested by a study of Jartoux, who in turn received it from European sources.



ARIMA'S PROBLEM OF INSCRIBED SPHERES

From Arima Raidō's *Shūki Sampō* of 1769. This shows the curious Japanese method of indicating a sphere by a circle with a lune on one side. In this work of Arima the value of π is given as

$$\frac{42822}{13630}\ \frac{45933}{81215}\ \frac{49304}{70117},$$


which is correct to twenty-nine decimal places

Among those who were intrusted with the secrets of the Seki school, since mathematical knowledge was still held as a kind of Pythagorean mystery, was Matsunaga Ryōhitsu (died 1744), who received them from his teacher Araki Hikoshirō Sonyei (1640–1718), a pupil of Seki's. Matsunaga did much toward the improvement of the native algebra, and by means of the *yenri* method computed the value of π to fifty figures.

Among the other Japanese writers of importance there was Miyake Kenryū, who wrote a well-known book of problems (1716) and another work in which he treated of the prismoid. About the same time Baba Nobutake (1706) wrote a work on astronomy that had much influence on the study of spherics.

Nakane Genjun (1701–1761), together with his father, Nakane Genkei (p. 440), contributed to the study of mathematics and astronomy. Each was influenced by the Chinese writings of the time, which in turn reflected the European

sciences which the Jesuit missionaries had made known in the East. One of the pupils of the younger Nakane was Murai Chūzen, who wrote (1765) a work[1] on numerical higher equations. In a later work (1781) he used the Pascal Triangle in expressing the coefficients of the terms in the expansion of a binomial. Arima Raidō (1714–1783), Lord of Kurume, was the first to publish the secret theory of algebra developed in the Seki school—a theory known as the *tenzan* algebra. As was the usual custom, Arima set forth his contributions in the form of problems. These problems related to indeterminate equations, the various roots of an equation, the application of algebra to geometry, the inscription of spheres within spheres, maxima and minima, binomial series, and stereometry. The most prominent of his protégés was Honda Teiken, better known by his later name Fujita Sadasuke (1734–1807). He wrote a notable work on algebra (1779),[2] together with various other treatises which show that he was an excellent teacher but a mathematician of no originality. Among his contemporaries was Aida Ammei (1747–1817), who gave a number of interesting series which he used in connection with his work in mensuration. Another contemporary and a fai




JAPANESE PROBLEM OF SPHERES TANGENT TO A TETRAHEDRON

This is typical of the native Japanese problems and shows the great patience of the Japanese scholar. The illustration is from an undated MS. by one Iwasaki Toshihisa (*c.* 1775)

[1] *Kaishō Tempei Sampō.* [2] *Seiyō Sampō.*

better mathematician, Ajima Chokuyen (1739–1798), was a contributor to the theory of indeterminate equations and to geometry. Among his solutions is an algebraic one of the Malfatti Problem. He also advanced the *yenri* theory by taking equal divisions of the chord instead, as his predecessors had done, of the arc.



WESTERN MATHEMATICS REPLACES THE *WASAN*

From a Japanese MS. of the middle of the 19th century, when European and American mathematics was beginning to replace the *wasan*, the native mathematics of Japan

The most interesting feature of Japanese mathematics in the 18th century is the slow penetration of Western theories into the domain of the *wasan*, or native mathematics. Japan showed great ingenuity in problem solving, but she never developed a great theory. Her nearest approach to originality in this respect was in her work on the *yenri*, and even here she seems to have had her start through contact with China and the Jesuit missionaries.

The 19th century opened with a general geodetic survey of the whole empire. By this time the European sources were

beginning to be available, but the problems were still of the ingenious native variety. In 1856 an arithmetic written on the European plan was published, and in 1859 an American work on the calculus, by Elias Loomis (1811–1899), was translated. Japanese mathematics assumed a new character from that time on. For better or worse the *wasan* was dead, and European science reigned in its stead. Whether Japan will ever be able to adapt her remarkable ingenuity to Western methods, or vice versa, remains to be seen.

10. The Historians of Mathematics

Seventeenth Century. Before referring to a few of the writers on the history of mathematics in the 18th century a word should be said concerning two of their predecessors. One of these was Bernardino Baldi,[1] an abbot, a native of Urbino, who wrote on mechanics but whose chief contribution was on mathematical biography,[2] his *Cronica* serving as a source to which writers on the history of mathematics have been much indebted. He was a linguist of unusual attainments and thus was able to secure his materials from several languages.

The second of these historians was John Wallis (1616–1703), whose contributions to mathematics in general have already been mentioned. His erudite *Treatise of Algebra*[3] contains a wealth of historical material and constitutes the beginning of the serious study of the history of mathematics in England.

Eighteenth Century. The first work bearing the name of history of mathematics was written by Johann Christoph Heilbronner,[4] a man who had given considerable attention to

[1] Born 1553; died 1617. The family name was Cantagallina.

[2] *Cronica de Matematici*, Urbino, 1707. His manuscripts on the subject were printed in Boncompagni's *Bullettino*, V, 427; VII, 337; XII, 420; XIX, 335, 437, 521; XX, 197. G. Zaccagnini, *Della vita . . . Baldi*, Reggio Emilia, 1918.

[3] English ed., London, 1685; Latin ed., Oxford, 1693. Some of the "additional treatises" in the English edition have dates 1684.

[4] Born at Ulm, 1706; died at Leipzig, *c.* 1747. His first work was the *Versuch einer Geschichte der Mathematik und Arithmetik*, Frankfort, 1739. His leading work is the *Historia Matheseos Universae a mundo condito ad seculum post Chr. Nat. XVI*, Leipzig, 1742.

theology and the mathematical sciences. He did some teaching in Leipzig, but his only work of importance was in the line of the history of mathematics. His style is sometimes prolix, and our present knowledge of details is naturally better than that of any writer of the middle of the 18th century, but the work has much value even today on account of its erudition, its extracts from early writers, and particularly its list of manuscripts then to be found in various libraries and its list of early printed books.



JEAN ÉTIENNE MONTUCLA

After an engraving by P. Viel, from a miniature

To Jean Étienne Montucla[1] are due two noteworthy treatises on the history of mathematics.[2] Of these the one on the quadrature of the circle was the first publication of the kind; and while the subject has greatly changed since Montucla wrote, the work is still a classic on the early history. His larger work was the first modern history of mathematics that may be called a classic. Montucla was a man of erudition, he wrote an excellent style, and there are no early histories more highly esteemed than his.


Of less importance in this field is the work of Charles Bossut,[3] a Jesuit, mathematical examiner at the École du Génie, Mézières, and later an examiner at the École Polytechnique.

[1] Born at Lyons, September 5, 1725; died at Versailles, December 18, 1799.

[2] *Histoire des recherches sur la quadrature du cercle*, Paris, 1754; 2d ed., by Lacroix, Paris, 1831; *Histoire des mathématiques*, 2 vols., Paris, 1758; 2d ed. by Lalande, 4 vols., Paris, 1799–1802.

[3] Born at Tarare, near Lyons, August 11, 1730; died in Paris, January 14, 1814.

He was primarily a writer of textbooks, but he also wrote various monographs relating to geometry. His history[1] is of unequal value, the part relating to ancient mathematics being much better than that on modern times. He shows a prejudice in favor of English mathematics,—a fact which may account for the translation and publication (1803) of the work in London.



AUTOGRAPH LETTER OF MONTUCLA
Written May 24, 1789, ten years before his death

Of the various Italian writers of the same period the best known is Pietro Cossali,[2] who taught both physics and astronomy at the University of Parma and was later (1806) professor of mathematics at Padua. He wrote various memoirs on mathematical questions, chiefly algebraic, but is best known for his history of algebra,[3] a work of considerable scholarship and even yet of service to the student of this phase of the subject.

The German historian of mathematics in the same period was Abraham Gotthelf Kästner,[4] Dozent (1739) and professor (1746) of mathematics in the University of Leipzig, and later

[1] *Essai sur l'histoire générale des mathématiques*, Paris, 2 vols., 1802; 2d ed., Paris, 1810; English translation, London, 1803.

[2] Born at Verona, June 29, 1748; died at Padua, December 20, 1815. G. Avanzini, *Elogio*, Modena, 1822.

[3] *Origine, trasporto in Italia, primi progressi in essa dell' Algebra*, 2 vols., Parma, 1797. On the contributions of the Italians to the more general history of mathematics see P. Riccardi, in the *Memorie della R. Accad. delle Scienze dell' Istituto di Bologna*, VI (5).

[4] Born at Leipzig, September 27, 1719; died at Göttingen, June 30, 1800.

(1756) at Göttingen. He was the first mathematician of prominence to write a work devoted entirely to the history of his subject.[1] Naturally, however, the best histories of mathematics have never been written by the best mathematicians, and Kästner's contribution in this line is not as important as his numerous though by no means brilliant works and memoirs on equations, geometry, hydrodynamics, and various other branches. The work is poorly arranged, contains no index, and shows the effect of hurried compilation from notes taken in the course of a wide reading.

Nineteenth Century. The 19th century has seen so many important works on the history of mathematics that it is possible to mention only a few of the earlier writers, together with some of the most prominent of the later ones who have already passed away.

Among the first in point of time to write upon the subject in this century was Pietro Franchini,[2] a priest whose life was devoted chiefly to the teaching of mathematics in various secondary schools of Italy. He was a mathematician of considerable power, writing several works on the various branches of the science and a number of essays of some originality on analysis. He wrote three works on the history of mathematics,[3] no one of which is of special importance to the student,[4] although each has some value in connection with the development of the science in Italy. Nothing of Franchini's, however, can be said to compare with Libri's work, which is mentioned later.

Arthur Arneth[5] was one of the minor German writers of the middle of the century. He was a teacher of mathematics and

[1] *Geschichte der Mathematik*, 4 vols., Göttingen, 1796–1800.

[2] Born at Partigliano, near Lucca, April 24, 1768; died at Lucca, January 26, 1837.

[3] *Saggio sulla storia delle matematiche*, Lucca, 1821, with *Supplementi al Saggio*, Lucca, 1824; *La storia dell' algebra e de' suoi principali scrittori*, Lucca, 1827, with a *Supplemento*, the same year; *Dissertazione sulla storia matematica dell' antica Nazione Indiana*, Lucca, 1830.

[4] G. Barsotti, *Nelle esequie fatte al Prof. Pietro Franchini*, with a list of his works (Lucca, 1837).

[5] Born at Heidelberg, September 19, 1802; died at Heidelberg, December 16, 1858.

physics in a Lyceum at Heidelberg and a Privatdozent in the University. He wrote a few memoirs of no special value, a work on geometry (1840), and a history[1] that may, at the best, be described as mediocre.

Hermann Hankel[2] might have been one of the greatest historians of mathematics had he enjoyed the ordinary span of life, for he not only knew mathematics remarkably well,[3] being a professor of the subject (1867) at the University of Leipzig, but he was versed in various Oriental languages and in those of ancient and modern Europe. He left a number of fragments on the history of mathematics which showed such power that they were collected after his death and published[4] by his father. Although made up of scattered notes, this work will well repay the student's attention.

Denmark produced in the 19th century one prominent writer on the history of mathematics, Hieronymus Georg Zeuthen.[5] He studied at Copenhagen, received the degree of doctor of philosophy in 1865, and became (1883) professor in the university. He was a member of various scientific societies and was one of the leading writers on the general history of Greek mathematics.

Of the British writers in the 19th century, George Johnston Allman[6] was one of the most scholarly. He became professor of mathematics in Queen's College, Galway, in 1853 and is well known for his *History of Greek Geometry from Thales to Euclid* (Dublin, 1889), a work unsurpassed in its line until the appearance of Sir Thomas Heath's history in 1921.

[1] *Die Geschichte der reinen Mathematik*, Stuttgart, 1852.

[2] Born at Halle, February 14, 1839; died at Schramberg, Schwarzwald, August 29, 1873.

[3] See his *Zur allgemeinen Theorie der Bewegung der Flüssigkeiten*, Göttingen, 1861; *Vorlesungen über die komplexen Zahlen und ihre Funktionen*, Leipzig, 1867 (Vol. I); *Die Elemente der projektivischen Geometrie*, Leipzig, 1875.

[4] *Zur Geschichte der Mathematik in Alterthum und Mittelalter*, Leipzig, 1874. He also wrote *Die Entwickelung der Mathematik in den letzten Jahrhunderten*, Tübingen, 1869; 2d ed., 1884.

[5] Born at Grimstrup, February 15, 1839; died at Copenhagen, January 6, 1920.

[6] Born at Dublin, September 28, 1824; died in 1904.

France, which led the world in the history of mathematics in the 18th century, produced few prominent writers on the subject in the century following. Libri, who wrote in France, was an Italian, and the only native writer of profound scholarship in the general field may be said to have been Paul Tannery.[1] Although he was "professor remplaçant" of Greek and Latin philosophy at the Collège de France (1892), he was never worthily recognized in academic circles. He was connected with government service, and devoted much of his leisure to writing upon the history of Greek mathematics, his essays easily ranking as the best produced in France at the close of the 19th century.

Of the earlier French writers Michel Chasles, whose name has already been mentioned (p. 498), was among the most important. Of the later biographers of mathematicians, Maximilien Marie (died May 8, 1891) was among the most pretentious writers. His work cannot, however, be recommended for its scholarship, nor can it be considered an accurate source of information.

Unquestionably the best-known work on the history of mathematics in the 19th century was that of Moritz Benedict Cantor.[2] He studied at Heidelberg and received his doctor's degree in 1851. His works include not merely the *Vorlesungen über Geschichte der Mathematik* (4 vols., 1880–1908), but also a work on Roman surveyors (1875), one on the mathematics of primitive peoples (1863), and a large number of monographs. He was also connected with various journals and was the founder and for a long time editor of the *Abhandlungen zur Geschichte der Mathematik* (Leipzig, from 1877) and editor of the "Historisch-literarische Abtheilung" of the *Zeitschrift für Mathematik und Physik* (from 1875). He began his work as Dozent in the University of Heidelberg in 1853 and became professor in 1877. The fourth volume of his *Geschichte* was chiefly the work of various contributors who carried out his plans.

[1] Born at Mantes, December 20, 1843; died November 27, 1904.

[2] Born at Mannheim, August 23, 1829; died at Heidelberg, April 10, 1920.

Of the Austro-German writers, Moritz Steinschneider[1] deserves special mention for his numerous essays on Hebrew and Arabic mathematics.

The best known of the Italian historians is probably Libri.[2] He was educated at Pisa and for some time was professor in the university of that city. He went to Paris as a political refugee in 1830, was naturalized in 1833, and became professor of analysis at the Sorbonne. His *Histoire des sciences mathématiques en Italie* was published at Paris (4 vols., 1838–1841). On the day when the printing of the first volume was completed he called at the printer's and carried away a few copies. An hour later all the rest of the edition was burned, and the work had to be reset.[3] Libri was accused (1847) of stealing valuable books and manuscripts, and indeed did, in some way not satisfactorily explained to his critics, acquire certain rare works for his large library. He fled to England but was (1850) found guilty in Paris and was sentenced to ten years' imprisonment. The latter part of his life was spent in exile.

Much more important as a contributor to the history of the science, however, Principe Baldassare Boncompagni[4] became well known through his *Scritti di Leonardo Pisano* (1857–1862), although he had already written on the same subject. His greatest contribution was his *Bullettino di Bibliografia e Storia delle Scienze matematiche e fisiche* (Rome, 1868–1887). In spite of the fact that this monumental series has a vast amount of infinitesimal detail and that pages were frequently changed in type as the printing progressed, it will always remain as a standard work of reference.

[1] Born at Prossnitz, Moravia, March 30, 1813; died at Berlin, January 24, 1907.

[2] Conte Guglielmo Bruto Icileo Timoleon Libri-Carucci dalla Sommaja. Born at Florence, January 2, 1803; died near Fiesole, September 28, 1869. G. Loria, "Guglielmo Libri come storico della scienza," *Atti della Soc. Ligustica di Sci. Nat. e Geog.*, XXVIII, No. 3, with excellent bibliography; A. Stiattesi, *Commentario storico-scientifico sulla vita e le opere del Conte Guglielmo Libri*, Florence, 2d ed., 1879.

[3] One of the original copies with Libri's corrections was given by him to Professor Jacoli of Venice and by him to the present writer.

[4] Born at Rome, 1821; died at Rome, 1894. See G. Codazza, "Il principe Boncompagni," *Politecnico*, Vol. XX (Milan).

The best known of the historians of special branches is Anton von Braunmühl,[1] who is naturally thought of as a German by birth because of his long professorship at Munich, although he was born on Russian soil. His history of trigonometry (Leipzig, 2 vols., 1900, 1903) is the standard work on the subject.

In addition to these writers there are many others who might properly be mentioned but whose names are not, in the main, so familiar to the general reader of mathematical literature. A few names will be added, however, for the purpose of showing the general interest taken in the subject during the 19th century.

Richard Baltzer (1818–1887), professor at Giessen, wrote various notes upon special points in mathematical history; David Bierens de Haan (1822–1895), professor at Leyden, contributed to the history of mathematics in the Netherlands; Gumersindo Vicuña (1840–1890), a native of Havana, wrote upon the history of Spanish mathematics; William Whewell (1794–1866) wrote upon the history of the inductive sciences (1837); Heinrich Suter (born in 1848) wrote upon the general history of mathematics, but his most valuable contribution was to the biography of Arab mathematicians; E. L. W. M. Curtze (born in 1837) published a large number of memoirs upon the history of the science; G. Milhaud (1858–1918) published several essays upon the achievements of the Greek scientists and left a work on Descartes which appeared posthumously (1921); Alexander Macfarlane (1851–1913) published a series of lectures upon ten of the leading British mathematicians of the 19th century; A. M. Rebière (1841–1900) wrote the lives of woman mathematicians; Paul Mansion (1844–1919), professor at Ghent and editor of *Mathesis*, published a history of the calculus; Max Simon (1844–1918) published various essays and works upon the history of mathematics; J. H. Graf (1852–1918) published a history of mathematics in Switzerland.

Some of these names have already been mentioned, and the brief list here given has been supplemented by many references in the footnotes. It serves to show, however, that the 19th century developed a new interest in the history of mathematics.

[1] Born at Tiflis, December 22, 1853; died at Munich, March 9, 1908.

TOPICS FOR DISCUSSION

1. Meaning of elementary mathematics and the approximate date at which, for most practical purposes, it was brought to its present state of development.

2. A comparison of the work of those mathematicians who ranked as infant prodigies with that of those who developed later in life.

3. Effect of Newton upon the mathematics of Great Britain.

4. Evidence of opposition to Newton's mathematical theories.

5. The contest in Great Britain between the fluxional and the differential notation.

6. A comparison of the general nature of the British mathematics in the 18th century with that of the French.

7. The general nature of the problems which were suggested by the calculus and which attracted the attention of mathematicians in the 18th century.

8. The three most noteworthy mathematicians of France whose work fell chiefly in the 18th century, with a sketch of the life and contributions of each.

9. The rise of the metric system.

10. A consideration of the statement that the close of the 18th century was an era of giants in the field of mathematics.

11. The new branches of mathematics developed in the 18th century, with the names of those chiefly involved in their development.

12. The rise of modern pure geometry.

13. The general nature of the work done by Italian mathematicians in the 18th and 19th centuries.

14. The three most noteworthy Italian mathematicians whose work fell chiefly in the 18th century, with a sketch of the life and contributions of each.

15. The contributions of Switzerland in the 18th century.

16. The general nature of the Oriental mathematics in the 18th century, and particularly that of Japan.

17. The contributions of Germany in the 19th century.

18. The general nature of the mathematics of the 19th century compared with that of the centuries immediately preceding.

19. The general nature of the standard works on the history of mathematics, with a comparison of their purposes, their scholarship, and their methods.

CHRONOLOGICAL TABLE

This table includes the more important mathematical names mentioned in this volume up to the year 1850.

Through the year 1800 the relative importance of the names, judged chiefly by their influence upon elementary mathematics, is indicated by three sizes of type, although it is evident that such a distinction is often a matter on which opinions might reasonably differ. In order to assist the student in placing mathematical events in their proper relation to world history, a number of well-known historical events (with their dates), through the year 1804, have also been inserted, italic type being used for this purpose.

In general the dates are only approximations, as is seen in the large number of items under centennial and semicentennial years. Students who wish the precise dates can easily find them, where they are known, by consulting the pages mentioned in the Index.

After each mathematical name there is given a brief statement of the major interest of the individual, particularly with respect to elementary mathematics, although it is evident that in the cases of men like Newton, Euler, Gauss, and Lagrange the statement cannot give any satisfactory idea of the range of work accomplished. Where the major interest is given as astronomy or physics, the individual applied mathematics in his work.

In this table, dates before the Christian Era are, in the left-hand column, indicated by a minus sign.

The significance of the styles of type will be understood from the following scheme:

1. *Historical names and events.*
2. Mathematical names and events.
3. **Important mathematical names and events.**
4. ***Mathematical names and events of greatest importance.***

-50000. *Early Stone Age begins. (Rough approximation of date.) Means of making fire discovered.*

-15000. *Middle Stone Age begins. Works of art appear.*

-5000. *Late Stone Age begins.*

-4700. Possible beginning of Babylonian calendar.

-4241. Egyptian calendar introduced.

-4000. *Metal discovered.*
-3500. *Writing in use.*
-3000. *Early Babylonia. Sargon I, 2750 B.C.; Hammurabi, 2100 B.C. Earliest stone masonry.* Egyptian reliefs refer to taxes.
-2900. *Great Pyramid built.*
-2852. *Fuh-hi, reputed first emperor of China.* Astronomical observations.
-2700. *Huang-ti reigns in China.* Astronomy and arithmetic.
-2400. Babylonian tablets of Ur record measures.
-2350. *Yau reigns in China.* Astronomy.
-2200. Date of many mathematical tablets found at Nippur.
-2100. *Hammurabi king of Babylon.* The calendar.
-2000. *Postal system in Asia.*
-1850. *Amenemhat (Amenemmes) III reigns in Egypt.* Surveying, leveling. Oldest astronomical instrument.
-1650. **Ahmes (Rhind) papyrus.**
-1500. Oldest Egyptian sundial.
Reliefs show tax lists.
Babylonians knew simple rules of mensuration.
Queen Hatshepsut reigns in Egypt.
-1350. Date of later mathematical tablets found at Nippur.
Rollin papyrus with elaborate problems about bread.
Seti I reigns in Egypt.
-1347. Rameses II (Sesostris) said to have redivided land in Egypt.
-1180. Harris papyrus with list of temple wealth in Egypt under Rameses III.
-1150. Won-wang may have written the I-king.
-1122. First historical period of Chinese mathematics.
Chow Dynasty begins with the reign of Wu Wang.
-1105. **The Chóu-peï, Chinese classic in mathematics, may have been written. Also (but possibly as early as the 27th century B.C.) the K'iu-ch'ang Suan-shu.**
-1055. *David becomes king of Israel.*
-1032. First historical record of rules for Chinese currency by weight.
-1000. *Vedic literature begins (c. 1000–c. 800 B.C.).*
-753. *Rome founded.*
-750. *Assyrian Empire (c. 750–606 B.C.).*
-670. Knife money appears as a Chinese coinage.
-660. Tradition of Japanese numeration to high powers of 10.
Coins appear in China in circular form.
-650. Coins struck in Lydia, Asia Minor.
-606. *Chaldean Empire (606–539 B.C.).*
-600. ***Thales. Demonstrative geometry.***
Solon. Calendar.
-575. Anaximander. Gnomon.

-561. *Reign of Nebuchadnezzar closes.*
-550. Ameristus. Geometry.
-542. Bamboo rods used in China for calculating.
-540. ***Pythagoras. Geometry, theory of numbers. Earth spherical.***
-539. *Cyrus captures Babylon.*
-530. Anaximenes. Astronomy.
-517. Hecataeus. Maps.
-500. Śulvasūtras (date very uncertain). Pythagorean numbers.
-485. *Xerxes begins to reign.*
-480. *Battle of Thermopylæ.*
-475. Dispersion of Pythagorean brotherhood.
-470. Agatharcus, Athenian. Perspective.
-465. **Œnopides of Chios. Geometry.**
-460. **Hippocrates of Chios. Quadrature.**
Parmenides. Astronomy.
-450. Zeno. Paradoxes of motion.
Herodotus the historian.
-444. *Pericles becomes supreme in Athens.*
-443. *Phidias begins the Parthenon*
-440. Leucippus. Atomic theory.
Anaxagoras. Geometry.
-432. Meton, Phæinus, Euctemon. Astronomy.
-430. **Antiphon. Method of exhaustion.**
-425. **Hippias of Elis. Quadratrix.**
Theodorus of Cyrene. Irrationality.
Philolaus. Gnomon.
Socrates. Induction and definition.
-410. **Democritus. Atomic theory, irrationals.**
-400. **Archytas. Proportion.**
-380. Leodamas. Analytic proof.
Plato. Foundations of mathematics.
-375. **Theætetus. Geometry.**
Callippus. Greek astronomer.
Chinese coins with weight or value stamped.
-370. **Eudoxus. Proportion.**
-350. **Menæchmus. Conics.**
Deinostratus. Quadratrix.
Philippus Medmæus. Geometry.
Theophrastus. History of mathematics.
Chinese compass.
Xenocrates. Theory of numbers. History of geometry.
-340. ***Aristotle. Applications of mathematics, logic.***
Speusippus. Proportion.

-336. *Alexander the Great begins his reign.*
-335. Eudemus. History of mathematics.
-332. *Alexandria founded.*
-330. Autolycus. Geometry.
-323. *Alexander the Great dies. Ptolemy Soter begins to reign in Egypt.*
-320. Aristæus. Solid loci. Conics.
Dicæarchus. Mensuration.
-300. ***Euclid. Geometry.***
-260. Aristarchus. Astronomy.
Conon. Astronomy. Spiral of Archimedes.
-250. Nicoteles. Conics.
Berosus introduces Chaldean astronomy into Greece.
-247. *Ptolemy Euergetes begins his reign in Egypt. Patron of learning.*
-230. **Eratosthenes. Prime numbers, geodesy.**
-225. ***Apollonius. Conics.***
Archimedes. Geometry, infinite series, mechanics.
Ch'êng Kiang Chen. Knotted cords.
-213. *Shī Huang-ti, emperor of China, burns all the books.*
-200. Ch'ang Ts'ang revises the "Nine Sections."
-180. **Hypsicles. Astronomy, number theory.**
Nicomedes. Conchoid.
Diocles. Cissoid.
Zenodorus. Isoperimetry.
-150. Perseus. Sections of an anchor ring.
-140. **Hipparchus. Astronomy, trigonometry.**
-106. *Pompey and Cicero born.*
-100. *Caesar (100–44 B.C.) born.*
Wu-ti (140–87 B.C.) opens communications with the West.
-77. Poseidonius. Geometry, cosmography.
-75. *Cicero discovers the tomb of Archimedes.*
-63. *Augustus (63 B.C.–14 A.D.) born.*
-60. Geminus. History of mathematics.
P. Nigidius Figulus. Astronomy.
Marcus Terentius Varro. Mensuration.
-51. *Caesar completes the subjugation of Gaul.*
-50. Dionysodorus. Geometry.
-46. Cæsar, assisted by Sosigenes, reforms the calendar.
-40. Cleomedes. Astronomy, arithmetic.
-20. Marcus Vitruvius Pollio. Applied mathematics.
-8. Diodorus of Sicily. History of mathematics.
-4. *Probable date of the birth of Jesus.*
1. *Rag paper known in China.*
10. Strabo. In his geography considerable history of mathematics.

25. Columella. Surveying.
50. **Heron of Alexandria. Geodesy, mathematics. (Possibly *c.* 200.)**
Serenus of Antinoopolis. Cylindric sections.
Sun-tzï wrote the Wu-ts'ao Suan-king.
64. *Buddhism introduced into China.*
66. Liu Hsing devises a new Chinese calendar.
75. Pliny the Elder. His Natural History is valuable for the study of Roman numerals.
Pan Ku. Bamboo rods in use in Chinese computation.
100. **Nicomachus. Theory of numbers.**
Menelaus. Spherics, anharmonic ratio.
Ch'ang ch'un-ch'ing. Commentary on the Chóu-peï.
Theodosius. Geometry, astronomy.
Balbus. Surveying.
Frontinus. Surveying.
120. Hyginus. Surveying.
125. **Theon of Smyrna. Theory of numbers. History of Pythagoras.**
Ch'ang Höng. Astronomy, geometry, $\pi = \sqrt{10}$.
150. **Claudius Ptolemaeus (Ptolemy). Astronomy, trigonometry, geodesy.**
Marinus of Tyre. Geodesy, geography.
166. *Marcus Aurelius sends an embassy to China.*
180. Nipsus. Surveying.
190. Ts'ai Yung. Chinese calendar.
200. Epaphroditus. Surveying, theory of numbers.
Domitius Ulpianus. Mortality table.
Quintus Sammonicus Serenus. General mathematics.
220. Sextus Julius Africanus. Encyclopedia with some history of mathematics.
235. Censorinus. Astronomy.
245. Wang Pi. On the I-king.
250. Siu Yo. Arithmetic.
Hsü Yüeh. Commentary on Siu Yo.
263. Liu Hui. Wrote the Hai-tau Suan-king.
265. Wang Fan. Astronomy; $\pi = \frac{142}{45}$.
275. ***Diophantus. Algebra, theory of numbers.***
Sporus of Nicæa. History of mathematics.
280. Anatolius. Astronomy.
Porphyrius. Life of Pythagoras.
289. Liu Chih. Possibly the one who gave $\pi = 3.125$.
300. **Pappus. Geometry.**
323. *Constantine becomes sole emperor.*
324. *Constantinople founded.*
325. **Iamblichus. Theory of numbers.**

325. Metrodorus. (Possibly, but see 500.)
340. Julius Firmicus Maternus. Astrology.
372. *Buddhism introduced into Korea.*
379. *Theodosius the Great, emperor of the East.*
390. Theon of Alexandria. Geometry.
400. Fa-hién, Chinese Buddhist in India. Hindu mathematics becomes known in China.
Sūrya Siddhānta written in the 4th or 5th century.
410. Hypatia of Alexandria. Geometry and astronomy.
Synesius. Astrolabe.
Sack of Rome by Alaric.
425. Wang Jong. Arithmetic.
Tun Ch'uan wrote the San-tòng-shu.
433. *Attila, king of the Huns.*
440. P'i Yen-tsung. Circle measure.
450. Ho Ch'êng-t'ien. Astronomy.
Wu, a Chinese geometer. $\pi = 3.1432+$.
Domninus. Theory of numbers.
Victorius. Computus.
455. *Sack of Rome by Genseric.*
460. **Proclus. Geometry.**
Capella. Encyclopedia.
470. Tsu Ch'ung-chih. $\pi = \frac{3}{1}\frac{1}{3}\frac{5}{5}$.
485. Marinus of Flavia Neapolis. On Proclus.
500. Metrodorus. Arithmetical epigrams in the Greek Anthology.
505. Varāhamihira. Hindu astronomy.
510. **Boethius. Geometry, theory of numbers.**
Āryabhaṭa the Elder. General mathematics. $\pi = 3.1416$.
Damascius. Geometry.
518. *Hui-sing, Chinese Buddhist, visits India.*
520. Cassiodorus. Computus, encyclopedia.
522. *Buddhism introduced into Japan (522–552).*
525. Dionysius Exiguus. Christian calendar.
Anthemius. Architecture, conics.
527. *Justinian's reign begins.*
529. *St. Benedict founds the monastery at Monte Cassino.*
535. Ch'on Luan wrote the Wu-king Suan-shu.
550. Hsia-hou Yang wrote his Suan-king.
Codex Arcerianus written probably about the 6th century. Surveying.
554. Korean scholars introduce Chinese mathematics into Japan.
560. Eutocius. History of geometry.
575. Ch'ang K'iu-kien. Arithmetic.
Men. $\pi = 3.14$.

600. Prince Shōtoku Taishi. Arithmetic.
602. Korean priests bring works on the calendar to Japan.
610. Stephen of Alexandria. Astronomy and general mathematics.
Isidorus. Encyclopedia.
615. *Arab ambassadors visit China.*
622. *Flight of Mohammed from Mecca. Beginning of the Hegira era.*
625. Wang Hs'iao-t'ung. Numerical cubic equations.
628. **Brahmagupta. Geometry, algebra.**
629. *Hüan-tsang goes to India. Translates Hindu works.*
635. Asclepias of Tralles. On Nicomachus.
636. *Priest from Rome visits China.*
640. Joannes Philoponus. Astrolabe, on Nicomachus.
642. *Library of Alexandria burned.*
650. Sebokht. Hindu numerals.
670. *Emperor Tenchi (Tenji) reigns, 668–672.* Observatory established.
Arithmetic.
710. **Bede. Calendar, finger reckoning.**
711. *Saracens invade Spain.*
713. *Arab ambassadors visit China and foreign ships sail to Canton.*
727. I-hsing. Chinese calendar, indeterminate equations.
732. *Battle of Tours. Charles Martel defeats the Saracens.*
750. Akhmim Papyrus written *c.* 7th or 8th century.
756. *Cordova made the seat of the western caliphate.*
762. *Bagdad founded by al-Mansûr, c. 762–763.*
766. The Sindhind translated into Arabic. Hindu numerals.
770. Geber. Alchemy, astrolabe.
771. *Charlemagne's sole reign begins (771–814).*
775. **Alcuin called to the court of Charlemagne. Mathematical problems.**
Ya'qûb ibn Ṭâriq. The sphere.
Abû Yaḥyâ. Translated Ptolemy.
Kia Tan. Geographer.
790. *Harun al-Rashid (reigned 786–808/9).* Patron of mathematics.
Al-Fazârî. Mathematical instruments.
800. Jacob ben Nissim. Theory of numbers.
Messahala. The astrolabe.
Al-Tabarî. Astronomy.
Harun al-Rashid sends an embassy to China.
820. **Mohammed ibn Mûsâ al-Khowârizmî. Algebra.**
Hrabanus Maurus. Computus.
Al-Nehâvendî. Astronomy.
Al-Hajjâj. Greek mathematics.
Al-Mâmûn (reigned 809–833). Patron of mathematics.

830. Al-'Abbâs. Greek mathematics.
Al-Aṣṭorlâbî. The astrolabe.
840. **Ḥonein ibn Isḥâq. Greek mathematics.**
Walafried Strabus (Strabo). Teacher.
850. **Mahāvīra. Arithmetic, algebra, mensuration.**
Sahl ibn Bishr. Astronomy, arithmetic, algebra.
Al-Arjânî. Greek mathematics.
Abû'l-Ṯaiyib Trigonometry.
860. **Alchindi. Astronomy, optics, proportion.**
Almâhânî. Trigonometry, cubic equation.
Al-Mervazî. Astronomy.
870. **Ṯâbit ibn Qorra. Conics. Greek mathematics.**
The Three Brothers. Geometry, astronomy.
871. *Alfred the Great begins his reign.*
880. Al-Ḥimṣî. Greek mathematics.
Albumasar. Astronomy.
890. Aḥmed ibn al-Taiyib. Algebra.
Aḥmed ibn Dâ'ûd. Algebra.
Tenjin. Japanese patron of mathematics.
900. **Abû Kâmil. Geometry, algebra.**
Isḥâq ibn Ḥonein ibn Isḥâq. Greek mathematics.
Remigius of Auxerre. On Capella.
Muslim ibn Aḥmed al-Leiṯî. Arithmetic.
Al-Qass. On Euclid.
Qosṭâ ibn Lûqâ. On Diophantus.
Al-Miṣrî. Geometry.
910. **Al-Nairîzî. Geometry.**
Al-Faraḍî. Arithmetic.
915. Sa'id ibn Ya'qûb. Greek mathematics.
920. **Rhases. Geometry.**
Albategnius. Astronomy.
Odo of Cluny (879–*c.* 942). Abacus.
925. Al-Ḥasan ibn 'Obeidallâh. On Euclid.
Aethelstan's reign begins in England. Learning fostered.
940. Al-Fârrâbî. On Euclid and Ptolemy.
950. Hasan. (Date very doubtful.) The calendar.
Bakhshālī manuscript. Algebra. (Date very doubtful).
960. Abû Ja'far al-Khâzin. Geometry.
970. Hrotsvitha, a nun. Number theory.
975. Al-Ḥarrânî. On Euclid.
980. **Abû'l-Wefâ. Trigonometry.**
Abbo of Fleury. Computus.
987. **Abû'l-Faradsh. The Fihrist.**

993. Bernward. Theory of numbers.
Al-Masîḥî. On Ptolemy.
1000. Mohammed ibn al-Leit. Geometry.
Al-Majrîṭî. Theory of numbers.
Ḥâmid ibn al-Khiḍr. Astrolabe, algebra.
Al-Ḥasan (al-Ḥaiṯam) of Baṣra. Algebra, geometry.
Manṣûr ibn 'Alî. Trigonometry.
Gerbert (Sylvester II). Arithmetic.
Byrhtferth. Calendar.
Ibn Yûnis. Astronomy.
Avicenna. Geometry, arithmetic.
Albêrûnî. On Hindu mathematics.
1020. **Al-Karkhî. Algebra.**
Bernelinus. Arithmetic.
Śrîdhara. Arithmetic.
Firdusi, Persian poet, dies.
1025. **Al-Nasavî. Greek mathematics.**
Ibn al-Ṣaffâr. Astronomical tables.
1028. Guido of Arezzo (Aretinus). Arithmetic.
1042. *Edward the Confessor becomes king.*
1050. **Hermannus Contractus. Arithmetic, astrolabe.**
Ch'on Huo. Astronomy.
Ibn al-Zarqâla. Astronomy.
Wilhelm of Hirschau. Teacher.
1066. *Norman Conquest.*
1075. Psellus. Quadrivium.
Franco of Liège. Arithmetic, geometry.
1076. *Turks capture Jerusalem.*
1077. Benedictus Accolytus. Rithmomachia.
1083. China prints Liu Hui's classic. Block book.
1084. China prints Ch'ang K'iu-kien's arithmetic.
1090. Fortolfus. Rithmomachia.
1095. *First Crusade proclaimed.*
1100. **Savasorda. Geometry.**
Omar Khayyam, the poet. Algebra, astronomy.
Abû'l-Ṣalt. Geometry.
Walcherus. Geometry, arithmetic, astronomy.
1115. China prints the Huang-ti K'iu-ch'ang.
1120. **Plato of Tivoli. Translates from the Arabic.**
Adelard of Bath. Translates from the Arabic.
1125. Radulph of Laon. Arithmetic.
1130. Jabir ibn Aflah. Trigonometry.
1137. Gerland of Besançon. Computus.

1140. **Abraham ben Ezra (Rabbi ben Ezra). Theory of numbers, magic squares, calendar.**
Avenpace. Geometry.
Johannes Hispalensis. Translates from the Arabic.
Robert of Chester. Translates from the Arabic.
1144. Rudolph of Bruges. Translates Ptolemy.
1146. *Second Crusade proclaimed.*
1150. **Gherardo of Cremona. Translates from the Arabic.**
Bhāskara. Algebra.
Fujiwara Michinori. Mensuration.
Gherardo da Sabbionetta. Translates from the Arabic.
N. O'Creat. Arithmetic.
1175. **Averroës. Astronomy, trigonometry.**
Maimonides. Astronomy.
Samuel ben Abbas. Arithmetic.
Al-Ḥaṣṣâr. Arithmetic.
1180. Ts'ai Yuan-ting. On the I-king.
1200. Ta'âsîf. On Euclid.
Ibn Yûnis. Conics.
Ibn al-Yâsimîn. Algebra.
Al-Râzî. Geometry.
Daniel Morley. Translates from the Arabic.
Alpetragius. Astronomy.
Ibn al-Kâtib. Geometry.
Al-Ṭûsî. Geometry, algebra.
1202. **Leonardo Fibonacci. Algebra, arithmetic, geometry.**
1220. *Chinghiz Khan's great expedition to Europe.*
1225. **Jordanus Nemorarius. Algebra.**
Michael Scott. Translates from Greek and Arabic.
Genshō. Arithmetic.
1230. Ye-lü Ch'u-ts'ai. Astronomy.
Barlaam. Algebra, Euclid.
1240. Jehuda ben Salomon Kohen. On Ptolemy.
Alexandre de Villedieu. Arithmetic.
Robert Greathead. Geometry computus.
John of Basingstoke. Translates from the Greek.
1250. **Sacrobosco. Numerals and the sphere.**
Naṣîr ed-dîn. Trigonometry.
Roger Bacon. Astronomy, general mathematics.
Ch'in Kiu-shao. Higher numerical equations.
Liu Ju-hsieh. Algebra.
William of Moerbecke. Translates from the Greek.
Li Yeh. General mathematics.

1250. Isaac ben Sid. Astronomical tables.
Albertus Magnus. Astronomy, physics.
Vincent de Beauvais. Quadrivium.
Guglielmo de Lunis. (Period very uncertain.) Translates algebra from the Arabic.
Prophatius. Translates Euclid and Menelaus.
Rise of European universities.
Europeans visit the Mongol court.

1260. **Campanus. Translates Euclid.**
Ibn al-Lubûdî. Algebra, Euclid.
Kublai Khan's reign begins.

1261. Yang Hui. On the "Nine Sections."

1265. Petrus de Maharncuria. Magnet, general mathematics.

1270. Bar Hebræus. On Euclid.
Witelo. Perspective.

1271. *Marco Polo begins his travels.*

1275. Oldest algorism in French.
Liu I. General mathematics.
Arnaldo de Villa Nova. Computus.
Alfonso X. Astronomical tables.
Cimabue, the Italian painter.
Guelphs and Ghibellines in Florence (1183–1295).

1280. John Peckham (*c.* 1230–1295). Perspective.

1286. Friar Odoric goes to Canton.
Western learning.

1290. Kóu Shóu-king. Mensuration.

1297. Bartolomeo da Parma. Geometry, astronomy.

1299. **Chu Shï-kié. Algebra.**

1300. **Albanna. Algebra, proportion.**
Pachymeres. General mathematics.
Cecco d'Ascoli. On Sacrobosco.
Hauk Erlendsson. Algorism.
Pietro d'Abano. Astrolabe.
Andalò di Negro. Arithmetic, astronomy.

1315. *Dante (1265–1321).*

1320. John Manduith. Trigonometry.
Kalonymos ben Kalonymos. On Nicomachus.

1325. Petrus de Dacia. Geometry.
Thomas Bradwardine. Geometry, arithmetic.
Walter Burley. Greek mathematician.

1330. Joannes Pediasimus. Geometry.
Levi ben Gerson. Arithmetic.
Isaac ben Joseph Israeli. Geometry.

1330. Richard of Wallingford. Trigonometry.
1340. Maximus Planudes. On Diophantus, arithmetic.
Johannes de Lineriis. Alfonsine tables, arithmetic.
Paolo Dagomari. Arithmetic.
Master Sven. The sphere.
1341. Nicholas Rhabdas. Arithmetic, finger symbols.
1345. Richard Suiceth. Coordinates.
1346. *Battle of Crécy. Attack on feudalism.*
1349. *Black Death destroys large per cent of European population.*
1350. Joannes de Muris. Arithmetic, calendar.
Chunrad von Megenberg. The sphere.
Ibn al-Shâṭir. Trigonometry.
Boccaccio (1313–1375).
Petrarch (1304–1374).
1360. **Nicole Oresme. Exponents, proportion, coordinates.**
Walter Bryte. Arithmetic.
Jacob Poël. Astronomy.
Imanuel ben Jacob. Astrolabe.
1365. Heinrich von Hessen. Geometry.
Albert of Saxony. Geometry.
1369. *Tamerlane's reign begins.*
1375. Simon Bredon. Geometry.
Jacob Carsono. Astronomy.
1380. Rafaele Canacci. Algebra.
Joseph ben Wakkar. Astronomy.
1383. Antonio Biliotti. Arithmetic.
1384. *Wycliffe's English Bible completed.*
1390. Uniform weights and measures in most of England.
1392. Moschopoulus. Magic squares.
1400. Ibn al-Mejdî. Trigonometry.
Matteo, Luca, and Giovanni da Firenze. Arithmetic.
Petrus de Alliaco. Computus.
Conrad von Jungingen. Geometry.
Biagio da Parma. Perspective.
1410. Prosdocimo de' Beldamandi. Algorism, geometry.
1420. *Era of the Medici in Florence.*
1424. Rollandus. Theory of numbers, algebra.
1425. Leonardo of Cremona. Trigonometry.
1430. **Johann von Gmünden. Trigonometry.**
Jacob Caphanton. Arithmetic.
1431. *Joan of Arc burned.*

1435. Ulugh Beg. Astronomy.
John Killingworth. Algorism, astronomy.
1440. *Donatello the artist, Florence (1386–1466).*
Al-Kashî. Geometry, arithmetic, astronomy.
1449. Jacob of Cremona. Translates Archimedes.
1450. **Nicholas Cusa. Geometry, theory of numbers.**
Jehuda Verga. Arithmetic.
George of Trebizond. Translates Ptolemy.
Printing from movable type.
1453. *Fall of Constantinople.*
1460. **Georg von Peurbach. Trigonometry.**
Benedetto da Firenze. Arithmetic.
1462. *Sack of Nassau affects printing.*
1469. *Lorenzo the Magnificent, Florence.*
1470. **Regiomontanus. Trigonometry.**
1475. **Al-Qalasâdî. Theory of numbers.**
Pietro Franceschi. Regular solids.
Georgius Valla. Geometry, arithmetic.
1478. First printed arithmetic, Treviso, Italy.
1481. Giorgio Chiarino. Commercial arithmetic.
1482. First printed edition of Euclid, Venice.
First printed German arithmetic, Bamberg.
1484. **Nicolas Chuquet. Algebra.**
Piero Borghi. Arithmetic.
1490. **Johann Widman. Algebra, arithmetic.**
1491. Calandri. Arithmetic.
1492. Pellos. Arithmetic.
Lanfreducci. Arithmetic.
Columbus discovers America.
1493. *Maximilian I, emperor of Germany.*
1494. **Pacioli. General mathematics.**
1500. **Leonardo da Vinci. Optics, geometry.**
Jacques le Fèvre d'Estaples. Geometry, arithmetic.
Georgius de Hungaria. Arithmetic.
Charles de Bouelles. Geometry, theory of numbers.
Johann Stöffler. Astronomical tables.
Elia Misrachi. Arithmetic.
Clichtoveus. On Boethius.
1503. Gregorius Reisch. Encyclopedia.
1505. Ciruelo. Arithmetic.
1506. Scipione del Ferro. Cubic equation.
Antonio Maria Fior. Cubic equation.

1510. **Albrecht Dürer. Geometry of curves.**
Raphael (1483–1520).
1512. Juan de Ortega. Geometry, arithmetic.
1513. Blasius (Sileceus). Arithmetic.
1514. Böschensteyn. Arithmetic.
1515. The Taglientes. Commercial arithmetic.
Gaspar Lax. Proportion, arithmetic.
Giel Vander Hoecke. Arithmetic.
1518. **Adam Riese. Arithmetic.**
1520. **Jakob Köbel. Arithmetic.**
Copernicus. Astronomy, trigonometry.
Feliciano da Lazesio. Arithmetic.
Estienne de la Roche. Arithmetic.
Ghaligai. Arithmetic.
1521. *Luther excommunicated.*
1522. **Tonstall. First arithmetic printed in England.**
1525. **Stifel. Algebra, arithmetic.**
Rudolff. Algebra, decimals.
Buteo. Algebra, geometry, arithmetic.
Erasmus. Thinker.
Oronce Fine. Geometry.
1527. Apianus. Pascal Triangle in print, astronomy, arithmetic.
1530. Zuanne de Tonini da Coi. Cubic equation.
Ringelbergius. Geometry, arithmetic.
Francesco dal Sole. Arithmetic.
Schoner. Arithmetic.
1534. Sfortunati. Arithmetic.
Claude de Boissière. Rithmomachia, arithmetic.
Jesuit order founded by Loyola.
1535. Jean Fernel. Proportion, astronomy.
Grammateus. Algebra, arithmetic.
Suryadaśa. Hindu algebra.
Ganeśa. Hindu algebra.
Giovanni Mariani. Arithmetic.
Glareanus. Geometry, arithmetic.
1536. *Calvin goes to Geneva.*
Tyndale is burned.
1540. **Gemma Frisius. Arithmetic.**
Melanchthon. General mathematics.
Camerarius. On Nicomachus.
1541. *De Soto discovers the Mississippi.*
1542. **Robert Recorde. Algebra, geometry, arithmetic.**
1543. Copernican system published.

1545. **Ferrari. Biquadratic equation.**
Tartaglia. Cubic equation, general mathematics.
Cardan. Cubic equation, general mathematics.
1550. **Rhæticus. Trigonometry.**
Maurolico. Geometry.
Johann Scheubel. Algebra.
Commandino. Greek mathematics.
Cosimo Bartoli. Geometry.
T'ang Shun-ki. On the circle.
K'u Ying-hsiang. Algebra, geometry.
Simon Jacob. Arithmetic
Ramus. Geometry, optics, arithmetic.
François de Foix-Candale. On Euclid.
Jacobus Micyllus. Arithmetic.
Nunes (Nonius, Núñez). Algebra, geometry, navigation.
Mohammed ibn Ma'rûf. Algebra, spherics, arithmetic.
Titian (1477–1576) and Michelangelo (1475–1564).
1558. *Elizabeth becomes queen of England (died 1603).*
1560. Peletier. Algebra, arithmetic.
1562. Juan Perez de Moya. Algebra, arithmetic.
1565. Trenchant. Arithmetic.
Richard de Benese. Surveying.
1566. Jerónimo Muñoz. Euclid, arithmetic.
1568. Humphrey Baker. Arithmetic.
1570. Billingsley and Dee. First English translation of Euclid.
Menher de Kempten. Arithmetic.
Neander. Metrology, spherics.
Xylander. On Diophantus.
Forcadel. Greek mathematics.
Benedetti. Theory of numbers.
Belli. Geometry.
Dasypodius. Euclid, lexicon.
1572. **Bombelli. Algebra.**
Digges, father (died 1571) and son (died 1595). Arithmetic, geometry.
1573. Otto. $\pi = \frac{3}{1}\frac{1}{1}\frac{5}{3}$ (old Chinese value).
1577. Herbestus. Polish arithmetic.
Gířjka Gŏrla z Gŏrlssteyna. Arithmetic.
1580. **François Viète. Algebra.**
Ludolf van Ceulen. On π.
Francesco Barozzi. On Proclus.
1583. **Clavius. Geometry, algebra, arithmetic, the calendar.**
Petrus Bongus. Mystery of numbers.
1587. Fyzi. Persian translation of the Lilāvati.

1590. **Cataldi. Continued fractions.**
Stevin. Decimal fractions.
Hsin Yun-lu. Calendar.
Van der Schuere. Arithmetic.
Thomas Masterson. Algebra, arithmetic.
1592. Mōri Kambei Shigeyoshi. Abacus.
1593. Adriaen van Roomen. Value of π.
Ch'êng Tai-wei. Arithmetic.
1594. Thomas Blundeville. Trigonometry, cosmography.
1595. **Pitiscus. Trigonometry.**
Magini. Geometry, astronomy, trigonometry.
1600. **Thomas Harriot. Algebra, analytic geometry.**
Jobst Bürgi. Logarithms.
Galileo. Geometry, astronomy, mechanics.
Behâ Eddîn. The sphere, arithmetic.
Ghetaldi. Geometry, algebra.
Bernardino Baldi. History of mathematics.
Shakespeare (1564–1616).
1603. **Matteo Ricci, Hsü Kuang-ching, and Li Chi Ts'ao translate Euclid into Chinese.**
James I proclaimed king of Great Britain.
1608. *Telescope invented.*
1610. **Kepler. Astronomy, geometry.**
1612. **Bachet de Méziriac. On Diophantus, recreations.**
1614. ***Napier. Logarithms.***
1615. **Henry Briggs. Logarithms.**
1618. Nicolò Longobardi and Giacomo Rho. European astronomy in China
1619. Savilian professorships (Oxford) founded.
1620. Gunter. Logarithms.
Paul Guldin. Geometry.
Faulhaber. Series.
Snell. Geometry, trigonometry.
Ursinus. Trigonometry, logarithms.
Daniel Schwenter. Recreations.
Francis Bacon, Novum Organum published.
1621. Raganātha. Hindu mathematics.
1630. **Mersenne. Greek mathematics, theory of numbers, geometry.**
Oughtred. Algebra, slide rule, logarithms.
Mydorge. Geometry, recreations.
Gellibrand. Logarithms.
Albert Girard. Algebra, trigonometry.
Denis Henrion. Logarithms.
Claude Richard. Greek mathematics.

1634. Hérigone. Algebra.

1635. ***Fermat. Analytic geometry, theory of numbers.***
Cavalieri. Indivisibles.
Yoshida Shichibei. General mathematics.
Richelieu founds the French Academy.

1637. ***Descartes. Analytic geometry.***

1639. Imamura Chishō. Geometry.

1640. **Desargues. Projective geometry.**
Florimond de Beaune. Cartesian geometry.
Torricelli. Geometry, physics.
Borelli. Greek mathematics.
Bernard Frénicle de Bessy. Geometry.
Antoine de la Loubère. Curves.
Roberval. Geometry.
Velasquez (1599–1660).

1649. *Charles I executed. England declared a commonwealth.*

1650. ***Pascal. Geometry, probability, theory of numbers.***
John Wallis. Algebra, series, history of mathematics.
Frans van Schooten. Edited Descartes and Viète.
Grégoire de Saint-Vincent. Geometry.
John Kersey. Algebra.
Wingate. Arithmetic.
Nicolaus Mercator. Trigonometry, logarithms.
John Pell. Algebra.
Athanasius Kircher. Instruments.
Smogolenski. Logarithms in China.
Sié Fong-tsu. Logarithms in China.
Milton and Hobbes.

1654. *Louis XIV crowned.*

1659. Ferdinand Verbiest. Astronomy in China.

1660. René François Walter de Sluze. The calculus, geometry.
Isomura (Iwamura) Kittoku. Problems.
Viviani. Geometry.
Dechales. On Euclid.
Brouncker. Series.

1663. Lucasian professorship (Cambridge) founded.
Muramatsu Kudayū Mosei. Geometry.

1665. Nozawa Teichō, Satō Seikō, and Sawaguchi Kazuyuki. Geometry and the native Japanese integration.
Neile. Geometry.

1670. **Barrow. Geometry.**
James Gregory. Series.
William Leybourn. Surveying.

1670. **Huygens. Geometry, physics, astronomy.**
Edward Cocker. Arithmetic.
Sir Christopher Wren. Geometry, astronomy, architecture.
John Locke (1632–1704) and Spinoza (1632–1677).
1671. Giovanni Domenico Cassini. Astronomy.
1675. Greenwich observatory founded.
Mei Wen-ting. Algebra, history of Chinese mathematics.
John Bunyan.
1680. **Seki Kōwa. The calculus.**
Sir Isaac Newton. Fluxional calculus, physics, astronomy entire field of mathematics.
Johann Hudde. Algebra.
Barrême. Arithmetic.
1681. *John Dryden.*
Pennsylvania granted to William Penn.
1682. ***Leibniz. The calculus.***
1690. **Marquis de l'Hospital. Applied calculus.**
Halley. Astronomy, life insurance, physics.
Jacques Bernoulli. Applied calculus, geometry, probability.
De Lahire. Geometry.
John Caswell. Trigonometry.
Tschirnhausen. Optics.
1696. Nakane Genkei. Japanese calculus.
1700. **Jean Bernoulli. Applied calculus.**
Michel Rolle. Equations.
Pierre Nicolas. Geometry.
Giovanni and Tommaso Ceva. Geometry.
Fatio de Duillier. Geometry.
Varignon. The calculus.
David Gregory. Optics, geometry.
Peter the Great (died 1725).
1702. *Anne becomes queen of England.*
1704. Charles Hayes. The calculus in English.
1710. **Roger Cotes. Geometry, analysis, the calculus.**
De Montmort. Probability, series.
Pierre Jartoux. Astronomy and analysis in China.
Humphrey Ditton. The calculus.
Saurin. Geometry.
De Lagny. Analysis.
Parent. Solid analytic geometry.
1715. Miyake Kenryū and Nakane Genjun. Problems.
Raphson. History of fluxions.

1720. **Brook Taylor. Series.**
De Moivre. Complex numbers, probability.
Nicolas (II) Bernoulli. Geometry.
Manfredi brothers. Geometry.
Christian von Wolf. General mathematics.
Pescheck. Textbooks.
Crousaz. Geometry.
Jacob Hermann. The calculus.
Fagnano. Curves, elliptic functions.
Guido Grandi. Geometry.

1722. Takebe. Geometry, π to 41 figures.

1730. Nicolas (I) Bernoulli. Differential equations, probability.
Saunderson. Algebra.
Van s'Gravesande. The calculus.
Nicole. Finite differences.
Maupertuis. Geodesy.
Matsunaga. Geometry, π to 50 figures.

1736. James Hodgson. The calculus in English.

1740. **Colin Maclaurin. Algebra, series, conics.**
Gabriel Cramer. Determinants, equations, curves.
George Berkeley. Attacks fluxional calculus.
Gua de Malves. Analytic geometry.
Frézier. Descriptive geometry.
Frederick the Great (Frederick II) becomes king of Prussia.

1745. Voltaire. On Newton.
Marquise du Châtelet. On Newton.

1750. ***Léonard Euler. Analysis, physics, astronomy.***
Montucla. History of Mathematics.
James Stirling. Geometry, series.
Robert Simson. Geometry.
Matthew Stewart. Geometry.
Jean (II) Bernoulli. Physics.
Riccati family. Differential equations.
Boscovich. Geometry, astronomy.
Daniel (I) Bernoulli. Physics.
Thomas Simpson. Algebra, geometry, the calculus.

1751. John Rowe. The calculus in English.

1760. **D'Alembert. Differential equations, astronomy, physics.**
John Landen. Elliptic integrals.
Alexis Claude Clairaut. Geometry, geodesy.
Seven Years' War (1756–1763). Lessing (died 1781), Burke (died 1797), Rousseau (died 1778), Voltaire (died 1778).

1765. Murai Chūzen. Equations.

1770. **Lambert. Hyperbolic trigonometry.**
Malfatti. Geometry.
Maria Gaetana Agnesi. Geometry.
Kästner. History of mathematics.

1775. Vandermonde. Algebra.
Bézout. Algebra.

1776. Pestalozzi. Arithmetic.
United States independence. Washington (died 1799), Jefferson (died 1826), Lafayette (died 1834).

1780. ***Lagrange. Theory of numbers, analysis, elliptic functions, astronomy.***
Condorcet. Analysis, probability.
Ajima Chokuyen. Indeterminate equations.
Aida Ammei. Series.
Fujita Sadasuke. Algebra.
Charles Hutton. Tables, dictionary, recreations.
John Wilson and Edward Waring. Theory of numbers.
Méchain. Metric system.

1789. *French Revolution.*

1790. Meusnier. Surfaces.

1795. École normale supérieure and École polytechnique founded about this time.

1800. ***Gauss. Theory of numbers, geometry, analysis, physics, astronomy, general field of mathematics.***
Laplace. Astronomy, physics, least squares.
Legendre. Elliptic functions, theory of numbers, geometry.
Carnot. Modern geometry.
Monge. Descriptive geometry.
Delambre. Astronomy, geodesy.
Lacroix. Analysis.
Mascheroni. Geometry of the compasses.
Pfaff. Astronomy, analysis.
Jean (III) Bernoulli. Probability.
Lhuilier. Geometry.
Ruffini. Algebra.
Bossut, Cossali, and Franchini. History of mathematics.
Trembley. The calculus.
James Ivory. Analytic methods.
Arbogast. Differential equations, variations, series.

1804. *Napoleon made emperor. Battle of Waterloo fought in 1815.*

1810. Hachette. Algebra, geometry.
Jean Robert Argand. Complex numbers.

1810. Fourier. Series, physics.
William Wallace. Hyperbolic functions.
Gergonne. Editor of Annales.
Woodhouse. Differential calculus.
Robert Adrain. Least squares.
1819. Horner. Numerical equations.
1820. Peter Barlow. Tables.
Poinsot. Geometry.
Sophie Germain. Elastic surfaces.
Bolzano. Series.
Poisson. Definite integrals, series, physics.
Crelle. Tables, editor of the Journal.
Brianchon. Geometry.
1825. Abel. Elliptic functions.
The Bolyais and Lobachevsky. Non-Euclidean geometry.
Nathaniel Bowditch. Celestial mechanics.
1830. Babbage. Calculating machine.
George Peacock. Differential calculus, algebra.
Möbius. Geometry.
Carl Gustav Jacob Jacobi. Elliptic functions.
Poncelet. Projective geometry.
Galois. Groups.
Cauchy. Functions, determinants, series.
Dupin. Geometry.
1840. Lamé. Elasticity, surfaces.
Jacob Steiner. Geometry.
Olivier. Descriptive geometry.
Arneth. History of mathematics.
J. F. W. Herschel. Astronomy, analysis.
MacCullagh. Surfaces.
1850. William Rowan Hamilton. Quaternions.
Chasles. Modern geometry.
Salmon. Geometry, algebra.
Biot. Physics, astronomy.
Grunert. Editor of the Archiv.
August. Mathematical physics.
De Morgan. History of mathematics, logic.
George Boole. Logic, differential equations.
Sylvester. Algebra.
Cayley. Invariants.
H. J. S. Smith. Theory of numbers.
Todhunter. History of mathematics, textbooks.
Kirkman. Analysis situs.

1850. Airy. Lunar theory.
Adams and Leverrier. Discover Neptune.
Liouville. Editor of the Journal.
Grassmann. Ausdehnungslehre.
Kummer. Series, surfaces.
Riemann. Surfaces, elliptic functions.
Eisenstein. Invariants.
Bellavitis. Geometry.
Gudermann. Hyperbolic functions.
Von Staudt. Geometry.
Plücker. Geometry.
Lejeune-Dirichlet. Theory of numbers.
Quételet. Statistics, geometry, history of mathematics.
Wronski. Philosophy of mathematics.
Benjamin Peirce. Algebra.
Steinschneider. History of mathematics.
Libri. History of mathematics.

INDEX

Since important names are often mentioned many times in the text, only such page references have been given as are likely to be of considerable value to the reader, the first reference after a proper name being to the biographical note in case one is given. Bibliographical references, in general, give only the page on which a book or essay is first mentioned, and on this page the full title appears, together with the abridged form subsequently used. In arranging the words and names alphabetically, ä, ö, and ü are taken as if written ae, oe, and ue respectively. The prefix Mc is indexed under Mac.

Aāhmesu. *See* Ahmes
Aa-user-Rā, 47
Abacus, 40, 141
'Abbâs. *See* Jauharî
Abbasides, 167
Abbo of Fleury (*c.* 980), 190, 195
Abbot Adrian, 185
Abbott, W. C., 195
Abdallâh. *See* Ibn al-Yâsimîn
Abdera, 59, 61
Abel, N. H. (*c.* 1825), 527
Abenbêder (*c.* 13th century), 211
Abraham bar Chiia (*c.* 1120), 206, 201
Abraham ben Ezra (*c.* 1140), 207, 170, 350
Abraham Judæus (*c.* 1120), 206
Abû 'Alî al-Chaiyât (*c.* 825), 209
Abû Bekr Mohammed. *See* al-Ḥaṣṣâr, al-Karkhî
Abû Ja'far al-Khâzin (*c.* 960), 175
Abû Kâmil Shojâ (*c.* 900), 177
Abû'l-Faradsh (*c.* 987), 177
Abû'l-Qâsim. *See* al-Majrîṭî
Abû'l-Ṣalt (*c.* 1100), 206
Abû'l-Ṭaiyib (*c.* 850), 172
Abû'l-Wefâ (*c.* 980), 175
Abû 'Oṭmân. *See* Sahl ibn Bishr
Abû Yaḥyâ (*c.* 775), 168
Abû Zakarîyâ. *See* Abû Bekr
Academy, Plato's, 88
Achilles problem, 78
Adam, C., 371, 375
Adams, J. C. (*c.* 1850), 469
Adelard of Bath (*c.* 1120), 203
Adémar, Monk (*c.* 1025), 186
Adrain, R. (*c.* 1808), 532, 502
Adriaan. *See* Adriaen
Adriaen Adriaenszoon (*c.* 1600), 340
Adriaen Anthoniszoon (*c.* 1620), 340
Adriaen Metius (*c.* 1620), 340
Adrianus. *See* Roomen and Adriaen
Æsop, 186
Aethelhard. *See* Adelard
Aethelstan (*c.* 925), 187
Africanus, Sextus Julius (*c.* 220), 132
Agatharchus (*c.* 470 B.C.), 79, 119
Agnesi, M. G. (*c.* 1770), 519
Aḥmed ibn 'Abdallâh al-Mervazî (*c.* 870), 174
Aḥmed ibn 'Abdallâh ibn 'Omar (*c.* 1050), 205
Aḥmed ibn al-Ṭaiyib (*c.* 890), 174
Aḥmed ibn Dâ'ûd (*c.* 890), 174
Aḥmed ibn Mohammed (al-Ṭaiyib) (*c.* 890), 174
Aḥmed ibn Mohammed ibn 'Imâd. *See* Ibn al-Hâ'im
Aḥmed ibn Mohammed ibn 'Oṭmân. *See* Albanna
Ahmed ibn Rajeb (*c.* 1425), 289
Ahmed ibn Yûsuf (*c.* 900), 172
Ahmes (*c.* 1650 B.C.), 47, 45
Aida Ammei (*c.* 1780), 537
Airy, G. B. (*c.* 1850), 469
Ajima Chokuyen (*c.* 1780), 538
Akhmim, 191
Akkad, 37
al-'Abbâs. *See* Jauharî
al-Arjânî (*c.* 850), 176
Alasia, C., 499
al-Aṣṭorlâbî (*c.* 830), 169
al-Ba'albekî. *See* Qosṭa
Albanna (*c.* 1300), 211
Albategnius (*c.* 920), 175, 201
al-Baṭrîq (*c.* 790), 168
al-Battânî. *See* Albategnius
Albert. *See* Albertus

Albertus Magnus (*c.* 1250), 228
Albertus de Saxonia (*c.* 1365), 242
Albêrûnî (*c.* 1000), 285, 145, 153, 157
Albumasar (*c.* 880), 174
al-Carsi (*c.* 1375), 241
al-Chaiyât (*c.* 825), 209
Alchindi (al-Kindî) (*c.* 860), 171, 174
Alcuin (*c.* 775), 185
Aldhelm, 185
al-Dimishqî. *See* Sa'îd ibn Ya'qûb
al-Dînavarî (*c.* 880), 174
Alexander the Great (*c.* 335 B.C.), 92, 93, 102, 144
Alexandre de Villedieu (*c.* 1240), 226
Alexandria, 59, 61, 102, 190
al-Faḍl. *See* al-Nairîzî
al-Faradî (*c.* 910), 192
al-Farġânî. *See* Alfraganus
al-Fârrâbî (*c.* 940), 175
al-Fazârî (*c.* 773), 168
Alfonsine Tables (*c.* 1250), 228, 210, 238, 259
Alfonso X, el Sabio, 228
Alfraganus (*c.* 833), 170, 209
Alfred the Great (*c.* 880), 187
Algebra, 40, 48, 62, 106, 114, 134, 158, 170, 177, 227, 270, 273, 278, 284, 300, 311, 320, 328
Algorism (algorithm), 170, 222, 225, 226, 238, 239, 260
al-Ḥaitam of Basra (*c.* 1000), 175
al-Ḥajjâj ibn Yûsuf (*c.* 820), 170
Alhakem II (*c.* 970), 192
al-Ḥarrânî. *See* Ibrâhîm ibn Hilâl
al-Ḥasan ibn al-Ḥasan. *See* al-Ḥaitam
al-Ḥasan ibn 'Obeidallâh (*c.* 925), 170
al-Ḥaṣṣâr, Mohammed ibn 'Abdallâh (*c.* 1175?), 210, 209
Alhazen. *See* al-Ḥaitam of Baṣra
al-Ḥimṣî (*c.* 880), 176
al-Ḥosein, ibn Sînâ. *See* Avicenna
'Alî ibn Abî Sa'îd. *See* Ibn Yûnis
'Alî ibn Aḥmed, al-Nasavî (*c.* 1025), 285
'Alî ibn Ibrâhîm. *See* Ibn al-Shâṭir
'Alî ibn Mohammed. *See* al-Qalasâdî
al-Jauharî. *See* Jauharî
al-Jorjânî (*c.* 1000), 174
al-Karkhî (*c.* 1020), 283
al-Kashî (*c.* 1430), 289
al-Khâzin (*c.* 950), 175
al-Khowârizmî (*c.* 825), 170, 168, 203, 329
al-Kindî. *See* Alchindi
al-Kushî. *See* al-Kashî
Allen, H. J., 25
Alligation, 32
Allman, G. J. (*c.* 1880), 543, xiv, 69
Almagest, 131, 160. *See* Ptolemy, C.
Almâhânî (al-Mâhânî) (*c.* 860), 171
al-Majrîṭî (*c.* 1000), 192
al-Mâmûn (*c.* 820), 169
al-Mansûr (*c.* 770), 167
al-Masîḥî (*c.* 1000), 174
al-Mervarrûdî (*c.* 830), 169
al-Mervazî (*c.* 860), 174
al-Misrî. *See* Aḥmed ibn Yûsuf
al-Nadîm (*c.* 900), 177
al-Nairîzî (*c.* 910), 170
al-Nasavî (*c.* 1025), 285
al-Nehâvendî (*c.* 820), 169
A-lo-pen (*c.* 640), 149
Alpetragius (*c.* 1200), 209, 210
al-Qalasâdî (*c.* 1475), 211
al-Qass (*c.* 900), 174
al-Qible (*c.* 900), 192
al-Qiwâm (12th century), 209
al-Rashîd (*c.* 800), 168, 149
al-Râzî. *See* Rhases, Râzî
al-Shadîb (14th century), 241
al-Tabarî (*c.* 800), 169
al-Ṭûsî (*c.* 1200), 288
Amenemhat (Amenemmes) III (*c.* 1850 B.C.), 45, 46
America. *See* United States
Ameristus (*c.* 550 B.C.), 77
Amisus, 59, 61
Ammei. *See* Aida
Anacreon, 70
Analytic method, 89, 91
Anatolius (*c.* 280), 133
Anaxagoras (*c.* 440 B.C.), 78
Anaximander (*c.* 575 B.C.), 68
Anaximenes (*c.* 530 B.C.), 77
Anchor ring, 118
Andalò di Negro (*c.* 1300), 232
Andrian, F. v, 14
Angle, trisection of, 82, 118, 171
Anharmonic ratio, 127, 137
Animal concepts, 5
Annotio, Perito. *See* Cataldi
Anthemius (*c.* 525), 191
Antilogarithms, 433
Antinoopolis (Antinoe), 59, 61, 126
Antiphon (*c.* 430 B.C.), 84
Antonio Biliotti (*c.* 1383), 233
Apamcia, 59, 61
Apepa I, 47
Apianus, P. (*c.* 1527), 333
Apollonius (*c.* 225 B.C.), 116, 182, 293, 302, 406; Arab translators of, 172
Appuleius (*c.* 150), 71

Apuleius (*c.* 150), 71
Aquitania, 59, 60
Arab achievements, 289
Arab writers, 210, 283
Arabia, 283
Arabic influence in China, 149
Arabic numerals. *See* Hindu-Arabic numerals
Arago, F. (*c.* 1840), 490, 494
Araki Hikoshirō Sonyei (*c.* 1690), 536
Arameans, 30, 99
Aratus (*c.* 270 B.C.), 92
Arbogast, L. (*c.* 1800), 500
Arcerianus, Codex, 183
Arcerius, J. (*c.* 1600), 183
Archibald, R. C., 107
Archilochus (*c.* 680 B.C.), 64
Archimedes (*c.* 225 B.C.), 111, 302
Archimedes, cubic of, 118, 122, 171
Archippus, 81
Architecture, 17
Archytas (*c.* 400 B.C.), 73, 84
Area, 40, 163. *See also* Circle, Triangle, Quadrilateral, Trapezium
Arenarius, 113
Arendt, 23, 24
Aretinus (*c.* 1025), 197
Argand, J. R. (*c.* 1810), 500
Arima Raidō (*c.* 1760), 536, 537
Aristæus (*c.* 320 B.C.), 94
Aristarchus (*c.* 260 B.C.), 107, 302
Aristotle (*c.* 340 B.C.), 93
Aristoxenes (*c.* 350 B.C.), 74
Arithmetic, 57, 58, 66, 89, 128, 129, 158, 335
Arithmetic in the Five Classics, 150
Arithmetic in Nine Sections. *See K'iu-ch'ang*
Arithmetic complement, 394
Arithmetic series, 155
Arjânî (*c.* 850), 176
Arjûza, 288
Arnaldo de Villa Nova (*c.* 1275), 229
Arnauld, A. (*c.* 1650), 386
Arnauld de Villeneuve (*c.* 1275), 229
Arneth, A. (*c.* 1840), 542
Arnold, Sir E., 98
Arnsperger, W., 502
Aronhold, S. H. (*c.* 1880), 511
Ars Magna, 297
Artabasda. *See* Rhabdas
Artillery science, 298
Āryabhaṭa the Elder (*c.* 510), 153, 152, 154, 155
Āryabhaṭa the Younger, 156
Āryabhaṭīya (*Āryabhaṭīyam*), 155
Āryāṣṭasata, 155
Ascham, R. (*c.* 1550), 318
Asclepias of Tralles (*c.* 635), 191
Assyria, 35, 39, 99, 146
Aṣṭorlâbî, al- (*c.* 830), 169
Astrolabe, 91, 169, 196, 197
Astrology, 39, 100, 146, 169, 218, 220, 221, 232
Astronomical instruments, oldest, 46
Astronomy, 17, 23, 37, 39, 67, 83, 95, 107, 118, 143, 158, *et passim*
Athelhard. *See* Adelard
Athelston (*c.* 925), 187
Athens, 59, 61, 87
Atomic theory, 80
Aubrey, J., 221
Augrim, 170
August, E. F. (*c.* 1850), 511
Augustine of Canterbury (*c.* 600), 185, 187
Augustine of Hippo (*c.* 400), 187
Aurispa (*c.* 1400), 243
Ausdehnungslehre, 507, 462
Austria, 255
Autolycus (*c.* 330 B.C.), 94, 302
Avanzini, G., 541
Avenpace (*c.* 1140), 208
Averroës (*c.* 1175), 208
Avicenna (*c.* 1020), 285
Awayama. *See* Ōyama
Aymar, Monk (*c.* 1025), 186
Azimabad, 153

Baba Nobutake (*c.* 1700), 536
Babbage, C. (*c.* 1840), 460
Babbage, H. P., 460
Babrius (*c.* 3d century), 186
Babylon, 35
Babylonia, 35, 37, 39, 100
Bach, J. N., 188
Bachet, C. G. (*c.* 1612), 379, 133
Bachuone, Arnald (*c.* 1275), 229
Bacon, Roger (*c.* 1250), 222, 81
Bagdad, 167, 172, 177, 283
Baker, H. (*c.* 1568), 321
Baker, M., 515
Bakhshālī manuscript, 164
Balbus (*c.* 100), 125
Baldi, B. (*c.* 1600), 539, 108
Ball, R. S. (*c.* 1900), 470
Ball, W. W. Rouse, xiv, 69, 381, 396, 402, 448
Baltzer, R. (*c.* 1860), 546
Bamberg arithmetic (1482), 260
Bamboo rods, 96, 141
Bapu Deva Sastri, 145

Baraniecki, M. A., 346
Barbieri, M., 70
Barbosa, 264
Bar Hebræus (*c.* 1270), 288
Barlaam (*c.* 1330), 232
Barlow, P. (*c.* 1820), 460
Barnard, F. P., 184
Barocius, F. (*c.* 1580), 302
Barozzi, F. (*c.* 1580), 302
Barrême, F. (*c.* 1680), 387
Barrow, I. (*c.* 1670), 396
Barsotti, G., 542
Barter, 22
Bartoli, C. (*c.* 1550), 303, 308
Bartolomeo da Parma (*c.* 1297), 220
Barton, G. A., 38
Basingstoke, John of (*c.* 1240), 222
Baṭrîq (*c.* 790), 168
Battaglini, G. (*c.* 1880), 518
Battânî. *See* Albategnius
Battierius, J. J., 428
Bauer, W., 69
Baumgartner, A., 70
Baur, L., 222
Bayer, J. W. (1704), 257
Bayle, P., 379
Bayley, Sir E. Clive, 153
Beale, T. W., 275
Beaune, F. de (*c.* 1640), 376
Beauvais. *See* Vincent
Bede the Venerable (*c.* 710), 184
Behâ Eddîn (*c.* 1600), 351
Behr. *See* Ursinus
Beldamandi (*c.* 1410), 246
Belgium, 526. *See also* Netherlands
Bell. *See* Schall von Bell
Bellavitis, G. (*c.* 1850), 517
Belli, S. (*c.* 1570), 304
Beltrami, E. (*c.* 1880), 518
Ben. *See* Ibn
Benares, 154
Benedetti, G. (*c.* 1570), 302
Benedetto da Firenze (*c.* 1460), 246
Benedict. *See* Benedetto, Benedictus
Benedict, St. (*c.* 529), 244
Benedictus. *See* Benedetto, Benedict
Benedictus Accolytus (*c.* 1077), 198
Benedictus Herbestus (*c.* 1560), 346
Benese, R. de (*c.* 1565), 322
Ben Ezra. *See* Abraham ben Ezra
Benî Mûsâ (*c.* 870), 171
Ben Mûsâ. *See* al-Khowârizmî
Benoy Kumar Sarkar, 153
Bensaude, J., 206, 349
Berkeley, G. (*c.* 1740), 469
Berlet, B., 337
Bernelinus (*c.* 1020), 197
Bernoulli, Christoph (*c.* 1840), 433
Bernoulli, Daniel (I) (*c.* 1750), 431
Bernoulli, Daniel (II) (*c.* 1790), 432
Bernoulli, Jacques (I) (*c.* 1090), 427, 417
Bernoulli, Jacques (II) (*c.* 1785), 433
Bernoulli, Jean (I), (*c.* 1700), 428, 417
Bernoulli, Jean (II) (*c.* 1750), 432
Bernoulli, Jean (III) (*c.* 1800), 432
Bernoulli, Jean Gustave (*c.* 1850), 433
Bernoulli, Nicolas (I) (*c.* 1730), 432
Bernoulli, Nicolas (II) (*c.* 1720), 432
Bernoulli family, 426
Bernward (*c.* 993), 190
Berosus (*c.* 250 B.C.), 140
Bertins, A. F. des (*c.* 1760), 474
Bertrand, J. L. F. (*c.* 1875), 310, 480
Bessel, F. W. (*c.* 1830), 511
Bessy. *See* Frénicle
Betti, E. (*c.* 1880), 518
Beugham, C. van (*c.* 1680), 426
Beverley. *See* John of Beverley
Bézout, É. (*c.* 1775), 481
Bhâskara (*c.* 1150), 275, 154, 161, 162
Biadego, G. B., 103
Biagio, Master (*c.* 1400), 233. *See also* Blasius
Bianchi, G., 515
Bianchini, Giovanni (*c.* 1450), 254
Bibliotheca Mathematica, xvi
Bierens de Haan, D. (*c.* 1870), 546, 330, 339, 423
Biernatzki, K. L., 32
Bigollo, Leonardo. *See* Fibonacci
Bigourdan, G., 38, 100
Bija Gaṇita, 278
Biliotti, Antonio (*c.* 1383), 233
Billingsley, H. (*c.* 1570), 324
Binary numerals, 25
Binomial expression, 40
Biot, E., 29
Biot, J. B. (*c.* 1840), 400, 30, 129, 140
Biquadratic equation, 294, 300
Birch, S., 52
Bithynia, 59, 61
Bjerknes, C. A., 527
Björnbo, A. A., 125, 106, 202
Blasius, J. M. (*c.* 1513), 345
Blasius Parmensis. *See* Biagio
Blundeville, T. (*c.* 1594), 324
Bobynin, V., 262, 435
Boccaccio, 230
Bôcher, M. (*c.* 1910), 533
Bodo, Henricus, 189
Böckh, A., 82

Böschensteyn, J. (*c.* 1514), 336
Boethius (Boetius) (*c.* 510), 178
Boissière (*c.* 1530), 309, 198
Boll, F., 119
Bolyai, F. (*c.* 1825), 528
Bolyai, J. (*c.* 1825), 529
Bolyai family, 527
Bolzano, B. (*c.* 1820), 531
Bombelli, R. (*c.* 1572), 300
Bonaccio, G., 215
Bonaini, F., 214, 216
Boncompagni, B. (*c.* 1870), 545, xvi, 201, 202, 209, 214, 251
Bongo. *See* Bongus
Bongus, P. (*c.* 1583), 304
Books, burned in China, 30, 32, 138; as parts of a volume, 105
Boole, G. (*c.* 1850), 463
Booth, J. (*c.* 1860), 469
Bopp, K., 386, 425
Borchardt, L., 43
Borel, P., 365
Borell (Borrel). *See* Buteo
Borelli, G. A. (*c.* 1640), 368
Borghi (Borgi), P. (*c.* 1484), 249
Boscovich, R. G. (*c.* 1750), 517
Bosmans, H., 303, 328, 340, 349, 424, 436
Bossut, C. (*c.* 1800), 540
Bouelles, Charles de (*c.* 1500), 307
Bougie, 215
Bouvelles. *See* Bouelles
Bowditch, N. (*c.* 1825), 532, 487
Boyer, J., 495
Bracciolini (*c.* 1414), 243
Bradwardine, Thomas (*c.* 1325), 236
Brahmagupta (*c.* 628), 157, 152, 154, 167
Brahmasiddhānta, 157, 167
Brancker. *See* Branker
Branker, T. (*c.* 1660), 412
Brassinne, E., 377
Braunmühl, A. von (*c.* 1900), 546, 202
Breasted, J. H., 38, 43, 49
Bredon, Simon (*c.* 1375), 237
Breier, F., 79
Brentari, O., 517
Bretschneider, C. A., xiv
Bretschneider, E., 140, 267
Brewer, J. S., 224
Brewster, D., 371, 398, 522
Brianchon, C. J. (*c.* 1820), 495
Bridferth (*c.* 1000), 196
Bridges, J. H., 224
Briggs, H. (*c.* 1615), 391
Brioschi, F. (*c.* 1880), 518
British writers. *See* Great Britain
Brouncker, W. (*c.* 1660), 410
Browne, G. F., 184, 186
Browning, R., 207
Brugmans, H., 425
Brunner, K., 10
Bryson (*c.* 450 B.C.), 84
Bryte, Walter (*c.* 1360), 237
Bubnov, N., 195
Buddha, 98, 154
Buddhism, 143, 151
Budge, E. A. W., 49
Bürgi, J. (*c.* 1600), 338, 433
Bürk, A., 98
Büttikoffer, J., 13
Bugia, 215
Bulengerus, J. C., 74
Bungus. *See* Bongus
Burgess, E., 145
Burkhardt, H., 516
Burlæus. *See* Burley
Burleigh. *See* Burley
Burley, Walter (*c.* 1325), 236
Bury, J. B., 243
Bussey, W. H., 302, 382
Buteo (Butéon), J. (*c.* 1525), 312
Buxerius. *See* Boissière
Byrhtferth (*c.* 1000), 196
Byzantium, 59, 61, 229. *See also* Constantinople

Cabala, 207
Cadiz, 59, 60
Cæsar (*c.* 46 B.C.), 122
Cairo, 191
Cajori, F., xiv, 78, 387, 392, 394, 531
Calandri (Calandrus, Calender) (*c.* 1491), 254, 255
Calculus, 430, 439, 448. *See also* Newton, Leibniz, Cavalieri, Barrow
Calendar, 37, 42, 100, 122, 138, 140, 151, *et passim*
Calendrini, L. (*c.* 1740), 520
Caliphs, 167
Calippus (*c.* 325 B.C.), 95
Callimachus (3d century B.C.), 71
Callippus (*c.* 325 B.C.), 95
Calo. *See* Kalonymos
Calsadilha (15th century), 263
Cambyses (*c.* 525 B.C.), 71
Camerarius, J. (*c.* 1540), 332
Campanus (*c.* 1260), 218
Canacci, R. (*c.* 1380), 232
Candale. *See* Foix
Candolle, A. de, 426
Canon Paschalis, 138. *See also* Calendar
Cantor, G. (*c.* 1900), 510

Cantor, M. (*c.* 1900), 544, xiv, 14, 98, 183
Capella (*c.* 460), 182
Capes, W. W., 185
Caphanton, J. (*c.* 1430), 263
Cardan, H. (or J.) (*c.* 1545), 295. *See also* Cardano
Cardano, F. (*c.* 1500), 224, 296. *See also* Cardan
Carette, A. M., 516
Caritat. *See* Condorcet
Carmen de Algorismo, 226
Carnot, L. N. M. (*c.* 1800), 494
Caron, F. (*c.* 1654), 442
Carpini, Fra Plano (*c.* 1246), 268
Carrara, B., 195
Carsi. *See* Carsono
Carsono, J. (*c.* 1375), 241
Casorati, F. (*c.* 1880), 518
Cassini, C. F., de Thury (*c.* 1756), 369
Cassini, G. D. (J. D.) (*c.* 1671), 369
Cassini, Jacques (*c.* 1725), 369
Cassini, J. D., de Thury (*c.* 1800), 370
Cassini family, 368
Cassiodorus (Cassiodorius) (*c.* 520), 180
Casting out nines. *See* Nines
Caswell, J. (*c.* 1690), 413
Cataldi, P. A. (*c.* 1590), 303
Catenary, 424
Cattaldi. *See* Cataldi
Cauchy, A. L. (*c.* 1830), 496
Caussinus, N., 235
Cavalieri, B. (*c.* 1635), 362, 418
Caxton, 314
Cayley, A. (*c.* 1870), 465
Cecco d' Ascoli (*c.* 1300), 231
Celtes, C., 189
Censorinus (*c.* 238), 132
Ceulen. *See* Ludolf
Ceva, G. (*c.* 1700), 511
Ceva, T. (*c.* 1700), 511
Chabas, M. F., 50
Chaignet, A. Ed., 69
Chaiyât (*c.* 825), 209
Chalcedon, 59, 61
Chalcis, 59, 61
Chaldea, 35, 39, 100, 146
Chang. *See* Ch'ang, *infra*
Chang Ch'i-mei (*c.* 1616), 270
Ch'ang ch'un (*c.* 1220), 268
Ch'ang ch'un-ch'ing (*c.* 100), 141
Ch'ang Hông (*c.* 125), 141
Ch'ang K'ién (2d century), 140
Ch'ang K'iu-kien (*c.* 575), 150, 267
Ch'ang K'iu-kien Suan-king, 150, 267
Ch'ang-ti (*c.* 1259), 268
Ch'ang Ts'ang (*c.* 200 B.C.), 32, 139, 164
Chang Tsu-nan, 534
Charlemagne (*c.* 780), 185
Charles, E., 224
Chasles, M. (*c.* 1850), 498, 202, 544
Chastelet. *See* Châtelet
Châtelet, M^me^ du (*c.* 1745), 477
Chaucer, 160, 170
Chau Ju-kua, 267
Ch'ên Chin-mo (*c.* 1650), 436
Ch'ên Shih-jen (*c.* 1715), 535
Ch'êng Kiang Chen (*c.* 225 B.C.), 139
Ch'êng Tai-wei (*c.* 1593), 352
Chiarino, Georgio (*c.* 1481), 249
Chia-tsü (sexagesimals), 24
Chiia. *See* Abraham bar Chiia
Chijja. *See* Abraham bar Chiia
Ch'i-ku Suan-king, 151
Child, J. M., 398 417, 418
Ch'in Dynasty, 138
China, 22, 96, 138, 148, 152, 266, 351, 435, 534
Chinghiz Khan, 268, 287
Ch'in Kiu-shao (*c.* 1250), 269
Chios, 59, 61
Chiu-chï-li, 149
Chiung Lin-tai, 534
Chokuyen. *See* Ajima
Ch'ön Huo (*c.* 1050), 266
Ch'ön I, 149
Ch'ön Luan (*c.* 535), 150
Chords, table of, 119, 127, 131, 145
Chou. *See* Shu
Chóu K'ü-feï (*c.* 1178), 267
Chóu-Kung (*c.* 1105 B.C.), 31, 32
Chóu-peï Suan-king, 29, 141, 152
Christian scholars, early, 177–187
Chrysoloras, Manuel (*c.* 1397), 235
Chunrad von Jungingen (*c.* 1400), 241
Chunrad von Megenberg (*c.* 1350), 241
Chuquet, N. (*c.* 1484), 261, 313
Chu Shï-kié (*c.* 1299), 273
Chu Sung ting. *See* Chu Shï-kié
Chü-t'an Chüan, 149
Chûzen. *See* Murai
Cicero, 84, 88, 115
Circle, 40, 79, 84, 92, 98, 139, 197, 236, 242; segment of, 139, 164
Circumference of the earth, 110
Ciruelo, P. S. (*c.* 1505), 344
Cissoid, 118, 424
Cities, ancient mathematical, 59–61, 154
Clairaut, A. C. (*c.* 1760), 475
Clairaut, J. B. (*c.* 1740), 475
Clairaut family, 475
Claudius. *See* Ptolemy, C.

Claudius, emperor, 124
Clavius, C. (*c.* 1583), 334
Clazomenæ, 59, 61
Clebsch, R. F. A. (*c.* 1870), 511, 505
Cleomedes (*c.* 40 B.C.), 122, 118
Clepsydra, 69
Clichtoveus, J. (*c.* 1500), 342
Clifford, W. K. (*c.* 1870), 467
Cnidus, 59, 61
Cocker, E. (*c.* 1670), 415
Codex Arcerianus, 183
Cohen, B., 346
Coi. *See* Zuanne
Coins, 56, 96, 140
Colebrooke, H. T., 144, 280
Colla. *See* Zuanne da Coi
Colson, J., 404
Columella (*c.* 25), 123
Comirato, 513
Commandino, F. (*c.* 1550), 302, 170
Commercial mathematics, 49, 56
Compass, 97, 143
Compasses, geometry of the, 302, 516; single opening of the, 302
Computus, 138. *See also* Calendar
Conant, L. L., 14
Conchoid, 118, 171, 387
Condorcet, A. N. C., Marquis de (*c.* 1780), 481, 521
Confucius, 24, 70
Conic sections, 92, 116, 331
Conon (*c.* 260 B.C.), 107
Conrad von Jungingen (*c.* 1400), 241. *See also* Chunrad
Constantinople, 59, 61, 190, 229, 230, 233, 242
Constellations, 18, 91
Continued fractions, 156, 303
Continuity, 93, 416
Contractus. *See* Hermannus
Convergence, 497
Coordinates, 116, 129, 375, 378, 389
Copernicus (*c.* 1520), 346, 108
Coppernicus. *See* Copernicus
Cordier, H., 25
Cortés, 353
Cosa. *See* Coss
Cosecant, 175
Cosmic figures, 2
Coss, 328, 320
Cossali, P. (*c.* 1800), 541, 215
Cossic. *See* Coss
Cotes, R. (*c.* 1710), 447, 402
Counting, 6; scales of, 9, 141
Cousin, V., 137, 375
Couturat, L. (*c.* 1910), 500
Crafte of Nombryng (*c.* 1300), 238
Cramer, F., 69
Cramer, G. (*c.* 1740), 520
Crawford, John, 14
Crawford, M., 180
Crelle, A. L. (*c.* 1820), 506
Cremona, Jacob of. *See* Jacob
Cremona, L. (*c.* 1880), 517
Crete, 52, 59, 61
Cribrum (sieve), 109
Croker, J., 400
Cromberger, J., 354
Crotona (Croton), 59, 61, 72
Crousaz, J. P. (*c.* 1720), 519
Crouzas. *See* Crousaz
Cube, duplication of, 83, 85, 118, 136
Cubes, tables of, 40
Cubic equation, 114, 134, 151, 294, 295, 297, 301
Cuneiform numerals, 36
Cuneiform writing, 36, 39, 40
Cunningham, W., 320
Curriculum, medieval, 214
Curtze, E. L. W. M. (*c.* 1880), 546, 206
Curves, higher plane, 136, 327; of descent, 424. *See also* Cissoid and other names
Cusa, Nicholas (*c.* 1450), 258
Cycloid, 259, 366, 385, 424
Cyprus, 59, 61, 86
Cyzicus, 59, 61

Da Coi. *See* Zuanne
Dagomari, Paolo (*c.* 1340), 232
D'Alembert, J. L. (*c.* 1760), 479
Damascius (*c.* 510), 182
Danck, Johann (14th century), 238
Dansie, J., 391
Dante, 230
Darboux, J. G. (*c.* 1900), 500, 527
Dark Ages, 177
Darvai, M., 528
Darwin, G. H. (*c.* 1900), 470
Dasypodius, C. (*c.* 1570), 348
De Beaune. *See* Beaune
Dechales, C. F. M. (*c.* 1660), 386
Decimal point, 255, 331. *See* Fractions
Dedekind, J. W. R. (*c.* 1880), 509
Dee, John (*c.* 1575), 323, 170
Definitions, 75
De Haan. *See* Bierens de Haan
Deinostratus (*c.* 350 B.C.), 92, 82
Delambre, J. B. J. (*c.* 1800), 492, 92
De latitudinibus formarum, 236, 239, 242

Delatore, A. (*c.* 1489), 264
Delatte, A., 69
Democritus (*c.* 400 B.C.), 80
De Moivre, A. (*c.* 1720), 450
De Montgomery, J. E. G., 185
De Morgan, A. (*c.* 1850), 462, 103, 310, 324, 398, 522
Deniker, J., 14
Dennett, R. E., 14
Desargues, G. (*c.* 1640), 383
Desboves, A., 381
Descartes, R. (*c.* 1637), 371
Deschales. *See* Dechales
Descriptive geometry, 472, 491
Determinant, 440, 497
Dettonville. *See* Pascal
Diaz. *See* Diez
Dicæarchus (*c.* 320 B.C.), 94
Dickson, L. E., 381
Dickstein, S., 531
Diderot, 522
Diez, J. (*c.* 1550), 353
Differential triangle, 418
Digges, L. and T. (*c.* 1572), 321
Digits, 10
Dînavarî, al- (*c.* 880), 174
Dini, U. (*c.* 1900), 518
Dinostratus (*c.* 350 B.C.), 92, 82
Diocles (*c.* 180 B.C.), 118
Diodorus Siculus (*c.* 8 B.C.), 121, 45
Diogenes Laertius (2d century), 68
Diogo Mendes Vizinho (*c.* 1500), 263
Diogo Ortiz (15th century), 263
Dionisi, G., 517
Dionysius Exiguus (*c.* 525), 180
Dionysodorus (*c.* 50 B.C.), 122
Diophantus (*c.* 275), 133, 333, 343, 380
Dirichlet, P. G. Lejeune- (*c.* 1850), 507
Ditton, H. (*c.* 1710), 456
Division, 255
Domenico Maria Novara da Ferrara (*c.* 1490), 254
Domitius. *See* Ulpianus
Domninus of Larissa (*c.* 450), 136
Dozen, 12
Drevet, 424
Duality, 93
Du Bois-Reymond, P. (*c.* 1880), 511
Du Boulay, 242
Dudley, J., 452
Düker, H., 190
Dürer, A. (*c.* 1510), 326
Düx, J. M., 258
Duhem, P. (*c.* 1900), 500, 75, 294
Duillier. *See* Fatio
Dupin, C. (*c.* 1830), 500, 490
Duplication of the cube, 83, 85, 118, 136
Dupuis, J., 129
Dupuy, P., 498
Dusan, Stephen, 242
Dutt, R. C., 35

e, transcendence of, 500
Earth, sphericity of, 75, 81, 157; measure of, 109
East and West, intercourse between, 140, 143, 148, 267, 425, 442
Easter, 138. *See also* Calendar
Eclipse, 100
École normale, 485
École polytechnique, 485
Edelinck, 424
Edkins, J., 25
Eells, W. C., 14
Egypt, 41, 100; redivision of land by Rameses II, 51
Eisenlohr, A., 48, 50
Eisenstein, F. G. M. (*c.* 1850), 500
Elchatayn (elchataieym). *See* False Position
Elea, 59, 61, 78
Elements of Euclid, 105, 106
Elia Misrachi (*c.* 1500), 350
Elis, 59, 61
Ellipse, 92, 116, 163, 171
Ellipsoid, 114
Elliptic functions, 488, 527
Elphinstone, M., 34
Emanuel. *See* Imanuel
Embadorum, Liber, 206
Emmanuel. *See* Imanuel
Encyklopädie, xiv
Eneström, G., xvi, 117, 126, 412, 470
Engel, F., 480, 527
Engineers, early, 43, 124
England. *See* Great Britain
Epaphroditus (*c.* 200), 132
Equations, biquadratic, 40, 294, 300; cubic, 114, 134, 151, 294, 297, 298; fifth-degree, 527; functions of coefficients, 423; indeterminate, 119, 134, 150, 155, 156, 158, 159, 163; limits of roots, 377; linear, 48; *n* roots, 423; numerical, 270, 439; quadratic, 40, 134, 155, 159, 162, 280, 281; signs, 376; simultaneous, 91, 159. *See also* Pell Equation, Riccati Equation
Equinox, 37
Erasmus (*c.* 1525), 325
Eratosthenes (*c.* 230 B.C.), 108

Erlendssön, Hauk (*c.* 1300), 240
Estaples. *See* Faber
Étaples. *See* Faber
Eucleides. *See* Euclid
Euclid of Alexandria (*c.* 300 B.C.), 103, 169, 182, 187, 201, 203, 250, 251, 293, 302, 304, 323, *et passim*; Arab translators of, 168, 169, 171, 176, 201; bibliography, 309
Euclid of Megara, 88, 105
Euctemon (*c.* 432 B.C.), 83
Eudemus (*c.* 335 B.C.), 94, 68
Eudoxus (*c.* 370 B.C.), 91, 73
Euler, J. A. (*c.* 1775), 524
Euler, L. (*c.* 1750), 520
Euler's Equation, 522
Eutocius (*c.* 560), 182
Evolutes, 424
Exhaustion, method of, 84, 91
Exponents, fractional, 239, 408; negative, 408

Faber Stapulensis (*c.* 1500), 307, 199
Fabricius, J. A., 85
Facio (Faccio). *See* Fatio
Faḍl. *See* al-Nairîzî
Fagnano, G. F. (*c.* 1765), 515
Fagnano di Fagnani, G. C. (*c.* 1720), 514
Fa-hién (*c.* 400), 143, 154
Faizi. *See* Fyzi
Fakhrî, 284
False Position, 33, 216
Fan Ching-fu, 534
Faradî (*c.* 910), 192
Farġânî. *See* Alfraganus
Fârrâbî (*c.* 940), 175
Fatio de Duillier, N. (*c.* 1700), 519
Faulhaber, J. (*c.* 1620), 418
Favaro, A., 48, 299, 362, 364, 366
Favorinus (*c.* 125), 75
Fazârî (*c.* 773), 168
Federici, D. M., 240
Federigo of Montefeltro, 243
Feilberg, H. F., 14
Feizi. *See* Fyzi
Feliciano da Lazesio (*c.* 1520), 306
Fermat, P. de (*c.* 1635), 377, 133
Fermat, S. (*c.* 1679), 378
Fernel (Fernelius), J. (*c.* 1535), 308
Ferrara. *See* Domenico
Ferrari, L. (*c.* 1545), 300
Ferro (Ferri, Ferreo, Ferreus) (*c.* 1500), 295
Fibonacci, Leonardo (*c.* 1202), 214, 206
Fibonacci series, 217
Ficquet, 419
Figulus, P. Nigidius (*c.* 60 B.C.), 122
Figura cata, 172
Figurate numbers, 129
Fihrist, Kitâb al- (Book of Lists), 177
Finæus. *See* Fine
Fine, Oronce (*c.* 1525), 308
Fineus. *See* Fine
Finger reckoning, 185, 235
Fingers in counting, 10
Fincke, T. (Fink, Finck, Finchius) (*c.* 1583), 348
Fior, Antonio Maria (*c.* 1506), 295
Firmiani, 299
Fiske, T. S., 531
Fitzgerald, E., 286
Fitzstephen (*c.* 1180), 204
Five, scale of, 9
Flamsteed, J. (*c.* 1690), 405
Fleet, J. F., 155
Fleming, William (*c.* 1250), 229
Fleury, Claude, 228
Floridus. *See* Fior
Flos (Fibonacci), 217
Fluxions, Newton's, 404. *See also* Calculus
Foix-Candale (*c.* 1550), 309
Forcadel, P. (*c.* 1570), 313
Fortolfus (*c.* 1090), 198
Fortunatae Insulae, 130
Fourier, J. (*c.* 1810), 500
Fractions, Babylon, 38, 41; China, 150; continued, 156, 303, 411; decimal, 255, 290, 331; division, 162; Egypt, 45; as exponents, 239, 408; Greece, 119; India, 156, 162; sexagesimal, 24, 41, 119, 131, 332; unit, 38, 45, 48
France, 201, 226, 238, 260, 306, 370, 470
Franceschi, Pietro (*c.* 1475), 247
Francesco di Simone Stabili (*c.* 1300), 231
Francesco dal Sole. *See* Sole
Franchini, P. (*c.* 1810), 542
Franco of Liège (*c.* 1066), 197
Frankland, W. B., 103
Franklin, F., 463
Fraser, J., 14
Fraser, R. W., 154
Frederick the Great, 473, 478, 483, 501
Frénicle de Bessy, B. (*c.* 1640), 385
Frézier, A. F. (*c.* 1740), 472
Friedlein, G., 137, 178, 195
Frisch, 416
Frisi, P., 362, 364

Frisius. *See* Gemma
Frontinus (*c.* 100), 123
Fuchs, E. L. (*c.* 1870), 510
Fuh-hi (*c.* 2800 B.C.), 23, 28
Fujita Sadasuke (*c.* 1780), 537
Fujiwara Michinori (*c.* 1150), 273
Fuss, N., 521
Fuss, P. H. von, 427
Fyzi (*c.* 1587), 276, 279

Gades (Cadiz), 60
Gaging, 330
Gairdner, J., 314
Galenus. *See* Pediasimus
Galileo (*c.* 1600), 363
Galiot de Genouillac, 298
Galois, E. (*c.* 1830), 498
Gambier, G., 310
Gambioli, 514
Gaṇaca, 158
Ganeśa (*c.* 1535), 351
Gaṇita of Aryabhata, 155
Gaṇita of Mahāvira, 161
Garrett, J., 153
Gaskoin, C. J. B., 186
Gauging. *See* Gaging
Gauss, C. F. (*c.* 1799), 502
Geber, 168, 206. *See also* Jeber
Gebler, K. von, 363
Gelcich, E., 368
Gellibrand, H. (*c.* 1630), 392
Geminus (*c.* 77 B.C.), 121
Gemma Frisius (*c.* 1540), 341
Gemma Frisius, C. (*c.* 1570), 342
Gemma Regnier (Rainer, *etc.*). *See* Gemma Frisius
Gennadi (15th century), 262
Genouillac, 298
Genshō (*c.* 1225), 274
Gentleman's Diary, 447
Geography, 77, 78, 108, 129, 130
Geometria Culmensis (*c.* 1400), 241
Geometric forms, 62
Geometric ornament, 15
Geometric progression, 40
Geometry, 40, 59, 62, 67, 75, 90; analytic, 375, 472; descriptive, 472, 491; modern, 383, 496, 504, 525; non-Euclidean, 528, 529. *See also* Coordinates
George. *See* Georgius, Georgios
George of Hungary (*c.* 1500), 263
George of Trebizond (*c.* 1450), 263
Georgios Pachymeres (*c.* 1300), 229
Georgius de Hungaria (*c.* 1500), 263
Gerasa, 59, 61, 127
Gerbert (*c.* 1000), 195
Gergonne, J. D. (*c.* 1810), 495
Gerhardt, C. I., 234
Gerland of Besançon (*c.* 1137), 205
Germain, Sophie (*c.* 1820), 499
Germany, 226, 241, 255, 324, 416, 501
Gerson (Gerschom, Gersonides). *See* Levi ben Gerson
Ghaligai, F. (*c.* 1520), 306
Gherardi, S., 231
Gherardo Cremonense (of Cremona) (*c.* 1150), 202, 170, 171, 172
Gherardo da Sabbionetta (*c.* 1150), 202
Ghetaldi, M. (*c.* 1600), 368
Giambattista Benedetti (*c.* 1570), 302
Gibbon, 242
Gibbs, J. W. (*c.* 1885), 533
Giles, H. A., 23
Giles, J. A., 184
Gill, J., 317
Ginsburg, J., 166, 207
Giovanni Aurispa (*c.* 1400), 243
Giovanni Battista Lanfreducci (*c.* 1490), 245
Giovanni Bianchini (*c.* 1450), 254
Giovanni da Firenze (*c.* 1422), 245
Girard, A. (*c.* 1630), 423, 343
Gītikā, 156
Gitti, V., 253
Glaber, Raoul (*c.* 1040), 195
Glareanus, H. L. (*c.* 1535), 348
Gmünden, Johann von (*c.* 1430), 258
Gnomon, 31, 69, 98
Goethals, F. V., 343
Gola, 155
Golden Section, 4, 218
Golénischeff, 49
Gomperz, T., 71
Gondisalvi, D., 209
Gôrla z Gôrlssteyna, G. (*c.* 1577), 346
Gottsched, J. C., 502
Goutière, 486
Gow, J., xiv, 77
Graf, J. H. (*c.* 1900), 546, 525
Grammateus (*c.* 1535), 329
Grandi, L. G. (*c.* 1720), 514
Grassmann, H. G. (*c.* 1850), 507
Grassmann, R. (*c.* 1860), 507
Graves, F. P., 310
Gravesande. *See* Van s'Gravesande
Gray, G. J., 402
Great Britain, 184, 187, 203, 220, 235, 261, 314, 387, 446
Greathead, Robert (*c.* 1240), 222
Great Pyramid, 43, 44

Greece, 54
Greek algebra, 62
Greek Anthology, 133
Greek cities, 59–61
Greek mathematics, 63
Greeks, 55; in India, 144
Grégoire. *See* Saint-Vincent
Gregory, D. (*c.* 1700), 413
Gregory, J. (*c.* 1670), 409
Gregory, J. C., 14
Gregory XIII, 42
Griffith, F. L., 48
Groma, 124
Gromatici, 124
Grosse, H., 335
Grosseteste. *See* Greathead
Grotefend, 36
Group theory, 499
Gruma, 124
Grunert, J. A. (*c.* 1850), 511
Gua de Malves, J. P. de (*c.* 1740), 475
Gudermann, C. (*c.* 1850), 510
Günther, S., xiv, 111, 190, 234, 318, 458
Guglielmini, G. B., 214
Guglielmo de Lunis (*c.* 1250?), 220
Guido of Arezzo (*c.* 1028), 197
Guido Grandi (*c.* 1720), 514
Guijeno. *See* Blasius, J. M.
Guilielmus. *See* William, Guglielmo
Guilielmus Brabantinus (Flemingus) (*c.* 1250), 229
Guldin, P. (*c.* 1620), 433, 137
Gunter, E. (*c.* 1620), 394

Haan. *See* Bierens de Haan
Ḥabash al-Ḥâsib (*c.* 870), 174
Hachette, J. N. P. (*c.* 1810), 499
Hager, J., 74
Haid, J., 501
Ḥaitam, al-, of Basra (*c.* 1000), 175
Hai-tau Suan-king, 142, 152
Haithon, 269
Hajjâj, al-, ibn Yûsuf (*c.* 820), 176
Haldane, E. S., 371
Hallam, 184
Halley, E. (*c.* 1690), 405, 116
Halliwell, J. O., 187, 204, 226, 227
Halphen, G. H. (*c.* 1880), 500
Halsted, G. B., 529
Ḥâmid ibn al-Khiḍr (*c.* 1000), 285
Hamilton, W. R. (*c.* 1850), 461
Hammurabi (Hammurapi) (*c.* 2100 B.C.), 38
Hankel, H. (*c.* 1870), 543, xiv, 289, 522
Hanseatic League, 255
Hapgood, Isabel F., 530
Hare and hound problem, 32, 187
Hariot. *See* Harriot
Harmonic proportion, 75
Harpedonaptae, 81
Ḥarrânî. *See* Ibrâhîm ibn Hilâl
Harriot, T. (*c.* 1600), 388
Harris Papyrus, 52
Hartsingius, P. (*c.* 1654), 442
Harun al-Rashid (*c.* 800), 168, 149
Hasan, Jewish judge (*c.* 950?), 208
Ḥasan, al-, ibn 'Obeidallâh (*c.* 925), 176
Hase, H., 191
Haskins, C. H., 201, 203
Ḥaṣṣâr, Mohammed ibn 'Abdallâh (*c.* 1175?), 210, 209
Hatasu. *See* Hatshepsut
Hatono Sōha (*c.* 1675), 442
Hatshepsut, queen (*c.* 1500 B.C.), 49
Hauk Erlendssön (*c.* 1300), 240
Havet, J., 105
Hawkins, J. (*c.* 1678), 415
Hayes, C. (*c.* 1704), 448, 404
Heath, Sir T. L., xiv, 64, 92, 103, 111, 112, 114, 116, 133
Hecatæus (*c.* 517), 77, 78
Hehn, J., 14
Heiberg, J. L., 103, 111, 114, 126, 229, 230
Heilbronner, J. C. (*c.* 1740), 539, 221
Heinrich, G., 410
Heinrich von Langenstein (von Hessen) (*c.* 1365), 241
Helceph, 204
Hellmann, W., 231
Helmholtz, H. L. F. von (*c.* 1870), 511
Henricus. *See* Heinrich
Henrion, D. (*c.* 1630), 370
Henry, C., 204, 377
Herbestus, B. (*c.* 1577), 346
Hérigone, P. (*c.* 1634), 371
Hermann, J. (*c.* 1720), 520
Hermann the Lame. *See* Hermannus Contractus
Hermannus Augiensis. *See* Hermannus Contractus
Hermannus Contractus (*c.* 1050), 197, 199
Hermite, C. (*c.* 1870), 500
Herodotus (*c.* 450 B.C.), 81; on division of land, 51
Heron (Hero) (*c.* 50, or possibly *c.* 200), 125, 284, 302
Herschel, J. F. W. (*c.* 1840), 469
Hesse, L. O. (*c.* 1860), 510

Hexagram, Pascal's, 453
Hexagrams (*Pa-kua*), 25
Hilâl ibn Abî Hilâl. *See* Ḥimṣî
Hill, G. W. (*c.* 1900), 532
Hillebrandt, A., 34
Hiller, E., 129
Hilprecht, H. V., xv, 37, 40
Ḥimsî (*c.* 880), 176
Hindu-Arabic numerals, 144, 166, 234. *See also* Zero
Hipparchus (*c.* 140 B.C.), 119, 92
Hippias of Elis (*c.* 425 B.C.), 82, 92
Hippocrates of Chios (*c.* 460 B.C.), 82
Hire. *See* Lahire
Hirth, F., 23, 140
Hirth and Rockhill, 267
Hispalensis. *See* Johannes
Historians, 539
Hittell, J. S., 14
Ho and Hi (*c.* 2350 B.C.), 24
Hoche, R., 127, 137, 191
Ho' Ch'êng-t'ien (*c.* 450), 143
Hochheim, A., 283
Hock, C. F., 195
Hodgson, J. (*c.* 1636), 448
Hoecke. *See* Vander Hoecke
Hoernle, R., 164
Holland, 526. *See* Netherlands
Holliday, F. (*c.* 1740), 449
Holzmann, W. *See* Xylander
Hommel, F., 100
Honda. *See* Fujita
Ḥonein ibn Isḥâq (*c.* 840), 176
Hook, F. W., 185
Hôpital. *See* L'Hospital
Hopkins, E. W., 72
Horace, 85
Horner, W. G. (*c.* 1819), 459
Horner's Method, 270, 273, 459
Ḥosein. *See* Avicenna
Hospital. *See* L'Hospital
Ho-t'u, 28
Hoüel, 529
Hrabanus Maurus (*c.* 820), 188
Hrotsvitha (*c.* 970), 189
Hsia-hou Yang (*c.* 550), 150
Hsia-hou Yang Suan-king, 150
Hsin Yun-lu (*c.* 1590), 352
Hsü Kuang-ching (*c.* 1605), 304
Hsü Yüeh (*c.* 250), 143
Huang-ti, emperor (*c.* 2700 B.C.), 24, 32, 97
Huang-ti K'iu-ch'ang, 267
Hüan-tsang (*c.* 640), 149, 155
Huart, C., 170
Huber, D., 480
Huber, E., 38
Hudde, J. (*c.* 1680), 425
Hugenius. *See* Huygens
Hui-sing (*c.* 518), 148
Hulsius, L., 433
Hultsch, F., 125, 136
Hume (Humius), J. (*c.* 1636), 311
Hunt, A. S., 49
Hutton, C. (*c.* 1780), 458, 431
Huygens, C. (*c.* 1070), 423
Hyde, T., 289
Hyginus (*c.* 120), 124
Hypatia (*c.* 410), 137
Hyperbola, 92, 116
Hyperbolic functions, 458
Hyperboloid, 413
Hypsicles (*c.* 180 B.C.), 119

Iamblichus (*c.* 325), 135, 69
Ibn Abî Ya'qûb al-Nadîm (*c.* 987), 177
Ibn al-Hâ'im (*c.* 1400), 288
Ibn al-Kâtib (*c.* 1200), 210
Ibn al-Lubûdî (*c.* 1200), 288
Ibn al-Mejdî (*c.* 1400), 289
Ibn al-Ṣaffâr (*c.* 1025), 205
Ibn al-Ṣalâḥ (*c.* 1140), 286
Ibn al-Shâtir (*c.* 1350), 289
Ibn al-Yâsimîn (*c.* 1200), 288
Ibn al-Zarqâla (*c.* 1050), 205
Ibn Bedr (*c.* 13th century), 211
Ibn Man'a (*c.* 1200), 287
Ibn Mûsâ. *See* al-Khowârizmî
Ibn Râhiweih. *See* al-Arjânî
Ibn Yûnis (*c.* 1000), 175, 191
Ibn Yûnis the Younger (*c.* 1200), 287
Ibrâhîm ibn Hilâl (*c.* 975), 175
Ibrâhîm ibn Sinân (*c.* 940), 175
Ibrâhîm ibn Yaḥyâ (*c.* 1050), 205
Ibrim, 52
I-hsing (*c.* 727), 151
I-king, 23, 25, 142, 267
Imaginary numbers, 301, 423, 430, 450
Imamura Chishō (*c.* 1639), 437, 353
Imanuel ben Jacob Bonfils, 241
Indeterminate equations, 119, 150, 155, 156, 158, 159, 163
India, 33, 97, 144, 152, 274, 435
Indivisibles, 93, 362, 408
Induction, 302
Infinite products, 312
Infinite series. *See* Series
Infinitesimal, 416
Integral, 430
Intercourse between East and West, 140, 143, 148, 267, 435, 442

Invariants, 483
Isaac. *See* Ishâq
Isaac ben Joseph Israeli (*c.* 1330), 240
Isaac ben Salom (*c.* 930), 192
Isaac ben Sid (*c.* 1250), 210
'Îsâ ibn Yaḥyâ (*c.* 1000), 174
Isely, L., 426
I-se-ma-yin (*c.* 1267), 272
Isḥâq ibn Ḥonein (*c.* 900), 172, 176
Isidorus (*c.* 610), 183
Isochronous curve, 428
Isocrates, 71
Isomura Kittoku (*c.* 1660), 437
Isoperimetry, 136
Italy, 214, 220, 231, 245, 294, 361, 511
Ivory, J. (*c.* 1800), 469
Iwasaki Toshihisa (*c.* 1775), 537

Jabir ibn Aflah (*c.* 1130), 206
Jâbir ibn Ḥaiyân al-Ṣûfî. *See* Jeber
Jacob. *See* Ya'qûb
Jacob ben Machir (*c.* 1250), 210
Jacob ben Nissim (*c.* 800), 188
Jacob Caphanton (*c.* 1430), 243
Jacob Carsono (*c.* 1375), 241
Jacob of Cremona (*c.* 1449), 247
Jacob Poël (*c.* 1360), 241
Jacob, Simon (*c.* 1550), 338
Jacobi, C. G. J. (*c.* 1830), 506
Jacoli, F., 242, 366, 385
Ja'far ibn Mohammed (*c.* 880), 174
Jainas (Jinas), 161
Jaipur, 154
Japan, 144, 151, 273, 437, 535
Jartoux, P. (*c.* 1710), 534, 536
Jauharî, al- (*c.* 830), 176
Jean d'Estrées, 298
Jean de Lignères (*c.* 1340), 238
Jean de Meurs (*c.* 1350), 238
Jebb, R. C., 260, 314, 344
Jebb, S., 224
Jeber (Geber, *c.* 770), 168
Jehuda ben Salomon Kohen (*c.* 1240), 209
Jehuda Verga (*c.* 1450), 263
Jemshîd. *See* al-Kashî
Jenghiz Khan, 268, 287
Jerusalem, 61
Jews, 172, 206, 240, 263, 346, 350
Jindai monji, 144
Joannes. *See* Johannes, Johann, John, Jean, Giovanni
Joannes Grammaticus (*c.* 640), 191
Joannes de Monteregio. *See* Regiomontanus
Joannes de Muris (*c.* 1350), 238
Joannes Pediasimus (*c.* 1330), 234
Joannes Philoponus (*c.* 640?), 191
Johann van Kunsperck. *See* Regiomontanus
Johann von Gmünden (*c.* 1430), 258
Johannes. *See* Joannes, Johann, John, Jean, Giovanni
Johannes Hispalensis (Hispanensis) (*c.* 1140), 209, 170
Johannes de Lineriis (*c.* 1340), 238
Johannes Saxoniensis (14th century), 238
John. *See* Joannes, Johannes, Johann, Jean, Giovanni
John of Basingstoke (*c.* 1240), 222
John of Beverley (*c.* 690), 185
John of Luna (*c.* 1140), 209, 170
John of Seville (*c.* 1140), 209, 170
Johnson, G., 128
Jones, W., 34
Jordanus Nemorarius (of Namur, de Saxonia) (*c.* 1225), 226, 199
Jorjânî, al- (*c.* 1000), 174
Joseph ben Wakkar (*c.* 1380), 241
Josephus Problem, 207
Jourdain, P. E. B., 398
Juan Diez. *See* Diez
Jûḥannâ. *See* Bar Hebræus
Junge, G., 75
Junta dos Mathematicos, 263

Kästner, A. G. (*c.* 1770), 541
Kahun, 47
K'ai-yüan Chan-king, 151
Kâlakriyâ, 155
Kaldi, 39, 100
Kalonymos ben Kalonymos (*c.* 1320), 241
Kâmil, Abû (*c.* 900), 177
Kankah (*c.* 766), 167, 168
Kanroku, 151
Karamzin, N. M., 350
Karkhî (*c.* 1020), 283
Karpinski, L. C., xv, 98, 166, 170, 177, 238, 239, 262, 413
Kashî, al- (*c.* 1430), 289
Kaye, G. R., 98, 144, 145, 153, 164, 274
Kazi Zadeh. *See* Kashî
Keane, A. H., 14
Kelvin, Lord (*c.* 1880), 470
Kemâl ed-dîn ibn Yûnis (*c.* 1200), 287
Kenryū. *See* Miyake
Kepler, J. (*c.* 1610), 416
Kern, H., 153
Kersey, J. (*c.* 1650), 414

Ketensis. *See* Robert of Chester
Keyser, C. J., 531
Khayyam, Omar (*c.* 1100), 286
Khâzin (*c.* 950), 175
Khiḍr (*c.* 1000), 285
Khorâsân, 172
Khosrú (*c.* 525), 164
Khowârizmî (*c* 825), 170, 168, 203, 329
Khwarezm, 170
Ki-li, emperor (*c.* 1200 B.C.), 97
Killingworth, J. (*c.* 1435), 261
Kindî. *See* Alchindi
King, C. W., 70
Kingsley, C., 137
Kingsmill, T. W., 140
Kircher, A. (*c.* 1650), 422
Kirkman, T. P. (*c.* 1850), 469
Kitâb al-Fihrist (*Book of Lists*), 177
Kittoku. *See* Isomura
K'iu Ch'ang ch'un (*c.* 1221), 268
K'iu-ch'ang Suan-shu (*Arithmetic in Nine Sections*), 31, 96, 142, 152
Kiu-shu, 152
Kiu-szu, 152
Klau. *See* Clavius
Klos, T. (*c.* 1538), 346
Kluge, F., 196
Knobel, E. B., 131, 289
Knott, C. G., 391
Knotted cords, 139
Köbel, J. (*c.* 1520), 336
Koenig, S. (*c.* 1750), 474
Köpke, E. R. A., 189
Konen, H., 412
Korea, 144, 151
Kosta. *See* Qosṭâ
Kóu Shóu-king (*c.* 1290), 272
Kovalévsky, S. (*c.* 1885), 530
Kōwa. *See* Seki
Krauss, S., 138
Kroll, W., 103
Kronecker, L. (*c.* 1870), 511
Krukovsky. *See* Kovalévsky
Kublai Khan, 272
Kummer, E. E. (*c.* 1850), 507
Kushi. *See* al-Kashî
Kûshyâr ibn Lebbân (*c.* 1000), 176
Kuster, L., 69
Kusumapura, 153, 155
Kuṭakhâdyaka, 158
Kutta, W. M., 302
K'u Ying-hsiang (*c.* 1550), 351

Lachmann, K., 124
Lacouperie, A. T. de, 24, 25
Lacroix, S. F. (*c.* 1800), 492, 540
Ladies' Diary, 447
Lafon, M. A., 495
Lagny, T. F. de (*c.* 1710), 472
Lagrange, J. L. (*c.* 1780), 482
Laguerre, E. (*c.* 1880), 500
Lahire, P. de (*c.* 1690), 386
Lair, J., 186
Lalande (*c.* 1800), 540
Lalitavistara, 98
Laloubère. *See* Loubère
Lalouvère. *See* Loubère
Lambert, J. H. (*c.* 1770), 480
Lambo, Ch., 261
Lamé, G. (*c.* 1840), 500
Landen, J. (*c.* 1760), 457
Landi, F., 516
Lanfreducci (*c.* 1490), 245
Lange, J., 525
Laodicea, 59, 61
Lao-Tze, 70
Laplace, P. S. (*c.* 1800), 486
La-pu-tan (*c.* 1267), 272
Larissa, 59, 61
Larsam, 39
Lāṭa, 145
Latin mathematical vocabulary. *See* Veratti
Lauchen, G. J. von. *See* Rhæticus
Lavrovsky, 350
Lax, G. (*c.* 1515), 345
Lazesio. *See* Feliciano
Lazzarini, M., 215
Le. *See* Li
Leap year. *See* Calendar
Least squares, 448, 502
Le Besgue, V. A., 214
Lebon, E., 499
Lebrixa, 264
Legendre, A. M. (*c.* 1800), 487
Leibniz, G. W. Freiherr von (*c.* 1682), 417
Lejeune-Dirichlet. *See* Dirichlet
Leland, J., 204
Leodamas (*c.* 380 B.C.), 91
Leonardo of Cremona (*c.* 1425), 247
Leonardo Fibonacci (of Pisa) (*c.* 1202), 214, 206
Leonardo da Vinci (*c.* 1500), 294, 254
Lepsius, J., 480
Leslie, J., 445
Letters of the era of the gods, 144
Leucippus (*c.* 440 B.C.), 80
Levi, B., 98
Levi ben Gerson (*c.* 1330), 240
Lew. *See* Liu
Leybourn, W. (*c.* 1670), 414

L'Hôpital. *See* L'Hospital
L'Hospital, G. F. A. de (*c.* 1690), 384
Lhuilier, S. A. J. (*c.* 1800), 525
Liang I (two principles), 25
Liber abaci. See Fibonacci
Liber augmenti et diminutionis, 207
Liber embadorum, 206
Liber quadratorum, 217
Liberal arts, 180
Libraries: Alexandrian, 103; early Renaissance, 243; Laurentian, 243; Vatican, 243
Librarii, 244
Libri, G. (*c.* 1850), xv, 545, 377
Li Chi Ts'ao (*c.* 1605), 304
Lie, S. (*c.* 1890), 527
Liebhard. *See* Camerarius
Lieblein, J., 45
Lietzmann, W., 69
Life, advent of, 5
Lignères, Jean de (*c.* 1340), 238
Lilāvati, 276, 158
Lilawati. See Lilāvati
Lineriis, Johannes de (*c.* 1340), 238
Liouville, J. (*c.* 1850), 500
Li Ping (*c.* 250), 142
Li Shu (*c.* 2700 B.C.), 24
Little, A. G., 221, 222
Little, A. J., 22
Liu-chang, 152
Liu Chih (*c.* 289), 143
Liu Chih the Younger (*c.* 1360), 143
Liu Hsiao (*c.* 25), 141
Liu Hsing (*c.* 66), 141
Liu Hui (*c.* 263), 142, 25, 267
Liu I (*c.* 1250), 271
Liu Ju-hsieh (*c.* 1250), 271, 272
Liveriis. *See* Lineriis
Li Yeh (*c.* 1250), 270, 535
Lobachevsky, N. I. (*c.* 1825), 529
Loci, plane, 117; solid, 95
Lockwood, D. P., 201
Loftus, W. K., 39
Logarithmic spiral, 428
Logarithms, 338, 391, 394, 416, 433, 436, 534
Loggan, D., 407
Logistic, 57
Lokotsch, K., 285
Longobardi, N. (*c.* 1618), 436
Longomontanus, 411
Loomis, E. (*c.* 1860), 539
Loria, G., xv, 214, 343, 366, 482, 514
Loritus. *See* Glareanus
Lo-shu, 28
Loubère, A. de la (*c.* 1640), 385
Loxodrome, 423
Lubbock, Sir J., 14
Lubûdî, al- (*c.* 1260), 288
Luca di Borgo. *See* Pacioli
Luca da Firenze (*c.* 1400), 245
Lucas, E., 24
Lucas, H. (*c.* 1640), 395
Lucasian professorship, 395
Lucianus (2d century), 128
Ludolf van Ceulen (*c.* 1580), 330
Ludolph. *See* Ludolf
Luke. *See* Luca
Lullus, Raymundus (*c.* 1280), 229
Lunes of Hippocrates (*c.* 460 B.C.), 82
Luther, M., 325, 338
Lutz, H. F., 41
Luzzato, 207
Lyons, 306
Lysis, 81

Mabillon, J., 184
McClatchie, T., 25
McClenon, R. B., 217
McClintock, E. (*c.* 1890), 533
MacCullagh, J. (*c.* 1840), 469
Macfarlane, A. (*c.* 1900), 546, 461
McGee, W. J., 14
Maclaurin, C. (*c.* 1740), 452
Maclaurin's Theorem, 453
MacMahon, P. A., 463
Magic squares, 28, 209
Magini, G. A. (*c.* 1595), 367
Magna Græcia, 72
Magnin, Ch., 189
Magnitzky (*c.* 1700), 435
Mahābhārata, 35, 98
Mâhânî. *See* Almâhânî
Mahāvīra. *See* Mahāvīrācārya
Mahāvīrācārya (*c.* 850), 161, 152, 154
Mahler, E., 46
Mailly, E., 526
Maimonides (*c.* 1175), 208
Maire, A., 381
Majriṭî (*c.* 1000), 192
Malchus the Tyrian (*c.* 275), 135
Malfatti, G. F. G. (*c.* 1770), 515
Malfatti's problem, 515, 538
Malves. *See* Gua de Malves
Mamercus, 77
Mâmûn, al- (*c.* 820), 169
Mancini, G., 247, 252
Manduith, John (*c.* 1320), 236
Manfredi, Eraclito (*c.* 1720), 512
Manfredi, Eustachio (*c.* 1720), 512

Manfredi, G. (*c.* 1730), 512
Manfredonia, G. B. (*c.* 1475), 254
Mangili, G., 516
Mangu Khan, 268
Mankah, 167
Mann, A., 14
Manning, Mrs., 155
Mansion, P. (*c.* 1890), 546
Mansûr, al- (*c.* 770), 167
Manṣûr ibn 'Alî (*c.* 1000), 285
Mantovani, V., 295
Manuel. *See* Moschopoulus, Chrysoloras
Manzolo (Manzolino), 249
Maps, mathematical-historical, 60, 113, 154
Marco Polo (*c.* 1271), 269
Mares (Marros) (*c.* 1850 B.C.), 45
Margarita Phylosophica, 326, 213, 214
Mariani, G. (*c.* 1535), 306
Marie, M. (*c.* 1880), xv, 544
Marinus of Flavia Neapolis (*c.* 485), 138
Marinus of Tyre (*c.* 150), 129
Marolois, S. (*c.* 1600), 423
Marre, A., 211, 261, 313, 351
Marshman, J. C., 34
Martin, T. H., 125, 129
Martin, Th., 190
Martin, W. A. P., 31
Martinus (Martínez), J. *See* Blasius, J. M.
Marullo. *See* Maurolico
Mascart, J., xv
Mascheroni, L. (*c.* 1800), 516
Mâ-shâ'-allâh. *See* Messahala
Masîḥî, al- (*c.* 1000), 174
Maspero, H., 108
Master Nicholas (*c.* 1250), 218
Masterson, T. (*c.* 1590), 324
Mas'ûdî (*c.* 950), 149
Maternus, J. Firmicus (*c.* 340), 135
Mathematics, defined, 1; elementary, 444
Mathematum Rudimenta quaedam (anon.), 200
Mathew. *See* Matteo
Matsunaga Ryōhitsu (*c.* 1730), 536
Matteo da Firenze (*c.* 1400), 245
Matteo Ricci (*c.* 1603), 303, 435
Matthaei, O., 241
Matthew. *See* Matteo
Maupertuis, P. L. M. de (*c.* 1730), 473
Maurolico, F. (*c.* 1550), 301
Maurolycus. *See* Maurolico
Maurus, Hrabanus (*c.* 820), 188
Mauthner, F., 137
Maxima and minima, 424
Maximus Planudes (*c.* 1340), 233
Maxwell, J. C. (*c.* 1870), 470
Meadows of Gold, 149
Mean proportionals, 83
Méchain, P. F. A. (*c.* 1780), 491
Mechanics, 112, 137
Medieval curriculum, 214
Medma (Medmæus), 59, 61, 91
Megasthenes, 153, 154
Megiorini, P. (*c.* 1540), 257
Meier, R., 125
Mei Wen-ting (*c.* 1675), 436
Melanchthon, P. (*c.* 1540), 331, 328
Men (*c.* 575), 150
Menæchmian Triads, 92
Menæchmus (*c.* 350 B.C.), 92
Mendes, 59, 61
Mendoza, A. de, 354
Menelaus (*c.* 100), 126, 172, 302, 406
Menge, 103
Menher, V. (*c.* 1570), 342
Mensa Pythagorica, 129
Méray, C. (*c.* 1880), 500
Mercator, N. (*c.* 1650), 434
Merejkowski, D., 204
Merian, P., 427
Merlai. *See* Morley, D.
Merriman, M., 488
Mersenne, M. (*c.* 1630), 380
Mervarrûdî, al- (*c.* 830), 169
Mervazî, al- (*c.* 860), 174
Mesopotamia, 33, 37, 99, 146, 16
Messahala (*c.* 800), 169
Messina, 59, 61
Metal discovered, 22
Metius. *See* Adriaen
Meton (*c.* 432 B.C.), 83
Metonic Cycle, 83
Metric system, 484, 491, 494
Metrodorus (*c.* 500 ?), 133
Meurs. *See* Jean de Meurs
Meusnier de la Place, J. B. M. C. (*c.* 1790), 499
Mexico, 353
Meyer, K. F., 14
Meyer, W. A., 137
Méziriac. *See* Bachet
Michael Scott (*c.* 1225), 220, 210
Michizane. *See* Tenjin
Micyllus, J. (*c.* 1550), 332
Middle Kingdom, Egypt, 47
Midolo, P., 111

Mieli, A., 64
Migne, 197
Mikami, Y., xv, 23
Milanesi, G., 214
Miletus, 56, 59, 61
Milhaud, G. (*c.* 1900), 546, 98, 371
Miller, G. A., xv
Miller, W. J. C., 466
Millet, J., 371
Mishnath ha-Middoth, 174
Misrachi, Elia (*c.* 1500), 350
Miṣrî. *See* Aḥmed ibn Yûsuf
Miyake Kenryû (*c.* 1715), 536
Möbius, A. F. (*c.* 1830), 504
Moerbecke, William of (*c.* 1250), 229
Moeris, 45
Mohammed ibn 'Abdallâh ibn 'Aiyâsh. *See* al-Ḥassâr
Mohammed ibn 'Abderraḥmân. *See* Ibn al-Kâtib
Mohammed ibn Aḥmed, al-Bêrûnî. *See* Albêrûnî
Mohammed ibn Aḥmed ibn Mohammed ibn Roshd, Abû Velîd. *See* Averroës
Mohammed ibn al-Leiṯ (*c.* 1000), 285
Mohammed ibn Ibrâhîm, 168
Mohammed ibn 'Îsâ. *See* Almâhânî
Mohammed ibn Jâbir. *See* Albategnius
Mohammed ibn Ketîr. *See* Alfraganus
Mohammed ibn Ma'rûf (*c.* 1550), 351
Mohammed ibn Mohammed. *See* Abû'l-Wefâ, Averroës, al-Fârrâbî
Mohammed ibn Mohammed, Naṣîr ed-dîn. *See* Naṣîr ed-dîn
Mohammed ibn Mûsâ. *See* al-Khowârizmî
Mohammed ibn 'Omar, al-Râzî (*c.* 1200), 286. *See also* Abenbêder
Mohammed ibn Yaḥyâ. *See* Avenpace
Mohammed ibn Zakarîyâ. *See* Rhases
Mohl, J., 25
Moivre. *See* De Moivre
Moltzer. *See* Micyllus
Monge, G. (*c.* 1800), 490
Monier-Williams, M., 153
Monmort. *See* Montmort
Montalte. *See* Pascal
Monte Regal Piedmontois (*c.* 1585), 374
Montgomery, J. E. G. de, 185
Montmort, P. R. de (*c.* 1710), 471
Montucla, J. E. (*c.* 1750), xv, 540
More, T. (*c.* 1516), 316
Morgan, M. H., 123
Môri Kambei Shigeyoshi (*c.* 1592), 352
Morley, Daniel (*c.* 1200), 204
Morley, F. V., 388
Morley, H., 295
Morley, S. G., 14
Morse, H. B., 140
Mortality table, 131
Mortet, V., 132
Moschopoulus (*c.* 1300), 234
Moses ben Maimun. *See* Maimonides
Moses ben Tibbon (*c.* 1250), 209, 210
Moses Kapsali, 350
Moya, J. P. de (*c.* 1562), 345
Müller, C. F., 330
Müller, F., 241
Müller, Johann. *See* Regiomontanus
Müller, T., 327
Muhammed. *See* Mohammed
Mullach, F. W. A. (F. G. A.), 80
Multiplication tables, 40, 58, 129, 138, 314
Muñoz, J. (*c.* 1566), 345
Munyos. *See* Muñoz
Murai Chûzen (*c.* 1765), 537
Muramatsu Kudayû Mosei (*c.* 1663), 439
Muris. *See* Jean de Meurs
Mûsâ ibn Shâkir (*c.* 830), 171
Mûsâ ibn Yûnis (*c.* 1200), 287
Music, 75, 106
Muslim ibn Aḥmed al-Leiṯî (*c.* 900), 192
Mydorge, C. (*c.* 1630), 378
Myriad, 116, 144
Mysore, 154, 161
Mystery of numbers, 304
Mysticism, 16, 89

Nadîm (*c* 987), 177
Nagl, A., 105. 239
Naigeon, 486
Nairîzî, al- (*c.* 910), 176
Nakane Genjun (*c.* 1750), 536
Nakane Genkei (*c.* 1700), 440, 536
Nânâ Ghât, 154
Napier, J. (*c.* 1614), 389
Naples, 59, 61
Narducci, E., 182, 220, 229, 253
Nasavî (*c.* 1025), 285
Naṣîr ed-dîn (*c.* 1250), 287
Nau, F., 288
Nauck, A., 69
Nazâm ed-dîn Aḥmed, 351
Nazîf ibn Jumn (Jemen) (*c.* 980), 174
Neander, A., 185
Neander, M. (*c.* 1570), 332

Nebuchadnezzar (*c.* 550 B.C.), 100
Negative numbers, 40, 159, 162
Nehâvendî (*c.* 820), 169
Neile, W. (*c.* 1605), 412
Nelli, G. P. C. de', 364
Ne-mat-re, 45
Netherlands, 338, 422, 526
Neumann. *See* Neander
Newbold, W. R., 82
Newcomb, S. (*c.* 1880), 532
Newton, Sir Isaac (*c.* 1680), 398
Nicæa, 59, 61
Niccolò de' Niccoli (*c.* 1400), 243
Nicholas, Master, 218
Nicholas Cusa (*c.* 1450), 258
Nicholas Rhabdas (*c.* 1341), 235
Nicolas, P. (*c.* 1700), 387
Nicole. *See* Oresme
Nicole, F. (*c.* 1730), 472
Nicomachus (*c.* 100), 127, 179
Nicomedes (*c.* 180 B.C.), 118
Nicoteles (*c.* 250 B.C.), 107
Nigidius. *See* Figulus
Nilometer, 46
Nine Sections. See K'iu-ch'ang
Nines, casting out, 207
Nippur, 40
Nipsus, M. Junius (*c.* 180), 131
Nissim, Jacob ben (*c.* 800), 188
Nix, L. M. L., 172
Nobutake. *See* Baba
Nöldeke, T., 164
Noether, M., 463, 505, 527
Nonius. *See* Nunes
Notker Labeo (*c.* 1000), 196
Novara, D. M. (*c.* 1490), 254
Nozawa Teichō (*c.* 1665), 439
Nuffar, 40
Nugne. *See* Nunes
Number and language, 6
Numbers, perfect, 189, 307; mystery of, 304
Numerals, Russian, 262; Sumerian, 36; Syriac, 8
Nunes, P. (*c.* 1530), 348
Núñez. *See* Nunes
Nûr ed-dîn al-Beṭrûjî. *See* Alpetragius

O'Connell, Mme F., 531
O'Creat, N. (*c.* 1150), 204
Ocreatus. *See* O'Creat
Odo of Cluny (*c.* 920), 190
Odoric, Friar (*c.* 1300), 272
Œnopides (*c.* 465 B.C.), 79, 83
Oktai Khan (*c.* 1240), 287
Oliphant, Mrs., 249
Olivier, T. (*c.* 1840), 500
Olleris, A., 195
'Omar ibn Ibrâhîm al-Khayyâmî. *See* Omar Khayyam
Omar Khayyam (*c.* 1100), 286
Omeiya. *See* Abû'l-Ṣalt
Oppert, G., 35
Optics, 106
Opuntius (*c.* 375 B.C.), 91
Oresme (*c.* 1360), 239, 109, 233
Orient, 95, 138, 350, 435, 533
Oriental mathematics. *See* Babylon, Chaldea, China, India, Japan, Persia
Ornament, geometric, 15
Ortega, J. (*c.* 1512), 344
Ortiz, D. (15th century), 263
Osgood, W. F., 533
Otho. *See* Otto
Otto, V. (*c.* 1573), 340
Oughtred, W. (*c.* 1630), 392
Oyama Shōkei (*c.* 1728), 535

π, 44, 114, 141–144, 150, 156, 158, 163, 164, 198, 242, 247, 270, 275, 312, 331, 340, 400, 411, 436, 437, 536
Pachymeres, Georgios (*c.* 1300), 229
Pacioli (*c.* 1494), 251, 317
Paciuolo. *See* Pacioli
Pa-kua, eight trigrams, 25
Palibothra, 154
Pamphila, 67
Pañca Siddhāntikā, 156
Pan Ku (*c.* 75), 141
Paolo Dagomari, dell' Abaco (*c.* 1340), 232
Paolo Megiorini (*c.* 1540), 257
Paper, 140
Pappus (*c.* 300), 136, 302
Papyrus, 71
Parabola, 92, 116
Paraboloid, 114
Parchment, 105
Parent, A. (*c.* 1710), 471
Paris, 306
Parmenides of Elea (*c.* 460 B.C.), 81
Paros 59, 61
Pascal, B. (*c.* 1650), 381
Pascal's Triangle, 382, 329, 333, 537
Pastor. *See* Rey Pastor
Pāṭaliputra (Pāṭoliputra), 153
Pāṭaliputraka, 154
Patna, 153, 154
Paton, W. R., 133
Pauliśa Siddhānta, 145
Pauly-Wissowa, xv, 103
Peachamus. *See* Peckham

Peacock, G. (*c.* 1830), 459
Peccamus. *See* Peckham
Peckham, John (*c.* 1280), 224, 296
Pediasimus (*c.* 1330), 234
Peepin, A. N., 262
Peet, T. E., 47, 48
Peiper, R., 108
Peirce, B. (*c.* 1850), 532
Peirce, C. S., 218
Peking, instruments at, 272
Peletier (Peletarius), J. (*c.* 1560), 313
Pell, J. (*c.* 1650), 411
Pell Equation, 160
Pellos (Pellizzati) (*c.* 1492), 255
Péluse. *See* Monge
Penno, 52
Perez (modern Pérez). *See* Moya
Pérez, J. A. S., 211
Perga, 50, 61, 116
Pergamum, 59, 61
Pericles (*c.* 444 B.C.), 87
Peripatetic School, 93
Perito Annotio. *See* Cataldi
Permutations, 274. *See also I-king*
Perseus (*c.* 150 B.C.), 118
Persia, 143, 164, 283, 286
Perspective, 79, 123
Pescheck, C. (*c.* 1720), 502
Peschel, O. F., 14
Pestalozzi, J. H. (*c.* 1776), 524
Peter. *See* Petrus, Pietro
Peters, C. H. F., 131
Petrarch, 230
Petrie, W. M. Flinders, 47
Petrus. *See* Peter, Pietro
Petrus de Alliaco (*c.* 1400), 240
Petrus de Dacia (*c.* 1325), 238
Petrus de Maharncuria (*c.* 1265), 226
Petrus Peregrinus (*c.* 1265), 226
Petrus Philomenus (*c.* 1325), 238
Peurbach, G. von (*c.* 1460), 259
Pez, 107
Pfaff, J. F. (*c.* 1810), 510
Phaeinus (*c.* 432 B.C.), 83
Phidias (Pheidias) (*c.* 443 B.C.), 87
Philippus Medmæus (*c.* 350 B.C.), 61, 91
Phillips, P. L., 388
Philolaus (*c.* 425 B.C.), 81, 73, 76
Philoponus, Joannes (*c.* 640?), 191
Philosophical Transactions, 259
Pi. *See* π above
Picatoste y Rodríguez, F., 343
Picavet, F., 195
Piedmontois. *See* Monte Regal
Piero. *See* Pietro, Petrus
Pierpont, J., 445, 531
Pietro d'Abano (*c.* 1300), 220
Pietro Franceschi (della Francesca) (*c.* 1475), 247
Pithsanus. *See* Peckham
Pitiscus, B. (*c.* 1595), 331
P'i Yen-tsung (*c.* 440), 143
Plane numbers, 155
Planudes, Maximus (*c.* 1340), 233
Plato (*c.* 380 B.C.), 87; on arithmetic, 57, 80; on geometry, 90
Plato of Tivoli (Tiburtinus) (*c.* 1120), 201, 207
Platonic number, 89
Playfair, J., 292
Pleyte, W., 50
Pliny (*c.* 75), 123, 45
Plücker, J. (*c.* 1850), 505
Plumian professorship, 447
Plus and minus signs, 258, 330, 341
Plutarch, 68, 88, 115
Poël, Jacob (*c.* 1360), 241
Poggendorff, J. C., xv
Poggio Bracciolini (*c.* 1414), 243
Poincaré, H. (*c.* 1900), 499
Poinsot, L. (*c.* 1820), 499
Poisson, S. D. (*c.* 1820), 495
Poland, 346
Polar triangle, 423
Polemarchus (4th century B.C.), 95
Polycrates, 70
Polygonal numbers, 90, 119
Polyhedrons, regular, 73, 75, 95, 119
Pompeii, 59, 61
Poncelet, J. V. (*c.* 1830), 496
Porisms, 107
Porphyrius (*c.* 275), 135
Portugal, 348
Porus. *See* Sporus
Poseidonius (*c.* 100 B.C.), 118
Posidonius. *See* Poseidonius
Poudra, N. G., 383
Poullet-Delisle, A. C. M., 504
Predari, F., 362
Prime numbers, 109
Principia, Newton's, 402, 406, 447
Printing, origin of, 244; early, 249
Prior, O. H., 314
Probability, 382, 428
Problems, famous and fanciful, 50, 133, 159, 162, 163, 164, 187, 207, 280
Proclus (*c.* 460), 137, 67, 302
Progressions, 40, 119, 150
Projection, 79, 119
Prophatius (*c.* 1250), 210
Proportion, 90

Proportion, harmonic, 75
Prosdocimo de' Beldamandi (*c.* 1410), 246
Prowe, L., 347
Psammetichus I (*c.* 625 B.C.), 71
Psammetichus III (*c.* 525 B.C.), 71
Psellus (*c.* 1075), 197
Ptolemæus, Claudius. *See* Ptolemy
Ptolemais, 50, 61
Ptolemy, Claude (*c.* 150), 130, 145, 168, 169, 170, 228, 302
Ptolemy IV (Philopator), 116
Ptolemy Soter, 102
Pugliesi, O., 215
Pulverizer, 158, 280
Purbach. *See* Peurbach
Putna. *See* Patna
Putnam, G. H., 244
Puzzles, 133. *See also* Problems
Pyramids of Egypt, 43, 44
Pythagoras (*c.* 540 B.C.), 69, 254
Pythagorean numbers, 90, 97, 281
Pythagorean Theorem, 75, 30, 33, 97, 137

Qaisar ibn Abî'l-Qâsim. *See* Ta'âsîf
Qalasâdî (*c.* 1475), 211
Qâsim. *See* al-Majrîṭî
Qass (*c.* 900), 174
Qible (*c.* 900), 192
Qiwâm ed-dîn Yaḥyâ (12th century), 209
Qorra. *See* Ṯâbit
Qosṭâ ibn Lûqâ al-Ba'albekî (*c.* 900), 172
Quadratic equation, 134, 155, 159, 162, 280, 281
Quadratrix, 82, 92, 136
Quadrature, 83, 197, 236, 408. *See also* Circle
Quadrilateral, 158, 163
Quadrivium, 180
Quarter squares, 283
Quaternions, 461
Quetelet, L. A. J. (*c.* 1850), 526, 424

Ra-ā-us, 47
Rabanus. *See* Hrabanus
Rabbi ben Ezra. *See* Abraham ben Ezra
Radau, H., 37
Radix in counting, 8
Radulph of Laon (*c.* 1125), 205
Rafaele Canacci (*c.* 1380), 232
Raganātha (*c.* 1621), 435
Rahn, J. H. (*c.* 1660), 412
Raidō. *See* Arima
Rainer. *See* Gemma
Ralphson. *See* Raphson
Ramaka, queen, 49
Ramanujacharia, N., 274
Rambaud, A., 178
Ramée. *See* Ramus
Rameses II (*c.* 1347 B.C.), 51
Rameses III (*c.* 1180 B.C.), 52
Rameses IV (*c.* 1167 B.C.), 52
Rameses VI (*c.* 1150 B.C.), 52
Ramus, P. (*c.* 1550), 300, 260
Raṅgâcârya, M., 161
Raphael, 87, 170
Raphson, J. (*c.* 1715), 449
Rashid, Harun al- (*c.* 800), 168, 149
Rashid ed-din, 269
Ratdolt, 251
Ratzel, F., 14
Rawlinson, H. G., 36, 153
Rayleigh, Lord (*c.* 1900), 470
Râzî (*c.* 920). *See* Rhases
Râzî the Younger (*c.* 1200), 286
Reade, William (*c.* 1370), 237
Rebière, A. (*c.* 1890), 546, 478
Rechenmeisters, 250, 337
Rechenschule, 256
Reclus, E., 154
Recorde, R. (*c.* 1542), 317
Rectifications, 412, 424, 458
Redenti, F., 516
Reduction, method of, 83
Regiomontanus (*c.* 1470), 259
Regnier. *See* Gemma
Regula sex quantitatum, 127
Regula de tribus (detri). *See* Rule of Three
Reisch, G. (*c.* 1503), 326
Rekhmire, tomb of, 49
Religion, 16, 143, 194
Remigius of Auxerre (*c.* 900), 188
Rémusat, 271
Renaissance, 242
Renier. *See* Gemma
Retinensis. *See* Robert of Chester
Rey Pastor, J., 343
Reye, K. T. (*c.* 1880), 511
Rhabdas, Nicholas (*c.* 1341), 235
Rhæticus, G. J. (*c.* 1550), 333, 331
Rhases (*c.* 920), 175
Rhind, A. H. (papyrus), 48
Rho, G. (*c.* 1618), 430
Rhodes, 59, 61
Rhonius. *See* Rahn
Rhumb line, 423
Ricca, F., 517

Riccardi, P., 103, 249, 309, 541
Riccati, F. (*c.* 1750), 514
Riccati, G. (*c.* 1750), 513
Riccati, J. F. (*c.* 1720), 512
Riccati, V. (*c.* 1750), 513
Riccati Equation, 513
Ricci, Matteo (Matthew) (*c.* 1603), 303, 435
Richard, C. (*c.* 1630), 370
Richard of Wallingford (*c.* 1330), 235
Richardson, H. K., 143
Riemann, G. F. B. (*c.* 1850), 508
Riese, A. (*c.* 1518), 337
Ringelbergius (Ringelbergh), J. F. (*c.* 1530), 339
Rithmomachia, 198, 197
Ritzenstein, A., 137
Rixner and Siber, 299
Robert of Chester (*c.* 1140), 203, 329
Robert Greathead. *See* Greathead
Robert of Lincoln. *See* Greathead
Roberval, G. P. de (*c.* 1640), 385
Robinson, L. G., 114
Roche, Estienne de la (*c.* 1520), 313, 261
Rockhill, W. W., 140
Rodet, L., 153, 207
Rodríguez. *See* Picatoste
Roettier, 401
Roger Bacon. *See* Bacon
Rollandus (*c.* 1424), 261
Rolle, M. (*c.* 1700), 387
Rollin Papyrus, 50
Romain. *See* Roomen
Romanus. *See* Roomen
Rome, 59, 61, 120
Roomen, A. van (*c.* 1593), 339
Roots, approximations for, 284
Roper, M., 317
Rosen, F., 174
Roth, P. (*c.* 1610), 421
Rothlauf, B., 90
Rowe, J. (*c.* 1751), 448
Royal road to geometry, 92
Rubrouck (13th century), 268
Rudio, F., 84, 480, 509
Rudolff, C. (*c.* 1525), 328
Rudolph of Bruges (*c.* 1144), 202
Rudorff, A., 124
Ruffini, P. (*c.* 1800), 515
Rule of False Position. *See* False Position
Rule of Three, 32, 150
Rupartus Lincolniensis. *See* Greathead
Russ, J. *See* Rho
Russia, 262, 350, 435
Ryōhitsu. *See* Matsunaga
Sa'adia ben Joseph (*c.* 930), 192
Sabbionetta. *See* Gherardo
Sachau, E. C., 153, 285
Sacrobosco (*c.* 1250), 221
Sadasuke. *See* Fujita
Sadlerian professorship, 466
Sâḥib al-Qiṭ'ah (*c.* 900), 192
Sahl ibn Bishr (*c.* 850), 172
Sa'îd ibn Ya'qûb (*c.* 915), 176
Sa'îd ibn Yûsuf (*c.* 930), 192
Saintsbury, G., 320
Saint-Vincent, G. de (*c.* 1650), 424
Salhab ibn 'Abdessalâm. *See* al-Faraḍî
Sallusti, G. de', 300
Salmon, G. (*c.* 1850), 462
Samos, 59, 61, 70
Samuel ben Abbas (*c.* 1175), 209
Samû'îl. *See* Samuel
Sand Reckoner, 113
San-k'ai Chung-ch'a, 152
San-tŏng-shu, 143
Sargon (*c.* 2750 B.C.), 37, 38
Sarkar, B. K., 153
Sarton, G., 498
Satō Seikō (*c.* 1665), 439
Saunderson, N. (*c.* 1730), 454
Saurin, J. (*c.* 1710), 472
Savasorda (*c.* 1100), 206
Savile, H. (*c.* 1600), 395
Savilian professorship, 395
Savioli, G., 516
Sawaguchi Kazuyuki (*c.* 1665), 439
Scales of counting, 9, 41
Scandinavia, 240, 527
Scardeone, B., 246
Schall von Bell, J. A. (*c.* 1650), 436
Schanz, 258
Scheubel, J. (*c.* 1550), 329
Scheybl, Scheybel. *See* Scheubel
Schindel (Szcindel), J., 258
Schlegel, G., 18, 23
Schmidt, M., 14
Schmidt, M. C P., 92, 103, 121
Schmidt, W., 125
Schöner. *See* Schoner
Schoner, A. (*c.* 1560), 335
Schoner, J. (*c.* 1530), 335
Schonerus. *See* Schoner
Schoolhouse, oldest, 38
Schooten, F. van (*c.* 1650), 425, 311, 375, 377
Schooten, P. van (*c.* 1670), 425
Schooten, van, family, 425
Schreiber, H., 348. *See also* Grammateus
Schreyber. *See* Grammateus

Schroeder, L. von, 72, 98
Schuere. *See* Van der Schuere
Schultess, K., 195
Schultz, A. (*c.* 1600), 331
Schultz, W., 69
Schur, F., 480
Schurzfleisch, 189
Schwartzerd. *See* Melanchthon
Schwenter, D. (*c.* 1620), 420
Sciliceus. *See* Blasius, J. M.
Scipione del Ferro (*c.* 1506), 295
Score, 13
Scott, Michael (*c.* 1225), 220, 210
Scriptor. *See* Grammateus
Scultetus. *See* Schultz, A.
Sea Island Arithmetic Classic, 142
Sebokht (*c.* 650), 166
Secant, 175
Sédillot, L. P. E. Am., 30, 289
Sefer ha-Echad, 207
Sefer ha-Mispar, 207
Sefer Jezira, 188
Seki Kōwa (*c.* 1680), 439, 535
Senkereh, 39
Serenus, Quintus Sammonicus (*c.* 200), 132
Serenus of Antinoopolis (*c.* 50?), 126
Series, general, 48, 266, 271; infinite, 312, 410, 449, 533; logarithmic, 434; trigonometric, 409
Serret, J. A. (*c.* 1865), 500
Sesostris, 51
Seti I (*c.* 1350 B.C.), 50
Sexagesimal system, 24, 41. *See also* Fractions
Sextus. *See* Africanus
Sfortunati, G. (*c.* 1534), 306, 255
Shadib (14th century), 241
Shakespeare quoted, 74, 106
Shang Kao, 31
Shapiro, H., 174
Shâṭir, Ibn al- (*c.* 1350), 289
She Huang-ti (*c.* 220 B.C.), 138
Shiba Tatsu (554), 151
Shigeyoshi. *See* Mōri
Shi Huang-ti, emperor (*c.* 213 B.C.), 30, 138
Shirazi, J. K. M., 286
Shirwood, Bishop, 199
Shōtoku Taishi (*c.* 600), 151
Shu-king, 24
Shun, emperor (*c.* 2250 B.C.), 25
Shu-shu-ki-yi, 142
Siber, Rixner and, 299
Sicily, 59, 61
Siddhāntas, 145, 156, 157, 167, 168
Sié Fong-tsu (*c.* 1650), 436
Sieve of Eratosthenes, 109
Sighart, 228
Silberberg, M., 207
Siliceo. *See* Blasius, J. M.
Simon, J., 375
Simon, M. (*c.* 1890), 546, 83
Simone. *See* Francesco di Simone
Simony, C. de, 232
Simpson, T. (*c.* 1750), 457
Simson, R. (*c.* 1750), 456
Sinân ibn Ṯâbit ibn Qorra (*c.* 940), 172
Sindhind, 167
Sind ibn 'Alî. *See* Abû'l-Ṯaiyib
Sine, 156, 175
Siu Yo (*c.* 250), 142
Sixtus IV, 260
Sixty, scale of, 41
Skeat, W. W., 169
Slide rule, 340, 394
Sluse (Slusius). *See* Sluze
Sluze, R. F. W., Baron de (*c.* 1660), 425
Smith, C. F., 323
Smith, D. E., xv, 98, 200, 207, 332, 353, 409, 462
Smith, H. J. S., 467
Smith, R. (*c.* 1750), 448
Smith, T., 103
Smith, W., xvi
Smogolenski (*c.* 1650), 436
Smyrna, 59, 61
Smyth, C. P., 44
Snell, R. (*c.* 1600), 422
Snell, W. (*c.* 1620), 423, 331, 343
Snellius. *See* Snell
Snyder, Carl, 107
Socrates (*c.* 425 B.C.), 79
Sole, F. dal (*c.* 1530), 313
Solid numbers, 90, 155
Solids. *See* cube, polyhedrons
Solon (*c.* 600 B.C.), 65
Sonyei. *See* Araki
Sophists, 82
Soroban, 352
South-pointing vehicle. *See* Compass
Spain, 192, 205, 343, 434
Spencer, H., quoted, 3
Speusippus (*c.* 340 B.C.), 90
Sphere, volume of, 156; astronomical, 91, 94, 127, 136, 167
Spherics. *See* Sphere, astronomical
Spirals, 3, 44, 107, 114, 136, 387
Sporus of Nicæa (*c.* 275), 132
Square, diagonal of, 98
Square root, 155

Square roots, tables of, 40
Squares, tables of, 40
Śrīdhara (*c.* 1020), 274, 280
Stabili. *See* Francesco di Simone
Stäckel, P., 480, 521, 524, 528
Stageira (Stagira), 59, 61, 93
Staigmüller, H., 251
Stapulensis. *See* Faber
Stationarius, 244
Staudt, K. G. C. von (*c.* 1850), 505
Steele, R., 224
Steichen, M., 343
Stein, M. A., 140
Steiner, J. (*c.* 1840), 524
Steinschneider, M. (*c.* 1850), 545, 128, 174, 188, 207, 209, 211
Stellar polygons, 218, 236
Stephen of Alexandria (*c.* 610), 191
Stephen Dusan, 242
Sterck van Ringelbergh. *See* Ringelbergius
Stevin, S. (*c.* 1590), 342, 133
Stevinus (Steven, Stephan). *See* Stevin
Stewart, M. (*c.* 1750), 456
Stiattesi, A., 545
Stiefel. *See* Stifel
Stifel, M. (*c.* 1525), 327
Stirling, J. (*c.* 1750), 449
Stobæus (*c.* 500), 62
Stöffler, J. (*c.* 1500), 327
Stone Age, 15, 21
Stone masonry, 22
Stow, J., 204
Strabo (*c.* 10), 122
Strabus, Walafried (*c.* 840), 188
Strachey, E., 278
Strecker, 189
Strutt, J. W. *See* Rayleigh
Suan-king, 29
Suan-pan, 352
Substractio, 345
Suiceth, Richard (*c.* 1345), 236
Suisset. *See* Suiceth
Śulvasūtras, 97, 98
Sumario Compendioso, 353
Sumeria, 35, 37, 39, 146
Sundial, 49, 91, 122, 125, 146
Sunon, Master (*c.* 1340), 240
Sun-tzï (1st century ?), 141
Sun-tzï Suan-king, 141, 152
Surveying, 43, 52, 81, 124
Suryadaśa (*c.* 1535), 351
Sūrya Siddhānta, 34, 145, 167
Suter, H. (*c.* 1890), 546, 168, 177, 212, 242
Sven, Master (*c.* 1340), 240
Swineshead. *See* Suiceth
Switzerland, 348
Syene, 59, 61, 110
Sylow, L., 527
Sylvester, J. J. (*c.* 1850), 463, 532
Symbols, of derivatives, 425; of operation, 258, 330, 341, 394; of proportion, 394; of relation, 320, 389, 394; for unknown, 280, 312
Symonds, J. A., 244, 346
Synesius (*c.* 410), 136
Syracuse (Siracusa), 59, 61, 113
Syriac numerals, 8
Syrians. *See* Arameans
Sz' Siang (four figures), 25
Szŭ-ma Ta (554), 151

τ (Greek letter) for zero, 204
Ta'âsîf (*c.* 1200), 287
Tabarî (*c.* 800), 169
Ṯâbit ibn Qorra (*c.* 870), 171, 209
Ṯâbit the Younger (*c.* 940), 172
Tables, 40, 58, 129, 131, 138, 175, 314, 534. *See also* Chords, Mortality, Multiplication, Roots, Sine, Tangent
Tagliente (*c.* 1515), 304
Ta 'hbula, 207
Taikō (Toyotomi Hideyoshi) (*c.* 1590), 352
Tait, P. G. (*c.* 1880), 470
T'ai-yen calendar, 151
Ṯaiyib (*c.* 890), 174
Takahara Kisshu (*c.* 1600), 437, 353
Takebe Hikojirō Kenkō (*c.* 1722), 535
Talchis. *See* Albanna
Tallentyre, S. G., 404, 480
Tamerlane (*c.* 1400), 287
Ta-nao (*c.* 2700 B.C.), 24
Tang Dynasty (618–907), 149
Tangent, 175
T'ang shu, 149
T'ang Shun-ki (*c.* 1550), 351
Tannery, P., 544, xv, 83, 89, 103, 108, 125, 128, 132, 133, 136, 207, 234, 235, 375, 377, 499
Taqî ed-dîn (*c.* 1575), 351
Tarentum, 59, 61
Ṭâriq. *See* Ya'qûb
Tartaglia, Nicolo (*c.* 1545), 297
Tartalea. *See* Tartaglia
Taxes, 45, 46, 47, 49
Taylor, B. (*c.* 1720), 451
Taylor, J., 158
Taylor's Theorem, 451, 454
Teiken. *See* Fujita

Telescope, 365, 409
Ten, scale of, 10
Tenjin (*c.* 890), 152
Tenzan, 537
Terquem, O., 207, 214
Tetrabiblos. *See* Ptolemy, C.
Textbook writers, 413
Thabit. *See* Tâbit
Thales (*c.* 600 B.C.), 64
Thasos, 59, 61
Theætetus (*c.* 375 B.C.), 86
Thebes, 49
Theodore of Tarsus, 185
Theodorus (*c.* 1220), 217
Theodorus of Cyrene (*c.* 425 B.C.), 86
Theodosius of Bithynia (*c.* 50 B.C.), 125
Theodosius of Tripoli (*c.* 100), 125
Theon of Alexandria (*c.* 390), 136
Theon of Smyrna (*c.* 125), 129
Theophrastus (*c.* 350 B.C.), 83, 94
Theudius (*c.* 360 B.C.), 92
Thibaut, G., 98
Thibetan "Wheel of Life," 27
Thiele, G., 186
Thompson, R. C., 100
Thompson, W. *See* Kelvin
Thorndike, L., 135
Thureau-Dangin, F., 38
Thutmose III (*c.* 1500 B.C.), 50
Thymaridas (*c.* 380 B.C.), 91
Tibbon, Moses ben (*c.* 1250), 209, 210
Tibbon family, 210
Tifernas, Gregory (1458), 260
Timæus, 88
Timur. *See* Tamerlane
Tittel, C., 121
Todhunter, I (*c.* 1850), 468, 382
Toland, J., 137
Tonini. *See* Zuanne
Tonstall, C. (*c.* 1522), 316, 314
Topics for Discussion, 19, 53, 101, 147, 193, 265, 291, 357, 443, 547
Torquemada, J., 248
Torres, T. (*c.* 1500), 264
Torricelli, E. (*c.* 1640), 366
Tortolini, B. (*c.* 1860), 518
Torus, 118
Toshihisa. *See* Iwasaki
Trail, W., 456
Trajan, 146
Translators, periods of, 176, 200, 203
Trapezium. *See* Trapezoid
Trapezoid, 32, 40, 163
Trapezuntios. *See* George of Trebizond
Trebizond. *See* George of Trebizond
Trembley, J. (*c.* 1800), 525
Trenchant, I. (J.) (*c.* 1565), 314
Treutlein, P., 197
Treviso Arithmetic, 248
Triangle, 40, 156, 158; right-angled, 160, 163; Pascal's (*see* Pascal)
Trigonometry, 119, 126, 145, 156, 175, 176, 177, 272, 324, 331
Trigrams (*I a-kua*), 25
Triśatika, 274
Trisection of an angle, 118, 171
Trivium, 180
Tropfke, J., xv
Ts'ai Yüan-ting (*c.* 1180), 267
Ts'ai Yung (*c.* 190), 141
Tschirnhausen, E. W., Graf von (*c.* 1690), 417
Tsu Ch'ung-chih (*c.* 470), 143
Tu Chih-ching (17th century), 436
Türnau, D., 188
T'u-huo-lo, 149
Tun Ch'üan (*c.* 425), 143
Turner, E. R., 166
Turrecremata, J., 248
Ṭûsî (*c.* 1200), 288
Ṭûsî's staff, 288
Twelve, scale of, 11
Twenty, scale of, 12
Tylor, E. B., 14
Tyre, 59, 61

Ujjain, 154, 157
Ulpianus, Domitius (*c.* 200), 131
Ulugh Beg (*c.* 1435), 289
Umbra versa, 175
Unger, F., 256
Unit fraction, 45
United States, 531
Units of counting, 7
Units of measure, 7, 38
Unity, 74, 93
Universities, 188, 212, 231, 245
University of Paris, 188, 245
Ur, 38
Ursinus, B. (*c.* 1620), 420

Vacca, G., 302
Valla, Georgius (*c.* 1475), 247
Vallín, A. F., 343
Vanderbanck, 455
Vander Hoecke (*c.* 1515), 341
Vandermonde, A. T. (*c.* 1775), 481
Van der Schuere, J. (*c.* 1590), 342
Van Schooten. *See* Schooten
Van s'Gravesande (*c.* 1730), 526

Vansteenberghe, E., 258
Varahamihira the Elder (*c.* 200), 156
Varahamihira the Younger (*c.* 505), 145, 152, 154, 156
Varignon, P. (*c.* 1700), 470
Varro (*c.* 60 B.C.), 120
Vassilief, 529
Vassura, 366
Vaulezard, J. L. de, 311
Vaux, W. S. W., 164
Vedas, 35, 97
Vella, P. F. S., 304
Venturi, P. T., 304
Veratti, B., 178, 202
Verbiest, F. (*c.* 1659), 436
Verga, J. (*c.* 1450), 263
Verino, U., 246
Victorinus. *See* Victorius
Victorius of Aquitania (*c.* 450), 138
Vicuña, G. (*c.* 1880), 546
Vieta, F. (*c.* 1580), 310
Viète. *See* Vieta
Vigesimal scale, 12
Vijaganita. *See Bija Ganita*
Villani, F., 232
Vincent de Beauvais (*c.* 1250), 226
Vinci. *See* Leonardo da
Vissière, 142
Vitello (*c.* 1270), 228
Vitruvius (*c.* 20 B.C.), 123, 85
Vittorino da Feltre (*c.* 1430), 233
Viviani, V. (*c.* 1660), 367
Vizinho, D. M. (*c.* 1500), 263
Vogt, H., 75, 91, 98, 105
Voltaire (*c.* 1745), 477, 404
Volterra, V., 400, 514
Vossius, G. J., 221
Vyutkalita, 162

Wäschke, H., 234
Wakkar, Joseph ben (*c.* 1380), 241
Walafried Strabus (*c.* 840), 188
Walcherus (*c.* 1120), 205
Wallace, W. (*c.* 1810), 458
Wallinby, O. *See* Leybourn, W.
Wallingford, Richard of (*c.* 1330), 235
Wallis, J. (*c.* 1650), 406, 535, 539
Walsh, J. J., 212
Wang Fan (*c* 265), 142
Wang Hs'iao-t'ung (*c.* 625), 150
Wang Jong (*c.* 425), 143
Wang Pao-liang (*c* 554), 151
Wang Pao-san (*c.* 554), 151
Wang Pi (*c.* 245), 142
Wappler, E., 200
Ward, A. W., 339
Waring, E. (*c.* 1780), 469
Wasan, 538
Waters, W. G., 247
Wax tablet, 58
Weidner, E. F., 37
Weierstrass, K. (*c.* 1865), 509
Weissbach, F. H., 100
Weissenborn, H., 178, 195, 283, 417
Werner, J., 331
Werner, K., 184, 195
Wertheim, G., 207
West, A. F., 186
Weyr, E., 45
Whewell, W. (*c.* 1840), 546, 396, 402
Whish, C. M., 153
Whiston, W. (*c.* 1700), 451
Whitford, E. E., 412
Widman (Widmann), J. (*c.* 1490), 257
Wiedemann, A., 45
Wieleitner, H., xiv, 363
Wilhelm of Hirschau (*c.* 1050), 197
Wilkinson, J. G., 75
Wilkinson, L., 145
William. *See* Wilhelm, Guilielmus, Guglielmo
William of Champeaux (*c.* 1100), 188
William Fleming. *See* Fleming
William of Moerbecke. *See* Fleming
Williams, S. W., 140
Wilson, J. (*c.* 1780), 458
Wilson's Theorem, 450
Wing, V. (*c.* 1648), 415
Wingate, E. (*c.* 1650), 414
Winterberg, C., 253
Winterfeld, 189
Witelo (*c.* 1270), 228
Wittstein, A., 146, 515
Wön-wang (*c.* 1150 B.C.), 25
Woepcke, F., 117, 211, 284, 286, 289
Wolf, C. (*c.* 1720), 501
Wolf, R., 24, 433
Wolf, S., 137
Wood, A. à, 204
Woodhouse, R. (*c.* 1810), 459
Woodward, R. S., 487, 531
Woodward, W. H., 233
Wren, C. (*c.* 1670), 412, 394
Writing, advent of, 22
Wronski, H. (*c.* 1850), 531
Wu (*c.* 450), 144
Wu-king (Five Canons), 25
Wu-ti, emperor (*c.* 100 B.C.), 140
Wu-ti, a later emperor (*c.* 560), 150
Wu-ts'ao Suan-king (*c.* 1st century), 96, 141

Wu-ts'ao Suan-shu, 152
Wu Wang, emperor (*c.* 1122 B.C.), 22
Wylie, A., 31, 32, 272

Xenocrates (*c.* 350 B.C.), 92
Xylander, G. (*c.* 1570), 333, 133, 343

Yaḥyâ ibn Mohammed. *See* Ibn al-Lubûdî
Yang (male principle), 25
Yang Hui (*c.* 1261), 271
Ya'qûb ibn Ishâq. *See* Alchindi
Ya'qûb ibn Ṭârîq (*c.* 775), 167
Yaroslav, Duke, 268
Yâsimîn (*c.* 1200), 288
Yau, emperor (*c.* 2350 B.C.), 24
Yehuda ben Rakufial (*c.* 10th century), 208
Ye-lü Ch'u-ts'ai (*c.* 1230), 271
Ye-lü Hi-liang (*c.* 1260), 268
Yenri (circle principle), 535, 536, 538
Yi-ching. *See I-king*
Ying (female principle), 25
Yonge, C. D., 116
Yorozu, 144
Yoshida Shichibei Kōyū (*c.* 1640), 437, 353
Yu, emperor (*c.* 2200 B.C.), 29
Yüan Yüan (*c.* 1810), 535
Yuen Hao-wen (*c.* 1260), 271
Yuen-Shih, 271
Yûnis (Yûnos), 'Alî . . . ibn (*c.* 1000), 175, 191
Yûnis, Mûsâ ibn (*c.* 1200), 287
Yûsuf ibn Ibrâhîm (*c.* 875), 172

Zaccagnini, G., 539
Zaddik, Isaac (14th century), 241
Zeitschrift für Mathematik und Physik (Hl. Abt.), 105
Zeno of Elea (*c.* 450 B.C.), 78
Zenodorus (*c.* 180 B.C.), 118
Zero, 162, 207, 230, 270, 274, 277
Zeuthen, H. G. (*c.* 1900), 543, xv, 153
Ziegler, A., 260
Ziegler, T., 186
Zodiac, 23, 39, 100
Zoroaster, 71
Zosima (15th century), 262
Zuane. *See* Zuanne, Giovanni, John, Joannes, *etc.*
Zuane Mariani. *See* Mariani
Zuanne de Tonini da Coi (*c.* 1530), 295, 300
Zubler, L. (*c.* 1607), 361
Zumárraga, J. de, 354

HISTORY OF MATHEMATICS

VOLUME II

SPECIAL TOPICS OF
ELEMENTARY MATHEMATICS

BY

DAVID EUGENE SMITH

COPYRIGHT, 1925, BY DAVID EUGENE SMITH
ALL RIGHTS RESERVED

825.2

PREFACE

As stated in Volume I, this work has been written chiefly for the purpose of supplying teachers and students with a usable textbook on the history of elementary mathematics, that is, of mathematics through the first steps in the calculus. The subject has come to be recognized as an important one in the preparation of teachers and in the liberal education of students in colleges and high schools, showing, as it does, mathematics as constantly progressing instead of being a static mass of knowledge. Through a consideration of the history of the science the student comes to appreciate the fact that mathematics has continually adjusted itself to human needs, both material and intellectual; and thus he comes into sympathy with the effort to improve its status, either adding to its store through his own discoveries or bettering the methods of presenting it to those to whom it is taught in our schools.

In Volume I the reader found a general survey of the progress of elementary mathematics arranged by chronological periods with reference to racial and geographical conditions. In this volume he will find the subject treated by topics. The teacher of arithmetic will now see, in three or four chapters, a kind of moving picture of the growth of his subject,—how the world has counted, how it has performed the numerical operations, and what have been the leading lines of applications in which it has been interested. In geometry he will see how the subject arose, what intellectual needs established it so firmly, what influences led to its growth in various directions, and what human interest there is in certain of the great basal propositions. In algebra he will see, partly by means of facsimiles, how the symbolism has grown, how the equation looked three thousand years ago, the way its method of expression has changed from age to age, and how the science has so adjusted itself to world needs as now to be a necessity for the average citizen instead of a mental luxury for the selected

few. He will learn how the number concept has enlarged as new needs have manifested themselves, and how the world struggled with fractions and with the mysteries of such artificial forms as the negative and the imaginary number, and will thus have a still clearer vision of mathematics as a growing science. The terminology of the subject will arouse interest; the common units of measure will mean something more than mere names; the minutes and seconds of time and of angles will take on a kind of human aspect; and the calendar will cease to be the mystery that it is to the youth. Trigonometry will have a new interest to the teacher who reads what Plutarch tells of the shadow-reckoning of Thales, and of the independent origin of the trigonometry of the sphere; and the calculus, which the freshman or sophomore burns in accordance with time-honored tradition, will be seen to have a history that is both interesting and illuminating. To see in its genetic aspect the subject that one is teaching or studying, and to see how the race has developed it, is oftentimes to see how it should be presented to the constantly arriving new generations, and how it can be made to satisfy their intellectual hunger.

While the footnote is frequently condemned as being merely an apology for obscurity or as an exhibition of pedantry, it would be difficult, in a work of this kind, to dispense with its aid. There are two principal justifications for such a device: first, it enables an author to place the responsibility for a statement that may be open to question; and, second, it encourages many students to undertake further study, either from secondary sources or, what is more important, from the original writings of the men who rank among the creators of mathematics. With these two points in mind, footnotes have been introduced in such a way as to be used by readers who wish further aid, and to be neglected by those who wish merely a summary of historical facts. For the student who seeks an opportunity to study original sources a slight introduction has been made to this field. The text of the book contains almost no quotations in foreign languages, the result being that the reader will not meet with linguistic difficulties in the general narrative. In the notes, however, it is frequently desirable to quote the precise words of an author, and this has been done with reference to such European languages as are more or less familiar. It is not necessary to translate literally all these

extracts, since the text itself sets forth the general meaning. Students who have some general knowledge of Latin, French, or German will have little difficulty, and in many cases will have much interest, in seeing various statements in their original form. For special reasons a few notes have been given in Greek, but in every case the meaning is evident from the text.

In these footnotes and occasionally in the text there have, in this volume, been inserted a few names of minor importance which were purposely omitted in Volume I. These names refer to certain arithmeticians who contributed nothing to the advance of mathematics, but who, through popular textbooks, helped to establish the symbols and terms that are used in elementary instruction. In such cases all that has seemed necessary in the way of personal information is to give the approximate dates. In the case of names of particular importance further information may be found by referring to the Index.

The difficult question of the spelling and transliteration of proper names is always an annoying one for a writer of history. There is no precise rule that can be followed to the satisfaction of all readers. In general it may be said that in this work a man's name is given as he ordinarily spelled it, if this spelling has been ascertained. To this rule there is the exception that where a name has been definitely Anglicized, the English form has been adopted. For example, it would be mere pedantry to use, in a work in English, such forms as Platon and Strabon, although it is proper to speak of Antiphon and Bryson instead of Antipho and Bryso. When in doubt, as in the case of Heron, the preference has been given to the transliteration which most clearly represents the spelling used by the man himself.

In many cases this rule becomes a matter of compromise, and then the custom of a writer's modern compatriots is followed. An example is seen in the case of Leibniz. This spelling seems to be gaining ground in our language, and it has therefore been adopted instead of Leibnitz, even though the latter shows the English pronunciation better than the former. Leibniz himself wrote in Latin, and the spelled the name variously in the vernacular. There seems, then to be no better plan than to conform to the spelling of those German writers who appear to be setting the standard that is to be followed.

In connection with dates of events before the Christian era the letters B.C. are used; in connection with dates after the beginning of this era no distinguishing letters are added except in a few cases near the beginning of the period, in which the conventional letters A.D. have occasionally been inserted to avoid ambiguity. With some hesitation, but for a purpose which seems valid, dates are frequently given in parentheses after proper names. It is well recognized that a precise date, like 1202 after the name Fibonacci, is of no particular value in itself. It makes no difference, in ordinary cases, whether Fibonacci wrote his *Liber Abaci* in 1202, or in 1180, or in 1220, or whether *abacus* is spelled *abbacus*, as in some manuscripts, or in the more correct Latin form. On the other hand, two things are accomplished by a free use of such dates. In the first place, a reader is furnished with a convenient measuring instrument; he does not have to look in the Index or in a chronological table in order to see approximately where the particular writer belongs in the world's progress. The casual reader may well be pardoned if he does not recall where Bede, Alcuin, Gerbert, Jordanus, Fibonacci, and Roger Bacon stood chronologically with respect to one another, and in reading a technical history of this kind there is no reason why he should not be relieved of the trouble of consulting an index whenever he meets with such names as these. In the second place, it needs no psychologist to confirm the familiar principle that the mind comes, without conscious effort, to associate in memory those things which the eye has frequently associated in reading. At the risk, therefore, of disturbing the minds of those who are chiefly interested in the literary aspect of a general statement of the progress of mathematics, many important dates have been repeated, especially where they have not appeared in the pages immediately preceding.

The extent of a bibliography in a popular work of this kind is a matter of judgment. It can easily run to great length if the writer is a bibliophile, or it may receive but little attention. The purpose of lists of books for further study is that the student may have access to information which the author has himself used and which he believes will be of service to the reader. For this reason the sources mentioned in this work are such as may be available, in many cases are sure to be so, in the libraries connected with

our universities, while the original sources are those which are of importance in the development of elementary mathematics or which may be of assistance in showing certain tendencies.

The first time a book was mentioned in Volume I, the title, date, and place of publication were given, together, whenever it seemed necessary, with the abbreviated title thereafter used. In general this plan has been followed in Volume II, at least in the case of important works. To find the complete title at any time, the reader has only to turn to the Index to find the first reference to the book. The abbreviation *loc. cit.* (for *loco citato*, in the place cited) is used only where the work has been cited a little distance back, since any more general use of the term would be confusing. The symbolism "I, 7" has been used for "Vol. I, p. 7," in order to conserve space, although exceptions have been made in certain ambiguous cases, as in the references to Heath's *Euclid*, references to Euclid being commonly by book and proposition, as in the case of Euclid, I, 47.

The standard works are referred to as given on pages xiv–xvi of Volume I.

In the selection of illustrations the general plan followed has been to include only such as will be helpful to the reader or likely to stimulate his interest; it would be undesirable to attempt to give, even if this were possible, illustrations from all the important sources, for this would tend to weary him. On the other hand, where the reader has no access to a classic that is being described, or even to a work which is mentioned as having contributed to the world's progress in some humbler manner, a page in facsimile is often of value. It is evident that space does not permit of the use of such bibliographical illustrations as those which comprise a large part of the facsimiles in the author's *Rara Arithmetica*.

In general the illustrations have been made from the original books or manuscripts in the well-known and extensive library of George A. Plimpton, Esq., who has been very generous in allowing this material to be used for this purpose, or from the author's collection of books, manuscripts, mathematical portraits and medals, and early mathematical instruments.

The scheme of transliteration and pronunciation of proper names is set forth fully on pages xvii–xxii of Volume I. Since Arabic, Persian,

Hindu, Chinese, and Japanese names are used less frequently in this volume, it will sufficiently meet the needs of the reader if he refers to the scheme there given.

As in Volume I, a few topics for discussion or for the personal consideration of students are suggested at the close of each chapter. Specific questions have been avoided, the purpose being not so much to examine the reader on the facts set forth as to encourage him to pursue his reading in other works upon the subject. In most cases this reading will be done in such encyclopedias as may be available, and, preferably, in other histories also; but in any case the reader will have his attention called to a number of general lines for further study, and he will have the consciousness that the present work is merely an introduction to the general subject, in which, it is hoped, his interest has increased.

On account of the extent of the index to Volume I it has not been combined with that of Volume II. It should therefore be consulted in connection with the index to this volume, particularly with respect to biographical and bibliographical references. Since, in many cases, textbooks are mentioned so frequently as to render a complete list so long as to be burdensome to the reader, thus defeating its purpose, such works are included only when the author is not mentioned in Volume I and when the work is of such importance as to make the reference valuable.

The author wishes to express his appreciation of the aid rendered by various friends in reading the proofsheets of both volumes, and especially by the late Herr Gustaf Eneström of Stockholm, by Professor R. C. Archibald, by Mr. Jekuthiel Ginsburg of New York, and by Captain E. L. Morss of Boston.

DAVID EUGENE SMITH

CONTENTS

CHAPTER PAGE

I. DEVELOPMENT OF THE ARITHMETICA

1. General Survey 1
2. The Seven Liberal Arts 2
3. Early Writers on Number Theory 4
4. Names for Arithmetic 7
5. Elementary Classifications of Number . . . 11
6. Unity 26
7. Later Developments 29

Topics for Discussion 31

II. LOGISTIC OF NATURAL NUMBERS

1. Fundamental Operations 32
2. Reading and Writing Numbers 36
3. Addition 88
4. Subtraction 94
5. Multiplication 101
6. Division 128
7. Roots 144
8. Checks on Operations 151

Topics for Discussion 155

III. MECHANICAL AIDS TO CALCULATION

1. The Abacus 156
2. Finger Reckoning 196
3. Modern Calculating Machines 202

Topics for Discussion . . . . . 2[illegible]

CHAPTER PAGE

IV. ARTIFICIAL NUMBERS

1. Common Fractions 208
2. Sexagesimal Fractions 228
3. Decimal Fractions 235
4. Surd Numbers 251
5. Negative Numbers 257
6. Complex Numbers 261
7. Transcendental Numbers 268

Topics for Discussion 269

V. GEOMETRY

1. General Progress of Elementary Geometry . . 270
2. Name for Geometry 273
3. Technical Terms of Euclidean Geometry . . 274
4. Axioms and Postulates 280
5. Typical Propositions of Plane Geometry . . . 284
6. Typical Propositions of Solid Geometry . . . 291
7. The Three Famous Problems 297
8. Analytic Geometry 316
9. Modern Geometry 331
10. Perspective and Optics 338
11. Instruments in Geometry 344
12. The Problem of Earth Measure 368

Topics for Discussion 377

VI. ALGEBRA

1. General Progress of Algebra 378
2. Name for Algebra 386
3. Technical Terms 393
4. Symbols of Algebra 395
5. Fundamental Operations 416

CHAPTER PAGE

6. CONTINUED FRACTIONS 418
7. THE WRITING OF EQUATIONS 421
8. THE SOLUTION OF EQUATIONS 435
9. DETERMINANTS 475
10. RATIO, PROPORTION, AND THE RULE OF THREE . . 477
11. SERIES 494
12. LOGARITHMS 513
13. PERMUTATIONS, COMBINATIONS, PROBABILITY . . . 524

TOPICS FOR DISCUSSION 531

VII. ELEMENTARY PROBLEMS

1. MATHEMATICAL RECREATIONS 532
2. TYPICAL PROBLEMS 536
3. COMMERCIAL PROBLEMS 552
4. APPLICATIONS OF ALGEBRA 582
5. MAGIC SQUARES 591

TOPICS FOR DISCUSSION 599

VIII. TRIGONOMETRY

1. GENERAL DEVELOPMENT OF TRIGONOMETRY . . . 600
2. TRIGONOMETRIC FUNCTIONS 614
3. TRIGONOMETRIC TABLES 623
4. TYPICAL THEOREMS 628

TOPICS FOR DISCUSSION 633

IX. MEASURES

1. WEIGHT 634
2. LENGTH 640
3. AREAS 644
4. CAPACITY 644

CHAPTER PAGE

5. VALUE 645
6. METRIC SYSTEM 648
7. TIME 651

TOPICS FOR DISCUSSION 675

X. THE CALCULUS

1. GREEK IDEAS OF A CALCULUS 676
2. MEDIEVAL IDEAS OF THE CALCULUS 684
3. MODERN FORERUNNERS OF THE CALCULUS . . . 685
4. NEWTON AND LEIBNIZ 692
5. JAPAN 701

TOPICS FOR DISCUSSION 703

INDEX 705

HISTORY OF MATHEMATICS

SPECIAL TOPICS OF ELEMENTARY MATHEMATICS

CHAPTER I

DEVELOPMENT OF THE ARITHMETICA

1. General Survey

Nature of Arithmetica. As stated in the Preface, it is the purpose of this volume to set forth in considerable detail the important steps in the historical development of the several branches of elementary mathematics. One of these branches is now known as arithmetic, a name which, as commonly understood in the English-speaking world, has little or no relation to the arithmetic of the ancients. In recent times the word has acquired the meaning given by the Greeks and Romans to logistic, or the art of computation, a much more humble discipline than that which they called arithmetic.

In order to make the distinction clear, the present chapter will set forth a sufficient number of simple details of the ancient arithmetic to enable the student to form an idea of its general nature, and the second chapter will consider the development of that elementary art which now bears the ancient name. It will be seen that the science which formerly appropriated the title was not related to ordinary calculation but was a philosophical study dealing with such properties as might now find place in a course in the theory of numbers if the latter had not outgrown most of these simple number relations and become a subject for the university student.

Modern Theory. The modern theory of numbers has so little direct relation to elementary mathematics that its history need only be referred to briefly in this volume.[1] Certain features like prime and composite numbers, polygonal numbers (such as squares), and solid numbers (such as cubes) are still found in elementary mathematics, however, and these features render essential a brief statement concerning the ancient arithmetic. In order to explain the position of this science in the ancient scheme of learning, it is desirable to speak first of the general range of knowledge according to the Greek schools of philosophy, and to distinguish between arithmetica, the classical theory of numbers, and arithmetic, the modern art of computation.

2. The Seven Liberal Arts

The Sevenfold Division. As stated in Volume I, three and seven have been the chief among mystic numbers in all times and among all peoples. Many reasons have been assigned for this universal habit of the race, most of them manifestly fanciful, and possibly no reason can be adduced that will command the general approval of scholars. If, however, we omit the number five, which was often used as a primitive radix and thus lost its element of mystery, a fairly satisfactory explanation is found in the fact that three and seven are the first prime numbers,—odd, unfactorable, unconnected with any common radix, possessed of various peculiar properties, and thus of a nature to attract attention in the period of superstition and mysticism.

One of the many results of this veneration for these numbers is seen in the fact that the ancients numbered seven great branches of learning, just as they numbered the Seven Wonders of the World and the Seven Wise Men of Greece. They separated these branches into two groups, four studies making up the domain of science as recognized by the Pythagoreans, and three constituting the nonscientific domain. Plato[2] spoke of

[1] The student will find it elaborately treated in L. E. Dickson, *History of the Theory of Numbers*, 3 vols., Washington, 1919–1923; hereafter referred to as Dickson, *Hist. Th. Numb.*

[2] *Republic*, IX. See also Aristotle's *Politics*, VIII, 1.

the liberal arts and separated them into two groups, but he did not limit them to any definite number. The scientific group, consisting of arithmetic, geometry, spherics, and music, constituted the ancient domain of mathematics.

The Seven Liberal Arts. It was probably in the work of Capella (*c.* 460), that the seven liberal arts were first distinctly specified.[1] These seven arts were thenceforth looked upon as necessary to the education of free men (*liberi*). They were then separated into the quadrivium,[2] constituting the Pythagorean group, and the trivium,[3] made up of grammar, dialectics, and rhetoric.[4]

The names of the seven arts are fairly descriptive of the subjects represented, with the exception of spherics, which related to mathematical astronomy; music, which related only to the theory of harmony;[5] and arithmetic, which had little in common with the subject known in English by this name.

[1] Varro (1st century B.C.) wrote a treatise on the "nine liberal disciplines," but the work is not extant. Capella introduced the liberal arts as the bridesmaids at the marriage of Philology and Mercury. Cassiodorus (*c.* 470–*c.* 564) placed the limit definitely at seven because of the seven pillars in the Temple of Wisdom (Proverbs, ix, 1).

[2] In medieval Latin also written *quadruvium*, the *quadruplex via*, as some writers have it. The term in its literal meaning is found as early as Juvenal. In its technical educational meaning it is used by Cassiodorus.

[3] Also written *truvium*.

[4] As "Hugnitio natione tuscus, civis pisanus, episcopus ferrariensis," to quote a medieval record, has it: "Et uero quia gramatica dialecta rethorica dicuntur triuuium quadam similitudine quasi triplex uia ad idem idest ad eloquentiam arismethica. musica. geometria. astronomia. quadam simili similitudine dicuntur quadriuuium quasi quadruplex uia ad idem idest ad sapientiam." See also the well-known verse quoted in Volume I, page 180.

[5] As an old Latin MS. has it:

> Musicorum et cantorum magna est distantia:
> Isti dicunt illi sciunt quae componit musica.

The distinction is well set forth in B. Veratti, *De' Matematici Italiani anteriori all' invenzione della stampa*, p. 4 (Modena, 1860). See also P. Tannery, "Du rôle de la musique grecque dans le développement de la mathématique pure," *Bibl. Math.*, III (3), 161; E. Narducci, "Di un codice . . . dell' opera di Giorgio Pachimere: *περὶ τῶν τεσσάρων μαθημάτων*," *Rendiconti della R. Accad. dei Lincei*, Rome, VII (1891), 191.

3. Early Writers on Number Theory

Origin of the Theory. There is no definite trace of the study of the theory of numbers before the time of Thales (*c.* 600 B.C.). Tradition says that this philosopher, filled with the lore of the Egyptians and probably well informed concerning the mysticism of the Babylonians, taught certain of the elementary properties of numbers in the Ionic School, of which he was the founder. Such meager knowledge as he had he imparted to his brilliant disciple, Pythagoras (*c.* 540 B.C.), who thereupon resorted to the priests of Egypt and probably of Babylon for further light. In the school which he established at Crotona, in southern Italy (Magna Græcia), he elaborated the doctrines of his teachers, including ideas which are distinctly Oriental, and made the first noteworthy beginning in the theory of arithmetica.

Little by little, first among the Pythagoreans and then in other schools of philosophy, the subject grew, a little being added here and a little there, until the time finally became ripe for the appearance of treatises in which the accumulated knowledge could be systematically arranged.

Books on the Theory. The first successful effort in the preparation of an expository treatise on the subject was made by Euclid (*c.* 300 B.C.), who is often known only as a geometer but who showed great genius in systematizing mathematical knowledge in other important lines as well. In his *Elements* he devotes Books II, V, VII, VIII, IX, X (in whole or in part) to the theory of numbers or to geometric propositions closely related thereto, and includes such propositions as the following:

If four numbers are proportional, they are also proportional alternately (VII, 13).

If two numbers are prime to two numbers, both to each, their products also will be prime to one another (VII, 26).

If a square number does not measure a square number, neither will the side measure the side; and if the side does not measure the side, neither will the square measure the square (VIII, 16).

If an odd number measures an even number, it will also measure the half of it (IX, 30).

The next worker in this field was that interesting dilettante in matters mathematical, Eratosthenes (*c.* 230 B.C.), who worked on a method of finding prime numbers[1] by sifting out the composite numbers in the natural series, leaving only primes. This he did by canceling the even numbers except 2, every third odd number after 3, every fifth odd number after 5, and so on, the result being what the ancient writers called the sieve.[2]

His friend and sometime companion Archimedes (*c.* 225 B.C.) did little with the theory of arithmetica, but made an effort to improve upon the Greek system of numbers,[3] his plan involving the counting by octads (10^8), in which he proceeded as far as 10^{52}, and making use of a law which would now be expressed by such a symbolism as $a^m a^n = a^{m+n}$, although he made no specific mention of this important theorem.

It was to the commentary on the *Timæus* of Plato, written by Poseidonius (*c.* 77 B.C.), that the Greeks invariably went for their knowledge of the number theories of the Pythagoreans.[4] This is seen in the fact that the phraseology used by such writers as Theon of Smyrna (*c.* 125) and Anatolius (*c.* 280), in speaking of this subject, is simply a paraphrase of that used by Poseidonius.

Nicomachus, and Theon of Smyrna. The first noteworthy textbook devoted to arithmetica was written by Nicomachus (*c.* 100), a Greek resident of Gerasa (probably the modern Jerash, a town situated about fifty-six miles northeast of Jerusalem). He was not an original mathematician, but he did for the theory of numbers what Euclid had done for elementary geometry and Apollonius (*c.* 225 B.C.) for conic sections,—he summarized the accumulated knowledge in his subject. In his work are found such statements as the following: "Now fur-

[1] Vol. I, p. 109. [2] Κόσκινον (*kos'kinon*), Latin, *cribrum*.

[3] Vol. I, p. 113. See also his Ψαμμίτης (*psammi'tes*, Latin *arenarius*, "sand reckoner"), *Archimedis Opera Omnia*, ed. Heiberg (Leipzig, 1880–1915), with revisions.

[4] F. E. Robbins, "Posidonius and the Sources of Pythagorean Arithmology," *Classical Philology*, XV (1920), 309. On Plato's appreciation of the value of this kind of work see F. Cajori, "Greek Philosophers on the Disciplinary Value of Mathematics," *The Mathematics Teacher* (December, 1920), p. 57.

thermore every square upon receiving its own side becomes heteromecic; or, by Zeus, on being deprived of its own side."[1]

The next writer of note was Theon of Smyrna (*c.* 125). He added several new propositions to the theory, two of them being of special interest: (1) If n be any number, n^2 or $n^2 - 1$ is divisible by 3, by 4, or by both 3 and 4; and if n^2 is divisible by 3 and not by 4, then $n^2 - 1$ is divisible by 4. (2) If we arrange two groups of numbers as follows:

$$\begin{aligned} n_1 &= 1 + 0 \\ n_2 &= 1 + 1 \\ n_3 &= 2 + 3 \\ n_4 &= 5 + 7 \\ &\vdots \\ n_r &= n_{r-1} + d_{r-1} \end{aligned} \qquad \begin{aligned} d_1 &= 1 + 0 = 1 \\ d_2 &= 2 + 1 = 3 \\ d_3 &= 4 + 3 = 7 \\ d_4 &= 10 + 7 = 17 \\ &\vdots \\ d_r &= 2\,n_{r-1} + d_{r-1} \end{aligned}$$

then d^2 is of the form $2n^2 \pm 1$; for example, $d_1^2 = 1 = 2n_1^2 - 1$, $d_2^2 = 9 = 2n_2^2 + 1, \cdots$. The numbers d_r were called by Theon diameters. It is interesting to observe a fact unknown to him, namely, that the ratios $d_1 : n_1 = 1$, $d_2 : n_2 = \frac{3}{2}$, $d_3 : n_3 = \frac{7}{5}, \cdots$ are the successive convergents of the continued fraction

$$1 + \cfrac{1}{2 + \cfrac{1}{2 + \cfrac{1}{2 + \cdots}}},$$

and hence approach nearer and nearer the square root of 2.

Boethius. Boethius (*c.* 510) appropriated the knowledge of such writers as Euclid, Nicomachus, and Theon, incorporating it in his work *De institutione arithmetica libri duo* and producing a textbook that was used in all the important schools in the Middle Ages. It is the source with which a student may advantageously begin his study of this subject.

Later Writers. The most noteworthy writer on the subject in the medieval period is Fibonacci (1202), and with respect to him and subsequent writers, all of whom have been considered in Volume I, we shall later speak in detail as necessity arises.

[1] Introduction, XX. See G. Johnson, *The Arithmetical Philosophy of Nicomachus of Gerasa*, Lancaster, 1916, hereafter referred to as Johnson, *Nicomachus*. The meaning is that $x^2 \pm x$ is not a square but a heteromecic or oblong number.

4. Names for Arithmetic

Arithmetic and Logistic. The ancient Greeks distinguished between arithmetic,[1] which was the theory of numbers and was therefore even more abstract than geometry,[2] and logistic,[3] which was the art of calculating. These two branches of the study of numbers continued as generally separate subjects until the time of printing, although often with variations in their names; but about the beginning of the 16th century the more aristocratic name of "arithmetic" came to be applied to both disciplines. This use of the term was not universal, however, and even today the Germans reserve the word *Arithmetik* for the theoretical part of the science as seen in the operations in algebra, using the word *Rechnung* for the ancient logistic.[4] Various writers,[5] preserved the word "logistic" in the 16th century, but in the older sense it generally dropped out of use thereafter.

From the fact that computations were commonly performed on the abacus, the name of this instrument was used in the early Middle Ages as a synonym of logistic. Finally, however, the word "abacus" came to mean any kind of elementary arithmetic,[6] and this usage obtained long after printing was invented.[7]

In the Middle Ages the name "arithmetic" was apparently not in full favor, perhaps because it was not of Latin origin. Thus, in a manuscript attributed to Gerbert the word is spoken of as Greek, the Latin being "numerorum scientia."[8]

[1] Ἀριθμητική (*arithmetikē*), from ἀριθμός (*arithmos*), number. It passed over into Latin as *arithmetica*.

[2] "Est enim Arithmetices subjectum purius quiddam & magis abstractum, quam subjectum Geometriae" (J. Wallis, *Opera Mathematica*, I, 18 (Oxford, 1695)).

[3] Λογιστική (*logistikē*), which passed over into Latin as *logistica*.

[4] Compare also the French *calcul*.

[5] J. Noviomagus, *De Numeris libri duo* (1539); Buteo (Lyons, 1559); Schoner edition of Ramus (1586); "Logistica quam uulgo uocant algoristicam et algorismum" (MS. notes in the 1558 edition of Gemma Frisius, in Mr. Plimpton's library). For biographical information relating to such writers as are of particular importance, see the Index of Volume I.

[6] As in Fibonacci's *Liber Abaci* (1202).

[7] See *Rara Arithmetica* for many works bearing such titles as *Libro d' abacho*.

[8] "Graece Arithmetica, latine dicitur numerorum scientia," from the colophon of the "Liber subtilissimus de arithmetica." See C. F. Hock, *Gerberto o sia Silvestro II Papa . . . trad. del . . . Stelzi*, p. 206 (Milan, 1846).

Vicissitudes of the Term. The word "arithmetic," like most other words, has undergone many vicissitudes. In the Middle Ages, through a mistaken idea of its etymology, it took an extra *r*, as if it had to do with "metric."[1] So we find Plato of Tivoli, in his translation (1116) of Abraham Savasorda, speaking of "Boetius in arismetricis."[2] The title of the work of Johannes Hispalensis, a few years later (*c.* 1140), is given as "Arismetrica," and fifty years later than this we find Fibonacci dropping the initial and using the form "Rismetrica."[3] The extra *r* is generally found in the Italian literature until the time of printing.[4] From Italy it passed over to Germany, where it is not uncommonly found in the books of the 16th century,[5] and to France, where it is found less frequently.[6] The ordinary variations in spelling have less significance, merely illustrating, as is the case with many other mathematical terms, the vagaries of pronunciation in the uncritical periods of the world's literatures.[7]

[1] Greek μέτρον, a measure, as in "metre" and "metrology."

[2] *Abhandlungen*, XII, 16. For such abridged forms see the Index of Volume I.

[3] This is in one of the MSS. formerly owned by Boncompagni. See the sale catalogue of his library, p. 104. Fibonacci (1202) commonly used "abacus."

[4] *E.g.*, see the "Brani degli Annali Decemvirali posseduti dall' archivo Decemvirale di Perugia," in Boncompagni's *Bullettino*, XII, 432; E. Narducci, *Catalogo di Manoscritti*, 2d ed. (Rome, 1892), No. 56, p. 26; hereafter referred to as Narducci, *Catalogo Manosc.*

[5] *E.g.*, "Die Kunst Arismetrica die aller edelst vnder den sybē freyen künsten," Köbel, 1514. A MS. in Scheubel's (*c.* 1550) handwriting in the Columbia University Library has "de Arrismetris." There is also a MS. copy made *c.* 1515 in Rome, by a Swedish savant, Peder Månsson, from the *Margarita phylosophica* of Gregorius Reisch (1503), in which the form "Arismetrice" is given. See *Bibl. Math.*, II (2), 17.

[6] So in a MS. written by Rollandus *c.* 1424 (see *Rara Arithmetica*, p. 446) the form "arismetica" is usually given, but the form "arismetrica" also appears. In an unpublished MS. entitled "traicte d'Arismetricque . . . faite et compillé A paris en lan mil 475" (for 1475) there is this curious etymology: "Arismeticve est vne des sept ars liberaulx & la premiere des quatre ars Mathematique En la quelle est la vertus de nombrer. Et est dicte de ares Nom grec qui est en latin Virtus Et de menos aussi nom grec qui est en Latin numerus parquoy est dicte Vertus de Nombre." E. Narducci, *Catalogo Manosc.*, No. 603, p. 395.

[7] Thus, we have "arimmetica" throughout Zuchetta's work of 1600 (see *Rara Arithmetica*, p. 425); "eritmeticha" in a 17th century MS. (see Narducci's *Catalogo Manosc.*, No. 446, 1, p. 267); "aristmeticque" in an anonymous French work, Paris, 1540; "Alchorismi de pratica Aricmetica," in a MS. of Sacrobosco, Boncompagni sale catalogue, No. 645.

Origin of Algorism. From the fact that the arithmetic of al-Khowârizmî (*c.* 825) was translated into Latin as *liber Algorismi* (*the book of al-Khowârizmî*), arithmetic based on the Hindu-Arabic numerals, more especially those that made use of the zero, came to be called algorism as distinct from the theoretical work with numbers which was still called arithmetic.[1] Since *al* often changes to *au* in French, we have "augrisme" and "augrime,"—forms which were carried over to England as "augrim,"[2] later reverting to "algorism"[3] or the less satisfactory form of "algorithm."[4]

The prefix *al* was dropped from this word by most Spanish writers, giving such forms as "guarisma"[5] and "guarismo,"[6] and in other countries there were many variations that were quite as curious.[7]

The word troubled many of the early Latin writers, and various fanciful etymologies were suggested, the best conjecture being that of Sacrobosco that it came from Algus or Argus,

[1] Thus, that part of the Rollandus MS. (*c.* 1424) relating to the theory is referred to in the phrase "Arismetrice pars primo tractanda est speculatiua," while the other part is called "algorismus." See also M. Chasles, *Comptes rendus*, XVI, 162.

[2] "¶ ouer the wiche degrees ther ben nowmbres of augrym; . . . & the nombres of the degres of tho signes ben writen in Augrim." Chaucer's *Astrolabe*, ed. Skeat, p. 5.

"Although a sypher in augrim have no might in significacion of it-selve, yet he yeveth power in significacion to other." Chaucer, *The Testament of Love*, ed. Skeat, Bk. II, chap. vii.

[3] Thus, Recorde (*c.* 1542) in his *Grovnd of Arts* (as spelled in the 1646 edition): "Some call it Arsemetrick, and some Augrime. . . . Both names are corruptly written: Arsemetrick for Arithmetick, as the Greeks call it, and Augrime for Algorisme, as the Arabians found it." 1646 ed., p. 8.

[4] One eccentric English writer, Daniel Fenning (1750), attempted to distinguish algorithm, as first principles, from algorism, as the practice of these principles.

[5] As in the Spanish *Suma de Arithmetica* of Gaspard de Texeda, Valladolid, 1546. The separate word *al* or *el* (the) was prefixed, however, and the form *algoritmo* is still preserved.

[6] ". . . de vn Filosofo llamado Algo, y por aquesta causa fue llamada el Guarismo" (Santa-Cruz, a Spanish writer, 1594); but see *Rara Arithmetica*, p. 407.

[7] "Arismethique qui vulgayrement est appellee argorisme" (E. de la Roche, a French writer, 1520). We also find such forms as *alkauresmus* and *alchocharithmus* in various MSS. of the same period.

a certain philosopher, this being merely a corruption of al-Khowârizmî.[1] It was not until 1849 that the true etymology was again discovered.[2]

The Etymology early Recognized and Forgotten. Very likely the etymology of the term "algorism" was known to such early translators or writers as Johannes Hispalensis[3] (*c.* 1140) and Adelard of Bath[4] (*c.* 1120). By the following century, however, al-Khowârizmî was quite forgotten by such Latin writers as Sacrobosco[5] (*c.* 1250) and Bacon (*c.* 1250). From that time on we have the word loosely used to represent any work related to computation by modern numerals[6] and also as synonymous

[1] So we have Chaucer's expression,

> Thogh Argus the noble covnter
> Sete to rekene in hys counter.
>
> *Dethe Blaunche* (*c.* 1369)

This derivation was followed by various writers, such as Santa-Cruz (1594), Cataldi (1602), and Tartaglia (1556 ed., I, fol. 3, *r.*). Of the other fanciful etymologies the following may be of interest: *argris* (Greek) + *mos* (custom); *algos* (Greek for "white sand") + *ritmos* (calculation); *algos* (art) + *rado* (number); *Algorus*, the name of a Hindu scholar; *Algor*, a king of Castile. See A. Favaro, Boncompagni's *Bullettino*, XII, 115; M. Cantor, *Mathematische Beiträge zum Kulturleben der Völker*, p. 267 (Halle, 1863); C. I. Gerhardt, *Ueber die Entstehung . . . des dekadischen Zahlensystems*, Prog., p. 26, n. (Salzwedel, 1853); K. Hunrath, "Zum Verständniss des Wortes Algorismus," *Bibl. Math.*, I (2), 70; and see VIII (2), 74. P. Ramus (*Scholarum Mathematicarum Libri XXXI*, p. 112 (1569)) derived it from *al* (Arabic for "the") + ἀριθμός (*arithmos'*, number), and J. Schoner (1534 edition of the *Algorithmvs Demonstratvs*, fol. A [iij], *v.*) did the same.

[2] By the orientalist J. Reinaud (1795–1867). See *Mém. de l'Institut national de France des inscriptions et belles-lettres*, XVIII, 303; Boncompagni's *Bullettino*, XII, 116. Even as late as 1861, however, L. N. Bescherelle's well-known French dictionary (Paris, 1861) gave *al* (the) + *ghor* (parchment), and the variants *algarthme, algarisme*. See also Boncompagni's *Bullettino*, XIII, 557.

[3] "Incipit prologus in libro alghoarismi de pratica arismetrice. Qui editus est a magistro Johanne yspalensi." See F. Woepcke, *Journal Asiatique*, I (6), 519.

[4] Who uses such forms as *algoritmi* and *algorizmi*.

[5] "Hanc igitur scientiam numerandi compendiosam edidit philosophus nomine Algus, unde algorismus nuncupatur, vel ars numerandi, vel introductio in numerum." Halliwell ed., p. 1.

[6] "Ceste signifiance est appellée algorisme" (MS. of *c.* 1275); see C. Henry, Boncompagni's *Bullettino*, XV, 53. "Secondo Lalgorismo" (Ghaligai, 1521); ". . . calculandi artem, quam uulgus Algorithmum uocat" (Schoner, 1534). So the MS. of Scheubel (*c.* 1550), already mentioned as in the Columbia Uni-

with the fundamental operations themselves[1] and even with that form of arithmetic which makes use of the abacus.[2]

Names for Logistic. There have been various other names for logistic. The early Italian writers often spoke of a practical arithmetic as a *practica*,[3] *pratica*, or *pratiche*.[4] Many of the Latin writers of the Renaissance, particularly in the 16th century, spoke of it as the art of computing (*ars supputandi*).[5] The Dutch writers used the term "ciphering,"[6] particularly in the 16th and 17th centuries, and from this source, through New Amsterdam, came the common use of the word in the early schools of America.

In Italy, in the 15th century, logistic occasionally went by the name of the minor art,[7] and arithmetic and algebra by the name of the major art.[8]

5. Elementary Classifications of Number

Abstract and Concrete. The distinction between abstract and concrete numbers is modern. The Greek arithmeticians were concerned only with the former, while the writers on logistic naturally paid no attention to such fine distinctions. It was not

versity Library, has such phrases as "Algebrae fundamenta seu algorismus," "Algorismus de surdis," and "Algorithmus quantitatis," showing the broader use of the term. Stifel (1544) used the term in the same way.

[1] Thus, Thierfelder (1587) uses "Der Algorithmus" and "Die Species" (p. 51) as synonymous. Similarly, "ALgorithmus ist ein lehr aus der man lernet Addiren/Subtrahiren/Multipliciren vnd Diuidiren" (Stifel, 1545).

[2] As in the *Algoritmus* of Klos (1538), the first Polish arithmetic, which is purely a treatise on abacus reckoning. See S. Dickstein, *Bibl. Math.*, IV (2), 57. Similarly, there were several books entitled *Algorithmus linealis* published in Germany early in the 16th century, all dealing with the abacus.

[3] As in the Treviso arithmetic (1478).

[4] As in Cataneo's arithmetic (1546).

[5] Thus, Tonstall (1522) calls his work *De arte supputandi*, a title already used by Clichtoveus (1503) in the abridged form of *Ars supputādi*. Glareanus (1538) speaks of the "supputandi ars," and "supputation" (for computation) was a term in common use in England until the 19th century. For example, see W. Butler, *Arithmetical Questions*, London, 2d ed., 1795.

[6] Cyffering, cyffer-konst, cyffer-boeck, and the like.

[7] *L'arte minore.* [8] *L'arte maggiore*; or, in Latin, *Ars magna.*

until the two streams of ancient number joined to form our modern elementary arithmetic that it was thought worth while to make this classification, and then only in the elementary school.

The terms "abstract" and "concrete" were slow in establishing themselves. The mathematicians did not need them, and the elementary teachers had not enough authority to standardize them. In the 16th century the textbook writers began to make the distinction between pure number and number to which some denomination attached, and so we find Trenchant (1566), for example, speaking of absolute and denominate number, the latter including not only 3 feet but also 3 fourths.[1]

From that time on the distinction is found with increasing frequency in elementary works. Such refinements, however, as required the product to be of the same denomination as the multiplicand are, in general, 19th century creations of the schools. Thus Hodder[2] asserts that "Pounds multiplied by 20, are shillings," and every scientist today recognizes such forms as "20 lb. × 10 ft. = 200 foot-pounds."

Digits, Articles, and Composites. One of the oldest classifications of numbers is based upon finger symbolism.[3] The late Roman writers seem to have divided the numbers below a hundred into fingers (*digiti*), joints (*articuli*), and composites

[1] "L'absolu est celuy qui n'a aucune denomination: comme 2, 7, 5, tel nombre est abstrét, & de forme nue se referant à la Theorique. Le denommé: est celuy qui si prononce auec quelque denomination . . . & se refére à la Pratique." The latter included "le vulgairement denommé, comme 8 auñ," and also "le rompu, comme $\frac{3}{4}$," although he says that in practice $\frac{3}{4}$ is considered as abstract unless some denomination is given to it: "lequel en pratiquant est entendu absolu s'il n'a quelque denomination de suget, comme disant $\frac{3}{4}$ d'aun" (1578 ed., p. 16).

Similarly, Stifel: "Numeri abstracti proprie dicuntur, q̄ nullā prorsus denominationē habēt" (*Arithmetica Integra*, 1544, fol. 7, *v.*). Xylander (1577) used *ledige* and *benannte Zahlen*.

[2] 1672 ed., p. 56.

[3] See page 196. Th. Martin, "Les signes numéraux," *Annali di mat. pura ed applic.*, V, 257, 337, and reprint (Rome, 1864); hereafter referred to as Martin, *Les Signes Num.* Suevus (*Arithmetica Historica*, 1593, p. 3) speaks of the finger origin: "Digitus heist ein Finger zal/die unter zehen bedeut"; M. Wilkens, a Dutch arithmetician (*Arithmetica*, Groningen, 1630; 1669 ed., p. 1), says: "Dese zijn Digiti, dat's Enckel ofte vingergetalen"; and many other early writers have similar statements.

(*compositi*) of fingers and joints, the joints being the tens, and the composites being numbers like 15, 27, and so on. In a passage attributed, but doubtfully, to Boethius it is said that this threefold division is due to the ancients.[1] While the terms were probably known in early times, they were not used commonly enough to appear in the places where finger symbolism is mentioned.[2] So far as extant works are concerned, the classification is medieval.

Meaning of "Digit." Since there are ten fingers, it is probable that the digits were originally the numbers from one to ten inclusive; but so far as appears from treatises now extant they were the numbers from one to nine inclusive, not the figures representing these numbers; that is, they were the numbers below the first "limit." The division of numbers into limits or differences (in which 10, 20, ···, 90 were of the first order; 100, 200, ···, 900, of the second order, and so on) is found in the works of such writers as Alcuin (*c.* 780), Jordanus Nemorarius (*c.* 1225), O'Creat (*c.* 1150), and Sacrobosco (*c.* 1250), and was evidently common.[3] Since unity was not considered a number until modern times, it was sometimes definitely omitted, leaving only eight digits.[4]

[1] Since this is the first time the division appears, so far as known, the passage is important enough to be quoted in the original: "Digitos vero, quoscunque infra primum limitem, id est omnes, quos ab unitate usque ad denariam summam numeramus, veteres appellare consueverunt. Articuli autem omnes a deceno in ordine positi et in infinitum progressi nuncupantur. Compositi quippe numeri sunt omnes a primo limite id est a decem usque ad secundum limitem id est viginti ceterique sese in ordine sequentes exceptis limitibus. Incompositi autem sunt digiti omnes annumeratis etiam omnibus limitibus." Boethius, ed. Friedlein, p. 395. See also G. Enestrom, *Bibl. Math.*, XI (2), 116.

[2] Pliny, *Hist. Nat.*, 34, 7; 2, 23; Martin, *Les Signes Num.*, 51.

[3] G. Enestrom, "Sur les neuf 'limites' mentionnés dans l'"Algorismus' de Sacrobosco," *Bibl. Math.*, XI (2), 97. See also the 12th century MS. described by M. Chasles in the *Comptes rendus*, XVI (1843), 237; the *Compotus Reinheri*, p. 28; Boncompagni's *Bullettino*, X, 626; S. Günther, *Geschichte des math. Unterrichts*, p. 99 (Berlin, 1887) (for Bernelinus), hereafter referred to as Günther, *Math. Unterrichts*.

[4] *E.g.*, by Peletier (1549): "Le Nombre Entier se diuise en Simple, Article, & Composé. Le Simple est le Nombre plus bas que 10: ce sont les huict figures, 2, 3, 4, 5, 6, 7, 8, 9." He uses numbers and figures as synonymous, and uses "simple" for "digit."

Meaning of "Article" and "Composite." The articles were sometimes limited to nine in number (10, 20, ···, 90), but it was more common to take any multiple of ten. In the early printed books they were occasionally called decimal numbers,[1] and as such they finally disappeared.

The term "composite," originally referring to a number like 17, 56, or 237, ceased to be recognized by arithmeticians in this sense because Euclid had used it to mean a nonprime number.[2] This double meaning of the word led to the use of such terms as "mixed" and "compound" to signify numbers like 16 and 345.[3]

The oldest known French algorism (*c.* 1275) has the threefold division[4] above mentioned, as does also the oldest one in the English language (*c.* 1300), already cited. The latter work is so important in the history of mathematics in this language as to justify a further brief quotation:

> Some numbur is called digitus latine, a digit in englys. Somme nombur is called articulus latine. An Articul in englys. Some nombur is called a composyt in englys. . . .
>
> ¶Sunt digiti numeri qui citra denarium sunt.[5]

[1] Thus, Pellos (1492, fol. 4) speaks of "numbre simple," "nūbre desenal," and "nūbre plus que desenal"; and Ortega (1512; 1515 ed., fols. 4, 5) has "lo numero simplice," "lo numero decenale," and "lo numero composto."

[2] *Elements*, II, def. 13. For other Greek usage see Heath's *Euclid*, Vol. II, p. 286.

Lazesio (1526), among others, pointed out this twofold usage "secūdo sacro busco ī suo algorismo" and "secōdo el senso di Euclide" (1545 ed., fol. 2). See also Pacioli, *Sūma* (1494 ed., fols. 9, 19); Tartaglia, *General Trattato* (1556, II, fol. 1, *v.*); Santa-Cruz (1594; 1643 ed., fol. 2).

Trenchant (1566; 1578 ed., p. 223) speaks of "Nombre premier, ou incomposé," and "Nombre second, ou composé," a natural use of "second" as related to "premier" (prime), and the same usage was doubtless common at that time.

[3] "Alius aūt mixt⁹ siue ꝑpositus," in the *Questio haud indigna eiusq3 solutio ex aurelio Augustino*, *c.* 1507. So Hylles (1592; 1600 ed., fol. 7) says: "The third sort are numbers MIXT or compound"; Digges (1572; 1570 ed., p. 2) uses "compound" alone; and Hodder (10th ed., 1672, p. 5) has "A Mixt, or Compound." Dutch arithmeticians avoided the difficulty by using terms in the vernacular; thus, Mots (1640) gives "De enckel getallen" (digits), "Punct ofte leden-getallen" (articles), "t'samen-gevoeghde getallen (composites).

[4] "Tu dois savoir ki sont .3. manieres de nombres car li .1. sont degit li autre article. li autre compost." See Ch. Henry in Boncompagni's *Bullettino*, XV, 53.

[5] The anonymous writer here quotes from the *Carmen de Algorismo* of Alexandre de Villa Dei (*c.* 1240). The translation follows.

¶Here he telles qwat is a digit, Expone versus sic. Nomburs digitus bene alle nomburs þat ben with-inne ten, as nyne, 8.7.6.5.4.3.2.1. . . . Articulis ben ben alle þat may be deuidyt into nomburs of ten & nothynge leue ouer, as twenty, thretty, fourty, a hundryth, a thousand, & such oþer. . . . Compositys ben) nomburs þat bene componyt of a digyt & of an articulle as fouretene, fyftene, sextene, & such oþer.

Recorde (*c.* 1542) sums the matter up by saying:

A diget is any numbre vnder 10. . . . And 10 with all other that may bee diuided into x. partes iuste, and nothyng remayne, are called articles, suche are 10, 20, 30, 40, 50, &c. 100, 200, &c. 1000. &c. And that numbre is called myxt, that contayneth articles, or at the least one article and a digette: as 12.[1]

At best such a classification is unwieldy, and many of the more thoughtful writers, like Fibonacci (1202), abandoned it entirely. Others, like Sacrobosco (*c.* 1250), struggled with it but were obscure in their statements;[2] while Ramus very wisely (1555) dismissed the whole thing as "puerile and fruitless."[3]

All that is left of the ancient discussion is now represented by the word "digit," which is variously used to represent the numbers from one to nine, the common figures for these numbers, the ten figures 0, 1, . . ., 9, or the first ten numbers corresponding to the fingers.

Significant Figures. After the advent of the Hindu-Arabic figures into Europe (say in the 10th century) the difference between the zero and the other characters became a subject of comment. The result was the coining of the name "significant figures" for 1, 2, 3, · · ·, 9. At the present time the meaning


[1] 1558 ed. of the *Grovnd of Artes*, fol. C iij. Similar classifications are found in most of the early printed books of a theoretical nature, but less frequently in the commercial books.

[2] Thus Petrus de Dacia (1291) confessed that he could not quite understand Sacrobosco, saying, "ita credo auctorem esse intelligendum."

[3] "Puerilis et sine ullo fructu." See also Boncompagni, *Trattati d'Aritmetica*, II, 27 (Rome, 1857); J. Havet, *Lettres de Gerbert*, p. 238 (Paris, 1889); Boncompagni's *Bullettino*, XIV, 91; *Abhandlungen*, III, 136.

has been changed, so that 0 is a significant figure in certain cases. For example, if we are told to give log 20 to four significant figures, we write 1.301. Similarly, we write 0.3010 for log 2, and 7.550 for $\sqrt{57}$. The term is doubtless to be found in medieval manuscripts; at any rate it appears in the early printed arithmetics[1] and has proved useful enough to be retained to the present time in spite of the uncertainty of its meaning.

Odd and Even Numbers. The distinction between odd and even numbers is one of the most ancient features in the science of arithmetic. The Pythagoreans knew it, and their founder may well have learned it in Egypt or in Babylon. It must have been common to a considerable part of the race, for the game of "even and odd" has been played in one form or another almost from time immemorial,[2] being ancient even in Plato's time.[3] The game consisted simply in guessing odd or even with respect to the number of coins or other objects held in the hand.



The odd number was also called by the geometric name of "gnomon," the primitive form of the sundial. If such a figure

[1] *E.g.*, Licht (1500); Grammateus (1518), "neun bedeutlich figuren"; Riese (1522), "Die ersten neun sind bedeutlich"; Gemma Frisius (1540); Stifel (1544), "Et nouem quidem priores, significatiuae uocantur"; Peletier (1549), "Chacune des neufs premieres (qui sont appellees significatiues)" . . . ; Recorde (*c.* 1542), "The other nyne are called Signifying figures"; Trenchant (1566).

[2] This is seen in such expressions as ἀρτιασμός, ἄρτια ἢ περιττὰ παίζειν, ζυγὰ ἢ ἄζυγα παίζειν. This ζυγὰ ἢ ἄζυγα, "yokes or not-yokes," is similar to the Sanskrit "yuj" and "ayug" for even and odd. Horace couples it with riding a hobby horse as a childish diversion:


> Ludere par impar, equitare in harundine longa.
> *Satires*, II, 3, 248

See also E. B. Tylor, "History of Games," in the *Fortnightly Review*, May, 1879, p. 735.

[3] In addition to the references to the Greek theory of numbers given in Volume I and in this chapter, consult Dickson, *Hist. Th. Numb.*; F. von Drieberg, *Die Arithmetik der Griechen*, Leipzig, 1819; G. Friedlein, *Die Zahlzeichen und das elementare Rechnen der Griechen und Römer*, Erlangen, 1869; Heath, *History*, I, 67–117. Heath mentions a fragment of Philolaus (*c.* 425 B.C.) which says that "numbers are of two special kinds, odd and even, with a third, even-odd, arising from a mixture of the two."

is turned to the east in the morning and to the west in the afternoon, the hours can be read on the horizontal arm as in the Egyptian sun clock mentioned in Volume I, page 50. Thus we have the origin of the right shadow, the *umbra recta*, used in early trigonometry. By such an instrument we come to "know"[1] the time, and by facing it to the south we also come to know the seasons, the solstices, and the length of the year.[2]

It is apparent that the gnomon here shown in the shaded part of the figure is of the form $2n + 1$ and hence, as stated above, is an odd number.[3] It is also apparent that $\sum_{0}^{n} (2n + 1)$ is a square, that is, that the sum of the first n odd numbers, including 1, is a square,—a fact well known to the Greeks, as is shown by the works of Theon of Smyrna[4] (*c.* 125).



That there is luck in odd numbers is one of the oldest superstitions of the race, with such occasional exceptions as the case of the general fear of thirteen,—a fear that seems to have long preceded the explanation that it arose from the number present at the Last Supper.[5]

The general feeling that odd numbers are fortunate and even numbers unfortunate comes from the ancient belief that odd numbers were masculine and even numbers, always containing other numbers, were feminine. This led to the belief that odd numbers were divine and heavenly, while even numbers were human and earthly. The superstition was quite general among ancient peoples. Plato says: "The gods below . . . should receive everything in even numbers, and of the second choice, and

[1] Greek γνώμων (*gno'mon*), one who knows, from γιγνώσκειν, γνῶναι, know.

[2] Heath, *History*, I, 78.

[3] "Gnomon . . . quod Latini amussim seu normam vocant." J. C. Heilbronner, *Historia Matheseos Universae* (Leipzig, 1742), p. 173; see also page 193; hereafter referred to as Heilbronner, *Historia*.

[4] *Theonis Smyrnaei . . . expositio*, ed. Hiller, p. 31 (Leipzig, 1878). On this entire discussion see also Johnson, *Nicomachus*, and especially Heath, *History*, I, 77.

[5] Ernst Böklen, *Die Unglückszahl Dreizehn und ihre mythische Bedeutung*, Leipzig, 1913, with extensive bibliography.

ill omen; while the odd numbers, and of the first choice, and the things of lucky omen, are given to the gods above,"[1] and the phrase "Deus imparibus numeris gaudet" ("God delights in odd numbers") probably goes back to the time of Pythagoras.[2]

The superstition runs through a wide range of literature. Thus, Shakespeare, in the *Merry Wives of Windsor*, remarks that "there is divinity in odd numbers, either in nativity, chance, or death." Such beliefs naturally persist among the less advanced peoples and are common even today. For example, on the island of Nicobar, India, an odd number of vessels of water are dashed against the hut where a corpse is being laid out, and the stretcher that bears it must always contain an odd number of pegs.[3]

Further Classification. The Greeks not only recognized odd and even numbers,[4] but they carried the classification much farther, including what Euclid calls "even-times-even numbers," "even-times-odd numbers," and "odd-times-odd numbers." His definitions of the first two differ from those given by Nicomachus (*c.* 100) and other writers,[5] with whom an "even-times-even number" is of the form 2^n; an "even-times-odd number" is of the form $2(2n+1)$; and an "odd-times-odd number" is of the form $(2n+1)(2m+1)$. How far back these ideas go in Greek arithmetic is unknown, for they were doubtless transmitted orally long before they were committed to writing.

Since the product of two equal numbers represents the numerical area of a square, this product was itself called a square,—a word thus borrowed from geometry. The product of two unequal numbers was called a heteromecic (different-sided) number. Square and heteromecic numbers were called plane

[1] *Laws*, Jowett translation, V, 100.


[2] On the general number theory of Pythagoras, see Heath, *History*, I, 65.

[3] E. H. Man, "Notes on the Nicobarese," in the *Indian Antiquary*, 1899, p. 253.

[4] In Euclid's *Elements*, VII, 6, 7, ἄρτιοι and περισσοί.

[5] For particulars see Heath's *Euclid*, Vol. II, pp. 277, 281 seq. For the "odd-times-even number," which Euclid seems to have taken as synonymous with an "even-times-odd number," see *ibid.*, p. 283; on the general classifications by the Greeks, see K. G. Hunger, *Die arithmetische Terminologie der Griechen*, Prog., Hildburghausen, 1874.

numbers, while the product of three numbers was called a solid number, the cube being a special case. These are



30 ARITHMETICAE

NICOLAVS.

Pariter impar.

QVis eſt alter numerus par? IVST. Eſt pariter impar, uel à paribus impar, Græcis ἀρτιοπέριοσος, uel ἀρτιάκις περισσος. Eſt autem, cum primum diuiditur, mox fit indiuiſibilis, ut 14. 18. 22. NICOL. Quomodo eum finit horum numerorum exquiſitus magiſter Euclides? IVST. Sic: ἀρτιάκις περισσος ὅταν, ὁ ὑπὸ ἀρτίου ἀριθμοῦ μετρούμενος μὴ περισσον ἀριθμόν. Ita numerus diuidens par eſt, ſed diuiſorius mox impar exurget. NICOL. Cur id nominis illi inditum eſt?

Cur uocatur pariter impar

IVST. Ideo, quod qualibet eius ordinis numeri pares, facti ſunt per imparem multiplicationem: ut bis ter, ſenarium, bis quinq; denarium conficiunt. Verum ſi cui altius contemplari libet, eundem uocabit imparem in ſua quantitate, ſed parem in denominatione. Eſto exempli gratia, denarius, cuius altera pars eſt quinarius, qui quantitate, hoc eſt, monadum congregatione eſt impar, ſed quia à binario denominatur, par iudicabitur. Quæ ratio nominis ex Boethio colligitur: Alia autem Euclidi eſſe uidetur. NICOL. Sunt ne huic de illo aliquot theoremata?

Symbola ex pariter impari.

IVST. Quidni? Vnum eſt. Si numerus dimidium impar habuerit, pariter impar eſt tantum. Nam hic duntaxat extremum, quod maxi-

THEORY OF ODD AND EVEN NUMBERS

From the arithmetic of Willichius (1540). The page also illustrates the use of the catechism method in the 16th century

particular types of the figurate numbers mentioned later. The Boethian arithmetic made much of this classification;[1]

[1] *Boetii de institutione arithmetica libri duo*, ed. Friedlein, p. 17 (Leipzig, 1867); hereafter referred to as *Boethius*, ed. Friedlein.

For a full discussion see R. Bombelli, *L'antica numerazione Italica*, cap. x (Rome, 1876); hereafter referred to as Bombelli, *Antica numer.* For the status of the classification in the early printed books, see Pacioli, *Sūma*, 1494 ed., fol. [1], *v.* (= *A* [1], *v.*).

the medieval writers, both Arab[1] and Latin,[2] did the same; and early writers in the vernaculars simply followed the custom.[3]

Prime Number. Aristotle, Euclid, and Theon of Smyrna defined a prime number as a number "measured by no number but by an unit alone," with slight variations of wording. Since unity was not considered as a number, it was frequently not mentioned. Iamblichus says that a prime number is also called "odd times odd," which of course is not our idea of such a number. Other names were used, such as "euthymetric" and "rectilinear," but they made little impression upon standard writers.[4]

The name "prime number" contested for supremacy with "incomposite number" in the Middle Ages, Fibonacci (1202) using the latter but saying that others preferred the former.[5]

Perfect Numbers. Conventionally we speak of the aliquot parts of an integral number as the integral and exact divisors of the number, including unity but not including the number itself. A number is said to be deficient, perfect, or abundant according as it is greater than, equal to, or less than the sum of its aliquot parts.[6]

[1] On Savasorda (*c.* 1100), or Abraham bar Chiia, and his classification, see *Abhandlungen*, XII, 16. On al-Ḥaṣṣâr (*c.* 12th century), see *Bibl. Math.*, II (3), 17.

[2] Thus Jordanus Nemorarius (*c.* 1225): "Par numerus est qui in duo equalia diuidi potest. Impar est in quo aliqua prima pars est absq3 pari: additq3 supra parem vnitatē. Parium numerorū alius pariter par: alius pariter īpar: et alius impariter par. Pariter par est quē nullus impar numerat. Pariter īpar est quē quicunq3 pares numerāt. Imparit9 par est quē quidā par scdm parē ī quidā scdm imparē numerāt." 1496 ed., fol. b (3).

[3] *E.g.*, Chuquet, *La Triparty* (1484); see Boncompagni's *Bullettino*, XV, 619. Curtze found an early German MS. at Munich (No. 14,908, Cod. lat. Monac.) with such terms as "gelich oder ungelich," "glich unglich," and the like. See *Bibl. Math.*, IX (2), 39.

[4] Heath, *Euclid*, Vol. I, p. 146; Vol. II, pp. 284, 285.

[5] "Nvmerorum quidam sunt incompositi, et sunt illi qui in arismetrica et in geometria primi appellantur. . . . Arabes ipsos hasam appellant. Greci coris canon, nos autem sine regulis eos appellamus." *Liber Abaci*, I, 30.

[6] *E.g.*, 8 is a deficient number, since $8 > 1 + 2 + 4$; 6 is a perfect number, since $6 = 1 + 2 + 3$; 12 is an abundant number, since $12 < 1 + 2 + 3 + 4 + 6$. Various other names are given to abundant and deficient numbers, such as redundant or overperfect (ὑπερτελής, ὑπερτέλειος) and defective (ἐλλιπής). Heath, *History*, I, 10, 74.

This classification may have been known to the early Pythagoreans, but we have no direct evidence of the fact; indeed, their use of "perfect" was in another sense, 10 apparently being considered by them as a perfect number.

Euclid proved[1] that if $p = \sum_{0}^{n} 2^n$ and is prime, then $2^n p$ is perfect.[2] Nicomachus[3] separated even numbers into the classes above mentioned, and gave 6, 28, 496, and 8128 as perfect numbers, noting the fact that they ended in 6 or 8. Theon of Smyrna (*c.* 125) followed the classification of Nicomachus, but gave only two perfect numbers, 6 and 28. Iamblichus[4] (*c.* 325) did the same, but asserted that there was one and only one perfect number in each of the intervals 1 ... 10, 10 ... 100, 100 ... 1000, 1000 ... 10,000, and so on, and that the perfect numbers end alternately in 6 and 8,—statements which are untrue but which are found repeated in the arithmetic[5] of Boethius (*c.* 510). Subsequent writers in the Middle Ages and the Renaissance frequently followed Nicomachus or Iamblichus.

Fibonacci (1202) gave $\frac{1}{2} \cdot 2^2(2^2 - 1) = 6$, $\frac{1}{2} \cdot 2^3(2^3 - 1) = 28$, $\frac{1}{2} \cdot 2^5(2^5 - 1) = 496$ as perfect numbers, and so in general for $\frac{1}{2} \cdot 2^p(2^p - 1)$ where $2^p - 1$ is prime,—a rule which holds for the first eight perfect numbers but is not universal.[6] Chuquet (1484) gave Euclid's rule and repeated the ancient error that perfect numbers end alternately in 6 and 8.

The fifth perfect number, 33,550,336, is first given, so far as known, in an anonymous manuscript[7] of 1456–1461. Pacioli (1494) incorrectly gave 9,007,199,187,632,128 as a perfect number.[8]

[1] *Elements*, IX, 36.

[2] On all this work see Dickson, *Hist. Th. Numb.*, with bibliography, I, i; R. C. Archibald, *Amer. Math. Month.*, XXVIII, 140, with valuable references to American contributions.

[3] *Arithmetica*, I, 14, 15.

[4] 1668 ed., p. 43.

[5] *Arithmetica*, I, cap. 20, "De generatione numeri perfecti."

[6] Dickson, *Hist. Th. Numb.*, I, 13.

[7] Codex lat. Monac. 14,908. Dickson, *Hist. Th. Numb.*, I, 6.

[8] "Sia el nūero a noi ꝑposto. 9007199187632128. q̄le cōmo e ditto: e ꝑfecto." Fol. 7, *v*.

Charles de Bouelles (1509) wrote on perfect numbers[1] and asserted, without proof, that every perfect number is even. He stated that $2^{n-1}(2^n-1)$ is a perfect number when n is odd, which is substantially the incorrect rule of Fibonacci. This was also given by various other writers of the 16th century, including as good mathematicians as Stifel[2] (1544) and Tartaglia[3] (1556).

Robert Recorde[4] (1557) attempted to give the first eight perfect numbers, but three in his list were incorrect. Cataldi[5] showed that Pacioli's pretended fourteenth perfect number is in fact abundant, that the ancient belief that all perfect numbers end in 6 or 8 is unfounded, and that perfect numbers of the type given by Euclid's rule do actually end in 6 or 8.

Descartes thought that Euclid's rule covered all even perfect numbers and that the odd perfect numbers were all of the type ps^2, where p is a prime.[6]

Fermat (1636) and Mersenne (1634) paid much attention to the subject, and their investigations contributed to the theory of prime numbers.[7]

Euler at first (1739) asserted his belief that $2^{n-1}(2^n-1)$ is a perfect number for $n = 1, 2, 3, 5, 7, 13, 17, 19, 31, 41$, and 47, but afterward (1750) showed that he was in error with respect to 41 and 47. He proved that every even perfect number is of Euclid's type, $2^n\sum_0^n 2^n$, and that every odd perfect number is of the form $r^{4\lambda+1}p^2$, where r is a prime of the form $4n+1$.[8]

There are many references to perfect numbers in general literature,[9] in Hebrew and Christian writings on religious

[1] "De Numeris Perfectis," in his general work published at Paris in 1509–1510. See *Rara Arithmetica*, p. 89.

[2] *Arithmetica Integra*, fols. 10, 11 (Nürnberg, 1544); also *Die Coss Christoffs Rudolffs*, fols. 10, 11 (Königsberg, 1553).

[3] *La seconda Parte del General Trattato*, fol. 146, *v.* Venice, 1556.

[4] *The whetstone of witte*, fol. [4, *v.*]. London, 1557.

[5] *Trattato de' nvmeri perfetti*. Bologna, 1603.

[6] *Œuvres*, II, 429. Paris, 1898. [7] Dickson, *Hist. Th. Numb.*, I, 11–13.

[8] *Ibid.*, p. 18. For the later theory, consult this work, I, i.

[9] Thus Macrobius, in his *Saturnalia*, says that six "plenus perfectus atque diuinus est." *Satvrnaliorvm Liber VII*, cap. xiii, ed. Eyssenhardt, 1868, p. 446.

doctrines,[1] including Isidorus of Seville and Rabbi ben Ezra, and in the works of medieval and Renaissance mystics.[2]

Amicable Numbers.[3] Two integral numbers are said to be amicable[4] if each, as in the case of 220 and 284, is equal to the sum of the aliquot parts of the other. These two numbers, probably known to the early Pythagoreans, are mentioned by Iamblichus. They occupied the attention of the Arabs, as in the works of Ṯâbit ibn Qorra (*c.* 870). It was asserted by certain Arab writers that talismans with the numbers 220 and 284 had the property of establishing a union or close friendship between the possessors, and this statement was repeated by later European writers, including Chuquet (1484) and Mersenne (1634).

For a long time the only amicable numbers known were the two given above, 220 and 284, but in 1636 Fermat[5] discovered a second pair, $17{,}296 = 2^4 \cdot 23 \cdot 47$ and $18{,}416 = 2^4 \cdot 1151$, and also found a rule for determining such numbers.[6] A third pair was discovered by Descartes[7] (1638), namely, $9{,}363{,}584 = 2^7 \cdot 191 \cdot 383$ and $9{,}437{,}056 = 2^7 \cdot 73{,}727$. Descartes gave a rule which he asserted[8] to be essentially the same as Fermat's, but which various later writers, apparently ignorant of this assertion, assigned to Descartes himself.

Euler[9] (1750) made a greater advance in this field than any of his predecessors, adding fifty-nine pairs of amicable numbers of the type am, an, in which a is relatively prime to m and n, and contributing extensively to the general theory. Dickson

[1] *E.g.*, J. J. Schmidt, *Biblischer Mathematicus*, p. 20 (Züllichau, 1736).

[2] See Curtze's mention of a Munich MS. (No. 14,908, Codex lat. Monac.) in the *Bibl. Math.*, IX (2), 39, with five perfect numbers.

Thierfelder (1587, fol. A_4, *r.*) says: "Den in sechsz tagen hat Gott Himmel vnd Erden/ vnd alles was darinen ist/ gemacht/ das ist ein *Trigonal* oder dreyeckichte Zahl/ welche Zahlen für die heiligen Zahlen gehalten werden/ vnd ist darzu die erste perfect Zahl."

[3] Dickson, *Hist. Th. Numb.*, I, 36.

[4] The terms "amiable" and "agreeable" are also used.

[5] *Œuvres*, 1894, II, 72, 208.

[6] The rule is given in Dickson, *loc. cit.*, p. 37.


[7] *Œuvres*, 1898, II, 93.

[8] *Œuvres*, 1898, II, 148.

[9] *Opuscula varii argumenti*, 3 vols., II, 23 (Berlin, 1746–1751). See also *Bibl. Math.*, IX (3), 263; X (3), 80; XIV (3), 351; Cantor, *Geschichte*, III, 616.

(1911) has obtained two new pairs of amicable numbers and has also added to the general theory of the subject.[1]

Figurate Numbers. The Greeks were deeply interested in numbers which are connected with geometric forms and which therefore received the name of figurate numbers.[2] These are triangular if capable of being pictured thus:



FIGURATE NUMBERS


From the first printed edition of the arithmetic of Boethius (1488)




and are therefore of the form

$$\sum_{1}^{n} n = \tfrac{1}{2} n(n+1).$$

They are square if they can be represented by squares, such as ⁘, and are then of the form n^2. They are pentagonal if in the form of a square with a triangle on top, thus:



so that the form is $n^2 + \frac{1}{2}n(n-1)$. Similarly, there are hexagonal numbers and other types of polygonal numbers.[3]



In the Greek manuscripts they appeared in such forms as those here shown, the α's standing for 1's or possibly for ἀριθμός (*arithmos'*, number).[4]

Related to figurate numbers there are the linear numbers. Under this name Nicomachus (*c.* 100) included the natural numbers, beginning with 2; side and diagonal numbers[5]; area,

[1] See also his *Hist. Th. Numb.* and *Amer. Math. Month.*, XXVIII, 195.


[2] Boethius defined them as numbers "qui circa figuras geometricas et earum spatia demensionesque versantur." Ed. Friedlein, p. 86, l. 12. See also Heath, *History*, I, 76.

[3] Boethius, ed. Friedlein, p. 98 seq.

[4] These two forms are from a 10th century MS. of Nicomachus in Göttingen.

[5] The πλευρικοὶ καὶ διαμετρικοὶ ἀριθμοί of Theon of Smyrna (*c.* 125). See also Boethius, ed. Friedlein, p. 90.

or polygonal, numbers[1]; and solid numbers,[2] including cubic, pyramidal, and spherical numbers.[3] A relic of such numbers is



PYRAMIDAL NUMBERS

From Joachim Fortius Ringelbergius, *Opera*, 1531. The four layers of the two pyramidal numbers 35 and 30 are shown

seen in problems relating to the piling of round shot, still to be found in algebras. Indeed, it is not impossible that they may have been suggested to the ancients by the piling of spheres in

[1] See Nicomachus, *Introd.*, II, capp. 8–11.
[2] Στερεοί. Nicomachus, *Introd.*, II, 14.
[3] Boethius, *loc. cit.*, pp. 104, 121.

such games as the *Castellum nucum* to which Ovid refers in his poem *De Nuce*, where the pyramidal number is mentioned.[1]

Continuous and Discrete. The distinction between continuous and discrete magnitude is commonly referred to the Pythagoreans or even to Pythagoras (*c.* 540 B.C.) himself,[2] the continuous magnitude being geometric and the discrete being arithmetic. The distinction was recognized by various Greek and Latin writers,[3] appearing in the works of such medieval authors as Fibonacci (1202) and Roger Bacon (*c.* 1250).[4]

Cardinals and Ordinals. The distinction between cardinal and ordinal numbers is ancient, but the names are relatively modern. A cardinal number is a number on which arithmetic turns[5] or depends, and hence is a number of importance,[6] while ordinal number is one which denotes order.[7]

6. Unity

Unity. Not until modern times was unity considered a number. Euclid defined number as a quantity made up of units,[8] and in this he is followed by Nicomachus.[9] Unity was defined by Euclid as that by which anything is called "one."[10] It was generally defined, however, as the source of number, as in the

[1] Quattuor in nucibus, non amplius, alea tota est,
Cum sibi suppositis additur una tribus.
See also F. Lindemann, "Zur Geschichte der Polyeder und der Zahlzeichen," *Sitzungsberichte der math.-physik. Classe der K. Bayerischen Akad. der Wissensch. zu München*, XXVI, 625–757 (Munich, 1897).

[2] "Ogni quantità . . . secondo Pythagora, è o continua, ouer Discreta, la continua è detta Magnitudine, . . . & la discreta moltitudine." Tartaglia, *General Trattato*, I, fol. 1, *r.* (Venice, 1556).

[3] Boethius, ed. Friedlein, pp. 8, 16; Heath, *Euclid*, Vol. I, p. 234.

[4] *E.g.*, in the Sloane MS. fol. 94 of the *Communia*.

[5] Latin, *cardo*, a hinge.

[6] Compare cardinal, a prince of the Church. Glareanus recognized this metaphor: "Sunt enim quaedam, quae Cardinalia appellant, à cardine sumpta, ut opinor, metaphora, quòd ut in cardine ianua uertitur, ita huius artis primum ac praecipuum negocium in hisce consistat" (1538; 1543 ed., fol. 3, *r.*).

[7] On the history of these terms see E. Bortolotti, "Definizioni di Numero," *Esercitazioni Matematiche*, II, 253, and *Periodico di Matematiche*, II (4), 413.

[8] Ἀριθμὸς δὲ τὸ ἐκ μονάδων συγκείμενον πλῆθος. *Elements*, VII, def. 2. See also Heath, *History*, I, 69.

[9] *Introd.*, I, 7, 1.

[10] Μονάς ἐστιν καθ' ἣν ἕκαστον τῶν ὄντων ἓν λέγεται. *Elements*, VII, def. 1; ed. Heath, Vol. II, p. 279, with references to other Greek writers.

anonymous *Theologumena*,[1] a Greek work of the early Middle Ages. The dispute goes back at least to the time of Plato, for the question is asked in the *Republic*, "To what class do unity and number belong?"—the two being thus put into separate categories.

It is not probable that Nicomachus (*c.* 100) intended to exclude unity from the number field in general, but only from the domain of polygonal numbers.[2] It may have been a misinterpretation of the passage from Nicomachus that led Boethius[3] to add the great authority of his name to the view that one is not a number. Even before his time the belief seems to have prevailed, as in the case of Victorius (457) and Capella (*c.* 460), although neither of these writers makes the direct assertion.[4] Following the lead of Boethius, the medieval writers in general, such as al-Khowârizmî[5] (*c.* 825), Psellus[6] (*c.* 1075), Savasorda[7] (*c.* 1100), Johannes Hispalensis[8] (*c.* 1140), and Rollandus[9] (*c.* 1424), excluded unity from the number field.[10] One writer, Rabbi ben Ezra (*c.* 1140), seems, however, to have

[1] Μονάς ἐστιν ἀρχὴ ἀριθμοῦ, θέσιν μὴ ἔχουσα. *Theologumena*, I, 1.

[2] Ἡ μὲν μονὰς σημείου τόπον ἐπέχουσα καὶ τρόπον. See also Johnson, *Nicomachus*, p. 7.

[3] "Numerus est unitatum collectio." Ed. Friedlein, p. 13, l. 10. In the Latin version of the so-called Boethian geometry it is asserted: "Primum autem numerum id est binarium, unitas enim, ut in arithmeticis est dictum, numerus non est, sed fons et origo numerorum. . . ." Ed. Friedlein, p. 397, l. 19. See also H. Weissenborn, *Gerbert*, p. 219 (Berlin, 1888).

[4] "Unitas illa, unde omnis numerorum multitudo procedit." From the *Calculus* of Victorius; see Boncompagni's *Bullettino*, IV, 443.

"Nec dissimulandum est ex eo quod monas retractantibus unum solum ipsam esse. ab eaque cetera procreari. Omniumque numerorum solam seminarium esse. solamque mensuram et incrementorum. causam. statumque detrimentorum." From a fragment of Capella; see E. Narducci in Boncompagni's *Bullettino*, XV, 566.

[5] "Quia unum est radix uniuersi numeri, et est extra numerum." From the supposed translation of Adelard of Bath.

[6] "Principium itaque omnis numeri est Monàs, non-numerus fons numerorum." See the 1532 edition, p. 13.

[7] "Numerus est ex unitatibus profusa collectio" (Plato of Tivoli's translation, 1145). See C. H. Haskins, *Bibl. Math.*, XI (3), 332.

[8] "Unitas est origo et prima pars numeri . . . sed ipsa extra omnem numerum intelligitur." See B. Boncompagni, *Trattati d' Aritmetica*, II, 25 (Rome, 1857); hereafter referred to as Boncompagni, *Trattati*.

[9] See Volume I, page 261. "Vnitas non est numerus sed principia numerorum" (Plimpton MS., Pt. I, cap. 1).

[10] See also Boncompagni's *Bullettino*, XV, 126.

approached the modern idea. In his *Sefer ha-Echad* (*Book on Unity*) there are several passages in which he argues that one should be looked upon as a number.

Most of the authors of the early printed books excluded unity, as is seen in the works of Pacioli[1] (1494), Köbel[2] (1514), Tzwivel[3] (1505), and many others. Thus the English writer Baker (1568) remarks that "an vnitie is no number but the beginning and original of number."[4] In the 16th century, however, the more thoughtful writers began to raise the question as to whether this exclusion of unity from the number field was not like the trivial disputes of the schoolmen,[5] and by the end of the century it was recognized that the ancient definition was too narrow. Thus Hylles (1592), speaking of "an vnit or an integer (which sometimes I also cal an Ace)," is rather afraid to take a definite stand in the matter, but says that "the latter writers, as namely Ramus, and such as have written since his time, affirme not only that an vnite or one, is a number, but also that euery fraction or parte of an vnite, is a number. . . . I do accompt it after a sorte for the first or least number . . . euen as an egg, with[6] in power possibilitie containeth a bird though really and actually it is none." Stevin (1585), a much greater man, used the argument that a part is of the same nature as the whole, and hence that unity, which is part of a collection of units, is a number.[7] To this Antoine Arnauld, "le

[1] "Et essa vnita nō e numero: ma ben principio di ciascun numero" (1494 ed., fol. 9).

[2] "Daraus3 verstehstu das I. kein zal ist/ sonder es ist ein gebererin/ anfang/ vnnd fundament aller anderer zalen" (*Zwey rechenbuchlin*, Frankfort ed., 1537, fol. 26). It is also in his *Rechenbüchlin*, 1531 ed. dedication, and 1549 ed., fol. 26.

[3] "Unitas eñ numerus non est. sed fons et origo numerorum" (fol. 2).

[4] 1580 ed., fol. 1.

[5] So Gemma Frisius (1540) makes it a matter of authority: "Nvmerum authores vocant multitudinem ex Vnitatibus conflatum. Itaque Vnitas ipsa licèt subinde pro numero habeatur, propriè tamen numerus non erit" (1563 ed., fol. 5).

Also Trenchant (1566): " . . . l'vnité n'est pas nombre . . . Mais en la pratique, ou le nombre est tousiours adapté à quelque sugét . . . l'vnité est prinse pour nombre" (1578 ed., p. 9).

[6] *Sic*, for "which." From the 1600 edition.

[7] "La partie est de mesme nature que le tout. Unité est partie d'une multitude d' unitez . . . et par consequent nombre." See also the Girard edition, of 1634, p. 1, with slight change in wording.

grand Arnauld" (1612–1694), replied that the argument was worthless, for a semicircle is not a circle. Stevin also used the argument that if from a number there is subtracted no number, the given number remains; but if from 3 we take 1, 3 does not remain; hence 1 is not no number.[1] The school arithmetics kept the Boethian limitation until the close of the 18th century.[2]

Another common notion was that unity is, like a point, incapable of division,—an idea also due to the Greeks.[3]

7. Later Developments

Higher Domain of the Arithmetica. The later developments in the arithmetica do not belong to the domain of elementary mathematics. Their history has been treated with great erudition by Professor Dickson in his *History of the Theory of Numbers*.[4] As a matter of general information, however, a few of the theorems which have attracted wide attention will be stated.

Typical Theorems. In 1640 Fermat, in a letter to Bernard Frénicle de Bessy (*c.* 1602–1675), set forth the theorem that if p is any prime number and x is any integer not divisible by p, then $x^{p-1}-1$ is divisible by p. The special case of 2^p-2 being divisible by the prime p had long been known to Chinese scholars, but the general theorem is due to Fermat. Leibniz proved the proposition some time before 1683.

Euler stated Fermat's theorem in a communication to the Petrograd Academy[5] in the form: If $n+1$ is a prime dividing neither a nor b, then a^n-b^n is divisible by $n+1$.

As stated in Volume I, page 459, Wilson discovered (*c.* 1760) that if p is prime, then $1+(p-1)!$ is a multiple of p. The manuscripts of Leibniz now preserved at Hannover show that he knew the theorem before 1683, but he published nothing upon

[1] *Abhandlungen*, XIV, 227.

[2] *E.g.*, Ward's *Young Mathematician's Guide*, p. 4 (London, 1771).

[3] "Il punto nella Geometria, & l'vnità nell' Arimmetica non è capace di partimento. Proclo sopra Euclide lib. 2.c.xi." Ciacchi, *Regole Generali d' Abbaco*, p. 352 (Florence, 1675).

[4] See also A. Natucci, *Il Concetto di Numero*, Turin, 1923.

[5] Presented in 1732, published in 1738.

the subject. Lagrange published a proof of the theorem in 1771, deduced it from Fermat's Theorem, and proved its converse.

Fermat gave as his opinion that $2^{2^n}+1$ is always prime, but asserted that he was unable to prove it. Euler (1732)[1] showed that Fermat's opinion was not warranted, since

$$2^{2^5}+1 = 641 \cdot 6{,}700{,}417.$$

Fermat's connection with numbers of this form led to their being called "Fermat's Numbers."[2]

With respect to the sum and the number of divisors of a number there is an extensive literature.[3] For example, Cardan (1537) stated that a product P of k distinct primes has $1+2+2^2+\cdots+2^{k-1}$ aliquot parts; for example, that $3 \cdot 5 \cdot 7$ has $1 + 2 + 4$ aliquot parts. This rule was proved by Stifel in his *Arithmetica Integra* (1544). Frans van Schooten (1657) proved that a product of k distinct primes has $2^k - 1$ aliquot parts, which is only another expression for Cardan's rule.

Descartes (probably in 1638) showed that if p is a prime the sum of the aliquot parts of p^n is $(p^n-1)/(p-1)$, a law simply illustrated by the cases of 2^3 and 7^2.

Fermat proposed (1657) two problems: (1) Find a cube which, when increased by the sum of its aliquot parts, becomes a square, one example being $7^3 + (1 + 7 + 7^2) = 20^2$; (2) find a square which, when increased by the sum of its aliquot parts, becomes a cube. Problems of this general nature attracted the attention of men like Frénicle de Bessy, Lord Brouncker, Wallis, Frans van Schooten, Ozanam, and various later scholars.

Other Subjects of Investigation. Among other subjects investigated is that of the factors of numbers that can be expressed in the form of $a^n \pm b^n$; for example, to find all the prime factors of $2^{45} - 1$. There are also such questions as the infinitude of primes in general; the tests for primality; the number of primes between assigned limits; the curious properties connected with the digits of numbers; periodic fractions; and the general theory of congruent numbers.

[1] Published in 1738. [2] R. C. Archibald, *Amer. Math. Month.*, XXI, 247.
[3] Dickson, *Hist. Th. Numb.*, I, 51.

TOPICS FOR DISCUSSION

1. The numbers three and seven in folklore and in literature.

2. The history of the seven liberal arts.

3. Distinction between arithmetic and logistic in ancient and medieval times.

4. History of the word "algorism" in various languages, particularly with reference to its forms and significance.

5. Various names given to what is now called arithmetic in the period known as the Renaissance.

6. History of the distinction between concrete and abstract numbers. The present status of the question, including that of operations with concrete numbers.

7. History of the finger names assigned to numbers, and the probable reason why such names attracted more attention in early times than at present.

8. Rise of the idea of significant figures and the present use of the term.

9. Probable reasons for the superstitions in regard to odd and even numbers and for the properties assigned to them at various times and by various peoples.

10. The gnomon and its relation to numbers and to other branches of mathematics.

11. Nature of and probable reason for certain other classifications of number in ancient times.

12. Probable cause for the special interest in prime numbers expressed by the ancients.

13. The historical development of the interest in amicable numbers and the present status of the theory.

14. The historical development of the theory of perfect numbers and the present status of the theory.

15. The interest in figurate numbers among the Greeks and the traces of such numbers in modern times.

16. The history of the concept of unity and of the controversy with respect to its being a number.

17. Traces of ancient arithmetic and logistic in modern textbooks in arithmetic and algebra.

18. Questions relating to the theory of numbers and attracting the attention of mathematicians during the 19th and 20th centuries.

CHAPTER II

LOGISTIC OF NATURAL NUMBERS

1. Fundamental Operations

Number of Operations. In America at the present time it is the custom to speak of four fundamental operations in arithmetic, that is, in what the ancients called logistic. This number is, however, purely arbitrary, and it is quite possible to argue that it should be increased to nine or more,[1] or even that it should be decreased to one.

The *Crafte of Nombrynge* (c. 1300) enumerates seven:

¶Here tells þat þer ben .7. spices or partes of þis craft. The first is called addicion̄, þe secunde is called subtraccion̄. The thryd is called duplacion̄. The 4. is called dimydicion̄. The 5. is called multiplicacion̄. The 6. is called diuision. The 7. is called extraccion̄ of þe Rote.[2]

Sacrobosco (*c.* 1250) had already spoken of nine of these operations,—numeration, addition, subtraction, duplation, mediation, multiplication, division, progression, and the extraction of roots,[3]—and Michael Scott had done the same.[4] This was a common number among medieval writers and, indeed, in the

[1] On the general question of the operations see J. Tropfke, *Geschichte der Elementar-Mathematik*, I (2) (Leipzig, 1921), hereafter referred to as Tropfke, *Geschichte*; Suzan R. Benedict, *A Comparative Study of the Early Treatises introducing into Europe the Hindu Art of Reckoning*, Dissertation, Univ. of Michigan, 1914.

[2] R. Steele, *The Earliest Arithmetics in English*, Oxford, 1923. The old letter þ is our *th*. Since it slightly resembles our letter *y*, the old word þe (the) is often ignorantly written as *ye*, as in "ye editor."

[3] In his *Algorismus*. See Volume I, page 222.

[4] Santa-Cruz (1594) refers to this, saying: ". . . las especies del qual, segū Iuā de sacrovosco, y Michael Scoto, son nueue" (1643 ed., fol. 9).

early printed books.[1] Pacioli (1494), however, claimed credit to himself for reducing the number to seven.[2] In due time a further reduction was made to six,[3] then to five, as with most 16th-century writers, and then to four. When five operations were taken, numeration was usually the first, the topic properly including notation.[4] One of the first of the writers of any note to reduce the number to four was Gemma Frisius (1540),[5] and such was his influence that this number soon became common. There have been those, indeed, who gave only three fundamental operations, multiplication being included in addition as a special case. This number was given, for example, by Elia Misrachi (*c.* 1500).[6]

Duplation and Mediation. The four operations generally recognized at present will be considered later in this chapter; the two operations of duplation (doubling) and mediation (halving), with the reasons for their use, will be explained briefly at this time.

[1] *E.g.*, Widman (1489), Peurbach (*c.* 1460; 1st ed., 1492), Huswirt (1501), Tartaglia (1556), and Santa-Cruz (1594).

[2] He says that nine were given by "Giouã de sacro busco e Prodocimo de beldemandis da padua dignissimo astronomo e molti altri in loro algorismi. Ma noi le ditte noue reduremo a septe" (fol. 19, *r.*).

Since the names have some interest, the list is reproduced: "La prima sira ditta numeratiõe ouer representatiõe: cioe sapere cognoscere e releuare le figure e caratteri del nũero. La secõda sira ditta additiõe ouer recogliere: agiognere; sũmare e acozare. La terza sira ditta subtractiõe ouer abattere: sotrare: cauare e trare. La q̃rta fia ditta multiplicatiõe. La quinta sira ditta diuisiõe ouer partire. La sexta sira ditta ꝓgressiõe. La septima sira ditta delle radici extractione." *Ibid.*

[3] *E.g.*, Glareanus (1538): "Eius sex, ut in epitome, prosequemur species, numerationem, additionem, subtractionem, multiplicationem, diuisionem, ac progressionem" (1543 ed., fol. 9).

[4] Numeration (Latin *numeratio*, from *numerus*, a number) has lately been used to mean the reading of numbers. Since medieval writers often called the characters 1, 2, 3, . . . *notae* (compare the "notes" in music), the writing of numbers has been called notation. The distinction is one chiefly of the schoolroom. Ramus (1569; 1586 ed., p. 1) was one of the first prominent writers to make it: "In numero spectatur primum notatio, deinde numeratio."

[5] "Qvatuor omnino sunt Arithmetices species" (1563 ed., fol. 6).

[6] G. Wertheim, *Die Arithmetik des Elia Misrachi*, Prog., Frankfort a. M., 1893; hereafter referred to as Wertheim, *Elia Misrachi*. The first edition of Misrachi appeared at Constantinople, 1532 or 1533; the second, at Basel, 1546.

The Egyptians often multiplied by continued doubling,[1] thus saving the trouble of learning a multiplication table. This was particularly convenient in working on the abacus. On this account duplation was generally recognized as a separate topic until the 16th century. Moreover, the Egyptian tables of measure were commonly arranged so as to make doubling and halving operations of great importance.[2] This method continued as long as the abacus was in use, and persisted for some time after that instrument was generally abandoned. An interesting illustration of mediation is seen, for example, in official papers of Russia prior to the time of Peter the Great (1672–1725), the word "half" being repeated as many as ten times to indicate a certain division.[3]

The use of duplation and mediation is seen in many of the Arab works,[4] and this fact influenced such medieval translators as Johannes Hispalensis[5] (*c.* 1140) and Adelard of Bath[6] (*c.* 1120). The processes were common in the theoretical works of the 15th century[7] but not in the commercial arithmetics, at least in Italy. The early printed books of Germany were less progressive in this respect than those of other countries, partly

[1] Thus, $7 \times 15 = 2 \times 2 \times 15 + 2 \times 15 + 15$.

[2] See the common use of the fractions $\frac{1}{2}$, $\frac{1}{4}$, $\frac{1}{8}$, $\frac{1}{16}$, in the Edfu Survey, in H. Brugsch, *Thesaurus Inscriptionum Ægyptiacarum*, Leipzig, 1883–1891, Vol. III; H. Brugsch, *Numerorum apud Veteres Ægyptios Demoticorum Doctrina*, Berlin, 1849; T. E. Peet, *The Rhind Mathematical Papyrus*, London, 1923; hereafter referred to as Peet, *Rhind Papyrus*. For the survival of doubling and halving until the present time, see the tables in Mahmoud Bey, "Le système métrique actuel d'Égypte," *Journal Asiatique*, I (7), 69, 82.

[3] V. V. Bobynin, "Esquisse de l'histoire du calcul fractionnaire," *Bibl. Math.*, X (2), 97.

[4] *E.g.*, al-Nasavî (*c.* 1025), on whose work see F. Woepcke, *Journal Asiatique*, I (6), 496; al-Ḥaṣṣar (12th century), on whose work see M. Steinschneider, *Abhandlungen*, III, 10; al-Khowârizmî (*c.* 825), on whose work and al-Ḥaṣṣar's see Suter, *Bibl. Math.*, II (3), 12.

[5] Who, however, speaks of them merely as special cases of multiplication and division. See Boncompagni, *Trattati*, II, 38. Similarly, as to Gernardus (13th century?), see G. Enestrom, *Bibl. Math.*, XIII (3), 289, 292.

[6] Boncompagni, *Trattati*, I, 10.

[7] *E.g.*, in the Rollandus MS. (1424); see *Rara Arithmetica*, p. 446. Rollandus gives: "addere. sbtrhere. mediare. duplare. diuidere. m̄ltiplicare. et radices invenire" (fol. 2).

because of the continued use of the abacus in that part of Europe, and so these two processes are found in the works of Tzwivel (1505), Köbel (1514), Grammateus (1518), Riese (1522), Rudolff (1526), and various other writers of that period, often with a statement that they are special forms of multiplication and division.[1] Stifel (1545) uses them only apologetically, and Scheubel (1545) omits them entirely.[2] They are rarely found in any of the printed arithmetics of Italy,[3] Spain,[4] France, or England. It is a curious fact, however, that Recorde (*c.* 1542) omits them with integers but includes them with fractions,—a vagary that endured at least as late as the 1668 edition of his *Ground of Artes*. His example was followed by Baker (1568), who had the notion that only fractions should be used with fractions, saying: "If you will double anye broken number you shall divide y^{e} same by $\frac{1}{2}$," and giving triplation and quadruplation in the same way. Gemma Frisius (1540) did as much as any other Continental writer to show the absurdity of following those "stupid people" who would include these operations.[5]

Names of the Operations. The awkward expression "the four fundamental operations" is modern. Several others used in the past possess the merit of greater brevity, and some of these are still found in various languages. A common name is "species," a term of the 13th century and made popular in the 16th century by the works of Riese (1522) and Gemma Frisius (1540).[6] Ramus (1569) used both "parts" and

[1] "Dupliren heist zwifeltigen/ ist nichts anders dañ ein zal mit 2 multiplicirn. Medijren heist halb machen od' halbiren/ ist nichts anders/ dañ ein zal in 2 abteilen." Köbel (1514).

[2] "De duplatione porro & Mediatione, cum illa multiplicationis, haec uero diuisionis pars sit, scribere quicqȝ, necesse non fuit." "Tractatus secundus" of the 1545 edition of his *De Numeris*.

[3] *E.g.*, Pacioli (1494): "Ma noi le ditte noue reduremo a septe. Peroche la duplatiõe īplicita in la multiplicatiõe: ela mediatiõe nella diuisiõe" (fol. 19, *r.*).

[4] *E.g.*, Santa-Cruz (1594): "Y porq̄ el doblar no se distingue del multiplicar, ni el mediar del partir" (1643 ed., fol. 9).

[5] "Quid verò mouerit stupidos illos nescio" (1563 ed., fol. 12).

[6] In the Latin editions: "De speciebus Arithmetices"; "Vocamus autem species certas operandi . . ." In the Italian translation: "Delle Specie dell' Aritmetica."

"species,"[1] while most of the Spanish[2] and Dutch[3] arithmeticians of the 16th and 17th centuries used the latter only.

A common Italian name in the 16th century was "acts,"[4] although "passions" was also used.[5] When Clavius wrote his algebra (1608), he used the word *operationes*, and it is probable that this word worked down from algebra to arithmetic.

Sequence of Operations. Our present traditional sequence has by no means been generally recognized, particularly in relation to fractions. Although all writers place notation and a certain amount of addition first, there has been little further uniformity. Abraham bar Chiia (*c.* 1120), Rabbi ben Ezra (*c.* 1140), and Fibonacci (1202), for example, use this sequence: multiplication, division, addition, subtraction, fractions, proportion, and roots.[6] Grammateus (1518) used the order: addition, multiplication, subtraction, division,[7]—an order which has much to commend it.

2. Reading and Writing Numbers

Babylonian Numerals. Since the early Babylonians were without papyrus or parchment, they doubtless followed the custom of most other early peoples and wrote upon leather. Living on an alluvial plain, they had no convenient access to stone for the purpose of permanent inscriptions, except in the northern region, and so they also resorted to the use of clay. They wrote by pressing into the clay with a stylus, the result being wedge-shaped (cuneiform) characters. These tablets were then baked in the sun or in a kiln of some kind, and thus they

[1] "Alii faciunt arithmeticae partes vel species . . .," *Arithmeticae libri dvo*, p. 111.

[2] *E.g.*, Santa-Cruz (1594): ". . . las especies del qual . . . son nueue."

[3] Thus Stockmans (1589), Houck (1676), and others speak of the "vier specien."

[4] So Sfortunati (1534) speaks of the "Cinque atti dell' arithmetica."

[5] Tartaglia (1556) prefers "atti" but says that "altri gli dicone Passioni del numero" (1592 ed., fol. 5). The word comes from the Latin *passio*, used by late Latin writers to mean "phenomenon"; originally, something endured.

[6] M. Steinschneider, *Abhandlungen*, III, 107.

[7] 1535 ed., fol. A iii.

became fairly permanent records. For relatively small numbers the numerical system was simple, consisting of the following characters:



$= 1, \quad = 10, \quad = 100.$

These symbols had different numerical meanings, however. The stood not only for 1 but also for 60, 3600, 12,960,000, and in general for 60^n. The stood for $10 \cdot 60^n$, and hence for 10, 600, 36,000, · · ·. In every case the context was depended upon to determine which value was to be taken. Furthermore, we often find the units represented by horizontal strokes, 10 represented by a vertical crossed by a horizontal stroke (like a plus sign), 20 represented by a vertical crossed by two horizontals, and so on. In certain tablets 71 is represented by 1 (for 60), the above symbol for 10, and a horizontal stroke for the unit.



BABYLONIAN NUMERALS FROM 1 TO 9


The forms vary in shape, but this gives an idea of the simpler numerals in common use. For the correct forms as seen in the clay tablets, see page 39

In writing their numerals the Babylonians made a slight use of the subtractive principle with which we are familiar in connection with the Roman notation. For example, the XIX of the Romans is equivalent to XX − I, a device that was anticipated some two thousand years by the Babylonians, who wrote for 19, the symbol (*lal* or *lá*) meaning minus.[1] In this case, then, we have 20 − 1, or 19. It has been suggested that one reason for writing 19 as 20 − 1 instead of 10 + 9 is that

[1] There are numerous forms for this symbol, some of them very complex. See H. V. Hilprecht, *Mathematical, Metrological, and Chronological Tablets from the Temple Library of Nippur*, p. 23 (Philadelphia, 1906); hereafter referred to as Hilprecht, *Tablets*. See also G. Reisner, "Altbabylonische Maasse und Gewichte," *Sitzungsberichte der k. Preussischen Akad. der Wissensch.*, p. 417 (Berlin, 1896); G. Contenau, "Contribution à l'histoire économique d'Umma," *Bibliothèque de l'École des hautes études*, fascicule 219 (Paris, 1915), with excellent facsimiles of various numeral forms. See especially Plate XIV for the representation of 71 referred to in the text. The tablets date from *c.* 2300 to *c.* 2200 B.C.

it was an unlucky number. The nineteenth day of a lunar month was the forty-ninth day from the beginning of the preceding month, and this forty-ninth day was one to be specially avoided. To avoid writing 19, therefore, the Babylonians resorted to writing $20 - 1$. This does not, however, account for such common forms as $60 - \frac{1}{3}$ for $59\frac{2}{3}$ and, as we shall presently see, the existence of the subtractive principle is easily explained on other and more rational grounds.


Since the larger numbers were used by relatively few scholars, there was no compelling force of custom to standardize them. The variants in these cases are not of importance for our purposes, and simply a few of the numerals will serve to show their nature. These illustrations[1] date from *c.* 2400 B.C.



As mentioned later, the Babylonians also used a circle for zero, at least to the extent that they employed it to represent the absence of number, but it played little part in their system

[1] G. A. Barton, *Haverford College Library Collection of Cuneiform Tablets*, Part I, Philadelphia [1905]; Allotte de la Fuye, "En-e-tar-zi patési de Lagaš," in the *Hilprecht Anniversary Volume*, p. 121 (Chicago, 1909).

of notation. More commonly a circle simply stood for 10, particularly in the early inscriptions, as shown on page 38.



TABLET FROM NIPPUR

Contains divisors of 60^4, the quotients being in geometric progression. Date *c.* 2400 B.C. The top line reads $2(\cdot 60)$, $5(\cdot 60)$, and $12(\cdot 60)$. The left-hand figure is the original; the right-hand one is a drawing. Courtesy of the University of Pennsylvania

Chinese Numerals. The present forms of the Chinese numerals from 1 to 10 are as follows:

一 二 三 四 五

六 七 八 九 十

The number 789 may be written either from the top downwards or from left to right, as follows:

七 百 八 十 九

The second character in the number as written on the preceding line means hundred, and the forth character ten.

It will be observed that the figure for 4, probably four vertical marks in its original form, resembles the figure for 8 inclosed in a rectangle. On account of this the Chinese have given it the fanciful name of "eight in the mouth."[1]

Chinese merchants also use the following forms for figures from 1 to 10: 〡 〢 〣 𠂇 𠄎 𠃌 𠃑 𠄌 𠂆 十

and they have special symbols for 100, 1000, and 10,000, besides those in which a circle is used for zero.

These symbols are not the same as the ancient forms, but our knowledge of the latter is imperfect.[2] There are many variants of each of the characters given above, as when Ch'in Kiu-shao (1247) used 𠄌 for 5, and both 𠂆 and 𠄎 for 9.[3] The numerals on the early coins also show the variations that are found from time to time. In the second century B.C., for example, we find the 5 given in the so-called seal characters in the form 𠄌,—a form which was used for hundreds of years.[4]


Rod Numerals. There were also numerals represented by rods placed on the counting board,—a device which will be described in Chapter III. These numerals appear in the *Wu-ts'ao Suan-king*, which may have been written about the beginning of our era, or possibly much earlier, and are found

[1] L. Vanhée, in *T'oung-Pao*, reprint. On the general subject of the Chinese numerals in their historical development the standard work is that of F. H. Chalfant in the *Memoirs of the Carnegie Museum*, Vol. IV, No. 1, and Plate XXIX.

[2] We have, however, various records going back to the early part of the Christian Era. For example, the Metropolitan Museum, New York, has a land grant of 403 in which the forms of the numerals are almost the same as those now in use.

[3] Chalfant, *loc. cit.*; Y. Mikami, *The Development of Mathematics in China and Japan*, p. 73 (Leipzig, 1913); hereafter referred to as Mikami, *China*. On the general topic see S. W. Williams, *The Middle Kingdom*, New York, 1882; 1895 ed., I, 619; hereafter referred to as Williams, *Middle Kingdom*. J. Hager, *An Explanation of the Elementary Characters of the Chinese*, London, 1801; J. Legge, *The Chinese Classics*, 2d ed., I, 449 (Oxford, 1893).

[4] H. B. Morse, "Currency in China," *Journal of the North China Branch of the R. Asiat. Soc.*, Shanghai, reprint (n.d.). Valuable on the history of Chinese money and weights.




MONOGRAM FORMS OF CHINESE NUMERALS


The Japanese *sangi* were sticks used for representing numbers and were descended from the "bamboo rods" of the ancient Chinese. They gave rise to a *sangi* method of writing numbers. From a work by the Japanese mathematician Fujita Sadasuke (1779) in Chinese characters. The number in the first line at the top is 46,431

even as late as the 19th century. The oldest forms for the units are commonly arranged as follows:




In the tens' column the symbols usually appear as follows:




the arrangement thereafter alternating, the hundreds being like the units, and so on. Sometimes the 𝍠 was used instead of 𝍩 for 7 hundreds, and so on, and similarly for 8 and 9. By the common plan the number 7436 would appear as  . In this system the zero takes the form of a circle in the Sung Dynasty (950–1280), as is seen in a work of 1247,[1] where the subtraction 1,470,000 − 64,464 = 1,405,536 appears as



with two forms for 4.

These numerals were frequently written in the monogram form;[2] for example, 123,456,789 appears as  .

Hindu Numerals. The history of those Hindu-Arabic numerals which may have developed into our modern European forms is considered later. It should be said, however, that there are various other systems in use in India and neighboring countries. Of these the most interesting is the modern Sanskrit, the numerals being as follows:




These characters are evidently related to the early Brāhmī forms which are mentioned later.

[1] The *Su-shu Kiu-ch'ang* of Ch'in Kiu-shao. See A. Wylie, *Chinese Researches*, Pt. III, p. 159 seq. (Shanghai, 1897); L. Vanhée, in *T'oung-Pao*, reprint, thinks that the zero reached China from India somewhat earlier.

[2] A. Vissière, *Recherches sur l'origine de l'abaque chinois* (Paris, 1892), reprint from the *Bulletin de Géographie*; hereafter referred to as Vissière, *Abaque*.

Of the numerals of the same general character and in use in parts of Asia adjacent to India, the following are types:



Their history has no particular significance, however, in a work of this nature, since the forms are local and are relatively modern.[1]


An American Place Value. When Francisco de Cordoba landed the first Spanish expedition on the coast of Yucatan, in 1517, he found the relics of a highly developed civilization, that of the Maya, which had received its deathblow in the wars of the preceding century.[2] Within a few years after the European invasion the independence of the Maya was completely lost. In 1565 Diego de Landa, bishop of Merida, in northern Yucatan, wrote a history of these people,[3] so that our knowledge of their achievements goes back to about the beginning of the period of European influence. They had an elaborate calendar before the Spaniards arrived, and capable investigators have asserted that the Maya cycle began as early as 3373 B.C.

[1] One of the best general works on Eastern notation is that of A. P. Pihan, *Exposé des Signes de Numération usités chez les Peuples Orientaux Anciens et Modernes*, Paris, 1860, with many tables.

[2] S. G. Morley, *An Introduction to the Study of the Maya Hieroglyphs*, p. 6 (Washington, 1915). This work should be consulted for details respecting this entire topic. Authorities vary as to the plural form of Maya, some giving Mayas and others Maya. See also C. Thomas, "Numeral Systems of Mexico and Central America," *Annual Report of the Bureau of American Ethnology*, XIX (1897–1898), 853 (Washington, 1900).


[3] In his *Relacion de las Cosas de Yucatan*, a work not printed until 1864.

The Maya counted essentially on a scale of 20, using for their basal numerals two elements, a dot (•) and a dash (—), the former representing one and the latter five. The first nineteen numerals were as follows, reading from left to right:



There were numerous variants of these forms,[1] but these offer no special peculiarities which we need consider.

The most important feature of their system was their zero, the character  , which also had numerous variants. Since their scale was vigesimal, they wrote 20 as we write 10, using their characters for 1 and zero.[2] The following table shows the general plan that was used when they wrote on flexible material:

Numerals					
Our forms	1 0	1 17	15·20 0	360 0 0	19·360 13·20 13
Values	20	37	300	360	7113

We see here a fairly well developed place value, the lowest order being units from 1 to 19, the next being 20's from 1 · 20

[1] See Morley, *loc. cit.*, p. 89, which should be consulted for a description of the system.

[2] Their special hieroglyphic for 20, used for certain purposes, need not concern us. On the word "hieroglyphic" see page 45, note 2.

to 17 · 20; the next being 360's from 1 · 360 to 19 · 360; the next being 7200's, and so on,[1] representing a very satisfactory system. There is no evidence in any extant record that it was used for purposes of computation, its use in the texts being merely to express the time elapsing between dates. The fact, however, that the pebble and rod are apparently the basal elements in the writing of numbers leads us to feel that we have in these numerals clear evidence of the early use of an abacus. If, as many ethnologists believe, there is a connection between the Japanese and certain of our primitive Americans, the use of the rods may be traced back to Asia.



EARLY FORMS OF COMMON EGYPTIAN NUMERALS

From a piece of pottery of the First Dynasty, *c.* 3400 B.C. The symbols for 10 and 100 are repeated several times

Egyptian Numerals. The Egyptians had four materials upon which they could conveniently record events. One of these was stone, a medium supplied by the quarries along certain parts of the Nile. Another medium was papyrus, a kind of paper made from strips of the pulp of a water reed which was apparently more common at one time than it is at present. The other two common materials were wood and pieces of pottery. Leather does not seem to have been so commonly used as in other countries.


In writing on stone the Egyptians took time for the work and made their characters with great care. These characters are called hieroglyphics.[2] The hieroglyphic characters were commonly written from right to left, but also from left to right.

[1]For complete description see Morley, *loc. cit.*, pp. 129–133.

[2]The sacred inscriptions; from the Greek *ἱερός* (*hieros'*), sacred + *γλύφειν* (*gly'phein*), to carve.

In the earlier inscriptions they are often written from the top down. This accounts for the various ways of writing the simple numerals, a character often being found facing in different directions. For our present purposes it suffices to give the ordinary form of hieroglyphic numerals,[1] as follows:

1	2	3	4	5	6	7	8	9	10
11	12	20	40	70	100	200	1000	10,000	



EGYPTIAN NUMERALS

Numerals reading from left to right. From the walls of a temple at Luxor

[1] These are as given in A. Eisenlohr, *Ein mathematisches Handbuch der alten Aegypter*, 2d. ed., table following p. 8 (Leipzig, 1877); hereafter referred to as Eisenlohr, *Ahmes Papyrus*. See also Peet, *Rhind Papyrus*; J. De Morgan, *L'Humanité Préhistorique*, p. 115 (Paris, 1921).

There were higher numerals, but the above will serve to show the general nature of the characters employed.


While the hieroglyphic forms were used in writing inscriptions on stone and in elaborate treatises on papyrus, other forms were early developed for rapid writing on papyrus, wood, and pieces of pottery. There were two forms of this writing, the hieratic (religious) and the demotic (popular). The former was a cursive script derived from the hieroglyphic, and the latter was a somewhat later form of the hieratic, beginning in the 7th century B.C. After the demotic forms came into general use, the hieratic was reserved for religious purposes.



HIEROGLYPHIC FOR 6000 (ABOUT 500 B.C.)

The meaning is, "The Falcon King led captive 6000 men of the Land of the Harpoon Lake," there being a harpoon just below this in the original inscription

The hieratic writing usually proceeded from right to left, although in early times it is occasionally found running from the top down. The numerals to 10 were of the following forms:[1]



1 2 3 4 5 6 7 8 9 10

The demotic forms offer no peculiarities of special interest.[2]

Greek Numerals. The first numeral forms of the Greeks seem to have been such upright strokes as were used in all Mediterranean countries, and perhaps represented the fingers. These strokes were repeated as far as the needs of the primitive inhabitants required. For example, in a stele from Corinth, of about the 5th century B.C., there is the numeral ||||||||, referring


[1] These are taken from the Ebers Papyrus as copied by Eisenlohr. Naturally the forms varied with different scribes.

[2] For a careful study of these forms, with numerous facsimiles, see H. Brugsch, *Numerorum apud Veteres Ægyptios Demoticorum Doctrina*, Berlin, 1849.




EARLY CYPRIOTE NUMERALS

From a fragment of a temple record found on the island of Cyprus. In the last two lines the numeral for 6 (||| |||) appears twice. Courtesy of the Metropolitan Museum of Art, New York



EARLY CYPRIOTE NUMERALS


The lower part of the fragment shown above. The numerals are the same as those on the tablets found at Knossos, Crete, where ◇ is used for 1000, ○ for 100, — for 10, and | for 1. The number 4 (||||) is in the first line and the number 14 (||||—) in the line next to the last. The Phœnicians also used these symbols for ten and one. Courtesy of the Metropolitan Museum of Art, New York



EARLY CYPRIOTE NUMERALS

From a fragment of a receptacle in a sanctuary. The inscription reads, "Zeus's portion of wine is three measures." Courtesy of the Metropolitan Museum of Art

to a fine of eight obols for intruding on certain property.[1] Inscriptions illustrating this usage are found not only in Greece but in various islands of the eastern Mediterranean Sea, as shown in the illustrations of monumental records from Cyprus. By the time Greece had reached the period of her intellectual ascendancy there had developed a system of numerals formed from initial letters of number names. These forms appear in records of the third century B.C., and were probably in use much earlier, although the custom of writing large numbers in words seems to have been general. Many generations later the system was described so fully by Herodianus, a prominent grammarian of the latter part of the second century, that the symbols were thereafter known as Herodianic numerals, although this name has no worthy sanction. In recent times they have been known as Attic numerals, since they are the only pre-Christian number forms found in Attic inscriptions. The system is also known as the acrophonic (initial) system, the initials



GREEK NUMERALS OF THE PTOLEMAIC PERIOD

On an icosahedral die of the Ptolemaic Period in Alexandria, just before the Christian Era. Such dice are occasionally found, usually made of basalt or quartz. This one is basalt, whitened for the purpose of photographing. From the author's collection

[1] *American Journal of Archæology* (1919), p. 353.

of the several number names, as of πέντε (*pen'te*), five, being used, singly or in combination, in the following manner:

Γ, an old form for Π, the letter *pi*, initial of ΠΕΝΤΕ (*pen'te*), five, used as a numeral for 5;
Δ, the capital *del'ta*, initial of ΔΕΚΑ (*dek'a*), ten, used as a numeral for 10; it is often written like ○ in the Greek papyri, and an inscription at Argos has ⊙;
H, the old Attic breathing, like our *h*, later represented by ', initial of ΗΕΚΑΤΟΝ (*hekaton'*), hundred;
X, the capital *chi*, initial of ΧΙΛΙΟΙ (*chil'ioi*), thousand;
M, the capital *mu*, initial of ΜΥΡΙΟΙ (*myr'ioi*), ten thousand.

These numerals were frequently combined, thus:

ⓟ or ⓟ, *pente-deka*, was used for 50;
ⓗ, *pente-hekaton*, was used for 500;

and so on for other numbers.

The forms of the letters varied in different cities and states of Greece, but the variants need not concern us in this description.[1]

The following will show how the characters were used:

ΓΙ = 6 ⓗHHH = 800
ΔΓIIII = 19 MMMM = 40,000
ⓟΔΔΔΔ = 90 ⓜ = 50,000

but in the manuscripts the forms vary so much as often to be exceedingly difficult to decipher.

[1] G. Friedlein, *Die Zahlzeichen und das elementare Rechnen der Griechen und Römer*, Erlangen, 1869; F. G. Kenyon, *Palæography of Greek Papyri*, Oxford, 1899; E. S. Roberts, *Greek Epigraphy*, p. 96 (Cambridge, 1887); J. Gow, "The Greek Numeral Alphabet," *Journal of Philology* (1884), p. 278; S. Reinach, *Traité d'Épigraphie Grecque*, pp. 216, 218 (Paris, 1885); J. P. Mahaffy, "On the Numerical Symbols used by the Greek Historians," *Trans. of the Royal Soc. of Literature*, XXVII (2), 160; Heath, *History*, I, 29. The best modern treatment is that of M. N. Tod, "Three Greek Numeral Systems," *Journal of Hellenic Studies*, XXXIII, 27, and "The Greek Numeral Notation," *Annual of the British School at Athens*, XVIII, 98. On the numerals of Crete see Sir A. J. Evans, *The Palace of Minos*, p. 279 (London, 1921), and *Scripta Minoa*, p. 258 (Oxford, 1909).

To these may be added the following characters related to numerical work:

T = talent and also $\frac{1}{4}$ obol
⊦ = drachma
I = obol, with Ↄ or C for $\frac{1}{2}$ obol
ⱷ = stater, so that ⱷⱷⱷⱷⱷⱷⱷ = 7 staters
and HHΔΔΔΔΓⱷⱷⱷ = 248 staters
Γ⁻ = 5 talents, ⍑ = 10 talents, H = 100 talents

Contemporary with the development of the Ionic alphabet we find numerical values assigned to the letters, somewhat as we use letters to number the rows of seats in an assembly room. The oldest forms that we have are substantially as follows:

A = 1	H = 7	N = 13	T = 19
B = 2	Θ = 8	Ξ = 14	Y = 20
Γ = 3	I = 9	O = 15	Φ = 21
Δ = 4	K = 10	Π = 16	X = 22
E = 5	Λ = 11	P = 17	Ψ = 23
⊥ = 6	M = 12	ⱷ = 18	Ω = 24[1]

These were used very early, but the system was manifestly of no value for computation. A more refined alphabetic system appeared at least as early as the third century B.C., running parallel with the more primitive systems.

As seen above, the Greeks had twenty-four letters in their common Ionic alphabet, but for a more satisfactory system of numerals they needed twenty-seven letters. They therefore added the three forms F or Ⅎ (the old digamma), ϙ or sometimes Ϙ (the Phœnician *koph*), and ϡ[2] (perhaps the Phœnician

[1] S. Reinach, *loc. cit.*, p. 220.

[2] A modern name for the character is *sampi* (*σαν+πι, san'pi*), suggested because of its resemblance to π in its 15th century form. The form in the 2d century was ⋔, and it may go back to the ⊤ (s), which was used from the 5th to the 2d century B.C. See Roberts, *loc. cit.*, p. 10.

shin or tsadé), after which they arranged their system as follows:

Units	Α	Β	Γ	Δ	Ε	F	Ζ	Η	Θ
	1	2	3	4	5	6	7	8	9
Tens	Ι	Κ	Λ	Μ	Ν	Ξ	Ο	Π	Ϙ
	10	20	30	40	50	60	70	80	90
Hundreds	Ρ	Ϛ	Τ	Υ	Φ	Χ	Ψ	Ω	Ϡ
	100	200	300	400	500	600	700	800	900

To distinguish the numerals from letters, a bar was commonly written over each number, as in the case of $\bar{\text{Α}}$, although in the Middle Ages the letter was occasionally written as if lying on its side, as in the case of ≺.[1]

The capital forms were used, the small letters being an invention of a much later period. In a manuscript of the 10th century in Göttingen the small letters are found, and there are no accents when these numerals appear in tables. When, however, they appear in the text, there are bars superscribed to distinguish the numerals from words, thus: $\bar{\alpha}$, $\bar{\epsilon}$, $\overline{\theta}$, $\overline{\iota\theta}$, etc. In modern books the forms usually appear as α', β', γ', δ', and so on, the accents being used to distinguish the numerals from letters. The thousands were often indicated by placing a bar to the left, thus:

/Α, /Β, /Γ, ··· for 1000, 2000, 3000, ···,

these appearing in modern Greek type as $,\alpha$, $,\beta$, $,\gamma$, ···.

The myriads, or ten thousands (*μύριοι, myr'ioi*), were represented by such forms as the following:

Μ or $\overset{\gamma}{\text{M}}$, 10,000; $\overset{\beta}{\text{M}}$, 20,000; $\overset{\Gamma}{\text{M}}$, 30,000, and so on.

In late Greek manuscripts the symbol ◠ was used for myriad, as in the case of $\overset{\frown}{\text{Ι}}\overset{\frown}{\Delta}$ for 14 myriads (140,000). We also find such forms as $\overset{\epsilon}{\frown}$ for 5 myriads (50,000).[2]

[1] V. Gardthausen, *Die Schrift, Unterschriften und Chronologie im Byzantinischen Mittelalter*, 2d ed., p. 360 (Leipzig, 1913); hereafter referred to as Gardthausen, *Die Schrift*. See also F. E. Robbins, "A Greco-Egyptian Mathematical Papyrus," *Classical Philology*, XVIII, 328.

[2] Gardthausen, *Die Schrift*, p. 371.

In the early Christian period the three lines of letters representing units, tens, and hundreds respectively were called verses or rows, and the rectangular arrangement of the figures in these verses was probably of some value in computation.[1]

Hebrew Numerals. The Jewish scholars used the letters of their alphabet for numeral symbols in the same way as the Greeks did. We find this usage well established in the Maccabean period (2d century B.C.), but it is probably of an earlier date. In the Talmud the numbers above 400 are formed by composition, 500 being formed of the symbols for 400 and 100,[2] and 900 being a combination of the symbols for 400, 400, and 100.[3] Later writers, however, followed a plan introduced by the Massoretes,[4] in which certain final forms of letters were used for the hundreds above 400. These numeral forms as now recognized are as follows:

א	ב	ג	ד	ה	ו	ז	ח	ט
1	2	3	4	5	6	7	8	9
י	כ	ל	מ	נ	ס	ע	פ	צ
10	20	30	40	50	60	70	80	90
ק	ר	ש	ת	ך	ם	ן	ף	ץ
100	200	300	400	500	600	700	800	900

The thousands were represented by the same letters as the units. Since the number 15 would naturally be represented by 10 and 5, read from right to left, that is, by יה, and since these are the first two letters of the word יהוה (Jhvh, Jahveh, Jehovah), the Hebrews wrote 9 + 6 (טו) instead.

[1] "Primus igitur versus est a monade usque ad enneadem," etc. (Capella, VII, 745). Favonius Eulogius (*c.* 400) remarks: "Primi versus absolutio novenario numero continetur." See J. G. Smyly, in "Mélange Nicole," *Recueil de Mémoires de Philologie Class. et d'Archéol.*, p. 514 (Geneva, 1905).

[2] תק. [3] תתק.

[4] The scholars engaged in the work of *Massorah*, the establishing of the traditional pronunciation and accents of the Hebrew scriptures. The work extended over a long period, closing in the 10th century. See *Jewish Encyclopedia*, IX, 348 (New York, 1905). For the zero, see Smith-Karpinski, p. 60.

Gematria. The fact that the letters of various ancient alphabets had numerical values, and hence were used in computation, led to the formation of a mystic pseudo-science known as gematria, which was very popular among the Hebrews as well as among other peoples.

Although it had many modifications, its general nature may be explained by saying that the numerical value of a name could be considered instead of the name itself. If two names had the same numerical value, this fact showed some relation between the individuals. It is probable that 666, "the number of the beast" in Revelations, was the numerical value of some name, this name being known to those who were in the secret, but being now lost. It is not improbable that it referred to "Nero Cæsar," which name has this value when written in Hebrew. For nearly two thousand years attempts have been made to relate the number to different individuals, particularly to those of a religious faith differing from that of the one suggesting the relationship. Thus, it has been assigned to various popes, to Luther, and to Mohammed; but it has also been related to statesmen, to the Latin Church, and to various other classes and organizations. In some cases a man's name and its gematria number have both appeared upon his tombstone. An interesting illustration of gematria is also found in our word "amen." Written in Greek, the numerical values of the letters are as follows: A(α) = 1, M(μ) = 40, H(η) = 8, N(ν) = 50, the total being 99. On this account we find in certain Christian manuscripts the number 99 written at the end of a prayer to signify "amen."[1]

Roman Numerals. The theories of the origin of the Roman numerals are for the most part untenable. Priscian (6th century) believed that "I" was used for 1 because it was the initial of the Greek *ἴα*, a dialectic Greek word for unity,[2] although long before the Greeks had any written language it was used for this purpose in Egypt, Babylon, and various

[1] Gardthausen, *Die Schrift*, p. 309.

[2] See the 1527 (Venice) edition of Priscian, fol. 271, *r*. For the feminine of *εἷς* (*heis*) the Æolic Greeks used *ἴα*; the other Greeks, *μία*. Homer used both forms.

other parts of the ancient world. His other theories were equally unscientific except in the cases of C and M. These symbols he took to be the initials of *centum* (hundred) and *mille* (thousand), and there was enough historical evidence for the late adoption of these letters as symbols for 100 and 1000 to justify him in making this statement. There are also various theories connected with stick-laying, but for these there is no historic sanction.

In the 16th century Mattheus Hostus[1] asserted that the theory of the early grammarians *frivolum est*; and while his own theories were generally about as frivolous, he made the plausible suggestion that the V was derived from the open hand, the fingers with the exception of the thumb being held together. This led naturally to taking the X as a double V,—a view held by various later writers and receiving powerful support from Mommsen (1850), the great German authority on Latin history and epigraphy. The theory is not inconsistent with the fact that the V is occasionally inverted (Λ), since this form, although an early one, may have developed relatively late with respect to X and may thus have represented half of that numeral.

Mommsen's most important suggestion, however, was that C and M are not primitive forms but are late modifications of such forms, influenced by the initials of *centum* and *mille*. The primitive forms for 50, 100, and 1000 he stated to be the Greek aspirates X (*chi*), from which ⅃ was derived; Θ (*theta*), from which comes the C; and Φ (*phi*), which is the origin of the M. As to this theory there is positive evidence that one of the earliest forms for X (*chi*) was ↓, and this, with the later forms ⇃, ⊥, and ⅃, was used for 50 in the inscriptions of about the beginning of our era.

As to the use of Θ for 100, we have also the early forms ⊗, ⊕, ⊙, and ⊖. If the last of these were written rapidly with a stylus or a reed pen, the result might easily resemble C. We have not, however, any of these transition forms extant, although by analogy with ⅃ and M we might well accept this theory.

[1] *De numeratione emendata veteribus Latinis et Graecis usitata*, Antwerp, 1582.

The Φ was also written Ⅾ, and the symbol for 1000 is very commonly given on the ancient monuments as ↀ, ↀ, ↀ, and the like, so that this part of the theory is reasonable. The M as a numeral is unusual on the older monuments, although an expression like $\overline{\text{II}}$M for 2000, where M evidently stands for the word *mille*, is not uncommon. Generally the Romans used one of the modifications of Φ as stated above, or the symbol ∞, which is probably a cursive form of ↀ, with numerous variants such as ⋈ and ∼.

As to the X for 10, there is the further theory that it may have come from the crossing off of ten single strokes for 1 by a *decussare* line, either as IIIIIIIIII or as IIIIIIIII, which was abbreviated as X.[1] This is analogous to the possible Egyptian plan of grouping ten strokes by an arc and thus obtaining their symbol ∩. There is much to commend this *decussare* theory, for 20 was commonly written ⋔ or ⋕, and similarly for 30 and 40. If this is the origin of the X for 10, then the V and Λ were naturally taken as halves of X. On the whole, this seems quite as probable as the hand theory. It has also been thought that X represents the crossed hands, thus giving two fives.

In 1887 Karl Zangemeister[2] advanced the theory that the entire system was based on the single *decussare* principle. Briefly, a crossing line multiplies any number by ten. Hence we have I and X for 1 and 10 respectively; X and ⋈ for 10 and 100, from the latter of which the X finally dropped out, leaving (, which became our C under the influence of the initial letter for *centum*; and ⋈ for 1000, which finally became the common ∞. Although the theory is interesting, it has never been generally accepted by Latin epigraphists, and so we at present fall back on the Mommsen theory as the most probable of any thus far suggested. It is quite as reasonable, however, to believe that the symbols were arbitrary inventions of the priests.

[1] *Decussare* is the verb form. The word also appears as *decussatio*, *decussatim*, and *decussis*, according to the sentence construction.

[2] "Entstehung der römischen Zahlzeichen," *Sitzungsberichte der Königl. Preuss. Akad. der Wissensch.*, XLIX, 1011, with a bibliography on page 1013.

An examination of the many thousand inscriptions collected in the *Corpus Inscriptionum Latinarum*[1] fails to solve the problem of origin, but it shows the change in forms from century to century. This change is even more marked in the medieval manuscripts. The following brief notes will serve to show how these numerals have varied.

The I is always a vertical stroke, or substantially so. Horizontal strokes are used in writing certain fractions. In late medieval manuscripts the stroke appears as i or, as a final letter, j.

The V also appears on the early monuments as U or Λ, and is frequently found in such contracted forms as ⋋ for 15. In the medieval manuscripts it varies with the style of writing, appearing as V, v, U, and u. In the late Roman times the character ꝙ, with numerous variants, was used for 6, possibly from the Greek numeral. To represent eight, for example, this character was combined with II.[2]

The X also appears on the monuments as ⋌ or ⋋. It is frequently combined with other letters in such forms as ⋈ for 70. In the medieval manuscripts it is often written as a small letter.

The L very frequently appears on the monuments of about the beginning of our era in the older forms of ↓, ⊥, and ⊥. In the Middle Ages it often appears as a small letter, as in a case like Clxviij for 168.

The C has changed less than the other forms, appearing on the ancient monuments as a capital and frequently in the later manuscripts as a small letter.

The D is generally thought to be merely half of the CIↃ which stood for thousands. It is occasionally written CI and appears very commonly as IↃ even after the beginning of printing. In the Middle Ages it appears both as a capital and as a small letter. There is a possibility that the use of D to represent 500 is due to the fact that the Etruscans had no such letter in their early alphabet, and consequently took the Δ (*delta*) for this purpose, just as they took other

[1] Berlin, 1863 seq.

[2] L. A. Chassant, *Dictionnaire des Abréviations . . . du Moyen Age*, p. 114 (Paris, 1884).

Greek letters for numerical purposes. The *delta* was then changed, in the course of time, to the form with which we are now familiar.[1]

When the Romans used the M in representing numbers, it was commonly as the initial of *mille*, thousand. When written with other numerals, the thousand symbol was usually CIↃ, ↀ, cIↃ, ↀ, ∞, ⋈, ∼, or some similar form, as in the case of ∞CIII for 1103. In the medieval manuscripts the M, usually a capital, replaced the earlier forms, as in the number Mcccclxxxxiiij for 1494.

The subtractive principle is found in certain cases like that of IV for 4, that is, $5 - 1$. This principle was, as we have seen, used by the Babylonians in the 3d millennium B.C. It was also used by the Hebrews, at least in word forms, but apparently not before the Etruscans and Romans used it. The Etruscans[2] preceded the Romans in recognizing the principle and made a more extensive use of it. They commonly wrote their numerals from right to left, and so we have such forms as the following:[3]

XIIIXX, for $20 + (10 - 3)$, or 27;
XIIXXX, for $30 + (10 - 2)$, or 38;
↑III, for $50 - 3$, or 47;
IIↁX, for $(50 - 10) + 2$, or 42.

The Etruscans also used ↓ for X, and so we find such forms as ↓⅃ for XL; when read from right to left this means our LX (60), but when written ⅃↓ it means our XL (40).[4] Such forms as XIↁ[5] for the Roman LIX are also found.

Subtractive Principle Widespread. The subtractive principle was probably used by various other early peoples, for an immature mind finds it easier to count backwards by one or two from some fixed standard, like 5, 10, 15, 20, and so on, than to count forwards by three or four. Thus, the Romans found it easier to think of "two from twenty" (*duo de viginti*) than of

[1] B. Lefebvre, *Notes d'Histoire des Mathématiques*, p. 30 (Louvain, 1920).
[2] R. Brown, "The Etruscan Numerals," *Archæological Rev.*, July, 1889.
[3] W. Corssen, *Ueber die Sprache der Etrusker*, 2 vols., Leipzig, 1874, 1875. *Corpus Inscriptionum Etruscarum*, I, Nos. 23, 27, 32, 38, *et passim* (Leipzig, 1893–).
[4] *Ibid.*, 4615.

"eight and ten" (*octodecim*), and of "one from twenty" than of "nine and ten." This is especially the case with numbers above five, since the difficulty is hardly experienced until nine or fourteen is reached.

As an indication of the tendency of primitive peoples to use the subtractive principle the fact may be mentioned that the Zuñi Indians, whose number names refer to the fingers, speak of four as "all the fingers almost complete," and of nine as "almost all are held up with the rest," each containing the idea of subtraction.[1] They had a system of knot numerals which involved the same principle. A medium knot indicated 5, and this with a small knot before it indicated $5 - 1$, whereas if the small knot came after the medium one the number was $5 + 1$. Similarly, a large knot indicated 10, and a small knot was used either before or after it so as to indicate 9 or 11 respectively.

Further Cases of the Subtractive Principle. It is because of the fact that the difficulty is not evident with so simple a number as 4 that the Romans did not commonly use the subtractive principle in this case, preferring the form IIII to the form IV. They used the principle more frequently in the case of 9, but even here they wrote VIIII oftener than IX. In the case of 400 they usually wrote CCCC, but occasionally they used CD. Even as late as the 16th century we often find a number like 1549 written in some such form as Mcccccxxxxviiij.

Relics of the subtractive principle are seen in our tendency to say "ten minutes of (or "to") six" instead of "fifty minutes past five," and to say "a quarter of (or "to") six" rather than "three quarters of an hour past five."

There is a possibility that the Romans avoided IV, the initials of IVPITER, just as the Hebrews avoided יה in writing 15, as the Babylonians avoided their natural form for 19, and as similar instances of reverence for or fear of deity occur in other languages.

[1] F. H. Cushing, "Manual Concepts," *American Anthropologist* (1892), p. 289; Th. W. Danzel, *Die Anfänge der Schrift*, p. 55 (Leipzig, 1912); L. L. Conant, *Number Concept*, p. 48 (New York, 1896).

Even when the subtractive principle was used, no fixed standard was recognized. The number 19 was commonly written XIX, but not infrequently IXX.[1] We also find IIX for 8 and IIXX for 18, but these were not so common. It is quite rare to find CD for 400 or CM for 900, and forms like MCM and DCD were never used in ancient or medieval times. In general, therefore, it may be said that the Romans recognized the subtractive principle but did not make much use of it.

Occasionally this principle was used with the fraction $\frac{1}{2}$, for which the Romans wrote the letter S, initial of *semis* (half). Thus we find SXC for $89\frac{1}{2}$ and SXXC for $79\frac{1}{2}$.

Large Numbers. The Romans had relatively little need for large numbers, and so they developed no general system for writing them. The current belief that they commonly used a bar, or *vinculum*, over a number to multiply it by 1000 is erroneous. What they ordinarily did, if they used numerical symbols at all, was to take some such forms as the following:

For 100,000: CCCIↃↃↃ

For 10,000: CCIↃↃ

For 5,000: IↃↃ

To represent larger numbers, these forms were repeated. Thus, the symbol , used for 100,000, is repeated twenty-three times on the *columna rostrata*,[2] making 2,300,000.


In the Middle Ages, however, we find such forms as |X| or IXI for million and |M| for hundred million, that is, for ten hundred thousand and one thousand hundred thousand.

Use of the Bar. The Romans commonly placed a bar over a number to distinguish it from a word, as in the case of $\overline{\text{II}}$VIR for *duumviri* (two men) and $\overline{\text{III}}$VIR for the triumvirate. The

[1] In early inscriptions this form was sometimes used for 21, since the Romans occasionally wrote numbers from right to left, like the early Greeks.

[2] A Roman monument set up in the Forum to commemorate the victory of 260 B.C. over the Carthaginians. This is the earliest noteworthy example of the use of large numbers in a Roman inscription.

oldest example that we have of the bar to indicate thousands dates from about 50 B.C. Cicero (106–43 B.C.), or, more probably, some late copyist of his works, also used $\overline{XX}$.CD and CCIↃↃ CCIↃↃ CCCC as equivalent. The *vinculum* is found frequently in the works of Pliny (1st century), but it is not used




ROMAN NUMERALS

From the work of Freigius, a Swiss writer, published in 1582

with much uniformity and we are not sure how many of his numeral forms are due to later scribes. In the Middle Ages the *vinculum* was called a *titulus*,[1] but even then it was more commonly used to distinguish numerals from words than to indicate thousands.

[1] Thus Bernelinus: "Nam sicut prima unitas notatur per elementum I, ita millenarius primus per idem I, superaddito tantum titulo." A. Olleris, *Œuvres de Gerbert*, p. 360 (Paris, 1867).

The Romans did not use the double bar to indicate 1000 × 1000, as in $\overline{\overline{V}}$ for 5,000,000, but it is said to be occasionally seen in the late Middle Ages.[1]



ROMAN NUMERALS


From Bongo's work on the mystery of numbers, Bergamo, 1584–1585

Late Coefficient Method. In the later Roman times there arose a kind of coefficient method of representing large numbers. Thus, Pliny used XIIM for 12,000, and we have a relic of this method in our modern use of 10 M. In such cases, however, M was looked upon as abbreviation for *mille* rather than as a symbol for 1000, although the distinction is, of course, not noticeable. We find the same thing in the Middle Ages, as when O'Creat (*c.* 1150) writes $X^{es}M.$ milia for ten thousand thousand. A somewhat similar usage appears in the *Compotus Reinheri* (13th century), where $IIII^{or}$ milia. $ccc^{a}.l.vi$ appears for 4356. Even as late as the 16th century the same plan was followed, as when Noviomagus (1539) wrote IIIM for 3000 and MM for 1,000,000, and when Robert Recorde (*c.* 1542) used vj.C for 600, ixM for 9000, CCC.M for 300,000, and 230 M, MM, M for $230 \cdot 10^{12}$. The coefficients were often written above the M, as in $\overset{lxxx}{MM}$ for 80,000,000 and $\overset{C}{MM}$ for 100,000,000, in a manu-

[1] A. Cappelli, *Dizionario di Abbreviature*, 2d ed., p. lii (Milan, 1912).

script of *c.* 1442.[1] They were also written below, as in $\underset{I}{C}xxiij$ for 123 and $\underset{I}{C}\ \underset{xxiij}{MM}\ \underset{iiij}{C}\ \underset{lvj}{M}\ \underset{vij}{C}\ \underset{lxxxix}{\ }$ for 123,456,789, in the arithmetic of Bartjens.[2]

Epigraphical Difficulties. The Romans varied their numerals, often according to the pleasure of the writers, and it takes a skilled epigraphist to decipher many of those that appear upon the amphoræ stating the amount or the price of wine. For example, the following numbers, which were taken from wine jugs of about the 1st century, and which are by no means among the most difficult, would certainly not be understood by the casual observer:



for ∞ CCC = 1300

for IXCS = $89\frac{1}{2}$

Such forms concern chiefly the student of epigraphy, however. The medieval numerals are more interesting, since they involve new methods, and hence a few types will here be given:

$\overline{c} \cdot \overline{lxiiij} \cdot ccc \cdot l \cdot i$, for 164,351, Adelard of Bath (*c.* 1120).

$\overline{vi} \cdot dclxvi$, for 6666, Radulph of Laon (*c.* 1125).

II.DCCC.XIIII, for 2814, Jordanus Nemorarius (*c.* 1225).

MↃCLVI, for 1656, a monument in San Marco, Venice.

ↃIↄ. Iↄ. Iↄ, for 1599, edition of Capella, Leyden, 1599.

xxyiii, for 28, edition of Horace, Venice, 1520.

$IIII_{xx}$ et huit, for 88, a Paris treaty of 1388.[3]

ⅭↃDCXL, for 1640, edition of Petrus Servius, Rome, 1640.

four Cli.M, two Cxxxiiii, millions, sixe ClxxviiiM. fiue Clxvii, for 451,234,678,567, Baker, 1568.

[1] A copy of Sacrobosco's arithmetic made *c.* 1442. See *Rara Arithmetica*, p. 450.

[2] A Dutch work of the 18th century, 1792 ed., p. 8.

[3] This is simply the French *quatre vingt* (4 × 20) and is common in medieval French MSS.

To the many other peculiarities of this system it is not possible to allow further space. The Roman forms persisted in use, especially outside of Italy, until printed arithmetics made our common numerals widely known. Even at the present time the fishermen of Chioggia, near Venice, use forms that closely resemble those of the early Etruscans, so persistent is custom in the humbler occupations of man.[1]

Our Common Notation. When we come to consider the origin of our common numerals, we are confronted by various theories, and the uncertainty is quite as marked as in the case of the Roman system. These symbols are generally believed to have originated in India, to have been carried to Bagdad in the 8th century, and thence to have found their way to Europe.[2] This is not certain, for various authors of scientific standing have attempted to show that these numerals did not originate in India at all,[3] but the evidence still seems much more favorable to the Hindu origin than to any other that has been suggested. The controversy has recently centered about the meaning of the word *hindasi*, which is often used by the Arabs in speaking of the numerals. It is asserted that the word does not mean Hindu, some claiming that it refers to Persia and others that it means that which is related to calculation. It is difficult, however, to explain away the following words of Severus Sebokht (*c.* 650), written in the 7th century and already quoted in Volume I:

I will omit all discussion of the science of the Hindus, a people not the same as the Syrians; their subtle discoveries in this science


[1] A. P. Ninni, "Sui segni prealfabetici usati . . . nella numerazione scritta dai pescatori Clodiensi," *Atti del R. Istituto Veneto delle sci. lett. ed arti*, VI (6), 679.

[2] Smith and Karpinski, *The Hindu-Arabic Numerals*, with bibliography, Boston, 1911 (hereafter referred to as Smith-Karpinski); G. F. Hill, *The Development of Arabic Numerals in Europe*, Oxford, 1915; J. A. Decourdemanche, "Sur la filiation des chiffres européens modernes et des chiffres modernes des Arabes," *Revue d'Ethnographie et de Sociologie*, Paris, 1912; G. Oppert, "Ueber d. Ursprung der Null," *Zeitschrift für Ethnographie*, XXXII, 122 (Berlin, 1900); G. N. Banerjee, *Hellenism in Ancient India*, p. 202 (Calcutta, 1919).

[3] *E.g.*, see Carra de Vaux, "Sur l'origine des chiffres," *Scientia*, XXI (1917), 273; but see F. Cajori, "The Controversy on the Origin of our Numerals," *The Scientific Monthly*, IX, 458.


of astronomy, discoveries that are more ingenious than those of the Greeks and the Babylonians; their valuable methods of calculation; and their computing that surpasses description. I wish only to say that this computation is done by means of nine signs.

Types of Early Hindu Numerals. The early numerals of India were of various types.[1] The earliest known forms are found in the inscriptions of King Aśoka, the great patron of Buddhism, who reigned over most of India in the 3d century B.C. These symbols are not uniform, the characters varying to meet the linguistic conditions in different parts of the country. The Karoṣṭhī forms, for example, are merely vertical marks, I II [III] IIII IIIII, and are not particularly significant. The Brāhmī characters found in some of these inscriptions are of greater interest. The only numerals thus far found in the Aśoka edicts are as follows:




1 2 4 6 50 50 200 200 200

The Nānā Ghāt Inscriptions. About a century after the Aśoka edicts certain records were inscribed on the walls of a cave on the top of the Nānā Ghāt hill, about seventy-five miles from the city of Poona. A portion of the inscriptions is as follows:




[1] For discussion and bibliography see Smith-Karpinski, p. 19.

The probable number forms contained in these inscriptions are as follows:




The next important trace of the numerals is found in the caves at Nasik, India. These are of the 1st or 2d century and are as follows:



The significant feature of these numerals is that they clearly resemble the Nānā Ghāt forms, and that in both we seem to have the progenitors of our present numerals.

It should be understood, however, that the interpretations of the inscriptions at Nānā Ghāt and Nasik are not universally accepted. All that we can say, in our present state of knowledge, is that these are the probable number forms as stated, that they resemble some of the numerals that were transmitted to Europe as of Hindu origin, that no zero appears in these early inscriptions, and hence that the place value, as we know it at present, had not yet been developed.

Variants of Hindu Forms. The variants of the Hindu forms preceding the invention of the zero may be seen in the table shown below.



As to the original significance of these forms we are wholly ignorant except in the cases of the first three. As to I, II, III, or —, =, ≡, there is, of course, no question. The vertical forms may have represented fingers used in counting or they may have been the marks that one naturally makes with a stylus or brush in keeping a numerical record. The horizontal forms may be pictures of computing sticks, like those which the Chinese used in remote times, and which appear in the Chinese numerals. Such sticks are naturally laid horizontally with respect to the eye. The earliest Sumerian forms of the numerals

[1]Karoṣṭhī numerals, Aśoka inscriptions, *c.* 250 B.C. For sources of information with respect to this table see Smith-Karpinski, p. 25.

[2]Same, Śaka inscriptions, probably of the first century B.C.

[3]Brāhmī numerals, Aśoka inscriptions, *c.* 250 B.C.

[4]Same, Nānā Ghāt inscriptions, *c.* 150 B.C.

[5]Same, Nasik inscription, *c.* 100.

[6]Kṣatrapa coins, *c.* 200.

[7]Kuṣana inscriptions, *c.* 150.

[8]Gupta inscriptions, *c.* 300 to 450.

[9]Valhabī, *c.* 600.

were horizontal,[1] and so the computing rod may have had its origin in Sumeria. The later Babylonian forms were vertical, and so the finger computation may have been in favor at that time. The Mediterranean lands adopted the vertical forms, and the Far East preferred the horizontal. From the vertical 𓏺𓏺 came the following Egyptian forms:

Hieroglyphic	𓏺𓏺
Hieratic	𓏺𓏺
Demotic	𓏺𓏺

From the symbol 𓏺𓏺 came also the Arabic ٢, which is our 2 if turned on its side. Indeed, our 2 is merely a cursive form of 𝌆 and our 3 is similarly derived from 𝌇, as is seen from a study of inscriptions and manuscripts.[2] These horizontal forms were early used by the Chinese and probably found their way from China to India.

Fanciful Theories. Numerous conjectures have been made as to the origin of the characters from four to nine, but no one of them has had any wide acceptance. We may dismiss at once all speculations as to their derivation from such combinations as ⊠ and ⊕ and from the number of sticks that might be laid down to make the figure 8. Such ideas are trivial and have no sanction from the study of paleography. There remain, however, various scientific theories, as that the forms are ancient initial letters of number words.[3] None of these theories, however, has stood the test of scholarly criticism, and today we have to confess that we are entirely ignorant as to the origin of the forms which began possibly in India in Aśoka's time and appear as the common numerals which we use.

[1] Sir H. H. Howard, "On the Earliest Inscriptions from Chaldea," *Proceedings of the Society of Biblical Archæology*, XXI, 301.

[2] An interesting fact in relation to the figure 2 is that the Romans often wrote $\frac{2}{2}$ as two lines, 𝌆, the twelfth being understood as we understand tenths when we write 0.2. They also wrote this character cursively, 𝍐, which is the character used for our 2 in several early printed books.

[3] For details of this theory consult the bibliography given in Smith-Karpinski, p. 30.

Origin of the Zero. The origin of the zero is quite as uncertain as the origin of our other numerals. Without it the Hindu numerals would be no better than many others, since the distinguishing feature of our present system is its place value. The earliest undoubted occurrence of a zero in India is seen in an inscription of 876 at Gwalior. In this inscription 50 and 270 are both written with zeros.[1] We have evidence, however, that a place value was recognized at an earlier period, so that the zero had probably been known for a long time. The Babylonians, indeed, had used a character for the absence of number, and they made use of a primitive kind of place value;[2] but they did not create a system of numeration in which the zero played any such part as it does in the one which we now use. There is also a slight approach to a place value in some of the late Greek works. For example, Diophantus seems, judging by certain manuscripts, to have used $\dot{\beta}\Lambda\Phi\Pi\varsigma$ for 23,587, the four points about the β being a late Greek symbol for myriads, and the position of Λ determining its value as 30 hundreds.[3]

The form of the zero may have been suggested by an empty circle, by the Greek use of omicron (O) to indicate a lacuna,[4] by the horned circle used in the Brāhmī symbols for ten, by the Hindu use of a small circle (o), as well as a dot, to indicate a negative, or in some other way long since forgotten. There is no probability that the origin will ever be known, and there is no particular reason why it should be. We simply know that the world felt the need of a better number system, and that the zero appeared in India as early as the 9th century, and probably some time before that, and was very likely a Hindu invention.

The Arabs represented 5 by a character that looked somewhat like the Hindu zero. In a manuscript of 1575 the numerals appear as ٩ ٨ ٧ ٦ ٥ ٤ ٣ ٢ ١. In other manuscripts we find such forms as ۴, ۵, and ۵. Because of the resemblance of their five to the circle the Arabs adopted a dot for their zero.

[1] For facsimiles see Smith-Karpinski, p. 52.

[2] For particulars see *ibid.*, p. 51.

[3] For other cases see Gardthausen, *Die Schrift*, p. 372.


[4] Being the initial of οὐδέν (*ouden'*), nothing. Thus, Archimedes might have used O to indicate the absence of degrees or minutes. See Heath, *Archimedes*, lxxii

For purposes of comparison the Sanskrit forms are here repeated and the modern Arabic forms are given:

Sanskrit, १ २ ३ ४ ५ ६ ७ ८ ९ ०

Arabic, ١ ٢ ٣ ٤ ٥ ٦ ٧ ٨ ٩ ٠

The various forms of the numerals used in India after the zero appeared may be judged from the table here shown.



[1] Bakhshālī MS. See Volume I, page 164; Smith-Karpinski, pp. 49, 50.

[2] The 3, 4, 6, from H. H. Dhruva, "The Land-Grants from Sankheda," *Epigraphia Indica*, Vol. II, pp. 19–24 with plates; date 595. The 7, 1, 5, from Bhandarkar, "Daulatabad Plates," *Epigraphia Indica*, Vol. IX, Part V; date *c.* 798.

[3] The 8, 7, 2, from "Buckhala Inscription of Nagabhatta," Bhandarkar, *Epigraphia Indica*, Vol. IX, Part V; date 815. The 5 from "The Morbi Copper-Plate," Bhandarkar, *Indian Antiquary*, Vol. II, pp. 257–258, with plate; date 804.


[4] The 8 from the above Morbi Copper-Plate. The 4, 5, 7, 9, and 0 from "Āsni Inscription of Mahipala," *Indian Antiquary*, Vol. XVI, pp. 174–175; date 917.

[5] The 8, 9, 4, from "Rashtrakuta Grant of Amoghavarsha," J. F. Fleet, *Indian Antiquary*, Vol. XII, pp. 263–272; date *c.* 972. See Bühler. The 7, 3, 5, from "Torkhede Copper-Plate Grant," Fleet, *Epigraphia Indica*, Vol. III, pp. 53–58.

[6] From "A Copper-Plate Grant of King Tritochanapâla Chanlukya of Latadeśa," H. H. Dhruva, *Indian Antiquary*, Vol. XII, pp. 196–205; date 1050.

[7] A. C. Burnell, *South Indian Palæography*, Plate XXIII, Telugu-Canarese numerals of the 11th century.

The following are later European and Oriental forms:



The Name for Zero. The name for zero is not settled even yet. Modern usage allows it to be called by the name of the letter *O*, an interesting return to the Greek name *omicron* used by Buteo in 1559. The older names are *zero*, *cipher*, and *naught*. The Hindus called it *śūnya*, "void," and this term passed over into Arabic as *aṣ-ṣifr* or *ṣifr*. When Fibonacci (1202) wrote his *Liber Abaci*, he spoke of the character as *zephirum*.[8] Maximus Planudes (*c.* 1340) called it *tziphra*[9] and this form was used by Fine (1530) in the 16th century. It passed over into Italian as *zeuero*,[10] *ceuero*,[11] and *zepiro*,[12] and in the medieval period it

[1] and [2] From a manuscript of the second half of the 13th century, reproduced in "Della vita e delle opere di Leonardo Pisano," Baldassare Boncompagni, Rome, 1852, in *Atti dell' Accademia Pontificia dei Nuovi Lincei*, anno V.

[3] and [4] From a 14th century manuscript.

[5] From a Thibetan MS. in the library of the author.

[6] From a specimen of Thibetan block printing in the library of the author.

[7] Śāradā numerals from *The Kashmirian Atharva-Veda, reproduced by chromophotography from the manuscript in the University Library at Tübingen*, M. Bloomfield and R. Garbe, Baltimore, 1901.

[8] ". . . quod arabice zephirum appellatur."

[9] From the Greek form τζίφρα (*tzi'phra*), used also by another writer, Neophytos, about the same time.

[10] Thus Jacopo da Firenze (1307), or Magister Jacobus de Florentia.

[11] As in the arithmetic of Giovanni de Danti of Arezzo (1370).

[12] As in a translation into Latin of the works of Avicenna.

had various other forms, including *sipos*, *tsiphron*, *zeron*, *cifra*, and *zero*. It was also known by such names as *rota*, *circulus*, *galgal*, *omicron*, *theca*, *null*, and *figura nihili*.[1]

Numerals outside of India. The first definite trace that we have of the Hindu numerals outside of India is in the passage already quoted from Severus Sebokht (*c.* 650). From this it seems clear that they had reached the monastic schools of Mesopotamia as early as 650.

The next fairly definite information as to their presence in this part of the world, and with a zero, is connected with the assertion that a set of astronomical tables was taken to Bagdad in 773 and translated from the Sanskrit into Arabic by the caliph's command. There is ground for doubt as to the assertion, but the translation is said to have been made by al-Fazârî (*c.* 773). It is probable that the numerals were made known in Bagdad at this time, and they were certainly known by the year 825. About that year al-Khowârizmî recognized their value and wrote a small book explaining their use. This book was translated into Latin, possibly by Adelard of Bath (*c.* 1120), under the title *Liber Algorismi de numero Indorum*.[2]

The Hindu forms described by al-Khowârizmî were not used by the Arabs, however. The Bagdad scholars evidently derived their forms from some other source, possibly from Kabul[3] in Afghanistan, where they may have been modified in transit from India. These numerals have been still further modified in some respects, and at present are often seen in the forms given on pages 70 and 71.

The Numerals move Westward. Owing to the fact that almost no records of a commercial nature have been preserved from

[1]For a full discussion see Smith-Karpinski, chap. iv.

[2]The Book of al-Khowârizmî on Hindu number. On this work see Smith-Karpinski, pp. 5 seq., 92 seq.

[3]It is curious that the old Biblical name of Cabul (1 Kings, ix, 13; Joshua, xix, 27) should be found in Afghanistan. Could it have been taken there by the ruling clan, the Duranis, who call themselves Beni Israel and who claim descent from the Israelites who fled to the Far East after the Assyrians devastated Samaria? If so, could these people, who also claim descent from Kish (1 Samuel, ix, 1), have taken the numerals from Egypt to Afghanistan?

the so-called Dark Ages of Europe, and that the number of scientific works that have come down to us is also very limited, we cannot say when the Hindu-Arabic numerals first found their way to the West. There are good reasons for believing that they reached Alexandria along the great pathway of trade from the East even before they reached Bagdad, possibly in the 5th century, but without the zero.[1] It would have been strange if the Alexandrian merchants of that time and later had not known the numeral marks on goods from India, China, and Persia. No system that did not contain a zero, however, would have attracted much attention, and so this one, if it was known at all, was probably looked upon only as a part of the necessary equipment of a trader with the East.

The Ġobâr Numerals. At any rate, numerals are found in Spain as early as the 10th century, and some of these numerals differ so much from the rest that they evidently came through a different channel, although from the same source. These were called the dust numerals,[2] possibly because they were written on the dust abacus instead of being represented by counters. It is worthy of note that Albêrûnî (*c.* 1000) states that the Hindus often performed numerical computations in the sand.

If these numerals reached Alexandria in the 5th century, they probably spread along the coasts of the Mediterranean Sea, becoming known in all the leading ports. In this case they would have been familiar to the merchants for purposes of trade and to the inquisitive for reasons of curiosity. The soothsayer and astrologer would have adopted them as part of the mysticism of their profession, and the scholar would have investigated them as possibilities for the advancement of science. In that case a man like Boethius (*c.* 510) would have been apt to know of them and perhaps to mention them in his writings.


All this is speculative. The *ġobâr* numerals exist as a fact, and this is their possible origin. In certain manuscripts of

[1] For bibliography and discussion see Smith-Karpinski.

[2] *Ḥurûf al-ġobâr.* The name appears in Tunis as early as the middle of the 10th century. There were also the *ḥurûf al-jumal*, or alphabetic numerals, used by the Jews and probably also by the Arabs.

Boethius there appear similar forms, but these manuscripts are not earlier than the 10th century and were written at a time when it was not considered improper to modernize a text. They do not appear in the arithmetic of Boethius, where we might expect to find them, if at all, but in his geometry, and their introduction breaks the continuity of the text. It therefore seems very doubtful that they were part of the original work of Boethius. Since any forms that reached Alexandria would probably have lacked the zero, and since a zero appears in the late Boethian manuscripts, there is the added reason for feeling that at least part and very likely all of the symbols were inserted by copyists.

These *ġobâr* numerals varied considerably but were substantially as shown in the following table:



Gerbert and the Numerals. The first European scholar who is definitely known to have taught the new numerals is Gerbert (*c.* 980), who later became Pope Sylvester II (999). He went to Spain in 967 and may have learned about them in Barcelona.[6] He probably did not know of the zero, and at any rate he

[1] For sources of information with respect to these numerals see Smith-Karpinski, p. 69.

[2] Al-Ḥaṣṣār's forms, H. Suter, *Bibl. Math.*, II (3), 15.

[3] The manuscript from which these are taken is the oldest (970 A.D.) Arabic document known to contain all the numerals.

[4] and [5] Woepcke, "Introduction au calcul Gobârî et Hawaî," *Atti dell' Accademia Pontificia dei Nuovi Lincei*, Vol. XIX.

[6] On this question see Smith-Karpinski, p. 110.

did not know its real significance. He placed upon counters the nine *caracteres*, as they were called by his pupils Bernelinus (*c.* 1020) and Richer, and used these counters on the abacus. Such counters, probably in the form of flattened cones, were called *apices*, a term also used in connection with the numerals themselves. These numerals were severally called by the names *igin, andras, ormis, arbas, quimas, calctis, zenis, temenias, celentis,* and *sipos.*[1] The origin and meaning of these terms have never been satisfactorily explained, but the words seem to be Semitic.[2]

The oldest definitely dated European manuscript that contains these numerals was written in Spain in 976. A Spanish



EARLY EUROPEAN NUMERALS

Oldest example of our numerals known in any European manuscript. This manuscript was written in Spain in 976

copy of the *Origines* of Isidorus, dated 992, contains the numerals with the exception of zero. Dated manuscripts of the Arabs have been found which give some of these numerals a century earlier, that is, in 874 and 888. They also appear in a Shiraz manuscript of 970 and in an Arabic inscription in Egypt dated 961. The earliest occurrence of these numerals in a date on a coin is found on a piece struck in Sicily in 1138.


There is good reason for believing that Gerbert obtained his knowledge of the numerals from studying in the convent of Santa Maria de Ripoll, a well-known center of learning near Barcelona;[3] indeed, it is not improbable that he saw the very

[1] There were variants of these forms. The *sipos* does not appear in the works of the pupils of Gerbert, but is found in a MS. of Radulph of Laon (*c.* 1125).

[2] Smith-Karpinski, p. 118.

[3] J. M. Burnam, "A Group of Spanish Manuscripts," *Bulletin Hispanique,* XXII, 229 (Bordeaux, 1920). With respect to the library in this convent see R. Beer, *Die Handschriften des Klosters Santa Maria de Ripoll,* Vienna, 1907.

manuscript of 976 above mentioned. There is considerable evidence to support the belief that the monks in this cloister obtained their knowledge of these numerals through mercantile




sources which were in communication with the East, rather than through any Moorish channels in Mohammedan Spain.

The changes in the forms of the numerals may be seen in the table above.[1] The forms as they appeared just before the


[1] This is from a table prepared by Mr. G. F. Hill of the British Museum, and is reproduced by his permission. His noteworthy article on the subject appeared in *Archæologia*, LXII (1910). This was elaborated in book form under the title, *The Development of Arabic Numerals in Europe*, Oxford, 1915.

invention of European printing may be seen in the annexed facsimile from a Latin manuscript written by Rollandus at Paris, *c.* 1424. After Europe began to print books, the forms varied but little, most of the changes being due simply to the fashions set by designers of type. For example, the figures 4 and 5 were changed to their present forms in the 15th century and have since then remained fairly well standardized.

Not only did the forms of the numerals change considerably during the Middle Ages, but the method of writing the ordinary numbers also varied from century to century. Some scribes always placed a dot before and after each figure, as in the case of a number like .2. Others adopted a somewhat similar plan in the case of numbers having several figures. For example, one writer of *c.* 1400[1] gives 5. 7. 8. 2. for 5782, and one of 1384 gives 1000. 300. 80. 4 for 1384, as shown in the following illustration from an anonymous *computus* written in Italy:



FROM THE ROLLANDUS MANUSCRIPT OF *c.* 1424



NUMERALS FROM A COMPUTUS OF 1384

The method of writing the date, 1000. 300. 80. 4, illustrates the difficulties in using the numerals. From Mr. Plimpton's library

[1] F. J. Studnička, *Algorismus prosaycus magistri Christani*, p. 9 (Prag, 1893). This Magister Christanus was Christanus Prachaticensis, or Christian of Prag (born 1368; died 1439).

Early English Algorism. An interesting illustration of the early use of the word "algorism" (algorym, augrim) in the English language may be seen in a manuscript now in the British Museum, dating from *c.* 1300.[1] The first page, which is here shown in facsimile, reads as follows:

Hec algorisms ars psens dicit^r in qua
Talibs indoꝫ fruim^r bis quīq figuris.[2]

This boke is called þe boke of algorym or Augrym after lewder use. And þis boke tretys þe Craft of Nombryng, þe quych crafte is called also Algorym. Ther was a kyng of Inde þe quich heyth Algor, & he made þis craft. And aft his name he called hit algory. Or els anoth cause is quy it is called Algorym, for þe latyn word of hit s. Algorismus coms of Algos grece q ē ars, latine, craft on englis, and rides q ē nm̄s, latine, A nombr on englys. inde dr algorismus ꝑ addicōnē huis sillabe ms & subtraccōnem d & E, qsi ars numāndi.[3]
¶ fforthermor[4] ye most undrstonde þt in þis craft ben usid teen figurys. as her ben writen for ensampul. ϕ . 9 . 8 . 7 . 6 . 5 . 4 . 3 . 2 . 1.
¶ Expone þe too vsus a for;[5] þis psent craft is called Algorisms, in þe quych we use teen figurys of Inde. Questio. ¶ Why ten fyguris

[1] It was first privately printed by the Early English Text Society (transcription by Robert Steele), London, 1894. It has already been referred to in Volume I, page 238, and in this volume, page 32.

[2] These are the two opening lines of the *Carmen de Algorismo*, of Alexandre de Villedieu (*c.* 1240). They should read as follows:


Haec algorismus ars praesens dicitur; in qua
Talibus Indorum fruimur bis quinque figuris.

It is translated a few lines later: "This present craft is called Algorismus, in the which we use ten figures of India."

[3] "Inde dicitur Algorismus per addicionem huius sillabe *mus* & subtraccionem *d* & *e*, quasi ars numerandi (Whence it is called Algorismus by the addition of this syllable *mus*, and the taking away of *d* and *e*, as if the art of numbering)." This idea had considerable acceptance in the 13th century.

[4] "Furthermore," the *f* being doubled for a capital. "Furthermore you must understand that in this craft there are used ten figures." The forms of the numerals given in the original were the common ones of the 12th and 13th centuries. The zero was not usually our form, but frequently looked more like the Greek *phi*. The 7, 5, and 4 changed materially in the latter part of the 15th century, about the time of the first printed books. The sequence here shown is found in most of the very early manuscripts, the zero or nine being at the left.

[5] "Explain the two verses afore."



Folio 1r of the manuscript

FIRST PAGE OF THE CRAFT OF NOMBRYNG

Egerton MS. 2622 in the British Museum, one of the earliest manuscripts in English which treat of any phase of mathematics

of Inde. Solucio.[1] for as I have sayd a fore þai wer fonde fyrst in Inde of a kyng of þat Cuntre þt was called Algor. ¶ Pma sigt uno duo vo scda[2] ¶ Tercia sigt tria sic pcede sinistre. ¶ Donc ad extmā venias que cifra vocar. ¶ Capm pmū de significacōe figurarm ¶ In þis verse is notifide þe significacōn of þese figuris. And þus expose þe verse the first signifiyth on. þe secūde signi[fiyth tweyn].[3]

Reading and Writing Large Numbers. One of the most striking features of ancient arithmetic is the rarity of large numbers. There are exceptions, as in some of the Hindu traditions of Buddha's skill with numbers,[4] in the records on some of the Babylonian tablets,[5] and in the *Sand Reckoner*[6] of Archimedes, with its number system extending to 10^{63}, but these are all cases in which the élite of the mathematical world were concerned; the people, and indeed the substantial mathematicians in most cases, had little need for or interest in numbers of any considerable size.

The Million. The word "million," for example, is not found before the 13th century, and seems to have come into use in England even later. William Langland (*c.* 1334–*c.* 1400), in *Piers Plowman*, says,

> Coueyte not his goodes
> For millions of moneye,

[1] "Answer."

[2] "The first means one, the second two, the third means three, and thus proceed to the left until you reach the last, which is called cifra." The author is quoting from the *Carmen* of Alexandre de Villedieu:

> Prima significat unum; duo vero secunda;
> Tertia significat tria; sic procede sinistre
> Donec ad extremam venias, quæ cifra vocatur.


[3] "Capitulum primum de significacione figurarum (Chapter I, On the meaning of the figures)."
"And thus explain the (Latin) verse: the first signifieth one."
"The second (secunde) signifieth twain."

[4] Sir Edwin Arnold speaks of this in *The Light of Asia*. See Smith-Karpinski, p. 16.

[5] Hilprecht, *Tablets*.

[6] Ψαμμίτης (*Psammi'tes*), translated into Latin as *Arenarius*. For the text, see *Archimedis opera omnia*, ed. Heiberg, with revisions, II, 242 (Leipzig, 1880–1913); Heath, *Archimedes*, p. 221. See also M. Chasles, in the *Comptes rendus*, April 11, 1842; the preface to the English translation of the work by Archimedes made by G. Anderson, London, 1784; Heath, *History*, II, 81.

but Maximus Planudes (*c.* 1340) seems to have been among the first of the mathematicians to use the word.[1] By the 15th century it was known to the Italian arithmeticians, for Ghaligai


¶ Como si fozmano milion

L million adoncha se dice formar per sette figure in questo modo. 1000000. perche la septima figura tien el luogo demiara demara: e per che mille miara fano uno million: et essendo in quel luogo la figura che ripzexenta uno pero bene edito uno milion. Ma. iquesto modo. 1 100000 diria uno milion e cento milia: perche oltra el million: in luogo de centenara de miara: sono la figura che ripzexenta uno si che bene edito uno million e cento milia. Ma. in questo modo. 1 1 10000. diria uno milion e cento e diexe milia perche oltra el million e cento milia: in luogo dele dexene demiar: sono la figura che ripzexenta uno: si che bene edito uno million e cento ediexe milia. Ma in questo modo. 1 1 1 1000. diria un million cento e undexe milia per che oltra el million cento e diexe milia: in luogo de numeri demiar sono la figura che ripzexenta uno: si che bene edito un million cento e undexe milia. Ma in questo modo. 1 1 1 1 100. diria un million cento e undexe milia e cento: perche oltra el million cento e undexe milia: in luogo de el simplice centenar: sono la figura che ripzexenta uno: si che bene edito un million cento e undexe milia e cento. Ma in questo modo. 1 1 1 1 1 10. diria un million. cento e undexe milla cento ediexe: perche oltra el million cento e undexe milia e cento: i luogo dele simplice dexene: sono la figura che ripzexenta uno. Ma in questo modo. 1 1 1 1 1 1 1. diria un million cento e undexe milia cento e undexe. per che anche in luogo dele simplice unita. sono la figura che ripzexenta uno. si che bene edito un million cento e undexe milia cento e undexe. et chosi procedédo per fina. 9999999. ponendo sempze a suo luogi quele figure repzexentante queli numeri oucro dexene ocenteuara. che si nomina et cetera. e questo basta cercha lo amaistramento del numerar. ben che in infinitum si pozia procedere. ma chom una zeneral figura misozcero dichiarir quanto potesse achader. et sarano questo sotto posta

1000000

1 100000

1 1 10000

1 1 1 1000

1 1 1 1 100

1 1 1 1 1 10

1 1 1 1 1 1 1

9999999

THE WRITING OF LARGE NUMBERS IN 1484

From Pietro Borghi's *De Arte Mathematiche*, Venice, 1484. This illustration is from the 1488 edition

(1521; 1552 ed., fol. 3) relates that "Maestro Paulo da Pisa"[2] read the seventh order as millions. It first appeared in a printed

[1] H. Wäschke translation, p. 4 n. (Halle, 1878); hereafter referred to as Wäschke, *Planudes*. The word simply means "great thousand" (from *mille* + *on*), just as *salon* means "great hall" (from *salle* + *on*) and *balloon* means "great ball."

[2] "La settima dice numero di milione." This Paul of Pisa may have been the Paolo dell' Abbaco (Dagomari, *c.* 1340) mentioned in Volume I, page 232.

work in the Treviso arithmetic of 1478. Thereafter it found place in the works of most of the important popular Italian writers, such as Borghi[1] (1484), Pellos[2] (1492), and Pacioli[3] (1494), but outside of Italy and France it was for a long time used only sparingly. Thus, Gemma Frisius (1540) used "thousand thousand"[4] in his Latin editions, which were published in the North, while in the Italian translation (1567) the word *millioni* appears. Similarly, Clavius carried his German ideas along with him when he went to Rome, and when (1583) he wished to speak of a thousand thousand he almost apologized for using "million," referring to it as an Italian form which needed some explanation.[5]

In Spain the word *cuento*[6] was early used for 10^6, the word *millon* being reserved for 10^{12}. When the latter word was adopted by mathematicians, it was slow in coming into general use.[7]

[1] "Il miar de milliara ò vuol dir il million" (1540 ed., fol. 5).

[2] His names beyond units are *desena, centenal, millier,* x^a *de* m^a, c^a *de* m^a, and *million* (fol. 2).

[3] He uses *milioni* (fol. 9) but no higher special names, although he repeats this word, as in "Migliara de miliõ de miliõ" (fol. 19, *v.*), adding: "Et sic ī sequẽtib⁹. ꝑseq̄re."

The spelling varies in the early books, sometimes appearing as *miglioni* (Pagani, 1591).

[4] *Millena millia.*

[5] In the Latin edition (1583): "Iam vero si more Italorum millena millia appellare velimus Milliones, paucioribus verbis & fortasse significantius" (*Epitome*, cap. i).

In the Italian edition (1586): "Hora se secõdo il costume d' Italia vorremo vn migliaio di migliaia chiamare millione, con manco parole, & forse piu significantemente" (p. 14).

In *De Cosmographia Libri IV* by Francesco Barozzi (*c.* 1538–*c.* 1587), a work published in Venice in 1585, it is stated that "septima (nota) pro Millenario Millenarii quem vulgus quidem Millionem appellant, Latini vero Milleno Millio."

[6] From *contar*, to count or reckon. Ciruelo, whose work was published in Paris in 1495, says: "Millies millena: quod vulgariter dicitur cuento: decies cuento/centies cuento/millies cuento/decies millies cuento/centies millies cuento [/millies millies cuento]/quod vulgariter dicitur millon." See 1513 ed., fol. a_2 and fol. A_3. He is not, however, uniform in the matter, using "millon" as synonymous with "cuento" in other places.

[7] As late as the 1643 edition of Santa-Cruz (1594) it was necessary to explain the word thus: "Millon que significa mil millares" (fol. 13, *r.*), the latter being the common form.

France early took the word "million" from Italy, as when Chuquet (1484) used it, being followed by De la Roche (1520), after which it became fairly common.

The conservative Latin writers of the 16th century were very slow in adopting the word. Even Tonstall (1522), who followed such eminent Italian writers as Pacioli, did not commonly use it. He seems to have been influenced by the fact that the Romans had no use for large numbers;[1] or by the fact that, for common purposes, it sufficed to say "thousand thousand," as had been done for many generations.[2] He simply mentions the word as a piece of foreign slang to be avoided.[3] Other Latin writers were content to say "thousand thousand."[4]

The German writers were equally slow in abandoning "thousand thousand" for "million," most of the writers of the 16th century preferring the older form.[5] The Dutch were even more conservative, continuing the old form later than the writers in the neighboring countries.[6] Indeed, for the ordinary needs of business in the 16th century, the word "million" was a luxury rather than a necessity.

[1]"Non me latet Romanos ueteres prisco more, suos nūmos Sestertios computātes, numerum trāscendentem centum millia . . . Latine nō enunciasse . . . ."

[2]Even as late as 1501, Huswirt, a German scholar, writes "quadraginta quattuor mille millia. quingēta millia quinquaginta nouē millia. octingenta. octoginta sex" for 44,559,886.

[3]"Septimus millena millia: uulgus millionē barbare uocat."

[4]So Stifel uses "millia, millies" (*Arithmetica Integra*, 1544 ed., fol. 1); Ramus uses "millena millia" (*Libri II*, 1569, p. 1); Glareanus has "mille millia" (1538; 1543 ed., fol. 9).

Thus, "tausant mal tausant" is used by such writers as Köbel (*Zwey Rechenbüchlin*, 1514; 1537 ed., fol. 14), Grammateus (1518; 1535 ed., p. 5), Riese (1522; 1529 ed., p. 3), and Rudolff (1526; 1534 ed., fol. 3). Rudolff, however, uses it together with the older form in his *Rechenbuch* (1526), and in his *Exempelbüchlin* (1530; 1540 ed., exs. 62 and 137) he says: "Vnd wirt ein million mit ziffern geschriben 1000000," and "ist zehenmal hundert tausent."

[6]Thus, "duysent mael duysent" is used by such writers as Petri (1567, fol. 1), Raets (1580, fol. A_3, with "duysentich duysent"), Mots (1640, fol. B_2), Cardinael (1659 ed., fol. A_3), Willemsz (1708 ed., p. 5), and Bartjens (1792 ed., p. 8). There were exceptions, as when Wentsel (Wenceslaus, 1599) used both "millioenen" and "millions" (p. 2), Stockmans (1589; 1679 ed., p. 8) occasionally used "millioen," and Starcken (1714 ed., p. 2) used "million" rather apologetically.

England adopted the Italian word more readily than the other countries, probably owing to the influence of Recorde[1] (*c.* 1542). It is interesting to see that Poland was also among the first to recognize its value, the word appearing in the arithmetic of Kłos in 1538.

The Billion. Until the World War of 1914–1918 taught the world to think in billions there was not much need for number names beyond millions. Numbers could be expressed in figures, and an astronomer could write a number like $9.15 \cdot 10^7$, or $2.5 \cdot 10^{20}$, without caring anything about the name. Because of this fact there was no uniformity in the use of the word "billion." It meant a thousand million (10^9) in the United States and a million million (10^{12}) in England, while France commonly used *milliard* for 10^9, with *billion* as an alternative term.

Historically the billion first appears as 10^{12}, as the English use the term. It is found in this sense in Chuquet's number scheme[2] (1484), and this scheme was used by De la Roche (1520), who simply copied parts of Chuquet's unpublished manuscript, but it was not common in France at this time, and it was not until the latter part of the 17th century that it found place in Germany.[3] Although Italy had been the first country to make use of the word "million," it was slow in adopting the word "billion." Even in the 1592 edition of Tartaglia's arithmetic the word does not appear. Cataldi (1602) was the first Italian writer of any prominence to use the term, but he sug-

[1] "203000000, that is, CCiii milliōs," "M. of millions," and "x.M. of millions" (1558 ed., fol. C_8).

[2] This plan is historically so important as to deserve being given in full. Chuquet gives the 6-figure periods, thus: 745324'8043000'700023'654321 (in which 8043000 should be 804300), and then says: "Ou qui veult le p̄mier point peult signiffier million Le second point byllion Le tiers poīt tryllion Le quart quadrillion Le cinq^e quyllion Le six^e sixlion Le sept^e septyllion Le huyt^e ottyllion Le neuf^e nonyllion et ansi des ault^{es} se plus oultre on vouloit ꝓceder ¶ Item lon doit sauoir que ung million vault mille milliers de unitez. et ung byllion vault mille milliers de millions. et tryllion vault mille milliers de byllions." From A. Marre's autograph copy of Chuquet. See also Boncompagni's *Bullettino*, Vol. XIII, p. 594.

[3] F. Unger, *Die Methodik der praktischen Arithmetik in historischer Entwickelung*, p. 71 (Leipzig, 1888), with the date of use of the word as 1681; hereafter referred to as Unger, *Die Methodik*.

gested it as a curiosity rather than a word of practical value.[1] About the same time the term appeared in Holland,[2] but it was not often recognized by writers there or elsewhere until the 18th century, and even then it was not used outside the schools. Even as good an arithmetician as Guido Grandi (1671–1742) preferred to speak of a million million rather than use the shorter term.[3]

The French use of *milliard*, for 10^9, with *billion* as an alternative, is relatively late. The word appears at least as early as the beginning of the 16th century as the equivalent both of 10^9 and of 10^{12}, the latter being the billion of England today.[4] By the 17th century, however, it was used in Holland[5] to mean 10^9, and no doubt it was about this time that the usage began to change in France.

As to the American usage, taking a billion to mean a thousand million and running the subsequent names by thousands, it should

[1] He generally used millions, thousand millions, million millions, and so on (p. 2); but he sometimes used *bilioni* for 10^9, although even then he preferred *duilioni*. His scheme of names is *millioni*, *bilioni* (or *duilioni*), *trilioni*, *quadrilioni* (or *quattrilioni*), *quintilioni*, for 10^6, 10^9, 10^{12}, 10^{15}, and 10^{18}. *Practica Aritmetica*, p. 5 (Bologna, 1602).

[2] Van der Schuere (1600) uses *millioen* (10^6), *bimillioen* (10^{12}), *trimillioen* (10^{18}), and *quadrimillioen* (10^{24}), but in a later edition (1634) of his arithmetic he gives *bimillion* and *billion* (10^9), *trimillion* and *trillion* (10^{12}), and so on to *nonemillion* and *nonilion*. Even as late as 1710 Leonhard Christoph Sturm (*Kurzer Begriff der gesamten Mathesis*, Frankfort a. d. Oder, 1710) used the words *trimillionen* and *bimillionen*.

[3] In his *Istituzioni di Aritmetica Practica*, p. 3 (Florence, 1740), he says: "millioni de' millioni (che possono dirsi Billioni) e li millioni di millioni di millioni (che si chiamano ancora Trillioni). . . . E così se fosse più lungo il numero, vi sarebbero ancora Quintillioni . . . Novillioni, ec. crescendosi ciascuno da ogni sei note."

[4] Thus, Trenchant (1566) uses *Miliars* (1578 ed., p. 14), and Peletier (1549) says: "Les François ont deux mots numeraux significatifs: l'un au septieme lieu, qui est Million, & l'autre au treizieme, qui est Milliart: c'est à dire, Million de Millions" (1607 ed., p. 15). Peletier states that the word was used by Budæus, and in the latter's *De Asse et partibus eius Libri quinq3* (1514; Paris edition of 1532, fol. 95, *v.*) the following appears: "hoc est denas myriadū myriadas, quod vno verbo nostrates abaci studiosi Milliartū appellāt, quasi millionū millionē." In Boissière's arithmetic of 1554 there is a statement similar to the one in Peletier's work. In E. Develey, *Arithmétique d'Émile*, 2d ed., Paris, 1802, only "billion" is used for 1000 millions.

[5] "milliart/ofte duysent millioenen," as Houck's arithmetic (1676, p. 2) has it.

be said that this is due in part to French influence after the Revolutionary War, although our earliest native American arithmetic, the Greenwood book of 1729,[1] gave the billion as 10^9, the trillion as 10^{12}, and so on. Names for large numbers were the fashion in early days, Pike's well-known arithmetic (1788), for example, proceeding to duodecillions before taking up addition.

Writing Large Numbers. Although it is nearly a thousand years since our common numerals appeared in any European manuscripts now extant, we have not even yet decided on the method of writing large numbers.

Influenced by the crosses placed on the thousands' and millions' lines of the abacus (see page 181) to aid the eye, the medieval writers often placed a dot above the thousands and above every third place beyond, but sometimes they placed one or more dots below, and these customs also appear in the early printed books. Thus, we have such a form as 6̇8̇54̇973,[2] with the occasional variant of a dot over the units' figure also.[3] Recorde (*c.* 1542) gives the rule as follows:

> Fyrst put a pricke ouer the fourth fygure, and so ouer the vij. And if you had so many ouer the x, xiij, xvj, and so forth, still leauing two fygurs betwene eche two pricks. And those roomes betwene the prickes are called Ternaries.[4]

Recorde also uses a bar (*virgula*) for separating the figures, saying:

[1] It was published anonymously, but, as is stated in the *Weekly News Letter* (Boston) of May 29, 1729, was written by Isaac Greenwood, sometime professor of mathematics at Harvard. As stated in Volume I, the first arithmetic printed in the New World appeared in Mexico in 1556; the first in what is now the United States was a reprint of Hodder's English arithmetic, Boston, 1719.

[2] Tonstall (1522, fol. C_1), Riese (1522; 1529 ed., p. 3), Rudolff (1526, fol. 3), Grammateus (1518; 1535 ed., p. 5), and many others. Widman (1489) recommends but does not use this plan: "Vnd setz vff ytlich tausent ain punct da by man mercken mag wie vil die letst figur mer tausent bedeut dann die vor ir" (1519 ed., fol. 5, *v.*).

[3] Thus Clavius (Italian edition of the *Epitome*, 1586, p. 14; Latin ed., 1583, p. 10). He recommends, however, the following: $4^{4}23^{3}29^{2}08^{1}95^{0}62800$.

[4] *Ground of Artes*, 1558 ed., fol. B_8. Similarly in Digges (1572; 1579 ed., p. 2), Baker (1568; 1580 ed., fol. 4), and Hodder (1672 ed.).

And some doo parte the nūbres with lynes after this forme 230|864|089|015|340, where you see as many lines as you made pricks.[1]

Some writers used this symbolism in grouping by sixes.[2] Besides placing one dot above a figure, the medieval writers often used such forms as $\ddot{2}43\dot{7}56\dot{2}93\dot{8}421$[3] and $\overset{\cdot\cdot}{2}437\overset{\cdot}{5}62\overset{\cdot}{9}38\overset{\cdot}{4}21$, and these occasionally appear in the printed works. Fibonacci (1202) used the arc, as in $\overset{\frown}{6}\overset{\frown}{7}8\overset{\frown}{9}3\overset{\frown}{5}7\overset{\frown}{8}4\overset{\frown}{1}0\overset{\frown}{5}296$,[4] but this was not a common form. A few of the other variations are given below:

$\overset{3}{7}\cdot53\overset{2}{8}.2\overset{1}{7}5.136$	Pellos (1492, fol. 4)
$\dot{4}.5.9.\dot{3}.6.2.\dot{9}.0.2.2$ or $\underset{4593629022}{\overset{a\ c\ b\ a\ c\ b\ a\ c\ b\ a}{}}$	Reisch[5] (1503, Lib. IIII, Tract. II, Cap. 4)
$25\overline{78}\overline{39}1\dot{0}6\overline{2}\overline{75}1\dot{2}3\overline{46}\overline{89}4352$	Barozzi[6] (1585)
$\overset{\cdot\cdot}{2}3.\overset{\cdot\cdot}{4}5\overset{\cdot\cdot}{6}.0\ddot{0}7.8\dot{4}0.0\ddot{0}0.3\dot{0}5.321$	Santa-Cruz (1594, fol. 12, *r.*)
1,234,567 or 1.234.567	Greenwood (1729)
$\dot{6}8\vert765\vert4\ddot{3}2\vert189\vert7\dot{1}6\vert789\vert132$	Blassière (1769)

The groups have been called by various names, such as periods,[7] regions,[8] and ternaries,[9] and occasionally, as with Trenchant (1566), there were four figures in the right-hand group.[10]

Spanish Method of Writing Large Numbers. One of the most interesting examples of the writing of large numbers found in the books of the 16th century is seen in the work of Texeda,

[1] Similarly in Gemma Frisius (1540), Trenchant (1566), and various others.

[2] *E.g.*, the Dutch arithmetic of Wilkens, 1669, p. 8.

[3] As in a 14th century algorismus in the Columbia University Library. This plan is also followed by Tartaglia (1556, I, fol. 7, *r.*).

[4] *Liber Abaci*, p. 1.

[5] These cases contain errors in printing in the first (1503) edition of the *Margarita phylosophica*, but they are corrected in the later editions.

[6] Francesco Barozzi, *De Cosmographia Libri IV*, Venice, 1585.

[7] "Haec prima est periodus," etc., Ramus (1569; ed. Schoner, 1586, p. 2).

[8] Santa-Cruz (1594), fol. 12.

[9] Recorde, as quoted above.

[10] As in 10,500,340,8020 on page 16.

a Spanish writer of 1546. In seeking to explain algorism he writes numbers in the Spanish style (*en Castellano*) and also in algorism (*en guarismo*). The following cases are typical:

c. lx. U 462 q̊s . : ix U 621[1]	160 U 462 q̊s 009 U 621
c. iij U. 75 q̊s c. ij U 300	103 U 075 q̊s 102 U 300
Dcccxcj Uccxxxiiij q̊s Dlx U.	891 U 234 q̊s 560/000
vij U . . . q̊s Dxlv Ucccclxijm̄	7 U 000 q̊s 545 U 462

It will be seen that in the Spanish forms, doubtless owing to the Arab influence, there is a tendency (not uniformly carried out) to use the dot for zero. Texeda also mixes his algoristic numerals with the Roman,—a custom not uncommon after the 12th century. The U stands for thousands, appearing in earlier times as ∪ with several variants, and being of uncertain origin. The q̊s stands for *quentos* (*cuentos*, millions). In the 16th century the Greek θ was also used instead of U, as in XXXVθCCCXXVI for 35,326, and 637θ500 for 637,500, and in the 18th century it often degenerated into a kind of inverted C.[2] In Portugal a symbol $ (*cifrão*) was used as early as the 16th century for the same purpose.

3. Addition

Terminology of Addition. The name of the operation which we call addition has had its vicissitudes. One writer of the 13th century, for example, used "aggregation" instead.[3] Writ-

[1] In the original the ix is misprinted x.

The numbers at the left are *en castellano*; those at the right, *en guarismo*. The illustrations are from fols. iij, *v*., to iiij, *v*.

The number is 160,462,009,621.

See F. Cajori, "Spanish and Portuguese symbols for 'Thousands,'" *Amer. Math. Month.*, XXIX, 201, who had not seen Texeda; he suggests that the U came from some variant of the Roman symbol for thousand.

[2] See the "Fragmentos del Archivo Particular de Antonio Perez, Secretario de Felipe II," *Revista de Archivos, Bibliotecas y Museos*, XXV (1920), 140 (Madrid, 1920). In the author's library are several Spanish manuscripts of *c.* 1725–1750 with the degenerate form of θ referred to in the text.

[3] "Agregare est quoslibet duos numeros uel plures in unum colligere" (B. Boncompagni, *Trattati*, II, 30). We still preserve the phrase "in the aggregate." The word is merely the Latin for προστιθέναι (*prostithen'ai*), used by Euclid and Diophantus, or συντιθέναι (*syntithen'ai*), as used by Heron and Pappus.

ing about the year 1200, Fibonacci used "composition" and "collection" as well as "addition."[1] Nearly a century after Fibonacci the earliest French algorism (*c.* 1275) used "assemble"[2] for "add," and two centuries later the first printed arithmetic used "join."[3] In the early printed books the word "summation" was a rival of "addition,"[4] and we still speak of summing up, and of summing certain numbers. Addition being the operation most frequently used, the operation probably gave rise to the expression "to do a sum," meaning to solve a problem. Various other names for the process have been used,[5] but they have no special significance. With the English tendency to brevity, there is little prospect of change in this language in the words "add" and "addition."

In such of the early printed arithmetics as were intended for popular use there was ordinarily no word corresponding to our term "addend."[6] On the other hand, the theoretical books, generally printed in Latin, spoke of the *numeri addendi*, that is, the "numbers to be added,"[7] and from this came the word *addendi* alone, as used by Fine (1530),[8] Gemma Frisius (1540), and later writers. From this we have our English "addends." Those who seek for a change have occasionally used the less familiar "summands."

[1] In the Latin, *compositio*, *collectio*, and *additio*.

[2] "Se tu veus assambler .1. nombre a autre" (Boncompagni's *Bullettino* XV, 53).

[3] *I.e.*, *jongere*. Treviso arithmetic, 1478

[4] "Addirn oder Summirn," in Rudolff's arithmetic of 1526 (1534 ed., fol. 3) Stifel's *Deutsche Arithmetica* (1545, fol. 1), Albert's arithmetic of 1534, and many others. Grammateus (1518) has "Additio oder Summierung," and Adam Riese (1522; 1550 ed.) has a chapter on "Addirn/Summirn/Zusamen legen," the last term derived from counter reckoning.

[5] *E.g.*, the German *Zusammenthuung*, the French *aiouster*, and the Italian *recogliere*, *summare*, and *acozzare*.

[6] *E.g.*, Recorde (*c.* 1542), Sfortunati (1534), Baker (1568), Digges (1572) Peletier (1549), Trenchant (1566), Pagani (1591), and Pacioli's *Sūma* (1494) The early Dutch arithmeticians rarely had such a word, and even the American Greenwood (1729) does not give one.

[7] As in Scheubel (1545, p. 13), Clavius (1583), Licht (1500, fol. 2), and many others, but curiously not in Tonstall (1522). There were also such terms as *termini addendi*, *numeri colligendi*, *numeri summandi*, and the like.

[8] Thus, he speaks of the *addendorum summa* as well as the *numeri addend* (1555 ed., fol. 3).

The word "addend" was frequently used to refer only to the lower of two numbers to be added, as in the following case from the *Margarita phylosophica* (1503 ed.):

4′6′7′9′	numerus cui debet fieri additio
3′2′3′2′	numerus addendus
7 9 i i	numerus ꝓductus

It was also used by many writers to refer to all the numbers to be added except the top one.[1]

The result obtained in addition has had a variety of names, although "sum" has been the favorite.[2] Next in order of popularity is "product," a term used for the result of any operation, but particularly in addition and multiplication. It was popular in Germany,[3] especially in early times,[4] and was also used in the Latin countries.[5]

Some of the Latin books of the 16th century also used *numerus collectus*, based upon the use of *collectio* for addition, and possibly we might now be using "collect" for "sum" if the Church had not appropriated the term.

The Operation of Addition. The operation of addition has not changed much since the Hindu-Arabic numerals began to be used. Even with the Roman numerals it was not a difficult process, and it is not probable that a Roman banker was com-

[1] George of Hungary (1499) calls only the lower of the two numbers the *numerus addendus*: "et numerus addendus, qui debet scribi in inferiori ordine" (Budapest reprint of 1894, p. 4). The same usage is found in an unpublished algorism of *c.* 1400 in the British Museum (Sl. 3281, fol. 4, *v.*).

[2] Thus, Chuquet (1484) uses *some*, and similar forms appear in many early printed books, including those of Pacioli (1494), Fine (1530), Tonstall (1522), Klos (1538), Sfortunati (1534), and Riese (1522).

[3] Günther, *Math. Unterrichts*, p. 316.

[4] Joannes de Muris (*c.* 1350) says in his *Quadripartitum*: "Propositis namque numeris addicionis, supra figuras cuiuslibet numeri calculis situatis adde singulam singulis, arcubus obseruatis, et productum signa per calculos atque lege." *Abhandlungen*, V, 144. This is also interesting because it describes the use of counters on a Gerbert abacus.

[5] An interesting case sometimes occurs, as in Savonne's work of 1563, where "sum" is used for addend, and "product" for the result: "Adiouster est mettre plusieurs nombres ou sommes ensemble pour en sçauoir le produit."

pelled to resort to the abacus in ordinary addition. This will easily be seen by considering a case like the following:

DCCLXXVII
CC X VI
DCCCCLXXXXIII

We might write this result CMXCIII, but a Roman would rarely if ever have done so. Even in the 16th century we find forms analogous to this, as in the work of Texeda (1546),[1] where we have the following parallel arrangement:

xxjUcxxvij	21U127
x vUccxviij	15U218
ijUccccliiij	2U454
jU.x.	1U010
xxxjUclxxxij	31U182
lxxUDccccxcj	70U991

By using their alphabetic numerals the Greeks were able to perform various operations without recourse to an abacus, although the work was somewhat more complicated than it is with our numerals.[2]

Hindu Method. Bhāskara (*c.* 1150) gives as the first problem in the *Lilāvati* the following:[3] "Dear intelligent Lilāvati, if thou be skilled in addition . . ., tell me the sum of two, five, thirty-two, a hundred and ninety-three, eighteen, ten, and a hundred, added together." In a commentary on this work, of unknown date, the following method is given:

Sum of the units,	2, 5, 2, 3, 8, 0, 0	20
Sum of the tens,	3, 9, 1, 1, 0	14
Sum of the hundreds,	1, 0, 0, 1	2
Sum of the sums, ...		360

[1] Fol. v, *v*. [2] For details as to the Greek methods see Heath, *History*, I, 52.

[3] H. T. Colebrooke, *Algebra with Arithmetic and Mensuration from the Sanscrit*, p. 5 (London, 1817); hereafter referred to as Colebrooke, *loc. cit.*, or to special topics under the heads *Āryabhaṭa*, *Brahmagupta*, *Bhāskara*, *Vija Ganita*, with these spellings.

The Hindus seem generally to have written the sum below the addends, beginning with units' columns as we do. They had at one time another method, however, which they designated as inverse or retrograde, the operator beginning at the left and blotting out the numbers as they were corrected.[1]

65391
3279
10420
~~7~~~~8~~~~9~~~~8~~0
909

Arab Method and its Influence. The Arabs, on the other hand, often wrote the sum at the top, putting the figures of the check of casting out 9's at the side.[2] This plan was adopted by Maximus Planudes (*c.* 1340), the form used by him being here shown.[3]

8030	2
5687	8
2343	3

How the traces of the Oriental sand table, with its easily erased figures, and the traces of the old counter-reckoning, showed themselves in early English works is seen in the following passage in *The Crafte of Nombrynge*[4] (*c.* 1300):

lo an Ensampull of all

326

216

Cast 6 to 6, & þere-of[5] wil arise twelue. do away þe hyer 6 & write þere 2, þat is þe digit of þis composit. And þen write þe articulle þat is ten ouer þe figuris hed of twene as þus

1

322

216

[1] This method is here indicated by canceling. The plan is one naturally adapted to the sand abacus. On the dispute as to whether the Hindus used this abacus, see Chapter III. See also C. I. Gerhardt, *Études historiques sur l'arithmétique de position*, Prog., p. 4 (Berlin, 1856) (hereafter referred to as Gerhardt, *Études*); J. Taylor, *Lilawati*, Introd., p. 7 (Bombay, 1816) (hereafter referred to as Taylor, *Lilawati*).

[2] H. Suter, "Das Rechenbuch des Abû Zakarîjâ al-Haṣṣar," *Bibl. Math.*, II (3), 15.

[3] Wäschke, *Planudes*, p. 6; Gerhardt, *Études*, p. 20. On such general early methods in the various operations see F. Woepcke, *Sur l'introduction de l'Arithmétique Indienne en Occident* (Rome, 1859).

[4] See pages 32 and 78.

[5] As stated on page 32, the old letter þ is our *th*.

Now cast þe articulle þat standus vpon þe figuris of twene hed to þe same figure, reken þat articul bot for one, and þan þere will arise thre. þan cast þat thre to þe neþer figure, þat is one, & þat wul be foure. do away þe figure of 3, and write þere a figure of foure. and let þe neþer figure stonde stil, & þan worch forth.

This is the oldest known satisfactory explanation of an example in addition in our language.[1]

Special Devices. In the way of special devices, Gemma Frisius (1540) gives one that is still used in adding long columns. It consists in adding each column separately, writing the several results, and then adding the partial sums as here shown. It will be observed that Gemma writes the largest number at the top, the object being to more easily place the various orders in their proper columns.[2]

9279
389
479
27
22
9
9
10147

In manuscripts of this period dots are sometimes used, as is the case today, to indicate the figure to be carried.

Carrying Process. The expression "to carry," as used in addition, is an old one and, although occasionally objected to by teachers, is likely to remain in use. It probably dates from the time when a counter was actually carried on the line abacus to the space or line above,[3] but it was not common in English works until the 17th century. Thus, we have Recorde (*c.* 1542) using "keepe in mynde," Baker (1568) saying "keepe the other in your minde," and Digges (1572) employing the same phraseology and also saying "keeping in memorie," and "keeping reposed in memorie." The later popularity of the word "carry" in English is largely due to Hodder (3d ed., 1664). In the 17th cen-

[1] On similar methods in the medieval manuscripts, not merely in addition but in the other operations, see L. C. Karpinski, "Two Twelfth Century Algorisms," *Isis*, III, 396.

[2] "Obseruandum igitur primò, vti maior numerus superiori loco scribatur, minores huic subscribantur" (1575 ed., fol. A 7).

[3] See Chapter III.

tury the expression "to carry" was often used in Italy.[1] Expressions like "retain," "keep in mind," and "hold" have, however, been quite as common.[2]

4. Subtraction

Terminology of Subtraction. As with addition, so with subtraction, the name of the process and the names of the numbers used have varied greatly and are not settled even now. Outside the school the technical terms of arithmetic are seldom heard. When we hear a statement like "*Deduct* what I owe and pay me the *rest*," we hear two old and long-used terms instead of the less satisfactory words "subtract" and "difference."

Terms Meaning Subtract. While the word "subtract," meaning to draw away from under,[3] has been the favorite term by which to indicate the operation, it has by no means enjoyed a monopoly. When Fibonacci (1202), for example, wishes to say "I subtract," he uses some of the various words meaning "I take."[4] Instead of saying "to subtract" he says "to extract,"[5] and hence he speaks of "extraction."[6] These terms, as also "detract,"[7] which Cardan[8] (1539) used, are etymologically rather better than ours. "Subduction"[9] has also been used for "subtraction," both in Latin[10] and in English. Digges (1572),

[1] As in "summa senza portare," "portare decine," and the like. See, for example, the arithmetic of G. M. Figatelli Centese, fol. 21 (Bologna, 1664).

[2] *E.g.*, "die ander behalt" (Riese, 1522; 1533 ed.), "behalt die ander in sinn, welche ist zu geben der nechsten" (Grammateus, 1518), "et alterā mente reconda" (Clichtoveus, 1503; *c.* 1507 ed., fol. D 2), "& secunda reservanda" (Ramus, ed. Schoner, 1586, p. 6), "ie . . . retien le nombre de diszeines" (Trenchant, 1566; 1578 ed., p. 24).

[3] *Sub* (under) + *trahere* (whence *tractum*) (to draw).

[4] *Tollo*, *aufero*, or *accipio*.

[5] *Extrahere* (to draw out or take away from).

[6] *Extra*[*c*]*tio*.

[7] *Detrahere* (to draw or take from).

[8] *Practica*, 1539, capp. 7–14.

[9] *Sub* (under) + *ducere* (to lead).

[10] *E.g.*, Tonstall (1522) devotes fifteen pages to *Subductio*. He also says: "Hanc autem eandem, uel deductionem uel subtractionem appellare Latine licet" (1538 ed., p. 23; 1522 ed., fol. E 2, *r.*). See also Ramus, *Libri duo*, 1569, 1580 ed., p. 3; *Schol. Math.*, 1569 ed., p. 115. Schoner, in his notes on Ramus (1586 ed., p. 8), uses both *subduco* and *tollo* for "I subtract." Gemma Frisius (1540) has a chapter *De Subductione siue Subtractione*, and Clavius (1585 ed., p. 26) says: "Subtractio est . . . subductio." In his arithmetic Boethius uses *subtrahere*, but in the geometry attributed to him he prefers *subducere*.

for example, says: "To subduce or subtray any sūme, is wittily to pull a lesse frō a bigger nūber." Our common expressions "to diminish" and "to deduct" have also had place in standard works, as in the translation of the *Liber algorismi*[1] and in the work of Hylles (1592).[2] Recorde (*c.* 1542) used "rebate" as a synonym for "subtract," and the word is used today in commercial matters in a somewhat similar sense.

In a manuscript written by Christian of Prag[3] (*c.* 1400) the word "subtraction" is at first limited to cases in which there is no "borrowing." Cases in which "borrowing" occurs he puts under the title *cautela* (caution), and gives this caption the same prominence as *subtractio*.[4]

The word "subtract" has itself had an interesting history. The Latin *sub* appears in French as *sub*, *soub*, *sou*, and *sous*, *subtrahere* becoming *soustraire* and *subtractio* becoming *soustraction*.[5] Partly because of this French usage, and partly no doubt for euphony, as in the case of "abstract," there crept into the Latin works of the Middle Ages, and particularly into the books printed in Paris early in the 16th century, the form *substractio*.[6] From France the usage spread to Holland[7] and England, and from each of these countries it came to America. Until the beginning of the 19th century "substract" was a common form in England and America,[8] and among those brought up in somewhat illiterate surroundings it is still to be found.

[1] Which uses both *diminuere* and *subtrahere*. See Boncompagni, *Trattati*, II, 32.

[2] He uses "abate," "subtract," "deduct," and "take away." [3] See page 77.

[4] The passage begins: "*Cautela* . . . si figura inferioris ordinis non poterit subtrahi a sibi supraposita."

[5] With such variants as *soubstraction*, *soubstraire*, and the like.

[6] It appears in the *Geometria* of Gerbert, but the MSS. used are of *c.* 1200; ed. Olleris, p. 430. As to the early printed books, Clichtoveus (1503), for example, generally uses *substractio*, although *subtractio* is occasionally found. See also his edition of Boethius, and see the 1510 edition of Sacrobosco. The word also appears in the work of George of Hungary (1499), along with *subtractio*, so that the usage was unsettled.

[7] Thus Wentsel (1599), Van der Schuere (1600), Mots (1640), and, indeed, nearly all Dutch writers before 1800. Petri (1567; 1635 ed.), however, uses *subtractio* in the Latin form and *subtraheert* in the Dutch, and Adriaen Metius (1633; 1635 ed.) also omits the *s*.

[8] Our American Greenwood (1729), for example, always used "substract" and "substraction," but dropped the *s* in "subtrahend."

The incorrect form was never common in Germany,[1] probably because of the Teutonic exclusion of international terms.[2]

Minuend and Subtrahend. The terms "minuend" and "subtrahend," still in use in elementary schools, are abbreviations of the Latin *numerus minuendus* (number to be diminished) and *numerus subtrahendus* (number to be subtracted).[3]

The early manuscripts and printed books made no use of our abridged terms. The minuend and subtrahend were called the higher and the lower numbers respectively, as in *The Crafte of Nombrynge* (*c.* 1300), the upper and under numbers,[4] the number from which we subtract and the subduced,[5] the total and less,[6] the total and abatement, and the total and deduction.[7] Among the most popular terms have been "debt" and "payment,"[8] but better still are the terms "greater" and "less."[9]

[1] As witness Köbel (1514; 1549 ed., fol. 110), Stifel (*Arithmetica Integra*, 1544, fol. 2), Albert (1534; 1561 ed.), Thierfelder (1587, p. 11), and many others.

[2] Their early writers used such forms as *abzihung* and *abzyhung*, instead of "subtraction," just as the Dutch used such terms as *Af-trekkinge* (Van der Schuere, 1600; 1624 ed., fol. 10). While the Italians used *abattere* and *cavare*, they also used *sottrare* and *trarre* (as in Cataneo, 1546; 1567 ed., fol. 5).

[3] See Boncompagni, *Trattati*, II, 33, on Johannes Hispalensis (*c.* 1140) and his use of *numerus minuendus*.

[4] "Die vnder zal sol nit übertreffen die ôbern" (Grammateus, 1518). Tonstall (1522) and other Latin writers have *numerus superior* and *numerus inferior*; the Italian edition of Clavius (1586) has *numero superiore* and *numero inferiore*.

[5] "Numerus ex quo subducitur" and "subducendus" (Gemma Frisius, 1540; 1563 ed., fol. 9).

[6] *Totalis, minor*, used by Tzwivel (1505), Clichtoveus (1503), and others.

[7] Hylles, 1600, fol. 19.

[8] Thus, the Dutch-French work of Wentsel (1599, p. 4) has:

Schult/Debte.	£. 15846
Betaelt/paye,	£. 5424
Reste	£. 10422

The Dutch names in the 16th and 17th centuries were generally *de Schult* and *de Betaelinghe*.

Similarly, we have the Italian *debito*, *pagato*, and *residuo* (as in the 1515 edition of Ortega), the French *dette* and *paye*, as well as *la superieure & inferieure* (Trenchant, 1566; 1578 ed., p. 30), and the Spanish *recibo* and *gasto* (Santa-Cruz, 1594; 1643 ed., fol. 20).

[9] Sfortunati (1534; 1544 ed., fol. 8), *il numero maggiore* and *il numero minore*; G. B. di S. Francesco (1689), *quantità maggiore* and *quantità minore*; Raets (1580), *Het meeste ghetal* and *Het minste ghetal*, with similar forms in other languages. See also Tartaglia, 1592 ed., fol. 9.

Name for Difference. The words "difference" and "remainder" have never been popular, in spite of the fact that they are commonly found in the textbooks of today. The popular term has been "rest," and in common parlance this is still the case, as when we say "Give me the rest," "Take the rest." It appears in the first printed arithmetic[1] and is found generally in the works of the Latin countries. Indeed, the verb "to rest" was not infrequently used to mean subtract.[2] In England, Tonstall (1522), writing in Latin, used sometimes *reliqua*[3] and sometimes an expression like "the number sought." Recorde (*c.* 1542) introduced "remayner" or "remainer," a term which Hylles (1592) also used, together with "remaynder," "remaynes," and "rest." The Latin writers commonly used *numerus residuus*,[4] *differentia*, *excessus*,[5] and *reliqua*. Of these terms we have relics in our language in the forms of "difference" and "excess," and another term commonly used by us is "balance."

An interesting illustration of the use of expressions which later resulted in technical terms is seen in the following from the *Margarita phylosophica* (1503 ed.):

9001386 numerus a quo debet fieri subtractio
7532436 numerus subtrahendus
1468950 numerus relictus

The Operation of Subtraction. The process of subtraction, unlike the processes of addition and multiplication, has never been standardized. There are four or five methods in common use today, the relative advantage of any one over the others not being decided enough to give it the precedence. A brief history of a few of the more prominent methods will be given.

[1] Treviso, 1478, p. 18.

[2] Thus, the Spanish writer Santa-Cruz (1594) uses *restar*; and Ortega (1512; 1515 ed.) begins a chapter *per sapere restare o / subtrahere*.

[3] Various other writers did the same. Thus, Glareanus (1538) has *relictum* and *reliquum*. Fibonacci (1202) used *residuum* and *reliquus*.

[4] *E.g.*, Fine (1530) and occasionally Clavius (1583). An unpublished algorism of *c.* 1400, now in the British Museum (Sl. 3281, fol. 4, *v.*), uses *a q" sbtrahi, subtrahēd*^{9}, and *residuū* for the three terms.

[5] Clavius speaks of *differentia siue excessus*. 1585 ed., p. 133.

1. The complementary plan is based upon the identity

$$a - b = a + (10 - b) - 10.$$

In particular, to find $13 - 8$ we may substitute the simpler process $13 + 2$ and then subtract 10. This plan is today used in the case of cologarithms and on certain types of calculating machines. It is not a modern device, however. Bhāskara (*c.* 1150) used it in the *Lilāvati*,[1] and no doubt it was even then an old one. It appears in *The Crafte of Nombrynge* (*c.* 1300), and the difficulty of the operation is apparent from the following extract:

lo an Ensampul.

2122
1134

take 4 out of 2. it wyl not be, þerfore borro one of þe next figure, þat is 2. and set þat ouer þe hed of þe fyrst 2. & releue it for ten. and þere[2] þe secunde stondes write 1. for þou tokest on[3] out of hym. þan take þe neþer figure, þat is 4, out of ten. And þen leues 6. cast[4] to 6 þe figure of þat 2 þat stode vnder þe hedde of 1. þat was borwed & rekened for 10, and þat wylle be 8. do away þat 6 & þat 2, & sette þere 8, & lette þe neþer figure stonde stille,

and so on with equal prolixity. The expression to "borro," used in this work, was already old. It was afterwards used by Maximus Planudes (*c.* 1340), acquired good standing in the works of Recorde (*c.* 1542) and Baker (1568), and has never lost its popularity.


The same method appears in the Treviso arithmetic[5] (1478),

[1] Taylor, *Lilawati*, Introd., p. 7. [2] For "where." [3] One.
[4] *I.e.*, add; a relic of the abacus. Compare our expression "cast accounts."
[5] The author adds 2 to 2, the result being 4, and then adds 1 to the next figure of the subtrahend, saying:
"al .4. tu die iongere .i. e levera .5. poi dira .5. da .5. che equale da equale: resta .0." (Treviso arithmetic, p. [19]). The *i* is used for 1, as on page 97 and as is the *j* in the following problem from Huswirt.


	452
	348
Lo resto	104

and Huswirt (1501) solved his first problem in subtraction by this means, saying:

5 from 4 I cannot. I take the distance[1] of the lower number, that is, 5 from 10, or 5, and this I add to the upper number, 4, and obtain 9, which I write directly under the bar and below the 5. I carry the j in mind or on the tablet,[2] first canceling the 4 and 5, and add it to the next number, that is, to 9. . . .




Among other authors of early printed books who favored the plan there were such writers as Petzensteiner (1483), Pellos (1492), Ortega (1512), Fine (1530), Gemma Frisius (1540), Ramus (1555), Albert (1534), Baker (1568), and Digges (1572). Savonne (1563) also used it and indicated the borrowing of ten by means of a dot, as shown in the annexed example from his arithmetic.



The early American arithmeticians looked with some favor on the plan. Thus, Pike[3] says:

If the lower figure be greater than the upper, borrow ten and subtract the lower figure therefrom: To this difference add the upper figure.


2. The borrowing and repaying plan, in which the 1 that is borrowed is added to the next figure of the lower number, is one of the most rapid of the methods in use today and has for a long time been one of the most popular. It appears in Borghi's (1484) well-known work, the first great commercial arithmetic to be printed. Borghi takes the annexed example and says, in substance: "8 from 14, 6; 8 from 15, 7; 10 from 13, 3; 3 from 6, 3." The plan was already old in Europe, however. Fibonacci[4] (1202) used it, and so did Maximus Planudes (*c.* 1340). These writers seem to have inherited it from the Eastern Arabs, as did the Western Arab writer al-Qalasâdî




[1] *Distantia*, for the complement.

[2] Very likely the wax tablet, still used in Germany at that time. See Chapter III. [3] 1788; 1816 ed., p. 12. [4] *Liber Abaci*, Boncompagni ed., I, 22.

(*c.* 1475). The arrangement of figures used by Maximus Planudes in the subtraction of 35843 from 54612 is here shown, the remainder being placed above the larger number, after the Arab and Hindu[1] custom. The top line was used only in the checking process. This method of borrowing and repaying was justly looked upon as one of the best plans by most of the 15th and 16th century writers, and we have none that is distinctly superior to it even at the present time.



3. The plan of simple borrowing is the one in which the computer says: "7 from 12, 5; 2 from 3 (instead of 3 from 4), 1." This method is also very old. It appears in the writings of Rabbi ben Ezra[2] (*c.* 1140), the computer being advised to begin at the left and to look ahead to take care of the borrowing. This left-to-right feature is Oriental[3] and was in use in India a century ago.[4] It was the better plan when the sand table allowed for the easy erasure of figures, but it had few advocates in Europe.[5]



When the computation began at the right, the borrowing plan was also advocated by such writers as Gernardus[6] (13th century?), Sacrobosco[7] (*c.* 1250), and Maximus Planudes[8] (*c.* 1340). The writers of the early printed arithmetics[9] were

[1] Taylor, *Lilawati*, Introd., p. 7.

[2] *Sefer ha-Mispar*, ed. Silberberg, p. 29 (Frankfort a. M., 1895); hereafter referred to as Silberberg, *Sefer ha-Mispar*.

[3] It is found in the works of al-Khowârizmî (*c.* 825), Behâ Eddîn (*c.* 1600), Albanna (*c.* 1300), and others. See H. Suter, *Bibl. Math.*, II (3), 15.

[4] See Taylor, *Lilawati*, Introduction.

[5] One of these was Ramus, who advocates "subductio fit á sinestra dextrorsum" (*Arith. Libri duo*, 1569; 1580 ed., p. 4; 1586 ed., p. 8).

[6] *Algorithmus demonstratus*, I, cap. ix. Formerly attributed to Jordanus Nemorarius (*c.* 1225). See G. Enestrom, *Bibl. Math.*, XIII (3), 289, 292, 331.

[7] See J. O. Halliwell, *Rara Mathematica* (London, 1838–1839), 2d ed., 1841, p. 7; hereafter referred to as Halliwell, *Rara Math.*

[8] With one or two other methods.

[9] *E.g.*, such writers as Tzwivel (1505), Clichtoveus (1510 edition of his Boethius, fol. 39), Köbel (1514; 1549 ed., fol. 120), Stifel (1544), Ghaligai (1521), Raets (1580), and Clavius (1583). Some of the more pretentious writers, like Pacioli (1494) and Tartaglia (1556), gave all three methods.

not unfavorable to it, although they in general preferred the borrowing and repaying method.

4. The addition method, familiar in "making change," is possibly the most rapid method if taught from the first. To subtract 87 from 243 the computer says: "7 and 6 are 13; 9 and 5 are 14; 1 and 1 are 2"; or else he says: "7 and 6 are 13; 8 and 5 are 13; 0 and 1 are 1," the former being the better. The method was suggested by Buteo (1559) and probably by various other early writers, but it never found much favor among arithmeticians until the 19th century. It has been called the Austrian Method, because it was brought to the attention of German writers by Kuckuck (1874), who learned of it through the Austrian arithmetics of Močnik (1848) and Josef Salomon (1849).

$$\begin{array}{r} 243 \\ 87 \\ \hline 156 \end{array}$$

5. Multiplication

General Idea of Multiplication. The development of the idea of multiplication and of the process itself is naturally more interesting than the evolution of the more primitive and less intellectual processes already described. Just as addition is a device for obtaining results that could be reached by the more laborious method of counting, so multiplication was developed as an abridgment of addition.[1] It was simply a folding together of many equal addends. This is expressed not merely in the Latin name[2] but in the corresponding names in various other

[1] Attention was called to this fact by various 16th century writers. Thus Ramus (1569) remarks: "Multiplicatio est qua multiplicandus toties additur, quoties unitas in multiplicante continetur, & habetur factus." Schoner, in his commentary, adds: "Ideoq3 multiplicatio est additio, sed ejusdem numeri secum, nō diuersorum" (1586 edition of the *Libri duo*, p. 12). Even as early as *c.* 1341 Rhabdas mentioned the same fact. See P. Tannery, *Notices et extraits des manuscrits de la Bibl. nat.*, XXXII, 155.

[2] From *multus* (many) + *plicare* (to fold); compare also our word "manifold." The term is simply the Latin form of the Greek πολυπλασιάζειν (*polyplasia'zein*), as used by Euclid, Pappus, and Diophantus, or πολλαπλασιάζειν (*pollaplasia'zein*), as used by Heron and Pappus, the latter using both forms. Such words as "three-ply" and "four-ply" illustrate this use of *plicare*.

languages.[1] The Latin writers of the Middle Ages and the Renaissance speak of leading a number into this multiplicity,[2] which explains our use of the expression "*a* into *b*," still retained in algebra but discarded in arithmetic.

Definition of Multiplication. The definition of multiplication has often disturbed teachers of arithmetic because of their failure to recognize the evolution of such terms. It gave no trouble in the ancient *arithmetica*, for the numbers there involved, in speaking of such a process, were positive integers; whereas in the ancient *logistica* no attention, so far as we know, was paid to any definitions whatever. When, however, the notion of the necessity of exact definition entered the elementary school, teachers were naturally at a loss in adjusting the ancient limitations to the multiplication by a fraction or an irrational number, and by such later forms as a negative or a complex number.

One of the best of the elementary definitions referring to integers, and at the same time one of the oldest in our language, is found in *The Crafte of Nombrynge* (*c.* 1300): "multiplicacion is a bryngynge to-geder of 2 thynges in on nombur, þe quych on nombur contynes so mony tymes on, howe mony tymes þere ben vnytees in þe nowmbre of þat 2."[3] The same definition is found in the arithmetic of Maximus Planudes (*c.* 1340),[4] in the first printed arithmetic (1478),[5] in the first noteworthy com-

[1] Compare the German works of the 15th and 16th centuries, with their *mannigfaltigen* and *vervielfachen*. Grammateus (1518) speaks of "Multiplicatio oder Merung."

[2] "¶ Si aliquis numerus . . . ducatur," as Jordanus Nemorarius (*c.* 1225) says (1496 edition of the arithmetic, fol. C 3, *et passim*). Similarly Clichtoveus (1503), "Duco .4. in .3. et fit .jz."; Gemma Frisius (1540), "Mvltiplicare, est ex ductu vnius numeri in alterum numerum producere, qui toties habeat in se multiplicatum, quoties multiplicãs vnitatem"; and many others. In the Latin edition of his arithmetic (1583; 1585 ed., p. 36) Clavius has "Multiplicatio est ductus vnius numeri in alium . . . Vt numerus 6. in numerum 5. . . . duci dicitur . . .," but in the Italian edition (1586, p. 35) he uses *per* for *in*, thus: "Moltiplicare vn numero per vn' altro."

[3] R. Steele's proof-sheet edition, p. 21 (London, 1894).

[4] Wäschke, *Planudes*, p. 13.

[5] "Che moltiplicare vno nũero per si ouero per vno altro: non e altro: che de do nnmeri ꝑpositi: trouere vno terzo numero: el quale tante volte contien vno de quelli numeri: quante vnitade sono nel altro. Exempio .2. fia .4. fa .8. ecco che .8. cõtiẽ in se tante .4. quante vnitade sono nel .2." Treviso arithmetic, p. [27].

mercial arithmetic (1484),[1] and in numerous other works.[2] Recorde (*c.* 1542) set the English standard by saying, "Multiplication is such an operaciō that by ij sumes producyth the thyrde, whiche thyrde sūme so manye times shall cōtaine the fyrst, as there are vnites in the second."[3]

A somewhat more refined definition, including the notion of ratio, was necessary for fractional multipliers, and this appeared occasionally in the early printed books, as in Huswirt (1501).[4] Its use in English is largely due to the influence of Cocker's popular arithmetic (1677), where it appears in these words: "Multiplication is performed by two numbers of like kind, for the production of a third, which shall have such reason [ratio] to the one, as the other hath to unite." The idea is Oriental,[5] appearing in various Arab and Russian works.[6]

The elementary teacher generally objects to such a form as 2 ft. × 3 ft. = 6 sq. ft., and on the ground of pedagogical theory there is some reason for so doing, but not on logical or historical grounds. With respect to logic, it all depends on how multiplication is defined; while with respect to history there is abundant sanction for the form in the works of early and contemporary writers. For example, Savasorda[7] (*c.* 1120) and Plato of Tivoli (1116)[8] broaden the definition in such a way as to allow a line to serve as a multiplier, and Baker (1568) remarks, "If you wil multiply any number by shillinges and pence," an expression commonly paralleled by children today. Few of our contemporary physicists would see anything to criticize in such an expression as 6 ft. × 10 lb. = 60 foot-pounds, and in due time such forms will receive more recognition in elementary arithmetic.

[1] Borghi, 1540 ed., fol. 6.

[2] *E.g.*, Tonstall, 1522, fol. G 1; Stifel, *Arithmetica Integra*, 1544, fol. 2; Sfortunati, 1534, 1544/5 ed., fol. 11; Tartaglia, 1556; Trenchant, 1566.

[3] 1558 ed., fol. G1. Digges (1572; 1579 ed., p. 4) gives the same form.

[4] "Multiplicatio est numeri procreatio, proportionabiliter se habentis ad multiplicandū sicut multiplicans ad vnitatem se habet" (fol. 3).

[5] See Taylor, *Lilawati*, Introd., p. 15.

[6] *E.g.*, Behâ Eddîn (*c.* 1600). It is still used in Russian textbooks on arithmetic.

[7] Abraham bar Chiia.

[8] In his translation of Savasorda: "Et primum quidem exponemus, quid significare velimus, cum dicimus: multiplicatio lineae in se ipsam."

Terminology of Multiplication. Of the terms employed, "multiplicand" is merely a contraction of *numerus multiplicandus*. In *The Crafte of Nombrynge* (*c.* 1300) it is explained as "Numerus multiplicandus, Anglice þe nombur þe quych to be multiplied." In most of the early printed Latin books it appears in the full form,[1] but occasionally the *numerus* was dropped, leaving only *multiplicandus*,[2] and this led the non-Latin writers to use the single term.[3] A few of the Latin writers suggested *multiplicatus*,[4] so that we had at one time a fair chance of adopting "multiplicate." In their vernacular, however, many writers tended to use no technical term at all, simply speaking of the number multiplied, as the Latin writers had done,[5] and to this custom we might profitably return. It is hardly probable that such terms as subtrahend, minuend, and multiplicand, signifying little to the youthful intelligence, can endure much longer.

The word "multiplier" has had a more varied career. *The Crafte of Nombrynge* (*c.* 1300) speaks of "numerus multiplicans. Anglice, þe nombur multipliynge," the former being the Latin name for "multiplying number." Since the word *numerus* was frequently dropped[6] by Latin writers, in the translations the technical term appeared as a single word, with such

[1] *E.g.*, Clichtoveus (1510 edition of Boethius, fol. 35), Tonstall (1522), Grammateus (1518), Scheubel (1545, I, cap. 4).

[2] *E.g.*, Pacioli (1494, fol. 26), Licht (1500, fol. 6), Huswirt (1501), Ciruelo (1495), Glareanus (1538), Fine (1530).

[3] Thus Trenchant used *multiplicande* (1566; 1578 ed., p. 35). On the Italian writers see B. Boncompagni, *Atti d. Accademia Pontificia di Nuovi Lincei*, XVI, 520; hereafter referred to as *Atti Pontif.*

[4] *E.g.*, Tzwivel (1505) used "nūerus multiplicādus siue multiplicatus," and Gemma Frisius (1540) used *multiplicandus* and *multiplicatus* interchangeably. So the Treviso arithmetic (1478) says: "Intendi bene. che ne la moltiplicatione sono prīcipalmente do numeri necessarii. zoe el nūero moltiplicatore: et el nūero de fir moltiplicato" (p. [27]), which is not quite the same usage.

[5] Thus Chuquet (1484) simply speaks of "le nombre multiplie," and similarly with Borghi (1484), Riese (1522; 1529 ed., p. 8), Sfortunati (1534; 1544/5 ed., fol. 11), and others. Digges (1572) speaks of it as "the other summe, or number to be multiplied."

[6] Thus Johannes Hispalensis (*c.* 1140); see Boncompagni, *Trattati*, II, 41. Clichtoveus (1510 edition of Boethius) uses *multiplicans* both as an adjective and as a noun. So also Huswirt (1501, fol. 3), Ciruelo (1495; 1513 ed., fol. A 6), Grammateus (1518; 1523 ed., fol. A 4), Gemma Frisius (1540), and many others.

variants as "multiplicans,"[1] "moltiplicante,"[2] "multiplicator,"[3] "multipliant,"[4] and "multiplier."[5]

The word "product" might with almost equal propriety be applied to the result of any other arithmetic operation as well as to multiplication. It means simply a result,[6] but it has some slightly stronger connection with multiplication on account of the use of the verb *ducere* in the late Latin texts.[7] It has, however, been used in the other operations by many writers, and its special application to the result of multiplication is comparatively recent. The tendency to simplify the language of the elementary school will naturally lead to employing some such term as "result" for the various operations.

The authors of the early printed books often took the sensible plan of having no special name for the result in multiplication. Certain of them used "sum"[8] or "sum produced,"[9] while *factus*, a natural term where *factor* is employed, had its advocates.[10] Finally, however, the *numerus* was dropped[11] from *numerus productus*, and "product" remained.[12]

[1] Pacioli, 1494, fol. 26.

[2] Cataldi, 1602, p. 21.

[3] *Multiplicador* (Pellos, 1492, fol. 8), *multiplicatore* (Ortega, 1512; 1515 ed., fol. 16), *multiplicatour* (Baker, 1568; 1580 ed., fol. 16). The word was common in English. Greenwood used it, with "multiplier," in the 1729 American arithmetic.

[4] Peletier, 1549; 1607 ed., p. 34.

[5] "The lesse is named the Multiplicator or Multiplyer." Digges, 1572; 1579 ed., p. 5.

[6] Latin *producere* (to lead forth); whence *productum* (that which is led forth).

[7] See pages 101, 102.

[8] *E.g.*, Pacioli (1494, fol. 26), Ortega (1512; 1515 ed., fol. 18), and Recorde (*c.* 1542; 1558 ed., fol. G 2). Fine (1530) uses it as well as *numerus productus*.

[9] Thus Hodder (10th ed., 1672, p. 25) speaks of "The Product, or sum produced." Similarly, Clichtoveus (1503) uses both *numerus productus* and *tota summa*, and Glareanus (1538) uses *summa producta*.

[10] Thus, Fibonacci (1202) uses "factus ex multiplicatione." He also speaks of the "contemptum sub duobus numeris." Ramus (1569) speaks of the *factus* in multiplication, and in his treatment of proportion he says: "Factus à medio aequat factum ab extremis."

[11] As in Licht (1500), Huswirt (1501), Gemma Frisius (1540), and Scheubel (1545).

[12] *Produit*, Trenchant (1566); *produtto*, Sfortunati (1534) and Cataldi (1602), or *prodotto* by later Italian writers. Unlike most of the Latin terms it found place in the early Teutonic vocabulary, as seen in Werner (1561) and such Dutch writers as Petri (1567), Raets (1580), and Coutereels (1599).

The Process of Multiplication. We know but little about the methods of multiplication used by the ancients. The Egyptians probably made some use of the duplation plan, 17 being multiplied by 15 as shown in the annexed scheme.[1] It is also probable that this plan was followed by other ancient peoples and by their successors for many generations, which accounts for the presence of the chapter on duplation in so many books of the Renaissance period. Indeed, even as good a mathematician as Stifel multiplied 42 by 31 by successive duplation, substantially as here shown.[2]

1	17
2	34
4	68
8	136
16	272
1	17
15	255

$1 \cdot 42 =$	42
$2 \cdot 42 =$	84
$4 \cdot 42 =$	168
$8 \cdot 42 =$	336
$16 \cdot 42 =$	672
$31 \cdot 42 =$	1302

There is also a contemporary example of the use of duplation and mediation, found among the Russian peasants today. To multiply 49 by 28 they proceed to double 28 and to halve 49, thus:

49	24	12	6	3	1
28	56	112	224	448	896

The fractions are neglected each time, and finally the figures in the lower row which stand under odd numbers are added, thus:

$$28 + 448 + 896 = 1372,$$

and this is the product of 28×49.[3]

Greek and Roman Methods. The abacus[4] was probably used so generally in ancient times that we need hardly speculate on the methods of multiplication used by the Greeks and Romans. It is quite possible, however, that the Greeks multiplied upon

[1] P. Tannery, *Notices et extraits des manuscrits de la Bibl. nat.*, XXXII, 125; *Pour l'histoire de la science hellène*, p. 82 (Paris, 1887); Heath, *History*, I, 52.

[2] See his *Rechenbuch* (1546), p. 12. He uses a similar process for division.

[3] In the above case we have

$\frac{49}{1}$	$\frac{49}{2}$	$\frac{49}{4}$	$\frac{49}{8}$	$\frac{49}{16}$	$\frac{49}{32}$
49	24	12	6	3	1
$1 \cdot 28$	$2 \cdot 28$	$4 \cdot 28$	$8 \cdot 28$	$16 \cdot 28$	$32 \cdot 28$

Here $49 \cdot 28 = (32 + 16 + 1) \cdot 28 = 896 + 448 + 28 = 1372$.

[4] See Chapter III.

their wax tablets about as we multiply, but beginning with their highest order;[1] and there is no good reason why the Romans should not have done very nearly the same. Indeed, in Texeda's arithmetic (1546) the "Spanish method" with Roman numerals is given side by side with the new method of "Guarism," thus:

ccclxxv———vijUD.vj	ȝ7ȝ———7U506
ijqsccljUDccc..	2251U800
DxxvUccccxx.	525U420
xxxvijUDxxx.	37U530
ijqsDcccxiiijUDccl.	2q̈s814U750

It should be observed, however, that the small tradesman has never had much need for this kind of work.[2]

Pacioli's Eight Plans. Our first real interest in the methods of multiplication starts with Bhāskara's *Lilāvati* (*c.* 1150), although we have a few earlier sources. Bhāskara gives five plans and his commentators add two more. These plans had increased to eight when Pacioli published his *Sūma* (1494), and these will now be considered.

Our Common Method of Multiplying. Our common form was called by Pacioli "Multiplicatio bericocoli vel scachierij," and appears in his treatise (fol. 26, *r.*) in the following form:


Multiplicandus. 9 8 7 6
Producentes.
Multiplicans. 6 7 8 9


		8	8	8	8	4
	7	9	0	0	8	
6	9	1	3	2		schachieri
5	9	2	5	6		Bericuocolo
summa	6	7	0	4	8	1 6 4

[1] P. Tannery, *Notices et extraits des manuscrits de la Bibl. nat.*, XXXII, 126; Heath, *History*, I, 54.

[2] An interesting witness to this fact is the first Bulgarian arithmetic (1833), described in *L'Enseignement Mathématique* (1905), p. 257.

He says that the Venetians called this the method "per scachieri" because of its resemblance to a chessboard,[1] while the Florentines called it "per bericuocolo" because it looked like the cakes called by this name and sold in the fairs of Tuscany.[2] In Verona it was called "per organetto,"[3] because of the resemblance of the lines to those of a pipe organ, and "a scaletta" was sometimes used because of the "little stairs" in the figure, as seen on page 107.[4]





This method is not found directly in the *Lilāvati*, but two somewhat similar ones are given. In the first of these[5] the multiplier is treated as a one-figure number and the work begins at the left, as shown above; in the second,[6] which is shown in the above computation at the right, the multiplier is separated as with us, but the work begins at the left as in the preceding case.

[1] *Scacchero*, the modern *scacchiere*. Our word "exchequer" comes from the same root. See page 188. The spelling varies often in the same book.

[2] ". . . el primo e detto multiplicare p Scachieri in vinegia ouer per altro nome per bericuocolo in firença . . . el primo modo di multiplicare chiamano. Bericuocolo: perch' pare la figura de q̄sti bricuocoli: o cōfortini che se vendano ale feste" (fols. 26, *r.*, 28, *v.*). A MS. in Dresden, dated 1346, has "lo modo di moltiplicare per ischachiere." B. Boncompagni, *Atti Pontif.*, XVI, 436, 439. An undated MS. in the Biblioteca Magliabechiana (Florence, C. 7. No. 2645) gives the name as *iscacherio*, *scacherio*, and *ischacherio*. Cataneo (1546; 1567 ed., fol. 10), although printing his work in Venice, calls the method *biricvocolo*. Those who do not have access to Pacioli may find the methods in facsimile in Boncompagni, *Scritti inediti del P. D. Pietro Cossali*, p. 116 (Rome, 1857), a work more likely to be found in university libraries.

[3] So Feliciano da Lazesio (1526) says: "Del multiplicar per scachier vocabulo Venitiano, ouer baricocolo uocabolo Fiorentino, ouer multiplicar per organetto uocabolo Veronese" (1545 ed., fol. 12). Similarly, Tartaglia (1556) says: "Del secondo modo di multiplicare detto per Scachero, ouer per Baricocolo, ouer per Organetto" (*General Trattato*, I, 23 (Venice, 1556)).

[4] "Multiplicatione a scaletta .&. aggregatione a bericocolo," in a MS. at Paris, described by Boncompagni, *Atti Pontif.*, XVI, 331.


[5] The *Swarupa gunanam*, "the multiplier as a factor." It is Bhāskara's first method. For the method of Mahāvīra (*c.* 850) see his *Gaṇita-Sāra-Sangraha*, Madras, 1912, p. 9 of the translation (hereafter referred to as *Mahāvīra*).

[6] The *St'hana gunanam*, "multiplication by places." It is his fourth method.

Since multiplication on the abacus required no symbol for zero, the earlier attempts with the Hindu-Arabic numerals occasionally show the influence of the calculi. This is seen in a Paris MS. in which the multiplication of 4600 by 23 is described in a manner leading to the form here shown.[1] It is possibly in forms like this that the chessboard method had its origin.

CM	XM	M	C	X	I
		4	6		
		1	8		
	1	2			
	1	2			
	8				
1		5	8		
				2	3

The name *scachiero* was used for a century after the chessboard form had entirely disappeared. The Treviso arithmetic (1478) does not attempt to mark off the squares, but the author uses the name,[2] as did various other Italian writers.[3] It was also used occasionally in Germany,[4] England,[5] and Spain,[6] but less in other countries.



Even after this method was generally adopted, the relative position of the figures was for a long time unsettled. In the oldest known German algorism[7] the multiplier appears above the multiplicand. In the Rollandus MS. (Paris, *c.* 1424)[8] the arrangement is as here shown. In the Treviso arithmetic the multiplier is sometimes placed at the right, as seen in the

[1] M. Chasles, *Comptes rendus*, XVI, 234 (1843).

[2] ". . . attēdi al terzo modo. zoe al moltiplicare per scachiero" (fol. 19).

[3] Borghi (1484) gives only this method, designating it "per scachier."

[4] *E.g.*, Petzensteiner (1483) says: "Also ich wil multipliciren in Scachir."


[5] Recorde (*c.* 1542) speaks of "one way that is wrought by a checker table" (1558 ed., fol. G8).

[6] Thus Texeda (1546) describes multiplication "escaqr o berricolo."

[7] A. Nagl, "Ueber eine Algorismus-Schrift des XII. Jahrh.," *Zeitschrift für Mathematik und Physik*, Hl. Abt., XXXIV, 129. This journal is hereafter referred to as *Zeitschrift* (Hl. Abt.).

[8] *Rara Arithmetica*, p. 446.

annexed facsimile; Widman (1489) gives the same arrangement in his second method; and this lateral position of the multiplier is preserved in our synthetic multiplication in algebra. The placing of the multiplier above the multiplicand[1] is possibly due to the fact that the writers of that period did not greatly concern themselves as to which of the two numbers was the operator, although the smaller one was more often chosen. The difficulty of settling down to a definite arrangement of figures is seen by a study of the various editions of the Taglientes' popular *Libro dabaco*,[2] eight of which[3] give the following examples:



SCACCHERO MULTIPLICATION

From the Treviso arithmetic, 1478. In this case the multiplier is placed at the right

(1515)	(1520)	(1541)	(1547)
456	456	456	456
23	23	23	23
1368	1368	136 8	1368
912	912	912	912
10488	10488	1048 8	10488

(1550)	(1561)	(1564)	(1567)
456	456	456	456
23	23	23	23
1368	1368	1368	1368
912	912	912	912
10488	10488	10488	10488

The various other editions give arrangements similar to the above. Some of these forms are doubtless due to printers' errors, but as a whole they go to show that a definite plan had

[1] *E.g.*, Sfortunati, 1534; 1545 ed., fol. 11.

[2] 1515, the work of two authors.

[3] The dates appear in parentheses.

not been agreed upon in the 16th century, although the general chessboard method was given the preference by most writers,[1] other methods being looked upon as mere curiosities. Thus Hylles (1600) says:

Also you shall vnderstand, that there are besides these sundrie other waies of Multiplication, aswell with squars as without which if you list to learne I referre you to M. Records ground of artes, where you may finde plentie of varietie.

The Castle Method. The second plan of multiplying laid down by Pacioli was, on account of the form of the work, known as "the castle" or, in Florence, "the little castle."[2] The significance of the name is best understood from the first example given by Pacioli.

	9876	6	Per .7
		6	
	6789	1	Proua
		61101000	
Castelucio		5431200	
		476230	[*sic*]
		40734	
Sūma		67048164	.1.

It will be observed that the figures are arranged somewhat like the wall and turret of a castle.[3] The scheme was merely a copy-book invention of the Italian schoolmasters and, although enduring until the close of the 17th century or later,[4] was always looked upon as a puerile method.[5]

[1] Thus Tartaglia (1556) calls it "vn modo generalissimo da nostri antichi pratici ritrouato, & piu di alcun' altro vsitato." Pagani (1591), although giving a list of methods like Pacioli, prefers this one, calling it "molto vago" (very pretty) and "molto sicuro" (very certain).

[2] "Del multiplicare per castello ouero castelluccio vocabulo Fiorentino" (Feliciano, 1526; 1545 ed., fol. 12). Pacioli (1494) says of it: "El secondo modo di multiplicare e detto castellucio" (fol. 26, *r.*), and a MS. of Benedetto da Firenze of *c.* 1460 calls it "elchasteluccio." In Spain (Texeda, 1546) it appears as "El .2. modo le dize castellucio."


[3] The figures at the right are the proof by casting out 7's.

[4] It appears in Ciacchi's *Regole Generali d' Abbaco*, fol. B3 (Florence, 1675).

[5] Thus Pagani (1591) says: "ma più tosto capriciose ch' vsitato, & vtile."

The Column Method. The third method given by Pacioli is known as the column or tablet plan.[1] By this method the computer refers to the elaborate tables, always in columns, like those used by the Babylonians, which are found in many of the 15th century manuscripts. It is essentially nothing but a step in the development of such elaborate and convenient multiplication tables as those of Crelle and others, which appeared in the 19th century.

Cross Multiplication. Pacioli's fourth method was that of cross multiplication, still preserved in our algebras and used, in simple cases, by many computers. To this he gave the name "crocetta" or "casella,"[2] adding that it is more fantastic and ingenious than the others.[3] His most elaborate illustration is that of 78×9876:



[1] "El terço e detto multiplicare ꝑ colōna ouer atauoletta" (1494 ed., fol. 26). Cataneo (1546; 1567 ed., fol. 8) gives it another name, "Del mvltiplicar a la memoria detto uulgarmente Caselle o Librettine," and the Taglientes (1515) speak of it as "ꝑ cholonella." Texeda (1546), who follows Pacioli very closely, speaks of it as "colona o taboleta." As "per colona" it is the first method given in the Treviso (1478) arithmetic, and this name is also used by Borghi (1484). Tartaglia (1556) calls it the oral or mental plan ("per discorso, ouer di testa") as well as "per colona" and "per colonella."

[2] "De .4°. mō multiplicandi dicto crocetta siue casella." The Treviso arithmetic calls it the method of the simple little cross: "Attendi diligētamente a lo segondo modo: zoe moltiplicare per croxetta simplice." The name *crocetta* was the more common one, for Feliciano (1526) speaks of it as "per crocetta o voi dire per casela"; Cardan (1539) gives it only the name "modus multiplicādi ꝑ cruceta"; and Tartaglia (1556) and Cataneo (1546) call it merely "per crosetta." *Crocetta* means a little cross, and *Casella* (a little house) is often used for "pigeonhole."

[3] ". . . piu fantasia e ceruello che alcūo d' glialtri." He admires it, however, as "bella e sotil e fo bel trouato" (fol. 28, *r.*).

The filling of the vacant places by zeros, in 0078, was not unusual among the Arabs. Thus in a manuscript of one of the works of Qosṭâ ibn Lûqâ[1] (*c.* 900) the multiplication of 21,600 by 4 appears in this form:



The plan is ancient, appearing in the *Lilāvati* (*c.* 1150) as the *tatst'ha* method, or method of the stationary multiplier, in distinction from the advancing multiplier, where the multiplying figures were advanced one place to the right after each partial product was found. This method is shown on page 118. The method given in the *Lilāvati* is fully explained by Ganeśa (*c.* 1535) in his commentary on Bhāskara.[2] It also appears in the arithmetic of Planudes (*c.* 1340), but in the forms here shown:[3]

840	114048	76842
24	432	1423
35	264	0054

While the first of these cases is simple, it is doubtful if the other two were practically used.

Some of the work of the Taglientes (1515) is related to cross multiplication, as may be seen from the illustration here given.



AN ITALIAN METHOD OF MULTIPLICATION

From the 1541 edition of the Taglientes' *Opera* of 1515

[1] The MS. is dated 1106 A.H. (1695 A.D.).

[2] Colebrooke, *Lilāvati*, p. 6 n.


[3] Wäschke, *Planudes*, p. 14.

On account of the difficulty of setting the crossed lines, printers often used the letter X between the multiplicand and the multiplier, and this may have suggested to Wright (1618) the multiplication sign (×) used by him and his contemporaries. In Pagani's arithmetic of 1591, for example, the work in cross multiplication appears as follows:

3		2		4		3		2
	X				I	X	I	X I
2		5		3		6		4
	800					157248		

He recognized that the method is not very practical with numbers of more than two figures.[1]

The Method of the Quadrilateral. The fifth method given by Pacioli is that of the quadrilateral.[2] It was really nothing but the chessboard plan with the partial products slightly shifted, as is seen in the illustration from the Treviso arithmetic (1478) here shown.



MULTIPLICATION PER QUADRILATERO

From the Treviso arithmetic, 1478

Gelosia Method. There seems to be no good reason why Pacioli should have postponed to the sixth in order the so-called gelosia, or grating, method, also known by the name of the quadrilateral, the square,[3] or the method of the cells,[4]

[1] "Il moltiplicare a crocetta di tre figure, è assai piu dificile de primo. . . . Il moltiplicare per crocetta di 4. figure è più dificile delli sopra nominati, & quanto più sono figure, tanto più sono dificile" (p. 17).


[2] "El quinto mō e detto ꝑ q̄drilatero" (fol. 26, *r.*). It appears in Pagani's work (1591) as "per quadrato." It is often merged with the *gelosia* method next mentioned, as when Tartaglia (1556) calls it "Per Quadrilatero, ouero per gelosia."

[3] "El sexto modo e detto ꝑ gelosia ouer graticola." Tartaglia (1556) says: "Il quinto modo di multiplicare è detto Quadrilatero qual è assai bello, perche in quello nō vi occorre a tener a mente le decene." Various other names are used, such as "modo de quadrato" (Feliciano, 1526), "ꝑ quadro" (16th century MS. of Gio. Dom. Marchesi), "per squadrado" (undated Bologna MS.).

[4] ". . . per le figure de le camerete," "dala fugura dela camerella," the cells also being called "camere triangulate." This is in an undated Turin MS. See B. Boncompagni, *Atti Pontif.*, XVI, 448.

and to the Arabs after the 12th century by such names as the method of the sieve[1] or method of the net.[2]

The method is well illustrated by two examples from the Treviso (1478) book here shown. It will be observed that the diagonals separate the tens and units and render unnecessary



FIRST PRINTED CASE OF GELOSIA MULTIPLICATION

From the Treviso arithmetic (1478), showing the form from which the Napier rods were developed

the carrying process except in adding the partial products. These diagonals sometimes slant one way and sometimes another, but in general the direction from the upper right-hand corner to the lower left-hand corner was the favorite.[3]

The method is very old and might have remained the popular one if it had not been difficult to print or even to write the net. It was very likely developed first in India, for it appears in Ganeśa's commentary on the *Lilāvati* and in other Hindu works.[4] From India it seems to have moved northward to

[1] In the writings of Albanna (*c.* 1300), possibly due to his commentator, al-Qalasâdî (*c.* 1475). See F. Woepcke, *Journal Asiatique*, I (6), 512.


[2] H. Suter, "Das Rechenbuch des Abû Zakarîjâ al-Ḥaṣṣâr," *Bibl. Math.*, II (3), 17.

[3] See the Marre translation of Behâ Eddîn's (*c.* 1600) *Kholâçat al Hissâb*, Paris, 1846; Rome, 1864, p. 13; hereafter referred to as Behâ Eddîn, *Cholâṣat*. Also see books as late as Giuseppe Cortese's *Aritmetica* (Naples, 1716).

The contrary direction is seen in MSS. of Ibn al-Hâ'im (*c.* 1400), dated 1132 A.H. (1720 A.D.), Albanna (*c.* 1300), and various other Arabic writers.

[4] Colebrooke, *Lilāvati*, p. 7. See also the introduction to Taylor, *Lilawati*, pp. 20, 33. It was called by the Hindus *Shabakh*.

China, appearing there in an arithmetic of 1593.[1] It also found its way into the Arab and Persian works, where it was the favorite method for many generations.




GELOSIA METHOD OF MULTIPLYING

From an anonymous manuscript written in Florence *c.* 1430. The author describes it as *Multiplicha p modo de Quadrato*

From the Arabs it passed over to Italy and is found in many manuscripts of the 14th and 15th centuries. In the printed books it appeared as late as the beginning of the 18th century,[2] but more as a curiosity than as a practical method.

As to the name *gelosia*, Pacioli's statement is more complete than that of any of his contemporaries:

> The sixth method of multiplying is called *gelosia* or *graticola* . . . because the arrangement of the work resembles a lattice or *gelosia*. By *gelosia* we understand the grating which it is the cus-



GELOSIA MULTIPLICATION AS GIVEN BY PACIOLI, 1494

Showing the same double arrangement of diagonals as in the Treviso book of 1478

[1] Libri, *Histoire*, I, 386, 389.

[2] *E.g.*, in the 1690 edition of Coutereels's *Cyffer-Boeck*; in Padre Alessandro's *Arimmetica* (Rome, 1714); and in Giuseppe Cortese's *Aritmetica* (Naples, 1716).

tom to place at the windows of houses where ladies or nuns reside, sc they cannot easily be seen. Many such abound in the noble city of Venice.[1]

The Repiego Method. Another method that was popular enough to survive and to have a place in some of our modern textbooks was called by the early Italians the "modo per repiego," that is, the method by composition, or, more exactly, by decomposition, of factors. For example, to multiply by 72, multiply by 9 and then by 8, thus saving the addition of partial products.[2] It is one of several methods inherited by the Italians through the Arabs, from Hindu sources.[3]

The Scapezzo Method. Pacioli's eighth method was commonly known among the Italians as *a scapezzo*, or multiplying by the parts, not the factors, of the multiplier.[4] Tartaglia (1556) gives as an illustration

$$26 \times 67 = (3 + 4 + 5 + 6 + 8) \times 67$$
$$= 201 + 268 + 335 + 402 + 536 = 1742;$$

but he could not have considered it as other than a curiosity although it was recognized by such writers as Ramus and Schoner.[5] It goes back to Bhāskara at least.[6]

[1] "Gelosia intendiamo quelle graticelle c̄h si costumono mettere ale finestre de le case doue habitano dōne acio nō si possino facilmē e vedere o altri religiosi Diche molto abonda la excelsa cita de uinegia" (fol. 28, *r.*). The word found its way into French as *jalousie*, meaning a blind, and thence passed into German and was carried even to the Far East, where it is met with today.

[2] Pacioli explains the term thus: "Repiego de vn numero se intende el producto de doi altri numeri che multiplicati vno nel laltro fanno quel tal numero aponto: del quale essi sonno ditti repieghi" (fol. 28, *v.*). It is Tartaglia's third method. See also Terquem's *Bulletin*, Vol. VI, and B. Boncompagni, *Atti Pontif.*, XVI, 404.


[3] It appears in Taylor's notes on the *Līlāvati* under the name *vibhaga gunanan* (submultiple multiplication). See translation, p. 8 n.


[4] "De octauo modo multiplicandi dicto aschapeçço." Pacioli, 1494, fol. 29, *r.* Tartaglia gives the name as "spezzato, ouer spezzatamente." *General Trattato* 1556, I, 26. In Texeda's Spanish work of 1546 it appears as *escapeço*.

[5] 1586 ed., p. 16.

[6] See Taylor's translation, p. 8. The name in his notes is *khanda gunanan* (parts multiplication).

Minor Methods of Multiplying. Besides the leading methods given by Pacioli there are many variations to be found in other early works. One of the most valuable is the left-to-right method, the "allo adietro" plan of Tartaglia,[1] still used to advantage in some cases but hardly worth teaching. The Arabs occasionally used it,[2] and the Hindus varied it by beginning with the lowest order of the multiplier and the highest order of the multiplicand.[3]





Another variant is seen in a cancellation method which went by various names. The Arabs called it the Hindu plan,[4] and Taylor[5] found the Hindus using it early in the 19th century. Al-Nasavî (*c.* 1025)[6] and other Arab writers thought highly enough of the method to give it a place in their works. It may be illustrated by the case of 76 × 43. The figures were first written as shown in the upper rectangle. The multiplication began with 7 × 4 = 28. As soon as a figure had served its purpose it was erased on the abacus and its place was taken by another. This procedure was modified in India, probably long after 200 B.C., when ink came into use,[7] the figures being canceled as here shown.[8]

In the 15th and 16th centuries there were numerous vagaries of the copy-book makers, the extensive discussion of which is

[1] *General Trattato*, 1556, I, 25, *v*. Calandri (1491) gives a page (fol. 8) to the method.

[2] *E.g.*, al-Karkhî, *c.* 1020. See H. Hankel, *Geschichte der Mathematik*, Leipzig, 1874, pp. 56, 188; hereafter referred to as Hankel, *Geschichte*.

[3] An elaborate example of the late use of the method is in Marten Jellen's *Rekenkundige Byzonderheden* (1779), p. 13.

[4] This translation of the term *Hindasi*, used by many Arab writers, is, however, disputed. It also means numerical, a translation that would have little significance in a case like this. See page 64.


[5] *Lilawati*, Introd., p. 9.

[6] F. Woepcke, *Journal Asiatique*, I (6), 497.

[7] See G. Bühler, *Indische Palaeographie*, pp. 5, 91 (Strasburg, 1896).

[8] This example is from an Arab arithmetician, Mohammed ibn Abdallâh ibn 'Aiyash, Abû Zakarîyâ, commonly known as al-Ḥaṣṣâr (*c.* 1175?). See H. Suter, *Bibl. Math.*, II (3), 16. Substantially the same plan is used by al-Qalasâdî (*c.* 1475), *ibid.*, p. 17. For various arrangements followed by the Hindus see Colebrooke, *Lîlâvati*, p. 7 n.


not worth while. Suffice it to say that certain teachers had their pupils arrange the partial products in the form of a rhombus, and even as good mathematicians as Tartaglia (1556) and Cataldi (1602) multiplied "per Rumbo" and "à Rombo." Others arranged the figures so that the outline of the work looked like a cup, chalice, or beaker.[1] The better writers, however, recognized that such work was time-consuming.[2]



THE FIRST EXAMPLE IN MULTIPLICATION PRINTED IN THE NEW WORLD

From the *Sumario cõpedioso* of Juan Diez, published in Mexico in 1556. The problem is to multiply 978 by 875 and the method is essentially *per copa*, that is, the method of the cup, so called because the figure resembles a drinking cup


Labor-Saving Short Methods. Besides the general methods already described there were many special devices for the saving of labor. Even when the multiplication table was learned, the medieval computers did not require it beyond 5×10, and various plans were developed for operating under this limitation when an abacus was not conveniently at hand. Of these methods, one of the best known is a complementary plan that is found in many of the 16th century books. To multiply 7 by 8, write the numbers with their complements to 10, as here shown. Then either $8 - 3$ or $7 - 2$ is 5, and $2 \times 3 = 6$, so that 56 is the product, the operation not requiring the multiplication table beyond 5×10 in any case. It is given by such writers as Huswirt (1501), Fine (1530), Riese (1522), Rudolff (1526), Stifel (1544), Recorde (*c.* 1542),



[1] Thus Tartaglia calls the plan "Per Coppa, ouer per Calice," and Cataldi says: "dalla forma loro si possono chiamare à Calice, Coppa, Tazza, ò Bicchiere." In Spanish (Texeda, 1546) it appears as "per copa" and is incorrectly given as "a la frãcesa" (a French method). It is essentially the method used by Juan Diez, Mexico, 1556, as shown in the illustration.

[2] So Tartaglia: "trouate piu per mostrar vn piu sapere, che per alcuna vtilità" (1592 ed., fol. 40). Other curious forms are given by Coutereels (1690 ed., p. 8) under the title "Vernakelijke Multiplicatie" (interesting multiplication).

Peletier (1549), and Baker (1568), while Peurbach (*c.* 1450; 1st ed., 1492) speaks of it as an ancient rule.[1] It was commonly used in connection with finger reckoning.[2] For example, to find 6×9, raise one finger on one hand and four fingers on the other hand, these representing the respective complements of 9 and 6. Then multiply the standing fingers for the units and add the closed fingers for the tens. The plan is still in use among the peasants in certain parts of Russia and Poland. As a variant of this method the following plan was until recently in use in certain towns in Russia: Number the fingers on each hand from 6 to 10, and give to each the value 10. Then the products of 8×8 and 7×9 are found as indicated below:



$8 \times 8 = 6 \times 10 + 2 \times 2 = 64$ $\qquad$ $7 \times 9 = 4 \times 10 + 2 \times 10 + 1 \times 3 = 63$

A few of the most common of the complementary methods will now be briefly indicated.

Various Arab and Persian writers[3] multiplied 8 by 7 by the relation $ab = 10b - (10 - a)b$, as here shown.[4]

In this case we have

$$\begin{aligned} 7 \times 8 &= 10 \times 8 - (10 - 7) \times 8 \\ &= 10 \times 8 - 3 \times 8 \\ &= 80 - 24 \\ &= 56. \end{aligned}$$

$$\begin{array}{r} 10 \times 8 = 80 \\ 3 \times 8 = \underline{24} \\ 56 \end{array}$$

The method was considered valuable by writers like Widman (1489), Riese (1522), Rudolff (1526), and Scheubel (1545).

[1] ". . . regulam illam antiquam."


[2] See pages 196 and 201.

[3] *E.g.*, Behâ Eddîn (c. 1600).

[4] G. Enestrom, *Bibl. Math.*, VII (3), 95.

Widman, Rudolff, and Grammateus (1518) used the relation

$$ab = 10b - 10(10 - a) + (10 - a)(10 - b)$$
$$= 10[b - (10 - a)] + (10 - a)(10 - b),$$



Trattatus

¶De multiplicatione Capl'm quartum

Multiplicatio eſt numeri procreatio.proportionabiliter ſe habentis ad multiplicandū ſicut multiplicatus ad vnitatem ſe habet.exempli gratia 3 ad 4 multiplicare eſt numerum 12 procreare. que ſic multiplicando videlicet 4 proportionantur quemadmodum multiplicans.ſcilicet 3 vnitati correſpondet.quia vtraq; eſt proportio tripla. Item multiplicatio prerequirit q; ſcitene multiplicationem digitorum inter ſe ſciat. Cuius talis datur regula Scribantur digiti ſubalterne.et cuilibet differentia a denario verſus dextram ponas.quas inter ſe multiplica.et productum inferius ſcribe.dein de differentiam vnius a digito alterius ſubtrahe et priori producto poſtpone.et prouenient ſumma.vt patet in figura

Exemplum. ſcies 8 quot ſunt multiplicationes eorū a denario.ſcilicet 4 et 2 in ſe.et erunt 8 que ſcribe.Deinde differentiam vnius a digito alterius ſubtrahe.et relictum priori producto poſtpone.vt patet in figura. ¶Alia regula de multiplicatione duorum numerorum infra 20 quorum quilibet duabus figuris ſcriptus eſt.propoſitis itaq; duobus numeris primas inferiorius cum prima ſuperiorius multiplica.et procreabitur numerus vna vel duabus figuris ſcribendus.ſi vna.ſcribatur.ſi duabus.primam harum ſcribe.ſecundam ſeruando in mente. Deinde iterum eisdem'ad ſe addas.et producto precedenti ſemper figuram in mente reſeruatam adiunge.et prouenient numerus duabus figuris vel vna ſcribendus Si vna.ſcribatur.et de vltimis figuris priorum numerorum eſt vnitas accipi debet.quam poſtpone prioribus et erit ſumma. Si autem duabus primam harum ſcribe.ſecundam vnitati a poſteriori b figuris accipiende adde.quas ſimul ſcribe.et erit ſumma

Exemplum

multiplica 4 in 7.et erunt 28 ſcribe 8. 2 ſeruādo i mente. Deinde adde primas figuras ad inuicem dicendo 4 ad 7 ſunt 11.et 2 ſeruata ſunt 13.ſcribe 3 poſt 8 vnā rem ſeruando in mente.quam addas ſigure vltimo accipiende.et erunt 2 que poſtpone 38 et erunt 238. ¶Pro dictorum dicendorum q; faciliori intellectu digitorum inter ſe multiplicatiuis tabula hic poſtea ponitur

COMPLEMENTARY MULTIPLICATION

From Huswirt's *Enchiridion nouus Algorismi summopere visus De integris*, Cologne, 1501. Much reduced

illustrated in the following multiplication of 8 by 7.

Here we have

$$7 \times 8 = 10 \times 8 - 10 \times (10 - 7) + (10 - 7)(10 - 8)$$
$$= 80 - 30 + 6$$
$$= 56.$$

$10 \times 8 =$	80
$- 10 \times 3 =$	− 30
$+ 3 \times 2 =$	6
	56

Further Algebraic Relations. The following relations also had their advocates:

$$ab = 10(a + b - 10) + (10 - a)(10 - b);^{1}$$

$$ab = 10(a - b + 2 \cdot \overline{b - 5}) + (10 - a)(10 - b);$$

$$ab = (10 - a)(10 - b) + 10(a + b) - 100;^{2}$$

$$(10a + a)(10b + b) = [(10a + a)b + ab]10 + ab;^{3}$$

$$(10a + b)(10a + c) = (10a + b + c)a \times 10 + bc;^{4}$$

$$(10a + b)(10a - c) = 100a^2 + 10ab - (10ac + bc);^{5}$$

$$ab = (a + b - 10) \times 10 - (a - 10)(10 - b),\ a > 10,\ b < 10;^{6}$$

$$(10 + a)(10b + c) = (ab + 10b + c) \times 10 + ac;$$

$$(3a)^2 = 10a^2 - a^2;\ (3a + 1)^2 = (3a)^2 + [(3a + 1) + 3a];$$

$$(3a - 1)^2 = (3a)^2 - [(3a - 1) + 3a];^{7}$$

$$a^2 = 10a - (10 - a) \times a;$$

$$a^2 = (\tfrac{1}{3}a)^2 \times 10 - (\tfrac{1}{3}a)^2,\ \text{where } a = 3m;$$

$$a^2 = (a - 1)^2 + [a + (a - 1)],\ \text{where } a = 3m + 1;$$

$$a^2 = (a + 1)^2 - [a + (a + 1)],\ \text{where } a = 3m + 2;^{8}$$

$$(a + b)(a - b) = a^2 - b^2;^{9}$$

$$(5a)^2 = 10a^2 \times 2\tfrac{1}{2};$$

[1] *E.g.*, Behâ Eddîn and Riese.
[2] The Petzensteiner arithmetic, Bamberg, 1483.
[3] Al-Karkhî (*c.* 1020), as in

$$22 \times 44 = (22 \times 4 + 8)10 + 2 \times 4 = 968.$$

[4] Al-Karkhî, Behâ Eddîn, Tartaglia. *E.g.*,

$$23 \times 27 = (23 + 7) \times 2 \times 10 + 3 \times 7 = 600 + 21 = 621.$$

[5] Al-Karkhî and Tartaglia. [6] An Arab writer, al-Kashî, *c.* 1430.
[7] Elia Misrachi (*c.* 1500) and Rabbi ben Ezra (*c.* 1140).
[8] Rabbi ben Ezra, who recognized that the limitations on a were unnecessary.
[9] Well known to the Greeks and given by Euclid.

$$(5a+1)^2=(5a)^2+[(5a+1)+5a];^1$$

$$(10+a)(10+b)=100+10(a+b)+ab;^2$$

$$ab=\left(\frac{a+b}{2}\right)^2-\left(\frac{a-b}{2}\right)^2.^3$$

The method of aliquot parts was also well known in the Middle Ages, both in Europe and among the Arabs, and the 16th century writers frequently gave our common rules of multiplying by numbers like 11 and 15. Beginning at least as early as the 14th century, multiplication by numbers ending in one or more zeros was commonly effected as at present.[4]

Contracted multiplication, the work being correct to a given number of significant figures, is a development intended to meet the needs of modern science. It began to assume some importance in the 18th century,[5] although a beginning had already been made by Bürgi (*c.* 1592) and Prætorius (*c.* 1599).[6]

The Multiplication Table. The oldest known arrangement of the multiplication table is by columns. This is the one always found on the Babylonian cylinders and the one commonly used by the Italian writers on mercantile arithmetic in the formative period of the subject. In general, no product appeared more than once; that is, after $2 \times 3 = 6$ was given, 3×2 was thought

[1] Elia Misrachi gives various rules of this kind.

[2] Huswirt (1501).


[3] This is the rule of quarter squares, which still has its advocates. It is probably due to the Hindus. See A. Hochheim, *Kâfî fîl Hisâb*, p. 7 (Halle a. S., 1878) (hereafter referred to as Hochheim, *Kâfî fîl Hisâb*); H. Weissenborn, *Gerbert*, p. 201 (Berlin, 1888). It is found in the *Talkhys* of Albanna (*c.* 1300), the work of al-Karkhî (*c.* 1020) mentioned above, and the works of Behâ Eddîn (*c.* 1600) and other Oriental writers. The preferred transliteration of the name of al-Karkhî's work is *al-Kâfî fi'l-Ḥisâb*, but the more familiar title as given in the European editions has been adopted in this work. See Volume I, page 283.

[4] *E.g.*, Maestro Paolo dell' Abbaco (*c.* 1340); see G. Frizzo's edition, p. 42 (Verona, 1883). Bianchini's correspondence with Regiomontanus (1462) contains it; see M. Curtze, *Abhandlungen*, XII, 197, 270. It is also in the Treviso arithmetic (1478), Pellos (1492), and other early works. There are, however, various cases in which it was not recognized in the 16th century.

[5] Greenwood's American arithmetic (1729) gives the reversed multiplier.

[6] M. Curtze, *Zeitschrift* (Hl. Abt.), XL, 7.

to be unnecessary, a view still taken by Japanese arithmeticians[1] and having much to commend it. The early Italian mercantile arithmetics gave, for purpose of easy reference, tables with the products of all primes to 47×47, or often to 97×97. Computers turned to these columns for the simpler products needed in multiplication *per colonna*. The Italians obtained the idea from the East, Rhabdas (1341) giving the column tables "which the very wise Palamedes taught me."[2]



MEDIEVAL MULTIPLICATION TABLE

Part of a table from an anonymous Italian MS. of *c.* 1456, but apparently a copy of an earlier work of *c.* 1420


The second arrangement was the square form generally used by nonmercantile writers and known as the Pythagorean Table,[3] whereof, as Hylles (1600) remarks, "Some affirme Pythagoras to be the first author." This mistaken idea was held by various early writers,[4] although the better ones seem to have recognized

[1] Smith and Mikami, *History of Japanese Mathematics*, p. 37 (Chicago, 1914); hereafter referred to as Smith-Mikami.

[2] P. Tannery's translation in *Notices et extraits des manuscrits de la Bibl. nat.*, XXXII, 167. For their use by Benedetto da Firenze, Luca dell' Abacho, and others, see *Rara Arithmetica*, p. 464 and elsewhere. They are also found in Pacioli (1494), Pellos (1492), Borghi (1484), and the Treviso book (1478), and in many other works. See also D. E. Smith, "A Greek Multiplication Table," *Bibl. Math.*, IX (3), 193.

[3] Table de Pythagore, Tabula Pythagorica, Mensa Pythagorae, Mensula Pythagorae, Tavola Pitagorica, Mensa Pythagorica, and other similar names are common.

[4] Thus Köbel (1514) speaks of "Der Pythagorisch Tisch oder Tafel" as "von dem Fürstē Pythagora geordnet" (1518 ed., fol. 17).




MULTIPLICATION TABLE (*c.* 1500)

The table as it appeared in an anonymous Latin MS. of *c.* 1500, being the same form as the one found in various MSS. of Boethius

that the later Pythagoreans were the inventors.[1] It is found in the arithmetic of Boethius[2] and in a work attributed to Bede (*c.* 710),[3] but the fact that Rhabdas (*c.* 1341) does not give it[4] suggests that the Greeks did not use it. It was common in the medieval works[5] and in the early printed books.[6] Some writers carelessly attributed it to Boethius,[7] while others arranged tables of addition, subtraction, and division on the same plan and gave to them the name of Pythagoras.[8]

The third standard form was the triangular array. It appears in a Prag manuscript[9] in the form here shown, but there are several variants. It is given in *The Crafte of Nombrynge* (*c.* 1300) as "a tabul of figures, where-by þou schalt se a-nonn) ryght what is þe nounbre þat comes of þe multiplicacion) of 2 digittes." Widman (1489) speaks of it as a Hebrew device,[10] and at any rate it is quite likely to be Arabic.[11] It was not so popular in the early textbooks as the columnar and square arrangements, although it was used by such writers as Widman (1489), Gemma Frisius (1540), Recorde (*c.* 1542), Baker (1568), and Trenchant (1566).



[1] Thus Boethius says: "Pythagorici . . . quam ob honorem sui praeceptoris mensam Pythagoream nominabant" (Friedlein ed., p. 396). See also A. Favaro in Boncompagni's *Bullettino*, XII, 148. Clavius (1583) says: "quòd Pythagoras eam vel primus excogitauerit, vel certe discipulos suos in ea mirifice exercuerit."

[2] Friedlein ed., p. 53. On the text see Boncompagni's *Bullettino*, XV, 139.

[3] *De arithmeticis numeris*, of doubtful authorship, where the "Pythagorica Mensa sive abacus numerandi" is given in full to 20×20, with the more important products as far as 1000^2.

[4] P. Tannery, *Notices et extraits des manuscrits de la Bibl. nat.*, XXXII, 121.

[5] *E.g.*, Jordanus Nemorarius (*c.* 1225), Rollandus (1424), and al-Kashî (*c.* 1430).

[6] *E.g.*, Tzwivel (1505) and such commentators as Faber Stapulensis and Clichtoveus.

[7] Thus Stifel (1545) says: "Dise tafel hat Boetius gesetzt."

[8] Possibly Ramus (1569) began this, for he gives these tables and says: "Hic Pythagoraeus additionis abacus est," and so for subtraction and division.


[9] See S. Günther, Boncompagni's *Bullettino*, XII, p. 149; very likely the MS. of Christian of Prag, already referred to on pages 77, 95.

[10] "Das erst ist eynn taffel geformiret auff den triangel geczogen aus3 hebraischer zungen oder iudischer."

[11] Behâ Eddîn (*c.* 1600) gives it in his *Kholâṣat al-ḥisâb*.

The extent to which the tables were carried varied considerably from time to time. Tables used for reference, as we might use Crelle's tables today, go back to ancient times, one of the 5th century giving the important products to 50×1000.[1] The medieval writers were usually content to stop with 20×20,[2] however.

For tables to be committed to memory it was sufficient, in the days of the medieval abacus, to go only to 5×10; even 4×9 was far enough for practical purposes.[3] Many of the 16th century writers outside of Italy found it necessary to urge their pupils



TRIANGULAR AND SQUARE FORMS OF THE MULTIPLICATION TABLE

From Widman's arithmetic (Leipzig, 1489), the edition of 1500

[1] "Victorii Calculus ex codice Vaticano editus a Godofredo Friedlein," in Boncompagni's *Bullettino*, IV, 443.

[2] As in a MS. written before 1284 and copied in 1385, described by Steinschneider in the *Bibl. Math.*, XIII (2), 40. See also Beldamandi's work (1410), printed in 1483, where the products extend to 22×22.

[3] See the devices for finding such products as 7×8, page 119. Thus Rudolff (1526; 1534 ed., fol. D 8) says: "Das ein mal eins . . . musten zum ersten wol in kopff fassen / doch nit weiter dañ bis auff 4 mal 9." Clavius, while recommending the learning of the table to 10×10, says: "Qvod si huiusmodi tabula in promptu nō sit, vtendum erit hac regula," namely, the one given on page 119.

very strongly to learn the table, showing that the custom was relatively recent in countries where the abacus had only just been abandoned or where its use was diminishing.[1] One Spanish writer says that it should be known as thoroughly as the Ave Maria,[2] and Digges (1572) encourages his pupils by saying: "This Table therefore first printe liuely in thy remembrance, and then boldly proceede farther, all difficultie I assure thee is past."

It may interest those teachers who feel that they must insist upon "two threes are six" instead of "two times three are six" to know that the former has at least some kind of remote sanction in a terse Latin form,[3] although in most languages the use of "times" has been general.

6. Division

Definition of Division. Division has generally been considered as the fourth of the fundamental operations,[4] the fifth when numeration is included, or the seventh when duplation and mediation are considered separately. In general the operation

[1] So Chuquet (1484) says: "¶Item plus est necesz^{e} de sauoir tout de cueur la multiplication dune chascune des .10. figures par soy mesmes et aussi par une chascune des aultres La quelle chose est appelle le petit liuret de algorisme." (From A. Marre's MS. copy in the author's library.) This "livret de algorisme" was a common name for the small multiplication table, "gli libretti minori" of the Italians, "gli libretti maggiori" referring to the table beyond 10 × 10. (Spelling as in the Dagomari MS. described in *Rara Arithmetica*, p. 435.) The couplet often found in 16th century books,

¶Lern wol mit fleisz das ein mal ein
So wirt dir alle rechnung gmein,

appeared first in print, so far as I have found, in Widman's work of 1489; 1508 ed., fol. 11.

[2] ". . . laquale tabula bisogna sapere ad memoria como la Aue Maria" (Ortega, 1512; 1515 ed., fol. 16). Thierfelder (1587) says: "Aber wer das ein mal eins nicht fertig lernet . . . wird nimermehr keinen fertigen Rechner geben" (p. 16). Metius is equally urgent: "Tabula Pitagorica, dieman wel vast in sijn memorie moet hebben" (1635 ed., p. 5).

[3] Thus Scheubel (1545, I, cap. 4) says: "Sixies septem sunt 42. septies quinq3 sunt 35," and so on, in which, however, the word "times" is concealed.

[4] The "quarto atto" of the Treviso arithmetic (1478).

has been known either as division[1] or as partition,[2] but many writers use both terms.[3] Thus Baker (1568) speaks of "Deuision or partition," and Digges (1572) says "To deuide or parte."[4]

As in the case of multiplication, no satisfactory definition, adapted to the understanding of beginners, is possible, since the concept is constantly extended as the pupil proceeds. To say that "diuision sheweth onlely howe often the lesse summe is conteyned in the bigger,"[5] or that

> Diuision doth search how oft the diuisor
> In Diuidend may be quoted or found
> Whereof the quotient is the decidor,[6]

is to exclude cases like 6 ft. $\div$ 2 or $3 \div 4$, although the latter was intentionally barred out by many writers[7] for the reason that a result like $\frac{3}{4}$ could not be "times" in the primitive use of the word. The early idea was manifestly that of an integral divisor and an integral quotient.[8]

A second definition which has had some sanction is that of finding a number which is contained as many times in the dividend as unity is contained in the divisor. It has long been used, being found in Maximus Planudes (*c.* 1340)[9] and the Treviso arithmetic.[10] An improvement upon this definition, and quite

[1] *E.g.*, with such medieval writers as Fibonacci (1202), *Liber Abaci*, p. 27. This is the idea of measuring, and so Euclid used the term μετρεῖν (*metrein'*) to mean both to measure and to divide.

[2] *E.g.*, the Treviso arithmetic, Huswirt (1501), Ghaligai (1521), Stifel (1544), Scheubel (1545), Cataldi (1602), Ortega (1512), Savonne (1563), and Santa-Cruz (1594). This form was preferred by Heron, Pappus, and Diophantus, all of whom used μερίζειν (*meri'zein*, to part).

[3] *E.g.*, Pacioli (1494), Tartaglia (1556), Trenchant (1566), Clavius (1583).

[4] So with some of the Dutch writers. Thus in the Dutch-French work of Wentsel (1599): "Deuisio: dat is deelinge," "Diuisio: e' esta dire, partir."

[5] Digges, 1572; 1579 ed., p. 8.

[6] Hylles (1600), the word "quoted" being interesting as related to "quotient."

[7] Thus Tzwivel (1505) says: "Officiuȝ diuisionis est cognoscere quotiens minor nūerus in maiore reꝑiat," and Peletier (1549) says: "c'est sçauoir combien de fois vn moindre nombre est contenu en vn plus grand."

[8] Thus Clavius: "Divisio est distributio propositi numeri in partes ab altero numero dato denominatas" (1583; 1585 ed., p. 48).

[9] Wäschke, *Planudes*, p. 23.

[10] "Trouare vno terzo nūero: el quale se troua tante volte nel mazore: quãte vnitade sono nel menore."

sufficient for pure number, is one that is based upon ratio,—the finding of a number which has to unity the same ratio as the dividend has to the divisor. It is often found in the 16th century books.[1]

It was natural in the Middle Ages, when division as performed on the abacus was often based upon subtraction, to base the definition also upon the latter operation.[2] This plan was followed by such writers as Ramus,[3] Schoner,[4] and Peletier[5] and has not wholly died out even yet.

Of all the elementary definitions the one most generally approved describes the operation as seeking a number which, multiplied by the divisor, is equal to the dividend, and it serves the purpose fairly well. It is perhaps the oldest definition extant[6] and it has the sanction of many scholarly writers.[7]

Two-fold Nature of Division. The above definitions do not, in general, distinguish between the two notions of division illustrated by the cases 6 ft. $\div$ 3 ft. $= 2$ and 6 ft. $\div 2 = 3$ ft., although the last definition includes both cases. Rudolff (1526) seems to have been the first to make this distinction perfectly clear,[8] and Stifel (1545) to have been the second.[9] Tartaglia[10] also gave it, and thereafter it was mentioned by various writers of the 16th and 17th centuries.

[1] "Diuisio est numeri ꝑcreatio ꝑportionabiliter se ad vnitatem habẽtis vt diuidẽdus ad diuisorẽ" (Huswirt, 1501, fol. 5).

[2] "Numerum per numerum diuidere est maiorem secundum quantitatem minoris partiri, uidelicet minorem de maiore tociens subtrahi, quociens in eo poterit inueniri." Johannes Hispalensis, *Liber Algorismi* (*c.* 1140), in Boncompagni's *Trattati*, II, 41.

[3] *Arith. libri duo*, 1569.

[4] *Tabulae Astronomicae*, 1536, fol. A 3*a*.

[5] 1549; 1607 ed., p. 48, as a secondary definition.

[6] J. P. A. Erman, *Life in Ancient Egypt*, p. 364, English translation by Tirard, New York, 1894 (hereafter referred to as Erman, *Egypt*), attributes it to the Egyptians.

[7] *E.g.*, Cataldi: "Il partire è modo di trouare vna quantità, quale moltiplicata per vna quantità proposta (ouero con la quale moltiplicando vna quantità proposta) produca vna quantità data" (1602, p. 32).

[8] "Diuidirn heisst abteilen. Lernet ein zal in die ander teilen/auff das man sehe/wie offt eine in d'andern beschlossen werde/oder wieuil auff einen teil kome" (1534 ed., fol. 8).

[9] *Deutsche Arithmetica*, 1545, fol. 1, where it is more clearly stated.

[10] *General Trattato*, 1556, I, fol. 27, *r*.

Terminology of Division. Early writers commonly gave names to only two of the numbers used in division, the *numerus dividendus* (number to be divided) and the *numerus divisor*.[1] These are, of course, not technical terms, and they appear as mere colloquial expressions in various medieval works. Gradually, however, the *numerus* was dropped and *dividendus* and *divisor* came to be used as technical nouns, as at present.[2] Such names as "answer" or "result" were commonly used for quotient[3] and were quite as satisfactory.

The names of the terms have undergone various changes. The divisor has frequently been called the "parter"[4] or the "dividens,"[5] but our present term has been the one most commonly used. The dividend has generally been called by this name, although there have been terms equivalent to "partend," with the usual linguistic variants.[6] The quotient has frequently been called the product,[7] the part,[8] the exiens,[9] and the outcome,[10] but the term used by English writers has been the favorite in most of the leading European languages.[11]

[1] Thus Clichtoveus, in his commentary on Boethius (1503; 1510 ed., fol. 36), says: "In divisione tres requiruntur numeri. Primus est numerus diuidēd^{9} & maior/ex hypothesi dandus. Secundus/numerus diuisor siue diuidens: etiam assignandus ex hypothesi. Tertius est numerus ex diuisione proueniens: & hic est querendus," no name being given for this quotient and no mention being made of a remainder.

[2] *E.g.*, in the Rollandus MS. (1424), where *quotiens* is also used. Joannes de Muris (*c.* 1350) used *dividendus* and *numerus quociens*, but not *divisor*. See *Abhandlungen*, V, 145.

[3] From *quoties*, how much.

[4] "L'autre qui le diuise, s'apele parteur, partisseur, ou diuiseur" (Trenchant, 1566; 1578 ed., p. 51). Chuquet (1484) calls it the *partiteur*, and Cataldi (1602) uses *il partitore*, following the Treviso book and other Italian works of the time. Pellos (1492), writing in a dialect mixture of French, Italian, and Spanish, called it the *partidor*. In the Teutonic languages it appeared in the 16th century as Theiler, Deyler, Teyler, Deeler, and deylder.

[5] ". . . nūer^{9} diuisor siue diuidēs," Tzwivel, 1505, fol. 6, and various other Latin works.

[6] Ortega (1512; 1515 ed.) calls it "la partitione," as he calls the multiplicand "la multiplicatione." Santa-Cruz (1594) calls it "suma partidera." Digges (1572) writes it "diuident."

[7] *E.g.*, Gemma Frisius (1540) and numerous other Latin writers.

[8] *E.g.*, the Treviso book (1478) gives "la parte."

[9] *E.g.*, Scheubel (1545).

[10] In Dutch, the *Uitkomst*.

[11] Of course with such variants as *quotiens* in the Latin books, *cociente* in the Spanish (Santa-Cruz, 1594), and so on.

For obvious reasons the name for the remainder has varied more than the others. The medieval Latin writers used *numerus residuus*, *residuus*, and *residua*, and various other related terms, and certain later authors employed the same word for the remainder as for the fraction in the quotient.[1]

The Process of Division. The operation of division was one of the most difficult in the ancient *logistica*, and even in the 15th century it was commonly looked upon in the commercial training of the Italian boy as a hard matter.[2] Pacioli (1494) remarked that "if a man can divide well, everything else is easy, for all the rest is involved therein." He consoles the learner, however, by a homily on the benefits of hard work.[3] So impressed was Gerbert (*c.* 980) by the difficulties to be overcome that he gave no less than ten cases in division, beginning with units by units, treated by continued subtraction.[4] Even as late as 1424 Rollandus gave only the simplest cases with small numbers, and nearly two centuries later Hylles (1600) recognized the difficulties when he said, "Diuision is esteemed one of the busiest operations of Arithmetick, and such as requireth a mynde not wandering, or setled vppon other matters."[5]

1	8
*2	16
$\frac{1}{2}$	4
*$\frac{1}{4}$	2
*$\frac{1}{8}$	1

Early Form of Division. Probably the oldest form of division is the one used by the Egyptians. This was based upon the processes of duplation and mediation. Thus, to divide 19 by 8 we may arrange the work as here shown. We take $2 \times 8 = 16$, $\frac{1}{2}$ of $8 = 4$, and so on, and select the numbers in the right-hand column which have 19 for their sum; for example, $16 + 2 + 1 = 19$. The quotient is therefore $2 + \frac{1}{4} + \frac{1}{8}$, the multipliers being marked here by asterisks.[6]

[1] *E.g.*, G. B. di S. Francesco (1689). In the case of $7 \div 3 = 2$, and 1 remainder, or $2\frac{1}{3}$, he uses *auanzo* for the 1 and also for the $\frac{1}{3}$. In a MS. of 1736 in the Woolwich Academy, England, "remainer" is used exclusively for "remainder."

[2] "Dura cosa e la partita" is a phrase often met.

[3] "Peroche nulla virtus est sine labore. E questo aferma el phylosopho q3 virtus cõsistit circa difficile" (fol. 32, *v.*).

[4] M. Chasles, *Comptes rendus*, XVI, 284.

[5] Fol. 37.

[6] Erman, *Egypt*, p. 365.

We are quite ignorant as to the way in which the Greeks and Romans performed the operation of division before the Christian Era. We have, however, a case described in the 4th century by Theon of Alexandria (*c.* 390), in which the literal numeral system of the Greeks is used and the work is not unlike our own, except that sexagesimal fractions are employed.[1] Since we know so little of the development of the operation among the ancients, we shall proceed at once to the history of the subject, showing particularly how long division was performed after the introduction of our modern numerals, say from about the year 1000.

Short Division. The simplest method, however, was the one which we call in English short division, which is based upon the recognition of the products in the columns of the multiplication table, and which has therefore been known as division by the column,[2] by rule, or by the table, as oral division, or as division in the head.[3] The method is illustrated in the Treviso book as follows:

Lo partitore	.2.	7624	o lauanzo,
La parte		3812	

which means that $7624 \div 2 = 3812$, with o remainder. The arrangement used by Sfortunati (1534) for a similar case is seen in the following example taken from his arithmetic:

P 14

1037382

$74098\frac{10}{14}$

[1] The details are given in Heath, *History*, I, 58.

[2] *Per colona* (Treviso, 1478); *per cholona* (Borghi, 1484).

[3] "Partire a regolo: ouer a tauoletta" (Pacioli, 1494; 1523 ed., fol. 32). Pacioli advises: "E comenza a partire sempre da lultima (more arabū)," that is, to begin at the left as the Arabs do. If anyone claims that the method is difficult, says Pacioli, "Bonum est difficile: malum autem facile . . . Stultorum infinitus est numerus" (fol. 32, *v.*). "Del primo modo de partire detto per colona, ouer di testa, ouer per discorso, ouer per toletta . . . aregolo, ouer alla dritta, ouer tauoletta." Tartaglia, *General Trattato*, 1556, I, fol. 29; 1592 edition of *Arithmetica*, fol. 43.

meaning that $1{,}037{,}382 \div 14 = 74{,}098\frac{10}{14}$.[1] As with us, the method was generally used only with a divisor of one figure[2] and until recently has not been very popular with teachers,[3] requiring as it did some attention to a division table.[4]

Gerbert's Method. Of the methods which make use of our common numerals in long division, one of the oldest is often attributed to Gerbert (*c.* 980), although it is uncertain whether he originated it and although he did not use the zero.[5] It may be illustrated by the simple case of $900 \div 8$. The process consists of dividing 900 by $10 - 2$, 2 being the complement of the divisor, and was essentially as follows:

$$
\begin{array}{l}
10-2)\,900\,(90+18+3+1+\tfrac{1}{2}=112\tfrac{1}{2} \\
\quad \underline{900-180} \\
\qquad 180 \\
\qquad \underline{180-36} \\
\qquad\quad 36 \\
\qquad\quad \underline{30-6} \\
\qquad\qquad 6+6=12 \\
\qquad\qquad\quad \underline{10-2} \\
\qquad\qquad\quad 2+2=4,\ \tfrac{4}{8}=\tfrac{1}{2}
\end{array}
$$

The form actually used by certain of the successors of Gerbert may be seen from an example in an anonymous manuscript of the 12th century now in Paris,[6] no zero appearing in the computation. The combination of Roman and Hindu numerals is

[1] 1544/5 ed., fol. 15, under "Partire per testa."

[2] "Si chiama Partire à Colonna, quando il Partitore sarà d'vn Numero solo." Gio. Batt. di S. Francesco, 1689, p. 29.


[3] Pike's very widely used arithmetic employs long division in the cases of $175{,}817 \div 3$ and $293 \div 8$. See the 8th edition, New York, 1816, pp. 18, 60.

[4] Some 16th and 17th century writers in Italy gave a division table, and Onofrio (1670) speaks of his as "di grandissima vtilta." The Japanese learn a peculiar division table for their *soroban* and the Chinese for their *suan-pan*. See Smith-Mikami, p. 40.

[5] H. Weissenborn, *Zur Geschichte der Einführung der jetzigen Ziffern*, p. 14 (Berlin, 1892); *Gerbert*, p. 169 (Berlin, 1888).

[6] M. Chasles, *Comptes rendus*, XVI, 235, 243.

frequently seen in this period. The long explanation in the manuscript may be summarized in the following solution:



This same method is one of three given by Adelard of Bath (*Regulae abaci*, *c.* 1120), who attributes it to Gerbert. These three methods are the *divisio ferrea*, as above; the *divisio aurea*, somewhat like our long division; and the *divisio permixta*.[2]

Division by Factors. A third method of division that was common in the late Middle Ages consisted in using the factors of the divisor, and was known as "per repiego."[3] By this method 216 ÷ 24 reduces to 216 ÷ 8 ÷ 3, the object being to

[1] The fraction ½ was neglected. The bracketed matter is not in the original.

[2] Of the "iron division" he says: "℔De ferreis quidem diuisoribꝫ [for "divisionibus," as in two MSS.] hec paucis dicta sufficiant. Tamen quia super his tractauit gibertus philosophꝰ vir subtilis ingenij diligenter et compendiose quidam eciam quem discipulum eius predicant quē guichardum nominant/diligenter et prolixe." See Boncompagni's *Bullettino*, XIV, 67.

[3] *Repiego* means "refolding." It appears with various spellings, often *ripiego*. In Texeda's Spanish arithmetic (1546) it appears as *repriego*. It was occasionally called "division by rule," a name also given to short division. Thus in a 14th century MS. in Mr. Plimpton's library: "Questo e partire per regola = cioe. Parti 9859 ꝑ 48 cioe ꝑ .6. & ꝑ .8. sua reghola. . . . fattiamo fino alpartmeo ꝑ Reghola." See also the repiego method of multiplication, page 117.

secure one-figure divisors that could be handled "per tavoletta." The illustration given by Pacioli (1494) is that of $9876 \div 48$. He first divides 9876 by 6, the result being 1646. He then divides 1646 by "the other number of the repiego"[1] and obtains $205\frac{6}{8}$ or $205\frac{3}{4}$. It is still used, although not commonly taught in school.

Division by Parts. If the divisor was a multiple of ten, the 16th century writers frequently resorted to "Partire per il scapezo," that is, "division by cutting up" the dividend. Thus, to divide 84,789 by 20, the dividend was cut by a bar, 8478|9, the first part being divided by 2 and the 9 being divided by 20, —a plan that is found essentially in our modern books.[2]

The Galley Method. By far the most common plan in use before 1600 is known as the galley, *batello*, or scratch, method and seems to be of Hindu origin. It may be illustrated by the case of $65{,}284 \div 594$, as given in the Treviso arithmetic (1478). To make the work clear, the first six steps are given separately as follows:


(1)	(2)	(3)
65284 | 1	1	16
594	0 5 2 8 4 | 1	0 5 2 8 4 | 1
	5 9 4	5 9 4

(4)	(5)	(6)
5	5	5
1 0 8	1 0 8	1 0 8
0 5 2 8 4 | 1	0 5 2 8 4 | 10	0 5 2 8 4 | 10
5 9 4	5 9 4 4	5 9 4 4 4
	59	5 9 9
		5

[1] " . . . dico che parta p laltro numero del repiego: cioe. p .8. neuen .205. sani: e auāza .6." (1494 ed., fol. 33, *r.*).

[2] It is given in *Le Regoluzze di Maestro Paolo dell' Abbaco* (14th century), ed. Frizzo, p. 43 (Verona, 1883). The relation of this to the decimal fraction is discussed on page 238. The plan is given by many writers, including Borghi (1484), Sfortunati (1534), Cataneo (1546), Baker (1568), Digges (1572), and Pagani (1591).

The completed work, the explanation for which occupies two and one-half pages, is as follows:



That is, 65,284 ÷ 594 = 109, with a remainder 538.

The method is by no means as difficult as it seems at first sight, and in general it uses fewer figures than our common plan. Maximus Planudes (*c.* 1340) throws some light upon its early history, saying that it is "very difficult to perform on paper, with ink, but it naturally lends itself to the sand abacus. The necessity for erasing certain numbers and writing others in their places gives rise to much confusion where ink is used, but on the sand table it is easy to erase numbers with the fingers and to write others in their places."[1] It thus appears that this method, which at first seems cumbersome, is a natural development of a satisfactory method used on the sand abacus. It was adopted by Fibonacci (1202), as here shown for the case of 18,456 ÷ 17.[2]

	6
	149
	18456
	17
	1085
$\frac{1}{17}$	1085


The names *galea* and *batello* referred to a boat which the outline of the work was thought to resemble.[3] An interesting

[1] From the French translation in the *Journal Asiatique*, I (6), 240. On the Hindu method, see Gerhardt, *Études*, p. 7.

[2] The Boncompagni edition (I, 32) gives no cancellation marks, and very likely Fibonacci made no use of them.

[3] As Pacioli says: "E q̄sto vocabulo li aduene a tale o*p*are *p* certa similitudine materiale che li respōde del offitio e acto de la galea materiale q̄le e legno marittimo acto al nauigare" (1494 ed., fol. 34). Tartaglia remarks: "È detto in Vinetia per batello, ouer per galea per certe similitudini di figure" (1592 ed., fol. 48). The spelling varied, as usual, giving such forms as *battello*, *vatelo*, *galera*, and *galia*. There was occasionally a distinction between the *galea* and *batello* forms, as in Forestani, *Pratica d' Arithmetica*, Venice, 1603.

illustration of this resemblance is seen in a manuscript of *c.* 1575, as here shown. Tartaglia[1] tells us that it was the custom of Venetian teachers to require such illustrations from their pupils when they had finished the work.




GALLEY DIVISION, 16TH CENTURY

From an unpublished manuscript of a Venetian monk. The title of the work is "Opus Arithmeticā D. Honorati veneti monachj coenobij S. Laurētij." From Mr. Plimpton's library


This method of dividing was used by the Arab writers from the time of al-Khowârizmî (*c.* 825), of course with variations. For example, al-Nasavî (*c.* 1025), in finding that $2852 \div 12 = 237\frac{8}{12}$, used the form on page 139. The advancing

[1] 1592 ed., fol. 53.

of the divisor one place to the right each time is here seen more clearly than in the usual Italian forms. The medieval Latin writers sometimes called this feature *anteriioratio*.[1] This advancing of the divisor was not universal, however, Rudolff (1526) telling us that the French and other computers often set the divisor down but once.[2]



As already stated, the galley method was the favorite one with arithmeticians before 1600, and it had many strong advocates up to the close of the 18th century.[3] It is found occasionally without cancel marks, probably owing in most cases to the lack of the necessary canceled types.[4] With or without this canceling, the method was preferred not merely by commercial computers but also by such scientists as Regiomontanus.[5] Even as good a mathematician as Heilbronner, in the middle of the 18th century, preferred it in all long examples.[6] One reason for this preference was, no doubt, that fewer figures were used; but even more important was the fact that the work was more compact,—an important item before



FIRST EXAMPLE IN LONG DIVISION PRINTED IN THE NEW WORLD

From the *Sumario Compendioso* of Juan Diez, Mexico, 1556. It illustrates the galley method, without canceled figures, as applied to the case of 114,400 ÷ 26 = 4400

[1] So Sacrobosco (*c.* 1250) uses this word and also the verb *anteriorare*. From this, no doubt, Chuquet (1484) was led to use *anteriorer*. See G. Eneström, *Bibl. Math.*, XIII (2), 54; Halliwell, *Rara Math.*, p. 17.

[2] "Frantzosen vnd etlich ander Nacion/welche den teyler nit mehr dann ein mal setzen/. . ." (1534 ed., fol. 11).


[3] Among those who preferred it to any other are Chuquet (1484), Widman (1489), Riese (1522), Tonstall (1522), Köbel (1514), Gemma Frisius (1540), Recorde (*c.* 1542), Baker (1568), Oughtred (1631), and certain Dutch writers even as late as Bartjens (1792).

[4] *E.g.*, Pellos (1492), Grammateus (1518), Albert (1534), and the Mexican work of 1556 as shown in the facsimile.

[5] See his correspondence with Bianchini in the *Abhandlungen*, XII, 197.

[6] *Historia*, pp. 776 *et passim*.

the days of cheap paper. Hodder, late in the 17th century, says that he "will leave it to the censure of the most experienced to judge, whether this manner of dividing be not plain, lineal, and to be wrought with fewer Figures than any which is commonly taught,"[1] and in this he follows the testimony of many of the best Italian writers for two centuries preceding.[2] The method is still taught in the Moorish schools of North Africa, and doubtless in other parts of the Mohammedan world.



Our Long Division. It is impossible to fix an exact date for the origin of our present arrangement of figures in long division, partly because it developed gradually. We find in various Arab and Persian works arrangements substantially like the one shown above for the case of $1729 \div 12 = 144$, and 1 remainder.[3] This resembles our method, although it has several points in common with the galley plan.

```
25)625(25
   4
  22
  10
 125
 100
  25
  25
```

In the 14th century Maximus Planudes gave what is called an Arab device. This is a step in advance of the one given above and yet is quite distinct from our method.[4] It appears in a form somewhat like the one here shown for the case of $625 \div 25$.


[1] 1672 ed., p. 54.

[2] Thus Pagani (1591): "Il partire a Galera e molto sicuro & legiadro ch' ogn' altro partire," and Pacioli is even more pronounced in his opinion.

[3] This is a composite of solutions in various MSS. examined, including several of the 16th century. See also the work of al-Kashî (*c.* 1430) as referred to in Taylor, *Lilawati*, Introd., p. 22; Gerhardt, *Études*, p. 14.


[4] Gerhardt, *Études*, p. 22.

The 15th century saw the method brought into its present form under the name *a danda* ("by giving"). This name came from the fact that when a partial product is subtracted we bring down the next figure and "give" it to the remainder.[1] An excellent illustration from a manuscript of *c.* 1460 is here shown, but it will be noticed that the remainder is repeated each time before the "giving." The name *danda*, or *dande* in parts of Tuscany, is still used to designate this method of dividing.[2] It has, however, been applied to forms quite different from the one shown above. For example, the case of $49{,}289 \div 23 = 2143$ appears in the form shown below in a 14th century Italian manuscript,[3] and the author speaks of an analogous solution as *a danda*.[4] The earliest printed book to give the method is Calandri's work of 1491, and the first example of the kind is shown on page 142. It next appeared as the third method of Pacioli,[5] and was given with increasing frequency in the following century, but rather as an interesting than as a particularly valuable device.[6] With the opening of the 17th century it began more effectively to replace the galley



EARLY EXAMPLE OF LONG DIVISION

One of the earliest examples of the present method. From an Italian MS. of *c.* 1460



[1] So Cataneo (1546) says: "È chiamato a danda il detto modo, perche à ogni sottration fatta nel operare se li da vna o piu figure dal lato destro" (1567 ed., fol. 15).

[2] Boncompagni's *Bullettino*, XIII, 252 n.


[3] *Rara Arithmetica*, p 437.

[4] "Questo sie' ilpartire adanda."

[5] "De tertio modo diuidendi dicto danda" (1494 ed., fol. 33).

[6] Pagani (1591) speaks of it thus: "Partire à danda è assai bello, & vago." Cognet (1573) mentions the advantage of not canceling: "Les Marchands Italiens, pour ne trencher aucune figure, divisent en la sorte qui s'ensuit"; and Trenchant (1566) remarks: "Il y a vne autre belle forme de partir, sans trencher aucune figure," or "sans rien couper."


method. Cataldi (1602) gives it as his first method, but with the quotient below the dividend, the first part of his work



FIRST PRINTED EXAMPLE OF MODERN LONG DIVISION
From Calandri's arithmetic, Florence, 1491. The problem is the division of 53,497 by 83


being as shown on page 143. In another example he places the quotient at the right, saying that this is the custom in

Milan.[1] In the galley method the most convenient place for the quotient was at the right; Cataldi's attempt at placing it below was awkward; the modern custom of placing it above the dividend in long division is the best of all, since it automatically locates the decimal point.



At the close of the 17th century the modern form of division was fairly well established, the galley method being looked upon more as a curiosity.[2]

There have been many variants of the *a danda* method, but the only one of any importance is that which omits the partial products as shown below. Cataldi (1602) calls it the abbreviated *a danda*.[3] It has had more or less vogue for three centuries, but it requires too much mental effort to become common. It was brought to the attention of American teachers by Greenwood (1729), who, speaking of the various methods, remarked that, as "most of the rest are at best an unnecessary Curiosity; I shall confine myself wholly to the Two ITALIAN Methods; which are the most usual," these two being *a danda* and the contracted form.



Of the various methods suggested in the 16th century one of the most interesting is that of Apianus (1527), particularly as it suggested the scheme of decimal fractions. To divide 11,664 by 48, Apianus first writes the aliquot parts of 48, with a corresponding series of numbers based on 48 as a unit, substantially as follows:


		48	corresponds to	1
$\frac{1}{2}$	of	$48 = 24$,		05
$\frac{1}{4}$	of	$48 = 12$,		025
$\frac{1}{8}$	of	$48 = 6$,		0125
$\frac{1}{16}$	of	$48 = 3$,		00625

[1] "Partire à Danda vsato in Milano."

[2] Thus Onofrio's *Aritmetica* (1670) gives it as "di poco ò nullo profitto."

[3] " . . . à Danda abbreuiato" (p. 88).

He then observes that $11 \div 48 \not> 1$, $11 \div 24 \not> 1$, $11 \div 12 \not> 1$, but $11 \div 6 > 1$. But $6 = \frac{1}{8}$ of 48, and hence the first part of the quotient is 0125. The rest of the work is substantially as follows:



$$11 - 6 = 5$$
$$5 \div 48 \not> 1$$
$$5 \div 24 \not> 1$$
$$\vdots$$
$$5 \div 3 > 1$$

hence we write $\frac{1}{16}$, or 00625, and so on. It is evident that Apianus had some idea of decimal fractions in his mind, although it was not developed in his treatise.

Clichtoveus (1503) gave a rule based upon the identity

$$\frac{10\left(a - \frac{c}{2}\right) + b}{c} + 5 = \frac{10a + b}{c}.$$

Thus, to find $29 \div 4$, take $a - \frac{c}{2} = 2 - \frac{4}{2} = 0$; then subtract 4 (or c) as often as possible from 9 (or b), thus finding that $9 \div 4 = 2\frac{1}{4}$. The final quotient[1] is then $2\frac{1}{4} + 5$, or 7^1.

Whatever method of dividing was used, a table of multiples of the divisor was early recognized as desirable. Such tables are found in many works, including those of Recorde (*c.* 1542), Fine (1530), Ramus (1569), Hylles (1592), and Greenwood (1729).

7. Roots

Finding Square and Cube Roots of Numbers. The Greeks found the square root of a number by a method similar to the one commonly set forth in the elementary algebras and arithmetics of the present time. It was shown geometrically by Euclid[2] that $(a + b)^2 = a^2 + 2ab + b^2$ (a fact that was prob-

[1] Edition of *c.* 1507, fol. D 4. He also gives a rule for the case of $a < \frac{1}{2}c$. For a few further notes on the history of division see E. Mathieu, "Méthodes de division en usage à la fin du siècle dernier," in *Journal de math. élément.*, V (4), 97.

[2] *Elements*, II, 4.

ably known long before his time), and by means of this relation Theon of Alexandria (*c.* 390), using sexagesimals, found the square root of a number by the following rule:

When we seek a square root, we take first the root of the nearest square number. We then double this and divide with it the remainder reduced to minutes, and subtract the square of the quotient; then we reduce the remainder to seconds and divide by twice the degrees and minutes [of the whole quotient]. We thus obtain nearly the root of the quadratic.[1]


By this rule he finds that $\sqrt{4500} = 67\ 4'\ 55''$ approximately.

From Greece the method passed over to the Arabs and Hindus, with no particular improvement. Thus Bhāskara (*c.* 1150) writes his number as follows:

	–		–	
8	8	2	0	9

and then proceeds much as Theon had done. He says:

Having deducted from the last of the odd digits[2] the square number, double its root; and by that dividing the subsequent even digit[3] and subtracting the square of the quotient from the uneven place,[4] note in a line the double of the quotient.[5]



One of the most interesting medieval examples of the finding of a square root is given by Maximus Planudes (*c.* 1340). To find $\sqrt{235}$ he arranges the work as here shown. This is quite

[1] For further details of the process see J. Gow, *History of Greek Mathematics*, pp 54–57 (Cambridge, 1884) (hereafter referred to as Gow, *Greek Math.*); K. Hunrath, *Ueber das Ausziehen der Quadratwurzel bei Griechen und Indern*, Prog., Hadersleben, 1883.

[2] That is, from 8, the third and last of the odd places denoted by a vertical line, counting from the right.

[3] Really, $(882 - 400) \div 40 = 9+$.

[4] Apparently meaning that 29^2 is subtracted from 882.

[5] For the rest of the rule see Colebrooke's *Lilāvati*, p. 9. For the work of the Arabs and Persians, see the Taylor translation, p. 23.

unintelligible without the accompanying explanation, which may be condensed as follows:[1]


$$\begin{array}{lll} & & 235(15 \\ & & \underline{1} \\ & 2 & 13 \\ 2\times 5= & \underline{10} & \\ & 30 & =\text{twice the root.} \\ \text{Hence} & 15 & =\text{the root.} \\ \text{But} & 15^2 & =225. \end{array}$$

Hence $\frac{235-225}{30}=\frac{1}{3}$, which must be added.

Hence the root is $15\frac{1}{3}$.

The early printed arithmetics generally used an arrangement of figures similar to the one found in the galley method of division. Thus Pacioli (1494) gives the following:[2]

Extractio radicū



that is, $\sqrt{99,980,001}=9999$.

Gradually, in the 16th century, the galley method gave way to our modern arrangement, although it was occasionally used until the 18th century.[3] Among the early writers to take an

[1] For examples of a more elaborate nature see Wäschke, *Planudes*.

[2] Fol. 45, *r*. For those who do not have access to original works a good illustration of this method may be seen in the *Abhandlungen*, XII, 201, 269. A problem of Chuquet's (1484) may also be seen in Boncompagni's *Bullettino*, XIII, 695.

[3] Among the better arithmeticians that used it in the 17th century was Wilkens, a Dutch writer (1630).

important step toward our present method was Cataneo (1546), who arranged the work substantially as follows:[1]

54756(234		
4	primo duplata	4
14	secondo	46
12		
27		
9		
185		
184		
16		
16		
0		

Among the first of the well-known writers to use our method in its entirety was Cataldi, in his *Trattato* of 1613.[2] Most early writers gave directions for "pointing off" in periods of two figures each, some placing dots above, as in $8\dot{2}4464^{8}$; some placing dots below, as in $11\dot{9}0\dot{2}5^{4}$ or as in $21\dot{1}7\dot{8}4\dot{0}4^{5}$; some using lines, as in $26\,000\,000\,000^{6}$; some using colons, as in $13:01:76:64:^{7}$; and some using vertical bars, as in $94|21|80|73|55$.[8] Many writers, however, did not separate the figures into groups.[9]

[1] *Le Pratiche*, Venice, 1567 ed., fol. 72.

[2] For various forms used by other writers see P. Treutlein, *Abhandlungen*, I, 64, 71.

[3] *E.g.*, Grammateus (1518), Scheubel (1545), Hartwell (1646 edition of Recorde's *Ground of Artes*), Wilkens (1630), and the American Greenwood (1729).

[4] *E.g.*, Gemma Frisius (1540), L. Schoner (1586), Peletier (1549), Santa-Cruz (1594), and Metius (1625). Cardan sometimes placed them above and sometimes below.

[5] This from the *Epitome* of Clavius (1583; 1585 ed., p. 309), although generally (as on page 310) he places the dots immediately below the figures.

[6] This in cube root, from the Rollandus MS. (1424).

[7] From Ortega (1512; 1515 ed., fol. 99). He also writes 3 : 6 : 0 : 8 : for the square root.

[8] This was very common and has much to commend it. It was given by Chuquet (1484), Pellos (1492), Finc (1530), Trenchant (1566), and many others.

[9] *E.g.*, the Arab al-Ḥaṣṣâr (*c.* 1175), Cataneo (see the example above), and Feliciano da Lazesio (1526).

In finding the square root and the cube root most of the early writers[1] gave the rules without any explanation, or at the most with merely a reference to the fact that $(a+b)^2 = a^2 + 2ab + b^2$. Thus Buteo (1559) proceeded no farther with cube root than to find the first figure, saying that it is better to use a table of cubes;[2] and more than a century later de Lagny[3] asserted that it would take most computers more than a month to find the cube root of 696,536,483,318,640,035,073,641,037. Although the ponderous work of Tonstall (1522) naturally included roots, Recorde (*c.* 1542) did not think the subject worthy of a place in his *Ground of Artes*.[4]

A conviction of the value of the reasoning involved in the subject led various writers in the 16th century to give clear explanations based on the geometric diagram.[5] The use of the blocks for explaining cube root was found somewhat later, and became fairly common in the 17th century.[6] In the 17th and 18th centuries the blocks are even used in finding the fourth root, x cubes being taken, each composed of x^3 cubes.[7]

[1] *E.g.*, Brahmagupta (*c.* 628) and Bhāskara (*c.* 1150), ed. Colebrooke, pp. 10, 279; al-Karkhī (*c.* 1020), ed. Hochheim, II, 13. Fibonacci (1202) described cube root, and it also appears in Sacrobosco's *Algorismus* (*c.* 1250) and in the *Carmen de Algorismo* of Alexandre de Villedieu (*c.* 1240).

[2] He gives such a table up to 40^3, and a rule which we may express by the approximation formula

$$\sqrt[3]{a^3 + r} = a + \frac{r}{3a^2 + 3a}.$$

[3] *Nouveaux Élémens d'Arithmétique et d'Algèbre*, Paris, 1697; A. De Morgan, *Arithmetical Books*, p. 55 (London, 1847) (hereafter referred to as De Morgan, *Arith. Books*).

[4] In Hartwell's edition, however, there is "An Appendix concerning the Resolution of the Square and Cube in Numbers, to the finding of their side," in which he speaks of the "Quadrat root, or the side of any Quadrat number," and gives the geometric diagram (1646 ed., p. 573).

[5] *E.g.*, Tonstall (1522, fol. TV_3), Trenchant (1566), L. Schoner (1586, p. 255), and Gemma Frisius (1540).

[6] A good illustration is found in Hartwell's edition of Recorde's *Ground of Artes* (1646 ed., p. 587).

[7] *E.g.*, Cardinael (1644; 1659 ed., fol. E 8). When Bartjens (1633; 1752 ed., pp. 242, 243) wishes to know "Hoe veel is de $\surd xx$ van 576" or "de Radix xx uit 3136 is," such being his two symbols for square root, he uses the diagram. He then uses the blocks in "kubicq-wortel" when he "trekt de $\surd x^3$ uit 5832" (p. 251), and in fourth root when he "trekt de $\surd x^4$ van 81450625." See also the 1676 edition, p. 242.

Higher Roots. The conviction of the value of the subject as a mental exercise led various writers to include some work in higher roots, the work being based upon a knowledge of the binomial coefficients. These coefficients were occasionally arranged in the triangular form subsequently known as Pascal's Triangle.[1] This arrangement was known to the Chinese[2] as early as 1303, and also to the Arabs,[3] and in Europe it appeared in print on the title-page of a work by Apianus published in 1527 and in a work[4] by Scheubel that appeared in 1545.

This arrangement of the binomial coefficients was first seriously considered in a printed book, in connection with higher roots, simultaneously by Stifel (1544)[5] and Scheubel (1545).[6] The latter finds the tenth root of 1,152,921,504,606,846,976, for example, to be 64, and he carries the work as far as to the finding of a 24th root. A little later it was used in France by such writers as Trenchant (1566)[7] and Peletier (1549),[8] and it appeared also in the works of various Dutch writers.[9]

[1] *Traité du triangle arithmétique,* published posthumously in 1665. The form used by Pascal is given later (p. 510).

[2] It appears in the *Szu-yuen Yü-kien* of Chu Shï-kié (1303), but as something already known. See Mikami, *China*, p. 90.

[3] Cantor, *Geschichte*, I (2), 645. [4] *Rara Arithmetica*, pp. 156, 236.

[5] In the *Arithmetica Integra*, fol. 44. As to their use in his *Coss* (1554), see *Abhandlungen*, I, 77; II, 43.

[6] *De Numeris*, in the *tractatus quintus*. See *Rara Arithmetica*, p. 236.

[7] "Doctrine generale pour extrére toutes racines." He also says: "Pour fondement de la quelle, i'ay formé ce trigône semé de nombres, s'imbolisans & s'engendrans les vns les autres par vn ordre de grandissime consideration" (1578 ed., p. 249). It will be observed that, by placing 1 at each end of each row, the successive rows give the coefficients in the expansion of $(a + b)^n$ for n equal to 2, 3, 4, ... This serves as a basis for the general rule for finding the nth root of any number. For example, to find the fifth root we observe that the arithmetic triangle gives the trial divisor as $5a^4$ and the complete divisor as $5a^4 + 10a^3b + 10a^2b^2 + 5ab^3 + b^4$, a principle well known to writers of the 16th century.

2
3 · 3
4 · 6 · 4
5 · 10 · 10 · 5
etc

[8] He speaks of it as a "Nouuelle maniere d'extraire les Racines, generale pour toutes extractions, jusques à infinité" (1607 ed., pp. 107, 178, 252).

[9] Thus Van der Schuere (1600) speaks of the "Drie-hoecks wijze" (triangle-like) arrangement. It is also used by Bartjens (1633), Cardinael (1644), and others.

Abbreviated Methods. Attempts at abbreviating the process are relatively late. One of the most popular rules for the abridgment of square root is attributed to Newton, and Greenwood (1729) gives it as follows:

SIR Isaac Newton takes notice of a very useful *Contraction*, in these Cases, viz. *That when a Root is carried on half way or above, the Number of Figures you intend it shall consist of; the remaining Figures may be obtained by Dividing the remainder by the double of the Radical Figures.*[1]

The Meaning of the Term. It should be stated in this connection that the use of "root" to mean the square root, common in Europe today, has historic sanction. Indeed, all the world still recognizes it by taking the symbol $\sqrt{a}$ instead of $\sqrt[2]{a}$ to indicate the positive square root of a. The usage, however, was not entirely general, many early writers specifying the square root as carefully as the cube root.[2]

The Arab writers conceived a square number to grow out of a root, while the Latin writers thought of the side of a geometric square. Hence the works translated from the Arabic have *radix* for a common term, while those inherited from the Roman civilization have *latus*.[3] Hence the Latin writers "found" the *latus* and the Arab writers "extracted," or pulled out, the root. Our arithmetics, based largely upon Arab sources, still use "extract," although the older usage of "find" is better. The fact that from *radix* we have both "radical" and

[1] P. 77. See Newton's *Arithmetica Universalis*, p. 33 (Cambridge, 1707): "Ubi vero radix ad medietatem aut ultra extracta est, cæteræ figuræ per divisionem solam obtineri possunt."

[2] Thus Suevus (1593), under his "Regvla qvadrata," gives "Extractio Radicis Quadratae," and Digges (1572) speaks of "the square *Radix*," "quadrat roote," and "quadrate root."

Among the early writers who used "root" for "square root" were al-Nasavî (*c.* 1025), L. Schoner (1586), Rollandus (1424), and probably Bhāskara (*c.* 1150; Taylor, *Lilawati*, introduction, p. 6).

[3] Schoner speaks of this in *De numeris figuratis liber*, appended to his 1586 edition of Ramus: "Sic 9 est aequilaterus, & latus ejus est 3. Hoc latus aequilateri ab Arabibus etiam dicitur Radix" (p. 3). Fibonacci (*Liber Abaci*, p. 353) uses "find" instead of "extract" with the word "root," having used "extract" for "subtract."

"radish" makes the use of "extract" more easily understood.[1] This use is found in various modern languages,[2] but is by no means universal. Thus Digges (1572) says, "To find the square *Radix*, or Roote of any number" (p. 13), although he also says, "to search or pull out the *Radix*, or roote cubical" (p. 16).[3]

8. Checks on Operations

Need for Checks. The fact that the intermediate steps in a long operation were erased on the various forms of the abacus rendered it impossible to review the work as may be done with our present methods. It was therefore necessary that some simple check should be used to determine the probable accuracy of a result. The inverse operation was generally too long to serve the purposes, and hence other methods were developed rather early.

Check of Nines. Of all these methods the check of nines is probably the best known. It is simple of application and serves to detect most of the errors that are likely to occur. The origin of the method is obscure. It is found in the works of various Arab writers, including al-Khowârizmî (*c.* 825), al-Karkhî (*c.* 1020), Behâ Eddîn (*c.* 1600), and others. Avicenna (*c.* 1020), however, in discussing the subject of roots, speaks of it as a Hindu method.[4] On the contrary, no Hindu writer is

[1] On the use of "root" see Wertheim's edition of Elia Misrachi (*c.* 1500), *Sefer-Ilamispar*, p. 20 (Frankfort a. M., 1893), and Tartaglia's *General Trattato*, II, fol. 53, *v.* (1556).

[2] *E. g.*, "Uyttreckinge der wortelen" (Cardinael, 1659 ed., p. 2), and "cavere la radice qvadra" (Ciacchi, 1675 ed., p. 335).

[3] Of other forms of expression the following are types: "7097, cuius tetragonicū latus inquirens . . .," Buteo (1559; 1560 ed., p. 71); " . . . sacar rayz quadrada . . .," Iuan Perez de Moya (1562; 1615 ed., fol. 223), *sacar* meaning to extract; "Del trare la radice de numeri quadrati" (fol. 15), but "Del trouare la radice Cubica" (fol. 18, *v.*), in the Italian translation of Fine (Venice, 1587), showing both "extract" and "find"; "Del modo di trar la radice quadra . . ." (fol. 182, *v.*), but "La estrattione delle radici cube" (fol. 187, *v.*), Forestani, *Pratica d' Arithmetica* . . ., Venice, 1603.

[4] "Fâ' l-tharîk al-hindacî," an expression that has been variously interpreted. See F. Woepcke, *Journal Asiatique*, I (6), 500; Carra de Vaux, "Sur l'histoire de l'arithmétique arabe," *Bibl. Math.*, XIII (2), 33.

known to have used it before the 12th century,[1] while the Arabs certainly used it early in the 9th century. Nevertheless, as careful a writer as Paul Tannery is convinced that the evidence at present points to its invention in India but to its first considerable use in the School of Bagdad.[2]

There is some interesting evidence of the recognition of the excess of nines in the number mysticism of one of the late Greco-Roman writers, Hippolytus, who seems to have lived in the 3d century and who wrote several theological treatises as well as a *canon paschalis*. He made no use of the principle, however, in the verification of computations, and so far as we know he was ignorant of this application of the theory.[3] What he did was to make use of gematria, as in estimating the relative ability of individuals by means of the numerical values of the letters of their names. Instead, however, of simply stating this value in the usual way, he stated it with respect to the modulus nine. For example, the numerical value of Hector (Ἕκτωρ) is 1225, but Hippolytus gave it as 1, which is the excess of nines in this number. He spoke of this plan as due to the Pythagoreans, meaning, no doubt, the Neo-Pythagoreans of a period much later than that of Pythagoras himself.

The check of nines seems to have come into general use in the 11th century, largely due to the influence of Avicenna (*c.* 1020) and his contemporary, al-Karkhî, and thereafter it is found in most of the other arithmetics of any importance for a period of about eight hundred years. Albanna (*c.* 1300) speaks of the Arab arithmeticians as giving proofs of their computations by the checks of 7, 8, 9, and 11, and as knowing of the checks by other numbers as well.

From the Arabs this method of checking passed over to the West, appearing in the works of the Hebrew-Arabic writer Kûshyâr ibn Lebbân (*c.* 1000), the Hebrew Rabbi ben Ezra

[1] G. R. Kaye, *Indian Mathematics*, p. 34 (Calcutta, 1915), hereafter referred to as Kaye, *Indian Math.*; Taylor's *Lilawati*, p. 7.

[2] P. Tannery, *Mémoires Scientifiques*, I, 185 (Paris, 1912). On the Arab writers see Boncompagni, *Trattati*, I, 12; *Bibl. Math.*, II (3), 17; XIII (2), 33; Hochheim, *Kâfî fîl Hisâb*, II, 10 n.

[3] P. Tannery, *Mémoires Scientifiques*, I, 185; Tropfke, I (2), 58.

(*c.* 1140),[1] the Hebrew-Christian Johannes Hispalensis (*c.* 1140), and the Christian writers Fibonacci (1202), Maximus Planudes (*c.* 1340), and their successors.

Fibonacci called the excess of nines the *pensa* or *portio*[2] of the number, and used it as a check in multiplication and division. Maximus Planudes arranged his work in multiplication as here shown, using 9 instead of 0 in the case of a zero excess, and apparently believing that the check was a complete one. Johannes Hispalensis and Fibonacci, however, recognized its limitations.

294849	9
543	3
543	3

In the early printed arithmetics the check is found quite generally. Pacioli (1494) speaks of it as "corrente mercatoria e presta,"[3] and Widman (1489) always concludes his operations by the query, "Wiltu probirn?" Scheubel (1545) considered the matter so important that he gave a table of multiples of nine for the convenience of computers.[4]

The failure of the check was considered at some length by Pacioli, but Clavius[5] was especially clear in his treatment of the case. So important was the whole matter considered that Santa-Cruz (1594) devoted twenty-two pages to the theory.[6] In the 17th century, owing to the general acceptance of the modern forms of computing, the revision of the operations became more simple, and hence some of the leading commercial arithmetics[7] discarded the check of nines. In England, however, the influence of Cocker[8] served to make it very popular, and such influence as Greenwood (1729) had in America was in the same direction. In the 19th century it dropped out of American arithmetics for the most part, but after 1900 it began to appear again.

So important did Tartaglia (1556) consider the check of nines, even in addition, that he gave a table of the excess of

[1] Silberberg, *Sefer ha-Mispar*, p. 94.

[2] *Liber Abaci*, I, 8.

[3] Fol. 20 [numbered 10], *r*.

[4] *De Nvmeris*, I, chap. 2, p. 12 (1545). He did the same for 7, 11, 13, and 19, using these numbers also for checking.

[5] *Epitome*, p. 22 (1583).

[6] 1643 ed., fol. 171. See also Sfortunati (1534; 1545 ed., fol. 8); Cataneo (1546; 1567 ed., fol. 18); and Pagani, p. 6 (1591).

[7] *E.g.*, Eversdyck's edition of Coutereels, p. 33 (1658); Mots (1640).

[8] *Arithmetick*, London, 1677, with later editions.

nines for each number from 0 to 90,[1] a waste of space that argues for the lack of appreciation of the ease with which one casts out the nines in any number, however large.

Checks with Other Numbers. Any other number besides nine may be used for checking, although nine is the most convenient. The use of other numbers is found in the works of various Arab writers, and Fibonacci[2] gives the checks for 7, 9, 11. Other medieval and Renaissance writers[3] also give such numbers as 2, 3, 5, 6, 13, and 19. Several of the early printed books show a preference for 7 on account of the diminished chance of error.[4] In general, however, they naturally give the proof by nines the preference.[5]

Inverse Operation. Although the check by the inverse operation took more time, it was more certain, and hence it found many advocates. It is so simple that its origin is probably remote, although it is not until the Middle Ages that we find it first stated definitely.[6] It appears frequently in the early printed books,—for example, in the works of Clichtoveus (1503), Albert (1534), and Thierfelder (1587). Tartaglia (1556) asserted that the method was illogical, since subtraction could not be used in checking addition, for the reason that it was taught after that subject,[7]—an objection that is of no practical significance.

[1] *General Trattato*, I, fols. 8, *v.*, and 9, *r.*

[2] *Liber Abaci*, pp. 8, 39, 45.

[3] B. Boncompagni, *Atti Pontif.*, XVI, 519. Rudolff (1526), Apianus (1527), Fischer (1549), Albert (1534), and Scheubel (1545) are particularly worth consulting.

[4] Thus Pellos (1492), comparing 7 with 2, says: "℄Item sapias che ꝑba de .7. es la plu segura ꝑba che pusca esser car la ꝑba de .2." (fol. 18). See also Borghi (1484).

[5] Thus Clavius (1583) prefers the proof "per abiectionem nouenarij" or, in the Italian edition, "col gettar via tutti li 9," to that "per abiectionem septenarij" or "col gettar via li 7"; and so with Chuquet, Pacioli, Buteo, Tartaglia, Cardan, and many others. The proof by other numbers than 9 and 11 is not often found after about 1600.

[6] For example, in the *Algorismus prosaycus magistri Christani* (*c.* 1400): "Et nota, quod subtraccio probat addicionem et addicio subtraccionem." Studnička ed., p. 9 (Prag, 1893).

[7] See the *General Trattato*, I, 8, *r.*

TOPICS FOR DISCUSSION

1. The number and the nature of the fundamental operations, and the reasons for the various classifications.

2. Significance of duplation and mediation in the development of logistic, particularly in early times.

3. Difficulties in adequately defining the fundamental operations as their nature expanded from time to time.

4. The leading principles determining systems of notation, with illustrations of each principle.

5. The leading systems of notation, with a study of their respective merits.

6. The significance and growth of the concept of place value in the writing of numbers.

7. The history of the Roman numerals, with a study of the variants from century to century.

8. The nature, history, and significance of the subtractive principle in the writing of numbers.

9. The history of our common numerals, with a study of the variants from century to century.

10. The reading and writing of large numbers at various periods and in various systems.

11. The terminology used from time to time in connection with the common operations.

12. Significant features of the work in addition and subtraction at different stages of the development of these operations and a study of the relative merits of the various methods.

13. A study of the different methods of multiplying, with a consideration of the relative merits of each and of the probable reason for the survival of the present common method.

14. A study of the different methods of division, with particular reference to the contest between our present plan (a modification of the *a danda* arrangement) and the galley method.

15. Traces of early methods of computations in our present operations with algebraic polynomials.

16. The historical development of the process of finding roots of numbers.

17. The historical development of the various methods of checking operations with integers.

CHAPTER III

MECHANICAL AIDS TO CALCULATION

1. The Abacus

Necessity for the Abacus. Since the numerals of the ancients were rather unsuited to the purposes of calculation, it is probable that some form of mechanical computation was everywhere necessary before the perfecting of the modern system. This probability becomes the stronger when we consider that all convenient writing materials were late developments in the history of civilization. Papyrus was unknown in Greece before the 7th century B.C., parchment was an invention of the 5th century B.C.,[1] and paper is a comparatively recent product,[2] while tablets of clay or wax were not suitable for calculation.

Meaning of the Term. In earliest times the word "abacus"[3] seems to have referred to a table covered with sand or with fine

[1] Pliny says, of the 2d century.

[2] It may have been brought into Europe in the 12th century by the Moors of Spain, but specimens dating from about the beginning of our era have been found on the eastern borders of China.

[3] The word comes from the Greek ἄβαξ (*a'bax*), probably from the Semitic אבק (abq), dust. Numerous other etymologies have been suggested. Among the most interesting is one given by Th. Martin (*Les Signes Num.*, p. 34) on the authority of Orion of Thebes, a lexicographer of the 5th century, and on that of several other scholars,—namely, that the word comes from α + βάσις (*a* + *ba'sis*, without base), referring to the fact that the computing tablet had no feet. A recent article by R. Soreau gives the improbable suggestion that ἄβαξ simply meant a numerical table, and came from α′, β′, + ἄξια (*a*, *b*, *ax'ia*, relating to value), meaning 1, 2, + αξ (indicating numerical values). See R. Soreau, "Sur l'origine et le sens du mot 'abaque,'" *Comptes rendus*, CLXVI, 67. The question was debated even in Pacioli's time, for he says (*Sūma*, fol. 19, *r.* (1494)): "e modo arabico e chiamase Abaco: ouer secōdo altri e dicta Abaco dal greco vocabulo." Of the various guesses, that of Joannes de Muris (*c.* 1350) is the most curious, that "abacus" is the name of the inventor: "Non est sub silencio transeundum de tabula numerorum, quam abacus adinuenit" (*Quadripartitum*, chap. xiv; in the *Abhandlungen*, V, 144).

dust, the figures being drawn with a stylus and the marks being erased with the finger when necessary. This at any rate is the testimony of etymology, and the dust tablet seems to have been the earliest form of the instrument.[1]

While all definite knowledge of the origin of the abacus is lost, there is some reason for attributing it to Semitic rather than to Aryan sources.[2]

The dust abacus finally gave place to a ruled table upon which small disks or counters were arranged on lines to indicate numbers. This form was in common use in Europe until the opening of the 17th century, and persisted in various localities until a much later date.

Meanwhile, and in rather remote times, a third form of abacus appeared in certain parts of the world. Instead of lines on which loose counters were placed there were grooves or rods for movable balls or disks, a form still found in Russia, China, Japan, and parts of Arabia.

We have, then, three standard types,—the ancient dust board, which probably gave the name to the abacus, the table with loose counters, and the table with counters fastened to the lines. These three, with their characteristic variants, will now be explained.[3]

The Dust Abacus. The dust abacus was merely a kind of writing medium of little greater significance in computation

[1] C. G. Knott, "The abacus in its historic and scientific aspects," *Transactions of the Asiatic Society of Japan,* Yokohama, XIV, 18; hereafter referred to as Knott, *Abacus.*

[2] Knott, *Abacus,* pp. 33, 44.

[3] The literature of the subject is extensive. The following are some of the general authorities consulted: Knott, *Abacus*; M. Chasles, *Comptes rendus,* XVI, 1409; F. Woepcke, *Journal Asiatique,* I (6), 516; Sir E. Clive Bayley, *Journal of the Royal Asiatic Society,* XV (N.S.); M. Hübner, "Die charakteristischen Formen des Rechenbretts," *Zeitschrift für Lehrmittelwesen und pädagogische Literatur,* II, 47; D. Martines, *Origine e progressi dell' aritmetica,* p. 19 (Messina, 1865) (hereafter referred to as Martines, *Origine aritmet.*); A. Terrien de Lacouperie, "The Old Numerals, the Counting Rods and the Swan-pan in China," *Numismatic Chronicle,* III (3), 297–340, reprinted in London in 1888 (hereafter referred to as Lacouperie, *The Old Numerals*). The most elaborate and scholarly work on the subject is F. P. Barnard, *The Casting-Counter and the Counting-Board,* Oxford, 1916 (hereafter referred to as Barnard, *Counters*).

than the clay tablet of the Babylonians, the wax tablet of the Romans, the slate of the Renaissance period, or the sheet of paper of today. In its use, however, is to be found the explanation of certain steps in the operations with numbers, and on this account it deserves mention.

The Hindus seem to have known this type in remote times but to have generally discontinued its use. Even in recent times, however, children have been instructed to write letters and figures in the dust or sand on the floor of the native school before being allowed to use the common materials for writing.[1] That the dust abacus was common a century ago is asserted by Taylor in the preface to his edition of the *Lilāvati*.

In the Greek and Roman civilizations the dust abacus was also well known. Figures were drawn upon it with a stylus, called by the Latin writers a *radius*,[2] much as they were drawn on the slate in recent times. The wax tablet, described later, was even more extensively used.


Nature of the Counter Abacus. As in the case of all such primitive instruments, the origin of the counter abacus is obscure.[3] We only know that in very early times there seems to have been a widespread knowledge of some kind of instrument in which objects (beads, disks, or counters) on one line indicated units, on the next line tens, on the next hundreds, and so on. Some general idea of this instrument may be obtained from the illustrations given on page 159. The first one shows the successive steps taken in the addition of numbers. The second illustration shows the use of the abacus in multiplication. Several variants of this type are given later.

[1] Sir E. Clive Bayley, *Journal of the Royal Asiatic Society*, XV (N.S.), 9 n., 15, and XIV (N.S.), Part 3 (in the reprint of the article "On the genealogy of ancient numerals" it appears in Part II, p. 71; see also Part I, p. 19, and Part II, pp. 50, 54); G. R. Kaye, "The use of the abacus in Ancient India," *Journ. and Proc. of Asiatic Soc. of Bengal*, IV (2), 293.

[2] "Ex eadem urbe humilem homunculum a pulvere et radio excitabo, qui multis annis post fuit Archimedes," Cicero, *Tusculan Disputations*, V, 23, 64; "Descripsit radio," Vergil, *Eclogues*, III, 41.

[3] On the history in general see A. Nagl, *Die Rechenpfennige und die operative Arithmetik*, Vienna, 1888.


There is some reason for believing that this form of the abacus originated in India, Mesopotamia, or Egypt. The whole



ADDITION ON THE ABACUS

An early computer, wishing to add 22 and 139, might have proceeded as follows: Place 2 pebbles on the units' line, as shown in the First Step. Then place 9 more, as shown in the Second Step. Then take away 10 of these pebbles and add one pebble to the tens' line, as shown in the Third Step. Then add 2 pebbles to the tens' line because of the 20 in 22, as shown in the Fourth Step. Then add 3 more because of the 30 in 139, as shown in the Fifth Step. Finally draw a line for hundreds, and on this place one pebble because of the 100 in 139. The answer is 161

matter is, however, purely speculative at the present time and it seems improbable that it will ever be definitely settled.



MULTIPLICATION ON THE ABACUS

Above the horizontal line in the middle it is easily seen that the number 4132 is represented. If we wish to multiply this by 2, we may simply double the objects (in this case the black dots) below the line, and the result is evidently 8264

The Abacus in Egypt. That the Egyptians used an abacus is known on the testimony of Herodotus, who says that they "write their characters and reckon with pebbles, bringing the hand from right to left, while the Greeks go from left to right." This right-to-left order was that of the Hieratic script, the writing of the priestly caste, and in this respect there is probably some relation between this script and the abacus.[1] No wall pictures thus far discovered give any evidence of the use of the abacus, but in any collection of Egyptian antiquities there may be found disks of various sizes which may have been used as counters.[2]

The Abacus in Babylonia. We have as yet no direct evidence of a Babylonian abacus. The probabilities are, however, that the Babylonians, like their neighbors, made use of it. Methods of computing were never chiefly confined to the learned class whose written records have survived. It was the trader first of all who used the abacus, and it was he who carried the customs and manners from country to country. Tradition not infrequently assigns the origin of the abacus to the Middle East, as in the writings of Iamblichus (*c.* 325), who not only states that Pythagoras introduced the instrument into Greece, but hints that he may have brought knowledge of this kind from Babylon.[3] The tradition that the primitive home of the abacus was in or near Babylon is also recorded by Radulph of Laon (*c.* 1125)[4] and other writers who had no special knowledge of the subject.

[1] On the Egyptian abacus see M. Cantor, *Geschichte*, I, chap. i; J. P. Mahaffy, *Old Greek Education*, p. 56 (New York, 1882), derives the Greek abacus from Egypt.

[2] In a papyrus of the time of Menephtah I (1341–1321 B.C., Lepsius) is a drawing which looks at first sight like an abacus (Cantor, *Geschichte*, I (1), 51), but which is more likely a record of the delivery of grain. Numerous similar illustrations are to be found in collections of Egyptian antiquities, as in the Archeological Museum at Florence (Egyptian coll., 2631 and 2652).

[3] *De Vita Pythagorae*, cap. v, § 22. "Primo itaque illum in arithmeticam et geometriam introduxit, demonstrationibus in abaco propositis. . . ." For further evidence as to the Babylonian origin see Volume I, page 40.

[4] "Et quum instrumenti hujus Assirii inventores fuisse perhibeantur." From a MS. in Paris, transcribed by F. Woepcke, *Journal Asiatique*, I (6), 48 n.

The Abacus in Greece. The abacus[1] and the counters[2] are mentioned several times in Greek literature. It is possible that one of the pictures on the so-called Darius vase in the Museum at Naples is intended to represent such an instrument, although what various writers have stated to be an abacus may be merely the table of the receiver of tribute. In the lowest line of figures in the illustration the king's treasurer may be seen as the figure next to the last one on the left. The other figures represent the bearers of tribute. On the table itself are the letters MΨHΔΠO<T, which are the ordinary numerals representing ten thousands, thousands, hundreds, tens, and fives, together with the symbols for the obol, half obol, and quarter obol. These symbols resemble those on the Salamis abacus mentioned below. The receiver of tribute holds a diptych, or two-leaved wax tablet, in his hand. Upon this tablet are the letters TAΛATA:H, which seem to stand for τάλα(ν)τα ἑ(κατόν) (*tal'anta hekaton'*, hundred talents). The receiver of tribute seems to be casting something on the table, the picture refers to the Persian wars of the time of Darius, these wars took place about 500 B.C., and coins were then known; hence he may have



THE DARIUS VASE

The collector of tribute mentioned in the text is the figure next to the left-hand one in the lowest row. He has a tablet in one hand, and there is a table in front. From the Museum at Naples

[1] "Αβαξ, ἀβάκιον (*a'bax, aba'kion*).


[2] Ψῆφοι (*pse'foi*).

been casting either coins or counters. The one thing that leads to the belief that the table is an abacus is the numerals, but there are no lines such as are found on the Salamis specimen. The date of the vase itself is unknown, but the style shows it to be of the best Greek period. It was found in 1851.[1]

Salamis Abacus. While there is some question as to the figure on the Darius vase, there seems to be little respecting an abacus found on the island of Salamis. It is of white marble, 1.49 m. long and 0.75 m. wide, and is broken into two unequal parts, but is otherwise well preserved and is now in the Epigraphical Museum at Athens.[2] Of the history of this specimen but little is known. It was found before the days of the careful keeping of records, and we are ignorant of its date and of the exact place in which it was discovered. It may have been the computing table in the counting house of some dealer in exchange, and in some of its features it is not unlike the tables used by bankers in the Middle Ages; or, as Kubitschek thinks, it may have been used in some school. The theory that it may have been used in scoring games of some kind seems to have no substantial foundation. In any case it was apparently used for the mechanical representation of numbers by means of counters. It should be observed that, although the crosses are at intervals of three spaces, the first is not on the fourth line as in the medieval European abacus.

[1] The vase is unusually large, being 1.3 meters high. For a good description see A. Baumeister, *Denkmäler des klassischen Altertums*, I, 408 (Munich, 1885). I have slightly changed the inscription from a personal examination of the vase. See Heath, *History*, I, 48; M. N. Tod, "Greek Numeral Notation," *Annual of the British School at Athens*, XVIII, 124.

[2] A description was first published by Rangabé in the *Revue Archéologique*, III, 295 seq., with a comment by A. J. H. Vincent, p. 401. Until 1899 all reproductions of the stone seem to have been derived from the drawing in Rangabé's article. In that year Dr. Nagl (*Zeitschrift* (Hl. Abt.), IX, 337–357, and plate) published an illustration of the abacus under the mistaken impression that it was different from Rangabé's specimen. In the same year W. Kubitschek set forth the facts and gave a satisfactory photograph in the *Wiener Numismatische Zeitschrift*, XXXI, 393–398, Plate XXIV. The author has had a cast taken from the original, and from this the above description is made. See also Harper's *Dictionary of Classical Literature and Antiquities*, p. 2 (New York, 1897); hereafter referred to as Harper's *Dict. Class. Lit.*; Heath, *History*, I, 49–51; Tod, *loc. cit.*, p. 116.



THE SALAMIS ABACUS

It will be seen that the marble slab is ruled as usual, so that counters could be placed on the lines. On three sides are Greek characters substantially as follows:

Ⱶ	1, drachma, a mutilated form of E, for ἕν
Γ	5, old form of Π, for πέντε
Δ	10, for δέκα
⟦Γ	50, for Π and Δ, five tens
H	100, for HEKATON, old form for ἑκατόν
⟦H	500, for Π and H, five hundreds
X	1000, for χίλιοι
I	the obol
C	the half obol
T	the quarter obol
χ	for χαλκοῦς, the eighth of an obol
⟦X	5000, for Π and X, five thousands
T	the talent of 6000 drachmas.

The lines at the top were for fractions. In the illustration the lines and symbols have been accentuated for the sake of clearness.

As to whether the Greeks commonly used loose counters or not we can only infer from this single extant specimen of an abacus, and possibly from the Darius vase. The former and possibly the latter lead us to believe that the loose counters were preferred to those sliding on wires or rods. We do not know any details as to the actual methods of computing, and in spite of the effort of Herodotus to be clear on the subject[1] we are uncertain whether the rows were horizontal or vertical with respect to the computer.[2] It seems probable that the Greeks made less use of the abacus than the Romans, the Greek numerals being better adapted to the purposes of computation, particularly of multiplication and division.[3]

[1] Liber II, cap. 36.

[2] H. Weissenborn, *Zur Geschichte der Einführung der jetzigen Ziffern*, p. 2 (Berlin, 1892), and authorities cited.

[3] J. G. Smyly, "The employment of the alphabet in Greek logistic," in the *Mélanges Jules Nicole*, p. 521 (Geneva, 1905); H. Suter, *Geschichte der math. Wissenschaften*, 2d ed., I, 11 (Zürich, 1873); Heath, *History*, I, 52.

The Abacus in Rome. There were at least three forms of abacus used by the Romans,—a grooved table with beads, a marked table for counters, and the primitive dust board.[1] In respect to each of these forms Latin writers give us considerable information. Horace, for example, speaks of the schoolboy with his bag and table hung upon his left arm, the table referring to the abacus or the wax tablet;[2] and Juvenal mentions both the table and the counters.[3] Cicero refers to counters when he speaks of the *aera* (bronzes), the computing pieces being then made of bronze,[4] and Lucilius the satirist, who lived a generation earlier, does the same.[5] The common name for these counters was, however, *calculi* or *abaculi*, and the material from which they were made was originally stone and later ivory and colored glass.[6] The word *calculus* means "pebble" and is the

[1] S. Hoüel, in his review of Friedlein's "Die Zahlzeichen," in Boncompagni's *Bullettino*, III, 78; Günther, *Math. Unterrichts*, p. 95 n.; A. J. H. Vincent, *Revue Archéologique*, III, 401; A. Kuckuck, *Die Rechenkunst im sechzehnten Jahrhundert*, p. 6 (Berlin, 1874). Although we have numerous references to the use of loose counters, it is curious that no ancient writer speaks definitely of the ruled table on which they are used; see Gerhardt, *Études*, p. 16. On the abacus as a gaming table, particularly for dice, see G. Oppert, *On the original inhabitants of Bharatavarṣa or India*, p. 329 (London, 1893); W. Ramsay and R. Lanciani, *Manual of Roman Antiquities*, 17th ed., p. 497 (London, 1901) (hereafter referred to as Ramsay and Lanciani). For a bibliography and description of the κύβοι (*kuʹboi*) and *tesserae* see J. Marquardt, *La vie privée des Romains*, French translation, II, 522 (Paris, 1893).

[2] Laevo suspensi loculos tabulamque lacerto.
Sat., I, 6, 74

[3] Computat . . . ponatur calculus, adsint
Cum tabula pueri; numera sestertia quinque
Omnibus in rebus, numerentur deinde labores.
Satire IX, 40

[4] "Si aera singula probasti." *Philosoph. Fragmenta*, V, 59.

[5] Hoc est ratio? perversa aera, summa est subducta improbe!
L. 886, ed. Marx; l. 740, ed. Lachmann

[6] Adeo nulla uncia nobis
Est eboris, nec tessellae nec calculus ex hac
Materia. Juvenal, XI, 131

Fragmenta teporata . . . fundi non queunt praeterquam abrupta sibimet in guttas, veluti cum calculi fiunt, quos quidem abaculos appellant aliquos et pluribus modis versicolores.—Pliny, *Hist. Nat.*, XXXVI, 26, 67

Capitolinus (*Pertinax*, I, 4), speaking of the boyhood of Pertinax (126–193), says: "Puer litteris elementariis et calculo imbutus."

Martial (II, 48) includes among his modest wants "tabulamque calculosque."

diminutive of *calx*, a piece of limestone (often referring to the special form of chalk, the name of which comes from the same root). It is therefore our word "marble" as applied to the small spheres with which children play games. From it came the late Latin *calculare*,[1] to calculate. Teachers of calculation were known as *calculones* if slaves, but *calculatores* or *numerarii* if of good family.[2] To calculate means literally, therefore, to pebble, and a calculator is a pebbler. The word *calculi* was transmitted by the Romans to medieval Europe and was in common use until the 16th century.[3]

We are not sure whether the small disks found in Roman remains were counters for purposes of calculation, counters for games (like American poker chips), or draughts. The games of backgammon and draughts are both very old,[4] and the former is our nearest approach, aside from such abaci as we still use, to the Roman and medieval abacus.[5]

The abacus in which the beads were allowed to slide in grooves or on rods is not mentioned by any early writer and seems to have been of relatively late invention. Indeed, in the 15th and 16th centuries it was commonly asserted that Ap-

[1] The Romans used *calculos subducere* instead of *calculare*. This word, in the sense of "to calculate," is first found in the works of the poet Aurelius Clemens Prudentius, who lived in Spain *c.* 400; see *Nouvelles Annales de Math.*, XVII, supplementary bulletin, p. 33.

[2] Tertullian, evidently with reference to the dust abacus, calls them "primi numerorum arenarii."


[3] Thus Clichtoveus, in his arithmetic of 1503 (1507 ed., fol. b, iij, *v.*), says: "Numeratio calcularis est cuiusq3 numeri suo loco et limite apta per calculos dispositio"; and Noviomagus (1539, fol. 9, *r.*) says: "Ut detur autem hac forma in calculis seu ut nunc fit nummis."

[4] They appear in various Egyptian, Greek, and Roman remains. For example, in the British Museum is an ancient model of an Egyptian barge on which a game of draughts is in progress, and A. Baumeister (*Denkmäler des klassischen Altertums*, I, 354 (Munich, 1885)) has reproduced an illustration of a similar game from an old Greek terracotta. There were two Roman games, the *ludus latrunculorum* and *ludus duodecim scriptorum*, on which pieces called *calculi* were used, but their exact nature is unknown. See Ramsay and Lanciani, p. 498; J. Marquardt, *La vie privée des Romains*, French translation, II, 530 (Paris, 1893); Harper's *Dict. Class. Lit.*, p. 562; J. Bowring, *The Decimal System*, p. 198 (London, 1854).

[5] It was probably the *ludus duodecim scriptorum* already mentioned, or the διαγραμμισμός (*diagrammismos'*), the late τάβλα (*ta'bla*), of the Greeks.

puleius invented this form of the instrument in the 2d century,[1]—a statement for which there is no standard authority.

Our knowledge of the grooved abacus is derived from a few specimens of uncertain date which have come down to modern times. One of these, formerly owned by Marcus Welser of Augsburg, was made of metal, is said to have been 4.2 cm. long



ROMAN ABACUS

Ancient bronze abacus of uncertain date, now in the British Museum

and 3.5 cm. wide, and had nineteen grooves and forty-five counters or buttons (*calculi*)[2]. Another was once owned by the reformer Ursinus (*c.* 1575), but is now lost. A third specimen, of bronze, is now in the Kircherian Museum at Rome. The general plan of the Roman abacus may be seen from the illustration here given, representing a specimen in the British Museum.

The symbols found on such specimens as are extant are usually the common Roman numerals from 1,000,000 down to 1,

[1] Unger, *Die Methodik*, p. 69.

[2] This was twice described before it was lost, once in Amsterdam in 1674 and once in Nürnberg in 1682. The measurements are questionable. See G. A. Saalfeld, "Der griechische Einfluss auf Erziehung und Unterricht in Rom," *Neue Jahrbücher für Philologie*, CXXVI, 371.

together with o (or θ) for *uncia*, or $\frac{1}{12}$ of the *as*; S for *semiuncia*; ɔ for the *sicilicus*, or $\frac{1}{4}$ *uncia*; and Z for the *duella*, or $\frac{1}{3}$ *uncia*.

The Abacus in China. At the present time the use of the abacus is universal in China. In banks, shops, and counting houses of all kinds the computations are performed on the *suan-pan*.[1] The computer works very rapidly, like an expert typist or pianist, and secures his results much more quickly than can be done by our common Western methods.[2] He learns its use by practical experience in business, probably as the Romans and Greeks learned it, and not in the village schools.[3]

The *suan-pan* is, however, a relatively late development of the abacus in China, appearing first, so far as we know at the present time, in the 12th century.[4] It is true that many writers[5] have placed its introduction much earlier, but there is no definite description of the instrument in Chinese before about 1175.[6]

[1] The term means computing plate or computing tray, often incorrectly translated as computing board. It is also called the *su-pan*, and there are other variants. It is called *suinbon* in Calcutta, where it is used by all the Chinese shroffs (computers, accountants, cashiers) in the counting houses. The common spelling is suan-pan, swan p'an, or swan pan. The instrument is also in common use in Siam and wherever Chinese merchants have determined business customs.

[2] See Knott, *Abacus*, p. 44; J. D. Bell, *Things Chinese*, p. 1 (New York, 1904); J. Goschkewitsch, "Ueber das chinesische Rechnenbrett," *Arbeiten der kaiserlich Russischen Gesandtschaft zu Peking*, I, 293 (Berlin, 1858); Smith-Mikami; R. van Name, "On the Abacus of China and Japan," *Journal of the Amer. Orient. Soc.*, X (*Proceedings*), p. cx; J. Bowring, *The Decimal System*, p. 193 (London, 1854).


[3] A. H. Smith, *Village Life in China*, p. 105 (New York, 1899).

[4] One of the most scholarly articles on the history of the *suan-pan* is the one already cited, by Lacouperie, *The Old Numerals*, pp. 297–340. It contains an excellent bibliography of the subject up to 1883.

[5] J. Hager, *An Explanation of the Elementary Characters of the Chinese*, p. x (London, 1801); H. Cordier, *Bibliotheca Sinica*, col. 509 (Paris, 1881–1895); L. Rodet, "Le Souan-pan et la Banque des Argentiers," *Bulletin de la Société Mathématique de France*, Vol. VIII (Paris, 1880). Chinese writers record that a work, *Su-shuh ki-i*, "Anecdotes of mathematics," written about 200, mentions various methods of computing, including "bead computation" and "hand computation," but the work gives no description of any process. See also an interesting early essay, Smethurst, "Account of the *shwan pan*," *Phil. Trans.*, XLVI (1749), 22.

[6] This occurs in two works, the *Pan chu tsih* and the *Tseu pan tsih*, which appeared in the Shun-hi dynasty, 1174–1190. They describe the *pan*, or tray, the word *suan-pan* not being then in use. Indeed, as late as the 16th century the name *pan shih* (board to measure) was used. See Lacouperie, p. 38.

As to the origin of the Chinese abacus, the evidence seems to point to Central or Western Asia. At the time of its appearance, China was largely under the domination of the Tangut or Ho-si state and of the Liao and Kin Tartars. The Tangutans were a mercantile race, and the Tartars were favorable to learning. Moreover, Arab and Persian traders are known to



MODERN CHINESE ABACUS

The *suan-pan*, known to have been used as early as the 12th century

have been in Canton in the 8th century, and the Nestorians were in contact with the northwest, so there was plenty of opportunity for such a simple device to make its way into China from Khorasan or some neighboring province. The fact that it seems to have reached Russia from Central Asia[1] adds to the belief that China may have received it from the same source.[2]

Before the time of the *suan-pan* the counting rods, often called the bamboo rods, had been used for more than a thousand years. They were known *c.* 542 B.C.[3] and are referred to as counting stalks in a statement of Hiao-tze, the ruler of Ts'in from 361 to 337 B.C. They are mentioned again about 215 B.C., and some specimens of this period were displayed in a museum


[1] Larousse, *Grand Dictionnaire Universel*, I, 636; Vissière, *Abacque*; A. Wylie, *Notes on Chinese Literature*, p. 91 (Shanghai, 1902).

[2] Lacouperie, *The Old Numerals*, p. 41.

[3] See Volume I, page 96.

of the Emperor Ngan (397–419). These were about 18 inches long, some made of bone and others of horn. In the reign of Wu-tí (140–87 B.C.) of the Han dynasty, it is related that an astronomer Sang Hung (about 118 B.C.) was very skillful in his use of the rods. In the third century of our era it is recorded that Wang Jung, a minister of state, spent his nights in reckoning his income with ivory calculating rods, and the expression "to reckon with ivory rods" is still used as an allusion to wealth. In the time of the Emperor Ch'eng (326–343) the counting rods were made of wood, ivory, or iron, and two centuries later the Emperor Siuen Wu (500–516) had counting rods cast in iron for the use of his people.[1]

The Chinese historian Mei Wen-ting (1633–1721), in his work on ancient calculating instruments,[2] states that about the beginning of the Christian era 271 rods constituted a set, or handful, and that they formed a hexagon that had nine rods on a side. This means that they were arranged in six groups of which the ends of each formed a triangular number of $1 + 2 + \cdots + 9$ units, or 45 in all. Six of these make 6×45, or 270, and these six were grouped about one central rod, making 271, thus affording an illustration of the use of figurate numbers in the East.




It seems from Mei Wen-ting's work that the rods were in general use until the 13th century. With respect to the *suan-pan*, he places the date somewhat later than other writers, saying, "If in my ignorance I may be allowed to hazard a guess, I should say that it began with the first years of the Ming dynasty," which would make the date about 1368. Subsequent writers are probably correct, however, in placing it a century or two earlier.

The Abacus in Japan. The primitive method of computing in Japan is quite unknown, but from the time of the Empress Suiko (593–628) the bamboo rods (*chikusaku*) were used.[3]

[1] Lacouperie, *The Old Numerals*, pp. 34–36 of reprint.
[2] *Ku-suan-k'i-k'ao.*
[3] Smith-Mikami, chap. iii, with bibliography.

These were round sticks about 2 mm. in diameter and 12 cm. in length, but because of their liability to roll they were in due time replaced by the *sanchu* or *sangi*, rectangular prisms about 7 mm. thick and 5 cm. long. The *soroban*, the name being probably the Japanese rendering of the Chinese word *suan-pan*, was developed but not generally adopted in the 16th century.[1]



THE SANGI BOARD IN JAPAN

Intended for computation with the *sangi* (rods). From Satō Shigeharu's *Tengen Shinan*, 1698


What may prove to be a relic of a very early Japanese system is seen in the tally sticks used in the Luchu (Liu Kiu, Riu Kiu) Islands, near Formosa, and known as *Shō-Chū-Mā*.[2]

The Abacus in Korea. The bamboo rods of China passed over to Japan by way of Korea, and in the latter country they re-

[1] Smith-Mikami, p. 19.

[2] B. H. Chamberlain, *Journal of the Anthropological Institute of Great Britain and Ireland*, XXVII, 383. For a brief mention of these tallies see the *Geographical Journal*, June, 1895.

mained in use long after they were abandoned elsewhere. The commercial class was acquainted with the *suan-pan* for a long




SANGI BOARD WITH NUMBERS INDICATED

From Nishiwaki Richyū's *Sampō Tengen Roku*, 1714. The *sangi* board was a board ruled as shown, the *sangi* being placed in the rectangles

time before the Japanese conquest, and now the *soroban* is common among the officials. But most of those who were edu-


cated in the native schools used the counting sticks until recent times, while those with but little education performed their



THE SANGI BOARD IN USE

From Miyake Kenryū's *Shōjutsu Sangaku Zuye*, 1716 (1795 ed.)

simple computations mentally or on their fingers. The counting sticks (*Ka-tji san*) were of bone, as in the illustration on page 174, or of bamboo split into long prisms. About a hun-



JAPANESE ABACUS


The *soroban*, known to have been used in Japan as early as the 16th century, and in universal use there at present

dred fifty were used in ordinary calculation, and these were kept in a bamboo case on the computer's desk. The sticks were laid as follows to represent the first twelve numbers:

I	II	III	IIII	X	XI	XII	XIII	XIIII	—	T	TT
1	2	3	4	5	6	7	8	9	10	11	12

In computing, the Koreans used the rods in substantially the same way as the Chinese and Japanese had used theirs. The

process was so cumbersome that it has recently given way to the Chinese and Japanese methods with the *suan-pan* and the *soroban*.[1] The Koreans also used pebbles and coins for the same purpose.[2]



KOREAN COMPUTING RODS

Computing rods made of bone. Until quite recently these were used in the schools of Korea. The numbers were represented as shown on page 173

The Abacus among the Mohammedans. The Arabs, Persians, Armenians, and Turks have a form of abacus which differs from that of the Far East and from the one used by the Romans, having ten beads on each line. Its early history is unknown, but since it resembles neither the abacus of China nor that of Western Europe, it probably originated among the Arab or Persian computers. The Turks call it the *coulba* and the Armenians the *choreb*.[3]

This form of the abacus does not seem to have been generally used by the Saracens in the Middle Ages. In

[1] On the mathematics of Korea in general, see P. Lowell, *The Land of the Morning Calm*, p. 250 (Boston, 1886). For the *Song yang hoei soan fa* or *Song yang houi san pep* (*Treatise on Arithmetic of Yang Hoei of the Song Dynasty*), which was for a long time a classic, see M. Courant, *Bibliographie Coréenne*, III, 1 (Paris, 1896). See also the *Grammaire Coréenne*, p. 44 (Yokohama, 1881), in which the description of the laying of the sticks recalls the Japanese method and differs from the one shown on page 173, which was given to the author by an educated Korean in Peking.

[2] *Grammaire Coréenne, loc. cit.*

[3] Pacioli speaks of this form of the abacus when he says that the orders of numbers increase from right to left "more arabū de simil arte pratica primi inuētori secōdo alcuni vnde ꝑ ignorātia et vulgo a corropto el vocabulo dicēdo la Abaco: cioe modo arabico. Che loperare suo e modo arabico e chiamase Abaco: ouer secōdo altri e dicta Abaco dal greco vocabulo." *Sūma*, fol. 19, *r.* (Venice, 1494).

that period the dust board was common and the numeral forms derived from being written on such a tablet were therefore, as already stated, called in the schools of the western Arabs the *ġobâr* (dust) numerals.[1] Thus the Moorish writer al-Qalasâdî (*c.* 1475), in his commentary on the *Talchîs* of Albanna (*c.* 1300), speaks of "a man of the Indian nation who took fine powder and sprinkled it on a table and marked on it the multiplications, divisions, or other operations, and this is the origin of the term *ġobâr*" (dust).[2] Further evidence of the rarity of any other form of the abacus among the Saracens in the Middle Ages is to be found in the silence of Maximus Planudes (*c.* 1340) upon the subject; for the contact with the East of one writing upon arithmetic in Constantinople would almost certainly have led him to speak of the bead abacus if it had been in common use among the Arabs of his time. It may be, however, that the dust abacus was used in some parts of the Mohammedan domain, and the bead abacus in other parts, the latter giving to Christian nations the line abacus. Some reason for this belief is found in the fact that certain medieval writers derived the word "abacus" from the Arabic,[3] while William of Malmesbury, although by no means a reliable chronicler, writing in the 12th century, says that Gerbert (*c.* 1000) obtained his idea of the instrument from the Saracens.[4] There is also a possible reference to the line abacus by Alchindi[5] (*c.* 860).

The Abacus in Russia. From the Mohammedan countries the bead abacus worked its way northward, and in comparatively

[1] H. Weissenborn, *Gerbert*, p. 235 (Berlin, 1888); *Zur Geschichte der Einführung der jetzigen Ziffern*, p. 7 (Berlin, 1892); Smith-Karpinski, p. 65.


[2] From an Arabic MS. in Paris, described by F. Woepcke, *Journal Asiatique*, I (6), 60.

[3] Thus a 12th century MS., *Regulae abaci*, published by M. Chasles in the *Comptes rendus* for 1843 (XVI, 218), asserts, "Ars ista vocatur abacus: hoc nomen vero arabicum est et sonat mensa."

[4] "Abacam certe primus a Sarazenis rapiens, regulas dedit, quae a sudantibus abacistis vix intelligentur." On the unreliability of this chronicler, see H. Weissenborn, *Gerbert*, p. 236 (Berlin, 1888).

[5] The reference is in the chapter "De numeris per lineas & grana hordeacea multiplicandis Liber I" of the Latin translation of his arithmetic. See H. Weissenborn, *Zur Geschichte der Einführung der jetzigen Ziffern*, p. 7.

recent times was adopted in Russia. It is still found in every school, shop, and bank of Russia proper, although in the former provinces of Finland and Poland it is seldom used. The computers handle it with much the same ease as the Chinese show in their use of the *suan-pan*, and there seems to be no reason why they should not continue to use it until it is replaced by more elaborate calculating machines. The Russians call their abacus the *s'choty*,[1] and the form is the same as that of the Armenian *chorcb* or the Turkish *coulba*. They occasionally speak of it as the Chinese abacus, so that there is this ground for the claim that it was introduced from China by way of Siberia, although the form of the instrument would go to show that it came from the South.



RUSSIAN ABACUS

The *s'choty* of the Russians. It is of the same form as the Armenian *chorcb* and the Turkish *coulba*

In the 16th century the German form of line reckoning was used in Poland,[2] and when this disappeared it was not replaced by the Russian abacus but by the algorism of Western Europe.

[1] Variously transliterated, but this form gives the pronunciation more nearly than the others.

[2] The third arithmetic printed in Poland, but the first in the Polish language, is that of Kĭos (1538). It is devoted almost entirely to this kind of computation. See the Baraniecki reprint (Cracow, 1889) and Dickstein's article in *Bibl. Math.*, IV (2), 57.

The Abacus in Western Europe. In medieval times in Western Europe the abacus had various names and forms. The followers of Boethius (*c.* 510) called it the Pythagorean table (*mensa Pythagorica*), a name also given to the square array of the multiplication table.[1] It was also known as the geometric table (*tabula geometricalis, mensa geometricalis*), table of the abacus (*tabula abaci*), and Pythagorean arc (*arcus Pythagoreus*), although *abax* or *abacus* was the common medieval name.[2] So common was this name that the verb "to abacus" became recognized,[3] and the arithmeticians of about the 11th century and later were occasionally called abacists.

The Dust Table. Of the various forms of abacus used in Europe, the dust table, already described as known in the Orient and in classical times, was one. We have evidence of its use at the close of the 9th century, when Remigius of Auxerre (*c.* 900), in his commentary on Capella's arithmetic, speaks of the table as being sprinkled with blue or green sand and the

[1] Cantor called attention to this distinction in his *Mathematische Beiträge zum Kulturleben der Völker*, Halle, 1863, p. 204 (hereafter referred to as Cantor, *Beiträge*); and Eneström did the same in the *Bibl. Math.*, I (2), 90. Adelard of Bath (*c.* 1120), in his *Regulae Abaci*, says: ". . . quidem mensam pithagoream ob magistri sui reuerentiam. sed postī tamē abacum dixerunt" (Boncompagni's *Bullettino*, XIV, 68). In this he apparently had in mind a passage in the *Ars Geometria* of Boethius (ed. Friedlein, p. 396): "Pythagorici vero . . . descripserunt sibi quandam formulam, quam ob honorem sui praeceptoris mensam Pythagoream nominabant . . . a posterioribus appellabatur abacus." Adelard even goes so far as to assert, with no foundation except tradition, that the abacus is due to Pythagoras himself: "Pythagorici vero hoc opus [abacum] composuerunt/ut ea que magistro suo pitagora docente audierant. ocul' subiecta retinerent: et firmius custodirent."

[2] Turchillus, writing on the abacus about 1200, says: "Ab antiquis mensa pytagorica, a modernis autem uel abax vel abacus nuncupatur" (Boncompagni's *Bullettino*, XV, 135). An anonymous MS. of the 12th century, in the Vatican, says: "(T)abula abaci quę pytagorea m̄sa uocatur" (*ibid.*, pp. 132, 154). See also M. Chasles, "Développements et détails historiques sur divers points du système de l'abacus," *Comptes rendus*, XVI, 1393, with references to other MSS.

[3] In a MS. in the Bibliothèque nationale in Paris there is some correspondence between one Radulph of Liège and Rogimbold of Cologne in the early part of the 11th century in which the writer says: "Hoc si abacizando probaveris." In the same MS. there is a letter addressed to Hermannus Contractus (*c.* 1050) in which the statement is made: "Ut meam abicizandi notem inscitiam." See Chasles, *loc. cit.*, p. 1417.

figures as being drawn with a *radius*.[1] A certain Papias, who wrote a *Vocabularium* in 1053, and who may be considered as representing the knowledge of his time, also speaks of the abacus as a table covered with green sand.[2]

The Wax Tablet. Allied to the dust table is the old wax tablet of classical times. This consisted of a tablet of wood or bone on which a thin coat of black wax was smeared, the figures being written with an iron stylus of which one end was pointed and the other was somewhat spoon-shaped, the latter being used for erasing by smoothing the wax down again.[3] This tablet passed into the medieval schools and counting houses, and specimens are extant which were in use as late as the 16th century.[4]

[1] "Abacus tabula est geometricalis super quam spargebatur puluis uitreus siue glaucus. Ibique cum radio uirgę formabantur figurę geometrię." See Boncompagni's *Bullettino*, XV, 572, and III, 84. In the same journal (X, 625) there is a description of a medieval MS. of unknown date in which the following passage appears: "Abacus vocatur mensa geometricalis que et in numeris et formis numerorum diuisa . . . ."


[2] "Abacus vel abax tabula: in qua uiridi puluē formae depinguntur." From the first printed edition of the *Vocabularium*, Milan, 1476, fol. 2, *v*. See also Boncompagni's *Bullettino*, XIV, 69.

It would seem that Adelard of Bath (*c.* 1120) referred to this form of abacus when he wrote: "Vocatur (Abacus) etiam radius geometricus, quia cum ad multa pertineat, maxime per hoc geometricae subtilitates nobis illuminantur." Radulph of Laon (*c.* 1125) had the same form in mind in writing the following: ". . . ad arithmeticae speculationis investigandas rationes, et ad eos qui musices modulationibus deserviunt numeros, necnon et ad ea quae astrologorum sollerti industria de variis errantium siderum cursibus . . . Abacus valde necessarius inveniatur." See Chasles, *loc. cit.*, p. 1414.

[3] One of the best specimens of this kind seen by the author is a 6th century Roman piece in the Rylands Library at Manchester, England. This is made of bone and is a diptych bound with iron and having an iron hinge. It has three iron styli, one end being pointed for writing and the other end being spoon-shaped for erasing.

[4] There is an elaborate set of Comptes de l'hôtel Saint Louis, written in 1256–1257, in the Musée des Archives, Paris. It consists of between ten and twenty placques. There is a 15th century specimen in the Germanic Museum at Nürnberg, from southern Germany; a piece apparently of the 16th century, from a church in Switzerland, may be seen in the British Museum; but the most interesting one of the medieval period that has come to the author's attention is in the Rathhaus at Goslar a. Harz,—a book of eight tablets bound together and making sixteen pages, two compartments to a page. This Goslar specimen is a Bürgerrolle of the 14th or 15th century, the numerals being Roman and the original stylus being annexed.

The Slate. In the later Middle Ages the slate replaced the wax tablet and sand table, and continued in use until the manufacture of cheap paper rendered it nearly obsolete at the close of the 19th century. The earliest printed reference to it is


Prosdocimi de beldamandis algo-
rismi tractatus perutilis ꝛ neceſſarius
foeliciter incipit. qui de generibus cal-
culationum ſpecie preteri. t nullaz. q̄ ſalte
neceſſaria ad hꝰ artꝫ ꝺgnitoz fuerat

Inueni inq̄z pluribus libris algoriſmi nūcupa
tis. mōs circa numeros opandi ſatis uarie:
atqz diuerſos. q licet boni exiſtere. atqz veri
erāt. tñ faſtidioſi: tū ꝓpt̄ ipaz regulaz mul-
titudinē: tū ꝓpt̄ eaz delectōes. tū etiā ꝓpter
ipaz operationū ꝓbaſōes: vtz. ſ. bone fuerint uel ue. Erāt ꝛ eti
am iſti modi inuñ faſtidioſi: q ſi in aliq̄ calculo aſtroloico erat
ꝺtigiſſz: calculatorē opatoz ſuam a capite incipere oportebat: da
to q erroꝛ ſuus adhuc ſatis ꝓpiquus exiſteret. ꝛ hoc ꝓpt̄ figu
ras in ſua opatōe deletas. Indigebat etiā calculatoꝛ ſemꝑ aliq̄
lapide uel ſibi ꝺformi. ſuꝑ quo ſcribere atqz faciliter delere poſſz
figuras cū ꝗbuſ opabat' in calculo ſuo. Et ꝗa hec oia ſuns fa:i

FIRST PRINTED REFERENCE TO A SLATE

From Beldamandi's work, 1483. See the word *lapide* in next to the last line

probably in the *Algorismus* of Prosdocimo de' Beldamandi, a work written in 1410 and first printed in 1483,[1] in which the author speaks of the necessity which a computer has for a slate from which he can easily erase what he has written.[2] The gain to the art of computation which resulted from this invention can hardly be realized at the present time.

[1] "Anno domini .1410. die .10. Iunij compilata." See *Rara Arithmetica*, p. 13.

[2] "Indigebat etiã calculator semꝑ aliq̄ lapide uel sibi ꝺformi, suꝑ quo scribere atqz faciliter delere possz figuras cũ ꝗbus opabat' in calculo suo" (1540 ed., fol. 2, *v.*). Compare also J. T. Freigius, *Pædagogvs*, Basel, 1582: "Numeri in abaco scribendi."

In this period, but we do not know precisely when, there came into general use the blackboard, arranged for hanging on a wall. This is frequently shown in the illustrations in the early printed books, as in the case of Böschensteyn's work of 1514.




EARLY ILLUSTRATION OF A BLACKBOARD

From Johann Böschensteyn's *Rechenbiechlin,* Augsburg, 1514


Gerbert's Abacus. To Gerbert (*c.* 1000) there is attributed the arc or column abacus.[1] If we could put one counter marked "4" on a line instead of putting four counters upon it, there would seem at first thought to be some gain. This apparent gain is offset, however, by the loss of time in selecting the counters and still more by the necessity for learning certain tables. The plan was followed by Gerbert, and possibly some of his succes-

[1] *Arcus Pythagoreus, tableau à colonnes.*

sors, the counters being the apices already mentioned on page 75. They represented the number 2,056,708, for example, as follows:



Over each triad of columns an arc was drawn to aid the eye, whence the name "arc abacus"; and in each column in which a number was to be represented a counter bearing that number was placed. As already stated, however, the gain over the older form was more apparent than real, for the computer was under the necessity of picking out the right counter each time. If Gerbert had understood the significance of the zero, he would not have used this device.



GENERAL PLAN OF A MEDIEVAL COMPUTING TABLE

This plan shows the arrangement of lines on the kind of computing table used in most parts of Europe in the Middle Ages

The Line Abacus. The most popular abacus of Western Europe consisted of a table ruled horizontally to represent different decimal orders, counters being placed upon the lines and in the spaces. It was often called a calculating table or simply a table,[1] and in England it received the name "counting table" or "counter."

The illustration of the line abacus, from Köbel's work of 1514, shows the form which was common in all Western Europe

[1] So Hudalrich Regius, in his *Epitome* (1536), says, "Abacus vulgo mensa dicitur calculatoria quibusdam distincta lineis"; and Radulph of Laon (*c.* 1125) asserts that "Greci enim Mensam abacum dicunt." See *Abhandlungen*, V, 96.

for several hundred years. The line nearest the computer represents units, the space above it, fives; the second line, tens; the second space, fifties; and so on.[1]

Representation of Numbers. On the lines and in the spaces, counters were then placed as shown in this illustration in the



COUNTER RECKONING IN 1514

From the title-page of Köbel's *Rechenbiechlin*, Augsburg, 1514

two columns on the table. In the right-hand column, which is the left-hand column of the computer who sits by the window, the number 26 is represented, 2 being on the tens line, 1 in the

[1] One of the best of the older authorities on this type is the work of T. Snelling, *A view of the origin, nature, and use of Jettons or Counters*, London, 1769. See also D. E. Smith, *Computing Jetons*, New York, 1921. The standard authority, however, is Barnard's work already mentioned.

fives space, and 1 on the units line. There should never be more than one counter left in a space or more than four counters on a line; for if there are five on a line, one is "carried" to the space above, and if there are two in a space, one is "carried" to the line above; whence our expression "to carry" in addition.[1] The thousands and millions lines were each marked by a small cross. This aided the eye in reading the numbers and is the origin of our system of separating the figures in groups of three by means of a comma.[2]

The intervals between the horizontal lines were commonly called "spaces" (*spacia*), and the divisions made by the vertical lines were called *cambien*, from the Italian *cambio* (exchange).[3]

Reckoning on the Lines. Computation on this form of abacus was called reckoning "on the lines," and many of the early German arithmetics include the expression "auf den Linien."[4] As a result, a boy who knew his abacus was said to "know the lines."[5] When he represented a number by means of counters on the lines, he was said to "lay" the sum;[6] and when he also knew the modern form of computing which had developed in Italy, he was said to be able to reckon "on the lines and with the pen."[7] He was often advised to "lay and seize" correctly, meaning that he must be careful to place the counter prop-

[1] Other illustrations from early printed books will be found in *Rara Arithmetica*. A good description of the common German abacus is given by A. Kuckuck, *Die Rechenkunst im sechzehnten Jahrhundert*, p. 7 (Berlin, 1874).

[2] "Dieselbe verzeichne mit einem Creutzlin," as J. Albert says in his arithmetic of 1534, mentioned below in note 4.

[3] Also "Cambien oder Bankir" by various other writers; e.g., Köbel, *Zwey rechenbüchlin* (1514; title of edition of 1517). Hudalrich Regius (1536) used the term *viculi*, and J. Albert (1534) speaks of the divisions as *Feldungen* as well as *Cambien* and *Cambiere*.

[4] For example, J. Albert, *Rechenbüchlein Auff der Federn*, Wittemberg, 1534 (title from 1561 ed.); A. Riese, *Rechenung auff der linihen vnd federn*, Erfurt, 1522.

[5] "Die Linien zu erkennen, ist zu mercken, das die underste Linie (welche die erste genent wird) bedeut uns, die ander hinauff zehen, die dritte hundert," etc. J. Albert, *loc. cit.*


[6] "Leg zum ersten die fl." J. Albert, *loc. cit.* This may be connected with our expression "to lay a wager."

[7] As in Riese's work of 1522.

erly and pick it up with the same care. Thus Albert (1534) tells him:


> Write right, lay right, seize right, speak right,
> And you will always get the answer right.[1]

Addition and Subtraction. The operation of addition can be understood by studying the illustration on page 185 from Recorde's work and by considering the following figure suggested by Albert's arithmetic:



(213 + 1450 + 2378 = 4041)

Subtraction was merely the reverse of the above operation, and the word "borrowing" had a more definite meaning than with us. The following figure is also suggested by Albert's work:




(1534 − 186 = 1348)

Multiplication and division were more complicated and are not of enough importance to warrant a description in this work.

Extent of Use of the Line Abacus. During the 15th century the line abacus furnished almost the only means of commercial computation throughout most of Western Europe north of the

[1] "Schreib recht/leg recht/greiff recht/sprich recht/
So kômpt allzeit dein Facit recht."

Alps. In the 16th century we find it given prominently in the printed arithmetics of Germany, Holland, Poland, and Austria, somewhat less prominently in England, and still less in France. We may say that those countries which were chiefly influenced



ADDITION.

Maſter.

The eaſieſt way in this arte, is to adde but two ſummes at ones togyther: how be it, you maye adde moze, as I wil tel you anone. therefore whenne you wylle adde two ſummes, you ſhall fyrſte ſet downe one of them, it forceth not whiche, and then by it draw a lyne croſſe the other lynes. And afterwarde ſette downe the other ſumme, ſo that that lyne maye be betweene them; as if you wouldde adde 1659 to 8342, you muſt ſet your ſümes as you ſee here.

And then if you lyſt, you maye adde the one to the other in the ſame place, or els you may adde them bothe tohither in a newe place: which way, becauſe it is moſt plaineſt I

PAGE FROM ROBERT RECORDE'S *GROUND OF ARTES*, C. 1542

This page shows the treatment of addition. It is from the 1558 edition

by the customs of the Italian merchants tended to abandon the abacus, while those which were in closer contact with the German counting houses continued to use it.

The popularity of the method may be seen in the fact that abacus-reckoning was a favorite subject of illustration in the title-pages of the arithmetics of Adam Riese, Gemma Frisius, and Robert Recorde, which were among the most widely circulated textbooks of the 16th century.

Origin of this Form of Abacus. It is not known when this form of the abacus first appeared. Indeed, there is a break of several centuries in the use of counters in any manner. We are ignorant as to how the Western world computed at the beginning of the Middle Ages, or what method Bede (*c.* 710) and Alcuin (*c.* 775) used in their calculations. In the 13th century counters were used for practical business computation, as they had been in the Roman days, but in the long interval the ancient scheme had changed, the vertical lines giving place to the horizontal. When or where this change took place there is at present no means of knowing.[1] Fibonacci (1202) names three methods of computing in use in his day,—finger reckoning, algorism (Hindu numerals), and the Gerbert abacus.[2] Of the use of ordinary counters he has nothing to say. Certain it is that counters were generally unknown in the 15th century in Italy, for we have the positive assertion of the Venetian patrician Ermolao Barbaro (*d.* 1495) that they were used only in foreign countries.[3]

The Counters in England. The most common of the English names for the small disks used on the line abacus was "counter," a word derived from the Latin *computare* through the French forms *conteor* and *compteur* and the Middle English *countere*, *cowntere*, and *countour*.[4] So in a work entitled *Know Thyself*, written about 1310, we are told to "sitte doun and take countures rounde.... And for vche a synne lay thou doun on Til thou thi synnes haue sought vp and founde," and in a work of 1496 mention is made of "A nest of cowntouris to the King." In the laws of Henry VIII (Act 32, cap. 14, 1540) we read, "Item for euery nest of compters .xviii.*s*," so that the expression was a common one and referred to the box or bag full of computing

[1] A. Nagl, *Die Rechenpfennige und die operative Arithmetik*, p. 8 (Vienna, 1888); A. Kuckuck, *loc. cit.*, p. 15.

[2] "Computatio manibus, algorismus, arcus pictagore."


[3] "Calculos sive abaculos . . . eos esse intelligo . . . qui mos hodie apud barbaros fere omnes servatur." Nagl, *loc. cit.*, p. 40.

[4] By a false etymology we have "comptroller," although this word is properly "controller," one who controls an account, from the Middle Latin *contrarotulum* (*contra* + *rotulus*), a counter roll, a check list.

pieces. Such a nest is probably referred to by Alexander Barclay (*c.* 1475–1552), in his *Egloges*, when he speaks of "The kitchin clarke . . . Jangling his counters."

In the Middle Ages in England it seems to have been the custom of merchants, accountants, and judges who had to consider financial questions to sit on benches (banks) with checkered boards and counters placed before them. Hence the checkered board came to represent a money changer's office, finally becoming a symbol for an inn, probably because innkeepers followed the trade of the money changer.

In Shakespeare's time abacus computation was in low repute, for the poet speaks contemptuously of a shopkeeper as a "counter caster." Counters apparently lost their standing only in the last half of the 16th century, for Robert Recorde, writing *c.* 1542, says: "Nowe that you haue learned the common kyndes of Arithmetike with the penne, you shall see the same arte in counters,"[1] and an anonymous arithmetic of 1546 has


THE GROVND OF
ARTES:
Teaching the woorke and practise of
Arithmetike, both in whole numbres
and fractions, after a more easyer
and exacter sorte, then anye lyke
hath hytherto beene
set forth: with di-
uers newe ad-
ditions.
Made by M. ROBERTE
RECORDE
Doctor of Physike.

COUNTER RECKONING

From the 1558 edition of Recorde's *Ground of Artes*

[1] From the 1558 edition of the *Ground of Artes*, in which Recorde devotes forty pages to this phase of the subject.

"An introduction for to lerne to reken with the pen, or with the counters accordying to the trewe cast of Algorisme." A century later Hartwell, in his appendix to the *Ground of Artes* (1646 ed.), speaks of ignorant people as "any that can but cast with Counters." Even in the first half of the 16th century people had begun to doubt the value of line reckoning, for Palegrave writes in 1530: "I shall reken it syxe tymes by aulgorisme or you can caste it ones by counters." That the abacus died out here before it did in Germany is also evident from the fact that German counters of the 15th and 16th centuries are very common in numismatical collections, while most of those used in England at this time were imported.[1]

From the use of "counter" in the sense described, the word came to mean an arithmetician. Thus we find in one of the manuscripts in the Cotton library the statement, "Ther is no countere nor clerke Con hem recken alle," and Hoccleve (1420) writes: "In my purs so grete sommes be, That there nys counter in alle cristente Whiche that kan at ony nombre sette."[2] The word also came to mean the abacus itself. Thus, in his *Dethe Blaunche* (*c.* 1369) Chaucer says: "Thogh Argus[3] the noble covnter Sete to rekene in hys counter."

Court of the Exchequer. Aside from the mere history of computation an interest attaches to the abacus in England because of its relation to the Court of the Exchequer, the *Chambre de l'échiquier* of the French.[4] In the *Dialogus de Scaccario*

[1] Barnard, *Counters*, p. 63.

[2] See Murray's *New English Dictionary*, II, 1057.

[3] The passage comes from the *Roman de la Rose*, in which this name, with also the spelling Algus, is given for al-Khowârizmî. Chaucer also speaks of the counters as "augrim (*i.e.*, algorism, from al-Khowârizmî) stones." On this subject see L. C. Karpinski, "Augrim Stones," *Modern Language Notes*, November, 1912 (Baltimore).

[4] The best original source of information as to the exchequer is the *Dialogus de Scaccario*, a work written by one Fitz-Neal in 1178–1179 (1181 according to Stubbs) and first edited by Madox in 1711. See also F. Liebermann, *Einleitung in den Dialogus de Scaccario*, Göttingen, 1875, and the Oxford edition of 1902. It is published in E. F. Henderson, *Select Historical Documents of the Middle Ages*, p. 20 (London, 1892). Consult also H. Hall, *The Antiquities and Curiosities of the Exchequer*, London, 1891 (reviewed somewhat adversely in *The Nation*, New

a disciple and his master discuss the nature of the exchequer as follows:

Disciple. What is the exchequer?

Master. The exchequer[1] is a quadrangular surface about ten feet in length, five in breadth, placed before those who sit around it in the manner of a table, and all around it has an edge about the height of one's four fingers, lest anything placed upon it should fall off. There is placed over the top of the exchequer, moreover, a cloth[2] bought at the Easter term, not an ordinary one but a black one marked with stripes, the stripes being distant from each other the space of a foot or the breadth of a hand. In the spaces moreover are counters placed according to their values.

The rest of the description is too long to be given, but it shows that a kind of abacus, although not the one above described, characterized this ancient court.

The counters finally came to be used to keep the scores in games,[3] as in the American game of poker and in the use of markers in billiards. They also remained in the schools for the purpose of teaching the pupils the significance of our number system, sometimes in the form of an abacus with ten beads on a


York, February 25, 1892, p. 157); R. L. Poole's Ford Lectures at Oxford in 1911, published under the title *The Exchequer in the Twelfth Century*; J. H. Ramsay, *The Foundations of England*, II, 323 (London, 1898); Martin, *Les Signes Num.*, p. 32; J. H. Round, *The Commune of London, and other Studies*, p. 62 (London, 1899); C. H. Haskins, "The abacus and the king's curia," *English Historical Review* (1912), p. 101. On the *Dialogus* as the earliest work on English government, consult J. R. Green, *Short History of the English People*.

[1]The word is a corrupt form of the Old French *eschequier* and Middle English *escheker*, based on the mistaken idea that the Latin *ex-* is taken with *scaccarium*. The term *scaccarium* for exchequer first appears under Henry I, about 1100. Before him, under William the Conqueror and William Rufus, we find the terms *fiscus* and *thesaurus*. "Exchequer" was later used to mean a chessboard, as in a work of 1300: "And bidde the pleie at the escheker"; and in Caxton's work on chess (*c.* 1475), p. 135, where it appears as *eschequer*.

[2]There is in the National Museum at Munich a green baize cloth embroidered in yellow with the ordinary arrangement of the medieval German abacus, intended to be laid on the computing table in the manner here described. For illustrations of such pieces see Barnard, *loc. cit.*, plates.

[3]"They were marking their game with Counters." Steele, in *The Tatler*, No. 15 (1709).

line, as seen in primary classes today, and sometimes for the purpose of teaching fractional parts. A specimen possibly intended chiefly for this purpose is seen in an abacus formerly used in the Blue Coat School in London, and here shown.



ABACUS FOR TEACHING FRACTIONS

Formerly used in the Blue Coat School, London. From Mr. Plimpton's collection

Counters in Germany. In no country was the line abacus more highly esteemed in the 15th and 16th centuries than in Germany. Its use had died out in Italy, the great commercial center of the world, but in the counting houses of Germany it was almost universal until the era of printed arithmetics. Indeed, even in the 16th century its superiority was stoutly maintained by various German Rechenmeisters,[1] and as late as the middle of the 18th century its use had not died out in a number of the towns.[2] Even after the leading merchants had learned the method of algorism the common people continued to do their simple sums by the aid of counters purchased from the itinerant peddler,[3] and the most popular of the

[1] Thus Apianus (*Eyn Newe . . . Kauffmansz Rechnung*, Ingolstadt, 1527) asserted: ". . . die Summirung der Register in gewicht mass vnd müntz durch die rechenpfenning auf der linie brauchsamer ist vnd vil schneller vnd füglicher geschicht dann durch die federn oder kreide."

[2] Heilbronner, in his *Historia*, p. 890, says that in his time the counters were still used "in pluribus Germaniae atque Galliae provinciis a mercatoribus," and speaks of computing on the line as "arithmetica calculatoria sive linearis est Scientia numerandi per calculos vel nummos metallicos."

[3] The priest Geiler of Kaiserberg (1445–1510) tells of peddlers' selling them in his day, and the custom doubtless continued. See *Bibl. Math.*, IV (3), 284.

early German arithmetics were based on "Rechnung auf Linien."[1] As late as 1587 Thierfelder testified to the fact that the use of counters was still common in Germany,[2] and in 1591 two arithmetics based on line computation were published.[3] Even as late as 1621 a textbook[4] on the subject appeared at Hildesheim. One of the last writers to describe the process fully was Leonhard Christoph Sturm, whose work was published in 1701.[5]

The common German name for the counter was *Rechenpfennig*,[6] although *Zahlpfennig*[7] and *Raitpfennig* were also used.

Counters in France. In the later Middle Ages France, like all the Latin countries, made less of counter reckoning than the Teutonic lands. The first printed description of counter reckoning in that country dates from about 1500,[8] and although

[1]So Köbel's well-known arithmetics, which went through various editions beginning with 1514, gave only the counter reckoning; Adam Riese's famous textbooks, beginning in 1518, favored it; and various other popular textbooks gave it prominent place. Even as good a mathematician as Rudolff introduced it immediately after the treatment of algorism in his *Kunstliche rechnung* of 1526.

The first arithmetics printed in Germany appeared at Bamberg in 1482 and 1483. Neither seems to have had anything to say about counters, although we have only a fragment of the earlier one, and perhaps this failure explains the lack of popularity of these books. The subject was so popular that Stifel (1544) calls it *Haussrechnung*. On the general subject, see H. Schubert, "Die Rechenkunst im 16. Jahrhundert," *Deutsche Blätter für Erziehenden Unterricht*, III, 69, 105.

[2]On the first page of his *Rechenbuch* he says: "Wie vil sind Arten oder Weisen/die im Rechnen am meysten gebraucht werden? Fürnehmlich zwo/ Die erst mit der Feder/oder Kreyden/durch die Ziffern. Die ander mit den Zahlpfenning auff den Linien."

[3]One (Swiss) by Mewrer in Zürich and the other by Kauder in Regensburg.

[4]The work was anonymous. The title is *Ein new Rechenbüchlein auf Linien und Ziffern*. See Nagl, *Die Rechenpfennige*, p. 27, for other cases.

[5]*Kurtzer Begriff . . . Mathesis.*

[6]Reckoning penny. The spelling varies. Stifel, for example, in his *Deutsche Arithmetica*, 1545, fol. 1, calls them *Rechenpfenning* (both singular and plural). *Rechenpfennig* was merely a translation of the medieval Latin name, as is seen in the arithmetic of Clichtoveus (1503): "quos denarios supputarios vocant," and in his commentary on Boethius (1510, fol. 33).

[7]Number penny. Rudolff, in his edition of 1534, uses this form, while Thierfelder, in his *Rechenbuch* of 1587, uses the form *Zahlpfenning*.

[8]In this anonymous and undated work, *De arte numerandi siue arismetice (perfectionis) summa quadripartita*, the author treats of the operations "per proiectiles," and says: "hec licet breuiter de proiectilibus sint dicta. negotianti tamen atque se exercenti per eos frequenter. abundantissime hec pauca sufficient." See Treutlein in the *Abhandlungen*, I, 24.

arithmetics on the subject are not so common as in Germany, they are sufficiently numerous to show that the system was well known.[1] The subject dropped out of the business textbooks in the last quarter of the 16th century, although the counter was used by women long after men had come to use algorism, writing not being so common among the former as among the latter.[2]

Because the counters are thrown upon the table the medieval Latin writers often called them *projectiles*.[3] The French translated this word, omitting the prefix, as *jetons*,[4] a word which still survives in France to mean a game counter, a small medal, or a token.

The older French jetons frequently bear such inscriptions as "pour les Comtes" and "pour les Finances,"[5] showing their use. Sometimes the legends are admonitory, thus: "Gectez, Entendez au Compte," "Gardez vous de Mescomptes," and "Jettez bien, que vous ne perdre Rien."[6]

The Tally and Related Forms. The subject of the abacus should not be dismissed without mention of the tally and certain other related forms. The tally was originally a piece of wood on which notches or scores[7] were cut to designate num-

[1] Among these books may be mentioned the following: Clichtoveus, *Ars supputādi tam per calculos q3 notas arithmeticas*, Paris, 1507; Clichtoveus, *De Mystica numerorum* (Paris, 1513, fol. 33, *r.*), subdividing *supputatio* into *calcularis* and *figuralis*, giving five common operations under the former and eight under the latter; Blasius, *Liber Arithmetice Practice* (1513); an anonymous *Le livre des Getz* (about 1500), in which is taught "la pratique de bien sçavoir conter aux getz comme à la plume"; Cathalan, *Arithmetiqve*, Lyons, 1555, in which the author explains "a Chiffrer & compter par la plume & par les gestz"; an anonymous *Arithmétique par les jects*, Paris, 1559; Trenchant, *Arithmetiqve*, 1566, the 1578 edition of which gives thirteen pages to "L'art et moyen de calcvler avec les Getons," the 1602 edition dropping the subject.

[2] Thus F. Legendre, in his arithmetic of 1729, says: "Cette manière de calculer est plus pratiquée par les femmes que par les hommes. Cependant plusieurs personnes qui sont employées dans les finances et dans toutes les jurisdictions s'en servent avec beaucoup de succès." [3] *Pro* (forward) + *jacere* (to throw).


[4] Also found in the following forms: *jettons, gects, gectz, getoers, getoirs, gettoirs, getteurs, jectoers, jectoirs, jetoirs, giets, gietons,* and *gitones*. Consult Snelling, *loc. cit.*, p. 2.

[5] Also "Getoirs de la chambre des comptes, Le Roi," "Ce sont les getoirs des ?tes [Comptes]. La Reinne."

[6] See also Snelling, *loc. cit.*, p. 3; Nagl, *loc. cit.*, p. 11; and Barnard, where many photographic plates may be consulted. [7] Scars; related to "shear."

bers. The word comes from the French *tailler* (to cut), whence our word "tailor." The root is also seen in the Italian word *intaglio* and is from the Latin *talea* (a slender stick).[1] The word has even been connected with the German *Zahl* (number) through the primitive root *tal.*[2]

The idea of keeping numerical records on a stick is very ancient, and in a bas-relief on the temple of Seti I (*c.* 1350 B.C.), at Abydos, Thot is represented as indicating by means of notches on a long frond of palm the duration of the reign of Pharaoh as decreed by the gods.[3]



TALLY STICKS OF 1296

Fragments found at Westminster in 1904. In the author's collection

In the Middle Ages the tally formed the standard means of keeping accounts. It was commonly split so as to allow each party to have a record, whence the expression "our accounts tally."[4] The root also appears in "tail," as when Piers Plowman, speaking of his gold, says that he "toke it by taille," meaning by count;[5] and in "tailage" (or "tallage") for toll or tax,[6]—a relic of the days when "our forefathers had no other books but the score and the tally."[7]

[1] Consult also Greenough and Kittredge, *Words and their Ways*, pp. 45, 266 (New York, 1901).

[2] G. Rosenhagen, "Was bedeutet Zahl ursprünglich?" *Zeitschrift für deutschen Altertum und deutsche Literatur*, LVII, 189.

[3] It is reproduced in G. Maspero, *Dawn of Civilization*, 3d ed. by Sayce, p. 221 (New York, 1897).

[4] Other expressions, like "keeping tally" at a game, "to tally up," and "stocks," are traced to this device.

[5] See also Chaucer, *General Prologue to the Canterbury Tales*, l. 570.

[6] The first poll tax is said to have been the "tailage of groats" levied by Parliament in 1377.

[7] Shakespeare, *2 Henry VI*, IV, vii, 38.

Usually a hazel stick was prepared by the "tally cutter," and the notches were cut before it was split, a large notch meaning M (£1000), a smaller one C (£100), a still smaller one X (£10), and so on down to pence. In earlier times the twentieth mark was a larger scar than the others, and the number was therefore called a score.[1] The system was used in the English Exchequer as late as 1812.

It is interesting also to note the relation of the tally stick to modern forms of investment. Formerly, if a man lent money to the Bank of England, the amount was cut on a tally stick.[2] This was then split, the bank keeping the "foil" (*folium*, leaf) and the lender receiving the "stock" (*stipes*), thereby becoming a "stock" holder and owning "bank stock."[3]

The tally was used in Germany for keeping accounts in the 13th and 14th centuries. Even at the beginning of the 15th century, and in as progressive a city as Frankfort am Main, the so-called *Kerbenrechnung* was common,[4] nor did the custom die out in Germany and Austria until the 19th century.[5]

[1] Teutonic *Stiege*, a word often used for twenty.

[2] The British Museum has a number of tally sticks, from 1348 to the "hop tally" still used in Kent and Worcestershire. There are six very perfect but not very old specimens in the Museum of Folklore at Antwerp. In cleaning the Chapel of the Pyx at Westminster in 1904 several specimens were found. These dated, as the inscriptions show, from 1296, and some of them came into the author's possession and are shown on page 193. Scotch "nick sticks" and Scandinavian calendar sticks belong to the same general class.

On the method of cutting and using tallies in England see the *Publications of the Pipe Roll Society*, Vol. III (London, 1884). On tally charts used by sailors, see *Zeitschrift für Ethnologie*, XXXV, 672 (Berlin, 1903). On their use as a means of communication see W. von Schulenberg, *Verhandlungen d. Berliner Gesellsch. für Anthropologie, Ethnol., und Urgeschichte*, XVIII, 384; *Zeitschrift für Ethnologie*, XIV, 370. On their other uses see Gyula v. Sebestyén, "Ursprung der Bustrophedonschrift," *Zeitschrift für Ethnologie*, XXXV, 755.

[3] Poole, *loc. cit.* See also C. H. Jenkinson, "Early wooden tallies," *Surrey Archæological Collection*, XXIII, 203.

[4] Günther, *Math. Unterrichts*, p. 287.

[5] F. Villicus, *Geschichte der Rechenkunst*, 3d ed., p. 15 (Vienna, 1897) (hereafter referred to as Villicus, *Geschichte*); R. Andree, *Braunschweiger Volkskunde*, p. 247 (Braunschweig, 1901), with several bibliographical notes of value; *Verhandlungen d. Berliner Gesellsch. für Anthropologie, Ethnol., und Urgeschichte*, XI, 763. With our "keeping tally" and baseball "scores" compare the German "Er hat viel auf dem Kerbholze." For interesting accounts of the use of the tally in Bohemia, see W. Wattenbach, *Das Schriftwesen im Mittelalter*, 3d ed., p. 95 (Leipzig, 1896).

In Italy the tally was evidently common in the 16th century, for Tartaglia (1556) gives a picture of one in his arithmetic, saying that one of the two parts was called by the Latin name *tessera*, a word often used to mean a counter.[1]

Knotted Cords. Related to the tally, in that they were used for recording numbers but not for purposes of calculation, are the knotted cords. These are used in various parts of the world and have such an extended history that only a passing reference can be given to them. Lao-tze, "the old philosopher," as the Chinese call him, in his *Tao-teh-king* of the 6th century B.C., referring to the earlier use of this device, says, "Let the people return to knotted cords (*chieh shing*) and use them."[2] Herodotus (IV, 98) tells us that the king of Persia handed the Ionians a thong with sixty knots as a calendar for two months, and a similar device of modern India may be seen in the museum at Madras. Indeed, in taking the census in India in 1872, the Santáls in the wilder parts of Santál Parganas used knots on four colors of cords, the black signifying an adult man, the red an adult woman, the white a boy, and the yellow a girl. The census was taken by the headmen, who, being unable to write, simply followed the popular method of keeping a numerical record.[3]

In the New World the knotted cord is best illustrated in the Peruvian *quipu*.[4] In each city of Peru there was, at the time

[1] "E laltro di questi dui pezzi lo chiamauano Tessera." *General Trattato*, I, fol. 3, *v*. (Venice, 1556).

[2] See Carus's English edition of the *Tao-teh-king*, pp. 137, 272, 323 (Chicago, 1898).

[3] *Proceedings of the Asiatic Society of Bengal*, p. 192 (Calcutta, 1872).

[4] The leading work on the general question of the *quipu*, with analyses of about fifty specimens, and with an extensive bibliography, is that of L. L. Locke, *The Quipu*, New York, 1923, published by the American Museum of Natural History. There is an article by the same author, entitled "The Ancient Quipu," in the *American Anthropologist* for 1912, p. 325. See also E. Clodd, *Storia dell' Alfabeto*, trad. del Nobili, cap. iii (Turin, 1903), or the English original; E. B. Tylor, *Early History of Mankind*, p. 160; *Westminster Review*, London, XI, 246; A. Treichel, *Verhandlungen d. Berliner Gesellsch. für Anthropologie, Ethnol., und Urgeschichte*, XVIII, 251; W. von Schulenberg, "Die Knotenzeichen der Müller," *Zeitschrift für Ethnologie*, XXIX, 491; H. G. Fegencz, "Kinderkunst und Kinderspiele," *Anzeiger d. Ethnolog. Abteilung d. Ungarischen National-Museums*, Budapest, XI, 103.

of the European invasion, a *quipucamayocuna* ("official of the knots") who may have performed duties not unlike those of a city treasurer today. At any rate, we have no evidence that the knots were used for any other purpose than the recording of numerical results, just as the Peruvian shepherd today uses them for keeping account of his herds.[1]



SPECIMEN OF QUIPU

The knotted cords of the ancient Peruvians

The knotted cords found in various forms of religious regalia may originally have recorded the number of prayers, pilgrimages, or sacrifices of the devotee. Examples of these are seen in the Lama rosary (*prenba*) and the rosaries of the Mohammedans, the Buddhists of Burma, and the Catholic Christians. Somewhat similar in use is the notched praying stick of the pilgrim, such as may be seen at the shrine of St. Fin Barr at Gouganebarra, Ireland.

2. Finger Reckoning


Finger Notation. The absence or rarity of suitable writing material led most early peoples to represent numbers by positions of the fingers,—a system not unlike the digital language of the deaf mutes of today. While this is manual rather than mechanical, it may properly be explained in this chapter. It is not improbable that the idea developed from the primitive method of counting on the fingers, usually beginning by point-

[1] They are also related to the wampum of the American Indian and possibly to the *lo-shu* and *ho-t'u* symbols of the ancient Chinese *I-king*.

ing at the little finger of the left hand with the second finger of the right, this being the result of holding the hands in a natural position for such a purpose. The person counting would thus proceed from right to left, and this may have influenced some of the early systems of writing numbers.[1]

The general purposes of digital notation were to aid in bargaining at the great international fairs with one whose language was not understood, to remember numbers in computing on an abacus, and to perform simple calculations.[2]

For the mere representing of the small numbers of everyday life the left hand sufficed. In this way it became the custom to represent numbers below 100 on the left hand and the hundreds on the right hand. Juvenal refers to this custom in his tenth satire, saying: "Happy is he indeed who has postponed the hour of his death so long and finally numbers his years upon his right hand."[3]



FINGER SYMBOLISM ABOUT THE YEAR 1140


One of a large number of drawings in a manuscript copy of Bede's works in the Biblioteca Nacional at Madrid, dating from *c.* 1140. The number 2000 is indicated. From a photograph by Professor J. M. Burnam

[1] On the general relation of the finger numbers to systems of counting and writing there is an extensive literature. See, for example, F. W. Eastlake, *The China Review,* Hongkong, IX, 251, 319, with a statement that the Chinese place the system before the time of Confucius; S. W. Koelle, "Etymology of the Turkish Numerals," *Journal of the Royal Asiat. Soc.*, London, XVI (N. S.), 141; Sir E. Clive Bayley, *ibid.*, XIV (reprint, part 2, p. 45 n.); M. Barbieri, *Notizie istoriche dei Mat. e Filosofi . . . di Napoli*, p. 10 (Naples, 1778); Villicus, *Geschichte*, p. 6; Bombelli, *Antica Numer.*, I, 108, 115 n., with bibliography and plates.

[2] P. Treutlein, *Abhandlungen*, I, 21; H. Stoy, *Zur Geschichte des Rechenunterrichts*, I. Theil, Diss., p. 31 (Jena, 1876).

[3] Felix nimirum, qui tot per saecula mortem
Distulit atque suos iam dextra computat annos.

That the system was familiar to the people is evident from a remark of Pliny[1] to the effect that King Numa dedicated a statue of two-faced Janus, the fingers being put in a position to indicate the number of days in a common year, and Macrobius testifies that the hundreds were indicated on the right hand.




FINGER SYMBOLISM IN THE 13TH CENTURY

From the *Codex Alcobatiensis* in the Biblioteca Nacional at Madrid, dating from *c.* 1200. From a photograph by Professor J. M. Burnam

The system was in use among the Greeks in the 5th century B.C., for Herodotus tells us that his countrymen knew of it. Among the Latin writers it is mentioned by Plautus, Seneca, Ovid, and various others.[2]

[1] See I. Sillig, edition of Pliny's works, V, 140 (Hamburg and Gotha, 1851), and the *Hist. Nat.*, XXXIV, vii, 16, 33.

[2] L. J. Richardson, "Digital Reckoning among the Ancients," *Amer. Math. Month.*, XXIII, 7; Bombelli, *Antica Numer.*, p. 102; A. Dragoni, *Sul Metodo aritmetico degli antichi Romani*, p. 10 (Cremona, 1811); Günther, *Math. Unterrichts*, p. 12. Possibly Juan Perez de Moya (1573) was correct in saying that the Egyptians used the system because they were "friends of few words,"—"los Egipcianos eran amigos de pocas palabras . . . destos deuio salir."



PACIOLI ON FINGER SYMBOLISM

From the *Süma* of Pacioli, Venice, 1494. The two columns at the left represent the left hand, the other two representing the right hand

Finger symbolism was evidently widely spread during many centuries, for there are also numerous references to it in both the Hebrew and the Arabic literature. Our precise knowledge of the subject is due chiefly, however, to a few writers,—to the Venerable Bede (*c.* 710), Nicholas Rhabdas (*c.* 1341), and a Bavarian writer, Aventinus (1522).[1] In the works of these writers the system is fully described, but brief summaries, often with illustrations, may be found in various books of the 16th century, including those of Andrés[2] (1515), Recorde[3] (*c.* 1542), Moya[4] (1562), Valerianus[5] (1556), and Noviomagus[6] (1539). The later literature of the subject is also extensive.[7]

The general scheme of number representations may be sufficiently understood from the illustrations given from the works of Pacioli and Aventinus. Bede gives a description of upwards of fifty finger symbols, the numbers extending through one million. No other such extended description has been given except the one of Rhabdas, but the works of Pacioli and Aventinus contain what are probably the best-known pictorial illustrations of the process.

[1] Bede, "De loquela per gestum digitorum," in his *Opera Omnia*, I, 686 (Paris, 1850); Nicholas Rhabdas, Ἔκφρασις τοῦ δακτυλικοῦ μέτρου; J. Aventinus, *Abacvs atqve vetvstissima, vetervm latinorum per digitos manusq3 numerandi . . . cōsuetudo*, Nürnberg, 1522 (title from Regensburg edition, 1532). See also St. Augustine, *Enarrationes in Psalmos*, xlix, 9, 1; *Sermones*, ccxlviii, ccxlix, cclii; and *Contra Iulianum*, iii, 11, 22; and M. Capella, *De Nuptiis Philologiae et Mercurii*, ii, 102, and vii, 729 and 746. On the Rhabdas symbolism see Heath, *History*, II, 551.

[2] Mossen Juan Andrés, *Sumario breve d' la prática de la Arithmetica*, Valencia, 1515.

[3] *Ground of Artes*, London, *c.* 1542.

[4] *Arithmetica Practica*, p. 627 (Salamanca, 1562).


[5] Joannes Pierius Valerianus Bellunensis, *Hieroglyphica*, p. 454 (Frankfort a. M., 1556; 1614 ed.).

[6] Cap. XIII of the 1544 edition of his *De nvmeris libri II*, Cologne.

[7] *E. g.*, see *Abhandlungen*, V, 91, 100; the Basel edition of St. Jerome's works, IX, 8 (1516); L. A. Muratori, *Anecdota*, Naples, 1776, with the "Liber de computo S. Cyrilli Alexandrini"; V. Requeno, *Scoperta della Chironomia*, Parma, 1797, with illustrations; M. Steinschneider, *Bibl. Math.*, X (2), 81; A. Marre, "Manière de compter des anciens avec les doigts . . .," Boncompagni's *Bullettino*, I, 309; Bombelli, *Antica Numer.*, cap. xiv, especially p. 109 n.; A. F. Pott, *Die quinäre und vigesimale Zählmethode bei Völkern aller Welttheile*, Halle, 1847, with an "Anhang über Fingernamen," p. 225; F. T. Elworthy, *The Evil Eye*, p. 237 (London, 1895), with illustrations; E. A. Bechtel, "Finger-Counting among the Romans in the Fourth Century," *Classical Philology*, IV, 25.

The representation of numbers below 100 was naturally more uniform, since they were in international use by the masses, while the representation of the higher numbers was not so well standardized.

Finger Computation. From finger notation there developed an extensive use of finger computation. This began, of course, with simple counting on the fingers, but it was extended to include particularly the simpler cases of multiplication needed



AVENTINUS ON FINGER SYMBOLS

From the *Abacvs* of Johannes Aventinus, Nürnberg, 1522 (Regensburg edition of 1532)

by the illiterate. For example, to multiply 7 by 8, raise two fingers on one hand and three on the other, since $5 + 2 = 7$ and $5 + 3 = 8$. Then add the numbers denoted by the raised fingers, $2 + 3 = 5$, and multiply those denoted by the others, $3 \cdot 2 = 6$, and the former result is the tens, 50, and the latter is the units, the product being 56. This depends, of course, upon the fact that $(10 - a)(10 - b) = 10\ (5 - a + 5 - b) + ab$.[1] The same principle is frequently seen in written work in arithmetic in the Middle Ages, since by its use it was unnecessary to learn the multiplication table above $5 \cdot 5$. In a somewhat

[1] See also pages 119 and 120.

similar way we may find the product of numbers from 10 to 15. For example, to find the product of 14 and 13, raise four fingers on one hand and three on the other, since $14 = 10 + 4$ and $13 = 10 + 3$. Then to 100 add ten times the sum of the number of fingers raised, and the product of the same numbers, the result being $100 + 10\,(4 + 3) + 4 \cdot 3 = 182$. The method is evidently general, since

$$(10 + a)\,(10 + b) = 100 + 10(a + b) + ab.$$

Such work is still to be seen in various parts of the world.

In the time of Fibonacci (1202) finger symbols were still used,[1] especially in remembering certain numbers in division.[2]

3. Modern Calculating Machines

Napier's Rods. The first important improvement on the ancient counter computation was made by Napier (1617). In his *Rabdologia*[3] he explains a system of rods arranged to represent the *gelosia* method of multiplication as seen in the illustration on page 203. The plan shows how crude were the methods of calculating even as late as the 17th century, although it would have had some value in connection with trigonometric functions if logarithms had not been invented. These rods were commonly known as Napier's Bones, as in Leybourn's *The Art of Numbring By Speaking-Rods: Vulgarly termed Nepeir's Bones*,[4] London, 1667. They attracted considerable attention, not merely in Europe but also in China and Japan.

[1] ". . . opportet eos qui arte abbaci uti uoluerint, ut subtiliores et ingeniores appareant scire computum per figuram manuum, secundum magistrorum abbaci usum antiquitus sapientissime inuentam." *Liber Abaci*, I, 5.

[2] So Fibonacci has a chapter, "De diuisione numerorum cordetenus in manibus per eosdem numeros," with such expressions as "ponens semper in manibus numeros ex diuisione exeuntes." *Ibid.*, I, 30.


[3] *Rabdologiae, Sev Nvmerationis Per Virgulas Libri Dvo*, Edinburgh, 1617; Leyden, 1626. Translations, Verona, 1623; Berlin, 1623. *Rabdologia* = late Greek ῥαβδολογία (*rhabdologi'a*), a collection of rods, from ῥάβδος (*rhab'dos*, rod) + λογία (*logi'a*, collection). Probably Napier took the word from the *Glossaria H. Stephani*, Paris, 1573, where the above meaning is given.

[4] W. Leybourn (*c.* 1670) derived *Rabdologia* from ῥάβδος (*rhab'dos*, rod) + λόγος (*log'os*, speech), and this etymology is still accepted by some writers.

Modern Machines. The essential superiority of the modern calculating machine over an instrument like the *suan-pan* is that the carrying of the tens is done mechanically instead of being done by the operator. For this purpose a disk is used which engages a second disk, turning the latter one unit after nine units have been turned on the former.[1]

The first of these instruments seems to have been suggested by a Jesuit named Johann Ciermans, in 1640,[2] but apparently nothing was done by him in the way of actually constructing such a machine.

The real invention may properly be attributed to Pascal (1642), who, at the age of nineteen and after many attempts, made an instrument of this kind, receiving (1649) a royal privilege for its manufacture,[3] and one particularly interesting specimen is still preserved in the Conservatoire National des Arts et Métiers at Paris. It is an adding machine adapted to numbers



NAPIER'S RODS IN JAPAN

The Napier Rods found their way into China at least as early as the beginning of the 18th century, and into Japan in the century following. This illustration is from Hanai Kenkichi's *Seisan Sokuchi*, of the middle of the 19th century

[1] For a succinct description of modern machines see F. J. W. Whipple, "Calculating Machines," in E. M. Horsburgh, *Handbook of the Exhibition at the Napier Tercentenary Celebration*, p. 69 (Edinburgh, 1914), hereafter referred to as Horsburgh, *Handbook*. For slide rules, *ibid.*, pp. 155 and 163.

[2] Kästner, *Geschichte*, III, 438; Cantor, *Geschichte*, II (1), 657.

[3] Cantor, *Geschichte*, II (1), 661; M. D'Ocagne, *Le Calcul simplifié par les procédés mécaniques et graphiques*, Paris, 1894; 2d ed., 1905; "Histoire des machines à calculer," *Bulletin de la Société d'Encouragement pour l'Industrie Nationale*, tome 132, p. 554, and other articles in the same number, with an extensive bibliography on pages 739–759.

of six figures and is one of the later attempts of Pascal. On the inside of the box is this inscription:

Esto probati instrumenti symbolum hoc; Blasius Pascal; arvernus, inventor. 20 mai 1652.[1]

In the same museum there are two other machines, apparently also of Pascal's make, one of which was verified and presented by a collateral descendant.

In 1673 Sir Samuel Morland (1625–1695), an English diplomat, mathematician, and inventor, made a machine for multiplying, and about the same time (1671) Leibniz constructed one in Germany. In 1709 the Marchese Giovanni Poleni (1683–1761), then professor of astronomy at Padua, made a similar attempt in Italy; and in 1727 there was described in Germany a machine constructed just before his death by Jacob Leupold (1674–1727), a Leipzig mechanic. These various attempts were recorded in 1735[2] by Christian Ludwig Gersten (1701–1762), then professor of mathematics at Giessen, in connection with a description of a machine invented by himself. It was not, however, until the 19th century that any great advance was made. In 1820 Charles Babbage began the construction of a machine for calculating mathematical tables, and in 1823 the Royal Society secured aid from the British government to enable him to continue his work. Babbage's progress not being satisfactory, this aid was soon withdrawn, but the work continued until 1856, when it was abandoned.[3] From the time when Babbage began to the present, however, the modern calculating machine has been constantly improved, first by Thomas de Colmar (1820), and various types are now in extensive use.[4]

[1] "Let this signature be the sign of an approved instrument. Blaise Pascal, of Auvergne, inventor. May 20, 1652."

[2] *Phil. Trans.*, Abridgment, 1747, VIII, 16.

[3] One of the best descriptions of this machine is given in *Babbage's Calculating Machine; or Difference Engine*, printed by the Victoria and Albert Museum, London, 1872; reprinted in 1907.

[4] There is a large collection in the Conservatoire National des Arts et Métiers at Paris.

Slide Rule. In 1620 Edmund Gunter designed the logarithmic "line of numbers," on which the distances were proportional to the logarithms of the numbers indicated. This was known as Gunter's Scale, and by adding or subtracting distances by the aid of compasses it was possible to perform multiplications and divisions. Thus the inventor worked out the principle of the slide rule, but instead of having the sliding attachment he used a pair of compasses.[1] This instrument was subsequently used in navigation.

In 1628 Edmund Wingate published at London his *Construction and Use of the Line of Proportion*, but this, like Gunter's Scale, was merely a rule in which the spaces on one side indicated numbers, while those on the other indicated the mantissas of these numbers.

About 1622 William Oughtred invented the slide rule,[2] but descriptions of his instrument did not appear in print until 1632. A pupil of Oughtred's, Richard Delamain, published at London in 1630 a small pamphlet entitled *Grammelogia; or the Mathematicall Ring*, in which he described a circular slide rule, apparently of his own invention. Oughtred, however, seems unquestionably to have invented the rectilinear logarithmic slide rule, and also, independently of Delamain, to have invented a circular one.

In the year 1654 a slide rule was made in which the slide moved between parts of a rigid stock, and a specimen of this type, now in the Science Museum at South Kensington, is inscribed "Made by Robert Bissaker, 1654, for T. W."[3] Who this T. W. was we do not know, but the invention was a notable step in the development of the modern type.

From that time on there were numerous inventors who improved upon the instrument. Among them were various obscure artisans, but there was also Newton, who devised a system of

[1] On this entire topic see F. Cajori, *A History of the Logarithmic Slide Rule*, New York, 1909; hereafter referred to as Cajori, *Slide Rule*.

[2] F. Cajori, *William Oughtred*, p. 47 (Chicago, 1916); hereafter referred to as Cajori, *Oughtred*.

[3] Horsburgh, *Handbook*, p. 163.

concentric circles for the solution of equations. In the 17th century the slide rule of the type now used attracted little attention, either in England or on the continent. In the following century, however, its value began to be recognized, and the instruments in use at that time resemble in several particulars those with which we are familiar. About 1748 George Adams made spiral slide rules that were carefully engraved and probably were of a higher degree of accuracy than those of his predecessors.

The first one to make a decided step in advance, however, was William Nicholson (1753–1815). He described (1787) the various types of rules then known, and suggested noteworthy improvements, particularly in the way of a rule which, through the device of a system of parallels, gave the effect of an instrument more than 20 feet in length. He also designed a spiral slide rule, apparently ignorant of the work done in this field by various predecessors. At about the same time various French and German writers contributed to the perfecting of the instrument, notably Jean Baptiste Clairaut (1720), who designed a new circular slide rule.

The most marked advance in the middle of the 19th century was made by Amédée Mannheim[1] (1831–1906), who (*c.* 1850) designed the Mannheim Slide Rule, which is still a standard, although modified in various particulars.[2] These modifications related (1) to increasing the length of the scales without increasing the size of the instrument; (2) to adapting the rule to specialized branches of science; and (3) to increasing the mechanical efficiency of the device.[3] Few such instruments have gained so much popularity in such a short time.

[1] *L'Enseignement Mathématique*, IX (1907), 169.

[2] For details as to other inventors see Cajori, *Slide Rule*. On the history of the planimeter, which may be classified among instruments relating to the calculus or among those having to do with calculation, see A. Favaro, "Beiträge zur Geschichte der Planimeter," Separat-Abdruck aus der *Allgemeinen Bauzeitung*, Vienna, 1873. The instrument seems to have been first designed *c.* 1814 by J. M. Hermann, but it attracted little attention. The first published description was that of an Italian inventor, Gonella; it appeared in 1825.

[3] Horsburgh, *Handbook*, p. 156.

TOPICS FOR DISCUSSION

1. Consider the difficulties of multiplying a number like 4275 by a number like 876, using only the Roman, Greek, Egyptian, or Babylonian numerals.

2. Reason for the persistence of the abacus in business calculations until the 17th century.

3. Reason for abandoning the abacus in Italy before it was abandoned in northern Europe.

4. The various etymologies of the term "abacus" as given in such dictionaries as are accessible.

5. Reason for believing that the origin of the line abacus may be Semitic.

6. Etymologies of the terms used in connection with the abacus, and the relation of the word "calculus" to other words in our language.

7. Words used in various languages to mean computing disks, with their etymologies.

8. Comparison of the various types of line abacus, with a discussion of their respective merits.

9. The various forms of the abacus used in the Far East, with a comparison of their merits.

10. A study of the evolution of the paper tablet used for computation by pupils at the present time, beginning possibly with the dust table or the wax tablet.

11. Gerbert's abacus and its chief defects.

12. History of the British Court of the Exchequer.

13. General use of counters in the countries of Western Europe.

14. History of the tally.

15. The history of finger symbols and finger computation, with special reference to the international character of the symbols themselves.

16. The general character of the quipu and of similar knot-tying devices in various parts of the world.

17. Relation of the "cat's cradle" to the knotted cords and possibly to the tying together of the stars to make the constellations of ancient astronomy.

18. Rise of the modern calculating machine.

19. Types and history of modern calculating machines and planimeters as described in current encyclopedias.

CHAPTER IV

ARTIFICIAL NUMBERS

1. Common Fractions

Origin of Artificial Numbers. The natural numbers seem to have served the purposes of the world until about the beginning of the historic period. Men broke articles and spoke of the broken parts, but even after weights came into use it was not the custom to speak of such a fraction as $\frac{3}{4}$ of a pound. The world avoided difficulties of this kind by creating such smaller units as the ounce and then speaking of the particular number of ounces. For example, the commercial fractions of Rome were referred to the *as*,[1] 16 *asses* making a *denarius*.[2] A twelfth part of the *as* was the *uncia*, whence the modern "ounce" and "inch." Hence the Romans used this scheme:

Multiples of the as

1 *as* $= \frac{1}{16}$ *denarius* $= \frac{1}{2}\frac{1}{4} + \frac{1}{4}\frac{1}{8} =$ *Denarii semuncia sicilicus*
2 *asses* $= \frac{1}{8}$ *denarius* $= \frac{1}{12} + \frac{1}{2}\frac{1}{4} =$ *Denarii uncia semuncia*
3 *asses* $= \frac{3}{16}$ *denarius* $= \frac{1}{6} + \frac{1}{4}\frac{1}{8} =$ *Denarii sextans sicilicus*
. . . . . .
15 *asses* $= \frac{15}{16}$ *denarius* $= \frac{11}{12} + \frac{1}{4}\frac{1}{8} =$ *Denarii deunx sicilicus*

Submultiples of the as

$\frac{11}{12}$, *deunx, i.e.*, $1 - \frac{1}{12}$, *de uncia*, $\frac{1}{12}$ taken away. The symbol is $S{=}{=}-$, meaning *semis* + $\frac{5}{12}$.

$\frac{10}{12}$, *dextans, i.e.*, $1 - \frac{1}{6}$, *de sextans*, $\frac{1}{6}$ taken away. The symbol is $S{=}{=}$, *semis* + $\frac{4}{12}$.

[1] Originally a pound of copper, but reduced by successive depreciations of coin until (191 B.C.) it weighed half an ounce.

[2] Originally a coin of 10 asses, but later of 16 asses, about 16 American cents.

$\frac{9}{12}$, *dodrans, i.e.,* $1 - \frac{3}{12}$, *de quadrans,* $\frac{1}{4}$ taken away. The symbol is $S=-$, *semis* $+ \frac{3}{12}$.

$\frac{8}{12}$, *bes, i.e., bi as* for *duae partes,* $\frac{2}{3}$. The symbol is $S=$, *semis* $+ \frac{2}{12}$.

$\frac{7}{12}$, *septunx, i.e., septem unciae.* The symbol is $S-$, *semis* $+ \frac{1}{12}$.

$\frac{6}{12}$, *semis,* half. The symbol is S, Σ, or (.

$\frac{5}{12}$, *quincunx, i.e., quinque unciae.* The symbol is $\equiv-$.

$\frac{4}{12}$, *triens,* one third. The symbol is $\equiv$.

$\frac{3}{12}$, *quadrans,* one fourth. The symbol is $=-$.

$\frac{2}{12}$, *sextans,* one sixth. The symbol is $=$.

$\frac{1}{12}$, *uncia,* ounce, inch. The symbol is $-$.

There were similar special names and symbols for $\frac{1}{24}$ (*semuncia,* Σ, ( ), $\frac{1}{48}$ (*sicilicus,* ↄ), $\frac{1}{72}$, $\frac{1}{144}$, $\frac{1}{288}$ (*scriptulum, scripulum, scrupulum,* ϑ, surviving in our "scruple"), and other fractions.[1]

It will be seen that the Roman merchant could speak of $\frac{3}{8}$ of a *denarius* as 6 *asses,* of $\frac{1}{24}$ of the *as* as a *semuncia,* and so on, without considering fractions at all, and this was the case with all ancient peoples. In fact, the origin of such compound numbers as 3 yd. 2 ft. 8 in. is to be sought in the effort of the world to avoid the use of fractions.

Gradually, however, the notion of a unit fraction developed; then came the idea of a general fraction; then the surd appeared; and so on through various types of fractions, irrational numbers, transcendental numbers, complex numbers, and other kinds of artificial numbers. Each was created to satisfy an intellectual need, and in due time each, excepting the latest creations, has satisfied important practical needs as well.

First Steps in Fractions. The first satisfactory treatment of fractions as such is found in the Ahmes Papyrus (*c.* 1550 B.C.).[2]

[1] For a brief discussion of Roman fractions, with bibliography, see Pauly-Wissowa, *Real-Encyclopädie,* II, 1114 (Stuttgart, 1896); hereafter referred to as Pauly-Wissowa. See also Ch. Daremberg and E. Saglio, *Dictionnaire des Antiquités Grecques et Romaines,* Paris, 1877. Of the early printed works on the subject the classical one is G. Budé (or Budaeus), *De asse et partibus ejus, Libri V,* Paris, 1516, with several later editions.

[2] F. Hultsch, *Die Elemente der ägyptischen Theilungsrechnung,* reprint from the *Abhandl. d. k. Sächs. Gesellsch. d. Wissensch.,* Bd. XXXIX; Eisenlohr, *Ahmes Papyrus*; Peet, *Rhind Papyrus,* where the date is put somewhat earlier.

Artificial numbers of this kind had already been used by the Babylonians,[1] but we have no noteworthy treatment of fractions prior to the work of Ahmes. The notion of the unit fraction was already old in Egypt, however, for the tables given by Ahmes bear evidence of a development through a long period.

The essential feature of the early Egyptian treatment is the unit fraction. The arithmeticians had long been able to conceive of $\frac{1}{10}$,[2] but they had no plural for it either verbally, or mentally. By the time of Ahmes, however, an idea akin to that of ratio had developed. The number 2 was divided, say into 43 equal parts, and what is essentially the ratio of 2 to 43, or twice $\frac{1}{43}$, was expressed, using modern symbols, as

$$2 : 43 = \tfrac{1}{42} + \tfrac{1}{86} + \tfrac{1}{129} + \tfrac{1}{301}.$$

Indeed, most of the ancient theory of fractions centered about the concept of ratio, and in such theoretical works as that of Boethius it lasted until the 16th century.

In the Ahmes Papyrus the fraction $\frac{1}{42}$, for example, is written  , where the dot is the unit-fraction symbol,  is the Ahmes hieratic symbol for 40, and  is used to denote 2.[3] It is a curious fact that the dot is occasionally found in modern times as a fraction symbol, as in the case of $\frac{\cdot}{2}$ and $\frac{\cdot}{4}$ for $\frac{1}{2}$ and $\frac{1}{4}$ in English copy-books of the 18th century.

How these unit fractions were derived we do not know. It is evident that more than one solution is possible, but it is not always evident why any given one should be preferred to any other. For example,

$$\begin{aligned} 2 : 43 &= \tfrac{1}{24} + \tfrac{1}{258} + \tfrac{1}{1032} \\ &= \tfrac{1}{30} + \tfrac{1}{86} + \tfrac{1}{645} \\ &= \tfrac{1}{36} + \tfrac{1}{86} + \tfrac{1}{645} + \tfrac{1}{1720} + \tfrac{1}{7744} \\ &= \tfrac{1}{40} + \tfrac{1}{860} + \tfrac{1}{1720} \\ &= \tfrac{1}{42} + \tfrac{1}{86} + \tfrac{1}{129} + \tfrac{1}{301},^{4} \end{aligned}$$

[1] D. E. Smith, "The Mathematical Tablets of Nippur," in the *Bulletin of the Amer. Math. Soc.*, XIII (2), 392; H. F. Lutz, "A Mathematical Cuneiform Tablet," *American Journal of Semitic Languages*, XXXVI, 240.

[2] *Re-met*, "mouth of ten." Erman, *Egypt*, p. 365. Compare the Hebrew *pe-esr*.

[3] For the complete work in facsimile, see the British Museum edition.

[4] Eisenlohr, *Ahmes Papyrus*, p. 12; Peet, *Rhind Papyrus*, p. 42.

and so on, to which, of course, may be added $\frac{1}{43} + \frac{1}{43}$. Of all these possibilities Ahmes and his predecessors took the form

$$2 : 43 = \frac{1}{42} + \frac{1}{86} + \frac{1}{129} + \frac{1}{301},$$

although $\frac{1}{24} + \frac{1}{258} + \frac{1}{1032}$ has the advantage that the first fraction is nearer the value of $\frac{2}{43}$ than it is in the others. Although there are numerous rules for forming the unit fractions, no one of them applies to all the cases. This shows that the treatise combined the results of earlier computers, each working by a secret rule of his own, or else that each solution was worked out laboriously by repeated trials.[1]

The Egyptians indicated a unit fraction by a fraction symbol with the denomination underneath. In hieroglyphics this symbol was  , but in the cursive hieratic writing it was merely a dot. Thus, $\frac{1}{5}$, $\frac{1}{10}$, and $\frac{1}{30}$ appear respectively as  , and $\frac{1}{42}$ appears as either  or  in hieroglyphic, but in the hieratic it appears as shown on page 210. For $\frac{1}{2}$, $\frac{1}{3}$, $\frac{1}{4}$, and $\frac{2}{3}$ there were special symbols, this having been rendered necessary by the frequent use of these fractions. Thus the symbol for $\frac{2}{3}$ was  .

The symbol  was also used with a different meaning. For example, in the Archeological Museum at Florence there is a marble cubit divided into parts marked with such characters as  ···  , representing 3, 4, 5, · · · 16 fractional parts, not $\frac{1}{3}$, $\frac{1}{4}$, $\frac{1}{5}$, · · · $\frac{1}{16}$. In the Louvre there is a similar measure made of wood with the symbols  ···  , ···  . It should be said, however, that the first of these symbols may be looked upon as meaning $\frac{1}{3}$ if we consider it as applying to $\frac{1}{3}$ of the subdivision of the cubit, say to $\frac{1}{3}$ of an inch, and similarly for the other fractions.

[1] *E. g.*, when $b + c = ka$, we have

$$\frac{a}{bc} = \frac{1}{b \cdot \frac{b+c}{a}} + \frac{1}{c \cdot \frac{b+c}{a}},$$

and this gives the Ahmes result in certain cases but not in others. Thus, $\frac{2}{15} = \frac{2}{3 \cdot 5}$, and $3 + 5 = 4 \cdot 2$. This fraction, therefore, is equal to $\frac{1}{12} + \frac{1}{20}$, but Ahmes gives $\frac{1}{10} + \frac{1}{30}$. See Eisenlohr, *Ahmes Papyrus*, p. 28; G. Loria, *Bibl. Math.*, VI (2), 97; VII (2), 84; Peet, *Rhind Papyrus*, p. 34.

Later Development of Unit Fractions. The separation of a fraction into partial fractions is an illustration of the force of tradition. The predecessors of Ahmes decomposed their quotients in this way, and so Ahmes did the same. Although the Greeks had meanwhile developed a fairly good system of fractions, Heron (*c.* 50?) followed the Egyptian tradition, adopting the standard set by Ahmes nearly two thousand years earlier.[1] Some six or seven centuries later, so the Akhmim Papyrus (*c.* 8th century) informs us, the identical method of Ahmes was still in vogue in the temple schools of Egypt. Even as late as the 10th century Rabbi Saʿadia ben Joseph al-Fayyumî[2] (died 941), a Hebrew writer living in Egypt, made much use of unit fractions in his computations relating to the division of inheritances.

Not all tables of fractions made by the Egyptians followed precisely the Ahmes type, as may be seen in one dating from about the 4th century and recently acquired by the University of Michigan.[3] This table gives the unit fractional parts up to tenths of the units from 1 to 9, of the tens from 10 to 90, of the hundreds to 900, and of the thousands to 9000. It then gives the elevenths, twelfths, and so on to the seventeenths of the units up to 11, 12, and so on to 17 respectively. For example, $\frac{1}{9}$ of 50 is given as $5\frac{1}{2}\ \frac{1}{18}$, and $\frac{1}{11}$ of 9 as $\frac{1}{2}\ \frac{1}{4}\ \frac{1}{22}\ \frac{1}{44}$.

Upwards of two centuries after Rabbi Saʿadia, Fibonacci gave a rule for separating fractions into partial fractions,[4] of which the separation into unit fractions is a special case. Until recently our textbooks in algebra have given similar directions, although the subject had no immediate application that the pupil could then understand.

In the Middle Ages unit fractions were sometimes called "simple fractions," the more general form being known as "composite fractions."[5] These "simple fractions" were not

[1] Professor Loria has called attention to the fact that Heron was not very accurate about it, for he gives $\frac{3}{5}\ \frac{1}{7}\ \frac{1}{5}$ as $\frac{1}{2} + \frac{1}{8} + \frac{1}{73}$, while the $\frac{1}{73}$ should be $1 : (73 + \frac{1}{7})$. *Bibl. Math.*, VII (2), 88. [2] *Traité des successions*, ed. Joël Muller. Paris, 1897.

[3] L. C. Karpinski, "Michigan Mathematical Papyrus, No. 621," in *Isis*, vol. iv.

[4] He called it a "regula uniuersalis in disgregatione partium numerorum." See *Liber Abaci*, p. 82.

[5] So the Rollandus MS. (*c.* 1424) says: "sicut $\frac{1}{2} \cdot \frac{1}{3} \cdot \frac{1}{4}$ que simplices fractões dicũt^{e}, sic $\frac{2}{3} \cdot \frac{3}{5} \cdot \frac{4}{5}$ que cõposite siue pregnãtes dicunt^{e}."

infrequently favored even by Renaissance mathematicians of some prominence, Buteo, for example, giving $1350534\frac{1}{2}\frac{1}{64}$ as the square of $1162\frac{1}{8}$,[1] even though he knew the other forms. As late as the 17th century Russian manuscripts on surveying speak of a "half-half-half-half-half-third" of a certain measure instead of $\frac{1}{96}$ of the measure,[2] and even today the unit fraction is used to some extent in the diamond trade in speaking of parts of a carat.

The difficulty met in early times in solving problems involving fractions is illustrated by an example from Ahmes: "A number together with its fifth makes 21 [; find the number]." Our solution would be $\frac{6}{5}x = 21$, whence $x = \frac{5}{6} \times 21 = 17\frac{1}{2}$; but Ahmes went through substantially this process: Multiplying 1 and $\frac{1}{5}$ by 5, we have 5 and 1, which make 6, and this is too small. To find how many times too small we divide 21 by 6, the result being $\frac{1}{2}$ and 3. Multiplying 5 by this result, the answer is $17\frac{1}{2}$.[3]

Development of the General Fraction. It seems probable that, except in very simple cases, the idea of a fraction with numerator greater than unity arose in Babylon. Although the unit fraction and possibly some idea of the sexagesimal fraction appear in the cuneiform records of *c.* 2000 B.C., the fraction forms also include special symbols for $\frac{2}{3}$, $\frac{2}{18}$, $\frac{4}{18}$, $\frac{5}{6}$, and other cases of a like degree of difficulty. No such elaborate treatment of the subject as that given by Ahmes, however, has been found as yet among the Babylonian remains.[4] In spite of this early use of the general fraction, our present forms are not due to Babylonian influences, at least not directly, but apparently to the Hindu arithmeticians.[5]

[1] *Ioan Bvteonis De Qvadratvra circuli Libri duo*, p. 39. Lyons, 1559.

[2] V. V. Bobynin, "Quelques mots sur l'histoire des connaissances mathématiques," *Bibl. Math.*, III (2), 104.

[3] Peet, *Rhind Papyrus*, p. 62, No. 27.

[4] For examples of the non-unit fractions among the Babylonians see Contenau, *loc. cit.* (cited on page 37), plates 3, 35, and 100; Peet, *Rhind Papyrus*, p. 28.

[5] R. C. Dutt, *History of Civilization in Ancient India*, I, 273 (London, 1893); hereafter referred to as Dutt, *History Civ. in Anc. India*. See also V. V. Bobynin, "Esquisse de l'histoire du calcul fractionnaire," *Bibl. Math.*, X (2), 100.

Greek Fractions. The Greeks followed the ancient plan of avoiding, by the use of submultiples, the difficulty of computing with fractions; but in due time the need for a fraction symbolism became so apparent that they developed a system that served their purposes fairly well. They designated $\frac{1}{3}$ (*τρίτον, tri'ton*) by the symbol $\overset{\circ\nu}{\Gamma}$, the Γ being the symbol for three. This was further abbreviated to $\overset{\prime\prime}{\Gamma}$. Similarly, for $\frac{1}{4}$ they used Δ (for four) with two accent marks, thus: $''\Delta$. In the same way they accented their other numerals, a method represented in modern typography by γ'', δ'', ϵ'', $\cdots$.

The more common fractional unit, one half (*ἥμισυ, he'misy*), had a special symbol, (, which was often written in a form resembling the Greek Σ or the Latin S. Two thirds (*δίμοιρον, di'moiron*) had various abbreviations, such as ς'', that is, $\frac{1}{2}+\frac{1}{6}$.[1]

Aristarchus (*c.* 260 B.C.) wrote the word for the numerator and the numeral for the denominator, as we might write "ten 71sts."[2] Various methods were afterwards used, such as writing the numeral for each term but doubling it for the denominator, as in the case of $2'5''5''$ ($\beta'\epsilon''\epsilon''$) for $\frac{2}{5}$; or writing the numerator, then the words "in part,"[3] and finally the denominator, as in the case of "3,069,000 in part 331,776," for $\frac{3\,069\,000}{331\,776}$.[4] Heron (*c.* 50?) and Diophantus (*c.* 275) used a symbol that naturally seems strange to the modern reader, namely, our common fraction reversed; that is, they wrote the equivalent of 1^{9}_{4} or 1^{9}_{4} for four nineteenths,[5] and similarly in other cases. Ordinarily, however, the unit fraction was preferred, $\frac{15}{16}$ being written as $\frac{1}{2}+\frac{1}{4}+\frac{1}{8}+\frac{1}{16}$.[6]

Roman Fractions. As already stated, the Romans, like their predecessors, avoided fractions to a great extent by the device

[1] For bibliographical references and for this general topic see Pauly-Wissowa, II, 1077; Heath, *History*, I, 42.

[2] *Δέκα οα''*. All the Greek symbols used hereafter in this section are modern.

[3] *Ἐν μορίῳ*, from *μόριον*, a piece, portion, or section, much as we should say "divided by."

[4] In modern Greek symbols, $\overline{\tau\varsigma}$. $\overline{\theta}$ μορ. $\overline{\lambda\chi\gamma}$. $\overline{\alpha\psi o\varsigma}$.

[5] *I.e.*, $\frac{\iota\theta}{\delta}(1^{9}_{4})$, or possibly $\frac{\iota\theta}{\delta}(1^{9}_{4})$, for 1^{4}_{9}.

[6] *I.e.*, $(\delta''\eta''\iota\varsigma''$. For a further discussion, see Pauly-Wissowa and Heath.

of compound numbers, although using a few convenient symbols. Even such names as *semuncia* (half-twelfth) were not numerous. Marcus Terentius Varro[1] (116–28 B.C.) mentions twelve such fractions, and Volusius Maecianus[2] (2d century) gives only two more. Of the later Latin writers, Isidorus (*c.* 610) mentions only eight and Papias (11th century) has eighteen.[3] Adelard of Bath (*c.* 1120) mentions twenty-four.

Chinese Fractions. The Chinese seem to have made use of fractions of considerable difficulty at a very early date.[4] The *Chóu-peï*, probably of about 1105 B.C. but possibly much earlier, has various problems involving such numbers as $247\frac{93}{1460}$, not stated, however, in numerical symbols but given in words. The work includes such divisions as that of 119,000 by $182\frac{5}{8}$, both of these expressions being multiplied by 8 before dividing. The unit fraction also entered into their work, as it did in all earlier civilizations. For example, in the *Nine Sections*, a work of very uncertain date but probably of the second millennium B.C.,[5] there is given the problem:

> There is a field whose length is one *pu* and a half, one-third *pu*, one-fourth *pu*, and one-fifth *pu*. If the area is 240 square *pu*, what is its breadth?

Present Writing of Common Fractions. It is probable that our method of writing common fractions is due essentially to the Hindus, although they did not use the bar. Brahmagupta (*c.* 628) and Bhāskara (*c.* 1150), for example, wrote $\frac{2}{3}$ for $\frac{2}{3}$.[6] The Arabs introduced the bar, but it was not used by all their writers, and when Rabbi ben Ezra (*c.* 1140) adopted the Moorish forms he generally omitted it.[7] It is ordinarily found

[1] *De Lingua Latina*, 1st ed. s.l.a., but Rome. Hain mentions six editions s.l.a. before 1501, and one dated 1474 and another 1498.

[2] *Assis Distributio*, 1st ed., Paris, 1565.

[3] *Vocabularium*, 1st ed., Milan, 1476. For a full discussion of these fractions see Boncompagni's *Bullettino*, XIV, 71, 109. Not the Papias of the 2d century.

[4] Y. Mikami, "Arithmetic with Fractions in Old China," *Archiv for Mathematik og Naturvidenskab*, Christiania, XXXII, No. 3.

[5] See Volume I, page 32.

[6] Taylor, *Lilawati*, Introd., p. 12; text, p. 24 n.; Villicus, *Geschichte*, p. 54.

[7] Silberberg, *Sefer ha-Mispar*, p. 104.

in the Latin manuscripts of the late Middle Ages, but when printing was introduced it was frequently omitted, doubtless owing to typographical difficulties. This inference is confirmed by such books as Rudolff's *Kunstliche rechnung* (1526), where the bar is omitted in all ordinary fractions like $\frac{2}{3}$ and $_1^8{}_2$ but is inserted in all fractions printed in larger type and in those having large numbers.[1] The same inference is drawn from his *Exempel-Büchlin* (1530), $\frac{1}{2}$ having the bar because that fraction was in the font, and the other fractions not having it because of the necessity for piecing them up. One of the interesting evidences of the troubles of early printers is seen in Ciacchi's *Regole generali d' abbaco* (Florence, 1675), where, in order to secure better alignment, every fraction in the book is set up like $\frac{2}{3}$, for $\frac{2}{3}$. The difficulties of the early printers probably account also for such forms as "Z3& septe octaui" for $23\frac{7}{8}$, and "Z3&V octaui" for $23\frac{5}{8}$, in Chiarino's work of 1481.

The omission of the bar was not, however, entirely a matter of typography. Hylles (1592), for example, omitted it after the first fraction in a case like $\frac{1}{2}$ of $\frac{2}{3}$ of $\frac{3}{4}$, writing this expression $\frac{1}{2} \cdot \frac{2}{3} \cdot \frac{3}{4}$ and saying:

> And here you see the first fractions to wit $\frac{1}{2}$ being a true fraction, written with his lyne as it ought to be. and the other two that is to say $\frac{2}{3}$ and $\frac{3}{4}$ to be written without any lyne as their vse and order is.[2]

Recorde (*c.* 1542) tells us that "some . . . expresse them thus in slope forme,"[3] as here shown: $\frac{\frac{3}{4}}{\frac{2}{3}}$ $\frac{1}{2}$

The common use of 2/3 for $\frac{2}{3}$ is the result of a desire to simplify written and printed forms.[4]


[1] Edition of 1534 examined.

[2] Fol. 11, *v*. Even as good a writer as Paolo Casati, *Fabrica et vso del compasso di proportione*, Bologna, 2d ed., 1685, however, omits the bar entirely.

[3] *Ground of Artes*, 1558 ed., fol. R iij, *v*.

[4] The questionable statement that 2/3 comes from 2 f 3, the f meaning *fratto* (fraction), is made by G. Frizzo, *Le Regoluzze del M. Paolo dell' Abbaco*, Bologna, 1857; 2d ed., enlarged, Verona, 1883, p. 45; but the manuscript was first published in G. Libri, *Histoire des Mathématiques*, III, 295.

Since the bar is an Oriental device, it was never used by the Greeks or Romans to indicate a fraction, at least in the way that we use it today. In Renaissance times, however, when Arab devices mingled with classical forms, we find the Roman numerals occasionally used[1] in cases like $\frac{IX}{XI}$, and the Greek numerals employed[2] in a similar manner.


I Dieſſe figur iſt vñ bedeūt ain fiertel von ainez
IIII ganzen/alſo mag man auch ain fünfftail/aym
ſechſtail/ain ſybentail oder zwai ſechſtail ꝛc. vnd alle
ander brüch beſchreiben/Als I/V | I/VI | I/VII | II/VI ꝛc.

VI Diß ſein Sechs achtail/das ſein ſechstail der
VIII acht ain ganz machen.

IX Diß Figur bezaigt ann newn ayilfftail das ſeyn
XI IX tail/der XI ain ganz machen.

XX Diß Figur bezaichet/zwenzigk ainundrey-
XXXI ſigk tail/das ſein zwenzigk tail der ains-
undreiſſigk ain ganz machen.

IIC Diß ſein zwaihundert tail/der Fierhun-
IIIIC.LX dert vnd ſechzigk ain ganz machen.

KÖBEL'S USE OF COMMON FRACTIONS

From Köbel's *Rechen biechlin* (1514; 1518 ed.), showing the attempt to use Roman numerals with common fractions

The Name "Fraction." The word "fraction" is from the Latin *frangere* (to break). It is a broken number and was often so called. Baker (1568), for example, speaks of "fractions or broken numbers," calling a fraction of a fraction a "broken of broken," and various other English writers did the same. The word "fragment" is from the same root and was not infrequently used for "fraction."[3]

[1] Köbel, *Ain New geordnet Rechen biechlin* (1514; 1518 ed.), fol. xxxiii, *r*.
[2] V. Strigelius (Strigel), *Arithmeticus Libellvs*, Leipzig, 1563.
[3] Thus, in the Italian edition (1586) of Clavius (p. 75) the word appears as *fragmēto*. The idea goes back to the Egyptians. See Peet, *Rhind Papyrus*, p. 15.

The use of this root has not, however, been universal. Boethius (*c.* 510) does not speak of fractions as such in his arithmetic, introducing instead an elaborate system of ratios; but in the geometry attributed to him there is a chapter *De Minutiis*,[1] so that if he spoke of fractions at all, other than as ratios, he called them minutes, and in this he was followed by various medieval writers.[2] In the 12th century, for example, Adelard of Bath used *minuciae*,[3] while about the same time Johannes Hispalensis preferred *fractiones*.[4] In the translation of al-Khowârizmî attributed to Adelard, however, *fraciones* is used.[5] There are many instances in the early printed books of the use of the two terms interchangeably, each signifying a common fraction.[6] Several reputable writers used "parts" as a synonym of "fractions."[7] In English the word "fraction" appeared early,[8] however, and has been the general favorite.

Since *ruptus*, like *fractus*, means broken, this has been the root of a name for fraction. In Italian it appears as *rotto* (plural, *rotti*),[9] in Spanish as *rocto*,[10] and in French in various forms.[11]

[1] Friedlein ed., pp. v, 425.

[2] We shall see that the term was also applied specifically to sexagesimals, although by no means generally.

[3] In his *Regulae abaci*. See Boncompagni's *Bullettino*, XIV, 109.

[4] In his *Liber Algorismi de pratica arismetrice*. See Boncompagni, *Trattati*, II, 49; *Abhandlungen*, III, 111. Fibonacci (1202) generally used *fractio*.

[5] For *fractiones*. *Minuta* is used (for *minutae*) to mean sixtieths. See Boncompagni, *Trattati*, I, 17.

[6] Thus Huswirt (1501): "Minutia siue fractio nihil aliud est q̄ pars integri" (fol. 11, *r.*); and Clavius (1583) expresses certain of his quantities "in numeris fractis, qui alio nomine Mintutiae, fractionesve dici solent vulgares" (p. 81).

[7] Thus Fine (1530): "De minutis, siue quotis eorundem integrorū partibus (quas uulgares appellant fractiones)"; and Gemma Frisius (1540): "Fractiones minutias aut partes." Gosselin, in his translation (Paris, 1578) of Tartaglia's arithmetic, uses *parties* more commonly than any other term for fractions. Hylles (1592) has the expression: "fractions of fractions (or as some men call them particles, that is as you would say parcels of parts)." Ramus (1555) speaks of "fractio sive pars."

[8] Thus Chaucer, in his *Astrolabe* (*c.* 1391), uses *fraccion*.

[9] Pacioli (1494) ordinarily speaks of *rotti*, although he also uses *fractioni* and *fracti* (fol. 48). Most of the 16th century Italian writers use *rotti*.

[10] Ortega (1512).

[11] Chuquet (1484), "nombres routz"; Savonne (1563), "roupt"; and later writers, "nombre rompu."

In the Teutonic languages the custom was followed of using vernacular expressions, and so the Latin *fractio* appeared as "broken number."[1]

Common Fraction. The expression "common fraction" was originally used to distinguish the fractions employed in trade from the sexagesimal fractions found in astronomy. It refers merely to the form of writing a fraction, $\frac{5}{10}$ being a common fraction, 0.5 being a decimal, and 30′ being a sexagesimal, although the values of the three are the same. In Latin the expression was *fractiones vulgares*, whence the "vulgar fractions" of the English. The adjective "common" is used at present in America, although this has not always been the case,[2] nor have the English uniformly followed their present usage.[3]

Definition of Fraction. In general a fraction has been defined as one or more parts, or equal parts, of a unit,[4] sometimes with the limitation that the numerator must be less than the denominator.[5] Occasionally the more scientific writers based the definition upon division, usually of a smaller number by a larger.[6] The idea of an improper fraction, like $\frac{3}{2}$, is a late development. Occasionally a 16th century writer like Recorde[7] (*c.* 1542), Gemma Frisius[8] (1540), or Tartaglia[9] (1556) mentioned this type of fraction as an expression of division, but little was done with it. Complex fractions, those in which a fraction appears

[1] Thus Riese (1522) speaks of "Ein gebrochene zal" (1550 ed., fol. 14, *v.*), and Grammateus (1518) has a chapter "Von Prüchen," speaking of a fraction as "ein iglicher pruch (welchen mañ in latein fraction nennet)." So in Dutch we find Raets (1580) speaking of "Die ghebroken ghetalen," and Mots (1640) and others speaking of "Ghebroken."

[2] Similarly in France, instead of *fraction ordinaire* Trenchant (1566) used *fraction vulgaire*. Our colonial arithmeticians usually followed the English use of "vulgar."

[3] Thus Digges (1572) speaks of "the vulgare or common Fractions."

[4] *E.g.*, Pacioli (1494): "Rotto e vno o vero piu parti de vno ītegro" (fol. 48, *r.*); Santa-Cruz (1594): "Quebrados es vna parte, ò partes dela cosa entera."

[5] *E. g.*, Pagani's arithmetic (1591).

[6] *E.g.*, Ramus (1555). The Dutch arithmetic of Raets (1580) defines a fraction as "een ghetal diuideert met een grooter." On the fusion of the notions of fraction and quotient, see V. V. Bobynin, *Bibl. Math.*, XIII (2), 81.

[7] See the 1558 edition of the *Ground of Artes*, fol. S vi, *v.*

[8] "Fractiones quae plus Integro valent."

[9] 1556 ed., I, fol. 107, *r.*

in either numerator or denominator, or in both, are older than might be expected. Rabbi ben Ezra, for example, has a problem involving the product of two such forms.[1]

Terms of a Fraction. The medieval Latin writers found it convenient to devise names for the terms of a fraction written after the Arab manner, and so they called the upper number by such names as *numerator* (numberer) and *numerus* (number),[2] while the lower number was called the *denominator* (namer). These terms are hardly destined to endure, but no others have been generally accepted. Among the medieval and Renaissance writers the numerator was often designated by such words as *nominator*,[3] "topterme," "top,"[4] *superior*,[5] and *denominato*,[6] and the denominator by such names as *base*, *inferior*, and *denominante*. Both the numerator and denominator took on vernacular forms with later Teutonic writers.[7] In the Latin languages, however, the favorite names were *numerator* and *denominator*, the former of which Tartaglia (1556) speaks of as being written above a *virgoletta* (little bar), and the latter as being written below it.[8]

[1] *Sefer ha-Mispar*, 39.

[2] So the Rollandus MS. (*c.* 1424) speaks of the "nūator et denomtor." See also the correspondence of Regiomontanus and Bianchini, *Abhandlungen*, XII, 287. In the 16th century Ramus (1555) speaks of the *superior terminus* as the *numerus sive numerator*.

[3] *E.g.*, Digges (1572), although he also used *numerator*. See pages 20, 24, 27 of the 1579 edition.

[4] Thus Hylles (1592): "Numerator which also for more shortnesse is sometimes called the Topterme or top onely: and that the lower term is vsually called the Denominator or Base."

[5] As in Gemma Frisius (1540), although he also uses *numerator*.

[6] Paolo dell' Abaco (*c.* 1340): "Sappi che ogni rotto si scrive con due numeri: il minore sta sopra la verga e chiamasi denominato; e il maggiore sotto la verga e chiamasi denominante" (ed. Frizzo, 1883, p. 45). The name was used by various 16th century writers, such as Sfortunati, *Nuovo Lume* (1534).

[7] *E.g.*, Widman (1489) has the Latin forms, but a little later the words *Zeler* and *Nenner*, with variants, came into general use. Occasionally a Dutch writer like Wentsel (1599) used the Latin forms, but most arithmeticians preferred "teller" and "noemer," with such variants as "telder" and "nommer," and similarly with the Scandinavian writers.

[8] ". . . l' uno di quali è detto numeratore (& questo si scriue sempre sopra vna virgoletta) l' altro è chiamato denominatore, e questo si scriue sempre sotto a quella tal virgoletta." *General Trattato*, I, 106, *v.*; 107, *r.*

Reduction of Fractions. Until recently the reduction of a fraction to lower or lowest terms was commonly known as abbreviation. Thus Digges (1572) says:

To abbreuiate any Fragment, is to bring a Fraction to his lest denomination. To make this abbreuiation, yee must diuide the Numerator of the Fraction, and so in the like maner the Denominator by the biggest number, that is some common part of them both.[1]

The word "depression" was also used, and, like "abbreviation," is more suggestive than "reduction,"[2] which sometimes had the special meaning of bringing fractions to a common denominator.[3]

Before the invention of decimals such fractions as $\frac{3}{4}\frac{2}{3}\frac{4}{2}\frac{5}{0}\frac{3}{8}\frac{1}{6}\frac{2}{4}$ were not uncommon,[4] and it was necessary to reduce them to lowest terms in order to operate with them. In general the cancellation of all common factors was not convenient, and hence the long form of greatest common divisor was essential. First, however, factors were canceled. A factor thus eliminated was called by the Italians a *schisatore*.[5] On account of the necessity for recognizing common factors, many of the early manuscripts and printed works gave the ordinary tests for divisibility by 2, 3, and 5, and even some kind of test for

[1] 1579 ed., p. 24. So Hodder (1672 ed.) says: "I would abreviate $\frac{24}{120}$," and J. Ward (1771 ed.) has a caption "To Abbreviare or Reduce Fractions into their Lowest or Least Denomination." The expression is much older than this, however, for Chuquet (1484) says: "Abreuier est poser ou escripre vng nombre rout par moins de figures . . ." (fol. 12, *r.*, of his MS.). Early Spanish writers used the same expression, as in Santa-Cruz (1594), "De abreuiar quebrados." The Dutch writers of the same period used various terms, including *abbreviatio, verminderinge, and vercontinghe*, and Van der Schuere (1600) says: "om ghebroken ghetallen te vercorten ofte minderen."

[2] Thus Pacioli (1494): "De vltima depssione fractorum siue modo schisandi dicto," adding "Che ī frāçese si chiama Abreuier" (fol. 48, *v.*).

[3] *E. g.*, Pellos (1492, fol. 21, *r.*), Chuquet (1484, fol. 10, *r.*), and others. This special meaning was not general, for Tartaglia uses it in the broader sense (1592 ed., I, fol. 169, *r.*).

[4] This and similar fractions are in the Treviso arithmetic (1478). Fractions like $\frac{9}{1}\frac{4}{4}\frac{3}{0}\frac{2}{6}\frac{5}{0}\frac{0}{0}$ and $\frac{1}{3}\frac{0}{0}\frac{0}{0}\frac{3}{8}\frac{2}{0}\frac{7}{0}\frac{6}{0}\frac{8}{0}\frac{3}{0}$ are given in the *Epitome* of Clavius (1583; 1585 ed., pp. 77, 124).

[5] A word suggesting canceling "across" (*schisa*), whence *schisare*, to reduce a fraction. So Pacioli (1494) speaks "De diuersis modis in ueniendi schisatorem" (fol. 49, *r.*), and Cataneo (1546) tells "Come si schisino i rotti" and speaks of "Lo schisamento" and "di schisare."

divisibility by 7.[1] When common factors were not readily seen, the greatest common divisor was resorted to at once, being found by the Euclidean method.[2] This is given in al-Karkhî's *Kâfî fîl Hisâb* (*c.* 1015),[3] and in various other Oriental works, manifestly all derived from Greek sources.

Greatest Common Divisor. The greatest common divisor went by various names in the early printed books.[4] The theoretical works usually gave a rule for finding it, although the mercantile works often omitted the subject entirely, the former making use of long fractions and the latter ignoring them. One of the earliest printed rules is stated by Pacioli (1494) and is credited to Boethius (*c.* 510).[5] In this the smaller term is continually subtracted from the larger, a smaller remainder from that, and so on, an evident modification of the Euclidean method. Several early writers used the latter method for "abbreviating," without mentioning the greatest common divisor as such.[6]

Sequence of Operations. By analogy to the sequence of operations in the case of integers, the sequence in fractions has generally begun with addition. Medieval[7] and Renaissance writers, however, often took the more sensible course of beginning with multiplication,—a course to which the primary schools have now returned. Recorde (*c.* 1542) was earnest in his advocacy of this method, saying:

[1] A good illustration of the use of these tests in the later works of the 16th century may be found in Van der Schuere's *Arithmetica* (1600).

[2] That is, the one given in the *Elements*. See Heath's *Euclid*, Vol. II, pp. 118, 299.

[3] Hochheim ed., p. 10.

[4] In the Latin books it usually appears as *maximus communis divisor*, and in the Italian works as *il maggior comune ripiego* (Cataneo's spelling, 1546) or *massima cõmune misura* (Cataldi's spelling, 1606).

[5] "Vn altro modo se elice da Boetio nel secondo della sua Arithmetica per trouare ditto schisatore" (fol. 49, *v.*). See Friedlein's *Boethius*, p. 77.

[6] *E.g.*, Chuquet (1484), under "Aultre stile de abreuir," and the Dutch writer Petri (1567). Somewhat similar treatments are given by Baker (1568), Raets (1580), Rudolff (1526), and others. The phraseology used by Grammateus (1518) is interesting ("¶Prüch kleyner zumachen"), and that of Rudolff is analogous to it ("Wie man gewiszlich erkennen mag/ob ein bruch müg noch kleiner gemacht werdẽ od' nit").

[7] For example, Abraham ben Ezra (*c.* 1140).

There is an other ordre to be folowed in fractions then there was in whole numbres. for in whole numbres this was the ordre, Numeration, Addition, Subtraction, Multiplyplication, Diuision and Reduction. but in fractions (to folowe the same aptnesse in procedyng from the easyest woorkes to the harder) we muste vse this ordre of the woorkes, Numeration, Multiplication, Diuision, Reduction, Addition, and Subtractiō.

The book is in the form of a dialogue, and upon the pupil's saying, "I desyre to vnderstond y^e^ reason," the master says:

As in the arte of whole numbres ordre woulde reasonablye begyn with the easiest, and so go forwarde by degrees to the hardest, even so reason teacheth in Fractions the lyke ordre.[1]

Addition and Subtraction. In adding or subtracting, early writers usually took for a new denominator the product of the given denominators, reducing the final result to lowest terms.[2] Because of the size of the common denominator thus found,[3] the early Rechenmeisters in Germany ordinarily added but two fractions at a time. Although the plan of reducing to the least common denominator before adding or subtracting was occasionally used by 15th and 16th century arithmeticians,[4] it was not until the 17th century that it began to be generally recognized,[5] and even then the name was slow of acceptance.[6]

[1] For further discussion see the 1558 edition of the *Ground of Artes*, fol. R iiii, *v*. The same order is followed by Pacioli (1494, fol. 51), Pagani (1591, pp. 34, 41), and others. Giovanni Battista di San Francesco, *Elementi Aritmetici* (Rome, 1689), even begins with division "that it may be better understood."

[2] Thus, $\frac{3}{4} + \frac{5}{6} = \frac{18}{24} + \frac{20}{24} = \frac{38}{24} = 1\frac{7}{12}$. The method is given by Bhāskara (*c.* 1150); see Taylor's translation, p. 24. It appears in many medieval MSS. and in such early arithmetics as those of Petzensteiner (1483), Pellos (1492), Riese (1522), Recorde (*c.* 1542), and Baker (1568).


[3] *E.g.*, the Dutch arithmetician Wilkens (1630) reduces $\frac{3}{4}$, $\frac{4}{5}$, $\frac{5}{6}$, and $\frac{7}{8}$ to 960ths before adding.

[4] Chuquet (1484) gives it (fols. 13 and 14), and it is found in such works of higher class as those of Tartaglia (1556) and Clavius (1583).

[5] So Cataldi (1606) reduces to the "minor commune denominatore"; the well-known Coutereels (1599), to "het minste ghetal"; and Wilkens (1630), to the "Kleynste gemeyne Noemer."

[6] The shorter name of "general denominator" was used by some writers. See Starcken's Dutch work of 1714, with "General Nenner." Ramus (1569) suggested "cognomen" for common denominator,—not a bad term.

The arrangement of an example in addition was somewhat uniform before the 17th century, and it may be understood from the following case of $\frac{5}{6} + \frac{3}{8}$ as given by Pacioli[1]:




Multiplication of Fractions. Although our present interest in the multiplication of fractions relates to such simple cases as $\frac{2}{3} \times \frac{4}{5}$, it is desirable to set forth some of the difficulties met by ancient writers. These difficulties appear in the works of the Egyptians and Greeks, but they are sufficiently evident in a single example given by Rabbi Sa'adia ben Joseph al-Fayyumî, a Hebrew scholar of the 10th century already mentioned on page 212. The problem, which shows the difficulties met with in the use of unit fractions, is to find the product of $61\frac{1}{3}\ \frac{1}{9}$ by $61\frac{1}{3}\ \frac{1}{9}$. The solution is substantially as follows: $61 \times 10 = 610$, $61 \times 20 = 1220$, $61 \times 40 = 2440$, $61 \times 60 = 3660$, the last three being found by doubling or by adding. Then $61 \times 61 = 3721$, evidently found by adding 61 to 3660. Then $\frac{1}{3} \times 61 = 20\frac{1}{3}$, and $20\frac{1}{3} + \frac{1}{9}$ of $61 = 26\frac{2}{3}$ (*sic*), the double of which is $53\frac{1}{3}$. Adding this to 3721, he obtains $3774\frac{1}{3}$, and this increased by $\frac{2}{3}$ gives the result, 3775. What he tries to do is to square $61\frac{1}{3}\ \frac{1}{9}$ by taking $61^2 + 2 \times 61 \times (\frac{1}{3} + \frac{1}{9}) + (\frac{1}{3} + \frac{1}{9})^2$ (a rule which was, of course, well known), but he fails in his computation.

With respect to the ordinary operation with simple fractions, the process of multiplication has not changed materially during the last few centuries except that cancellation was not generally used by early writers,[2] although a few of the better arithmeticians saw its advantages.[3]

[1] 1494 ed., fol. 51, with an error in the quotient as printed.

[2] *E. g.*, Calandri (1491) multiplied $\frac{4}{5}$ by $\frac{3}{4}$ thus:

Multiplica $\frac{3}{4}$ uie $\frac{4}{5}$



Fanno $\frac{3}{5}$

Even in the Greenwood American arithmetic (1729) this method is followed.

[3] Thus Rudolff (*Kunstliche rechnung*, 1526) says "das man ein ober vnd ein vnter gegen einander mag auffheben oder kleyner machen" during the operation.

In the matter of language, the schools have usually protested against the broadening of the meaning of any technical term. A teacher will object to saying "$\frac{3}{2}$ times 4" but will say "$1\frac{1}{2}$ times 4." The contest is an old one; thus Ortega (1512) would not write "$3\frac{1}{4}$ ducats," preferring the awkward expression "3 ducats and one fourth of a ducat";[1] but Rudolff (1526), at about the same time, did not hesitate to speak of "$\frac{1}{2}$ times" a number.[2]

Of the various special rules, most of which came from the Arabs, a single one may serve as a type. Expressed in modern symbols,

$$\frac{a}{b} \cdot \frac{c}{d} = \frac{\frac{ac}{b}}{d} = \frac{\frac{ac}{d}}{b},$$

that is,

$$\frac{2}{3} \cdot \frac{5}{7} = \frac{\frac{10}{3}}{7} = \frac{\frac{10}{7}}{3},$$

thus reducing the work to dividing a fraction by an integer and suggesting cancellation more strongly.[3]

Many of the early writers expressed concern over the fact that the product of a number by a proper fraction was less than the multiplicand. Borghi (1484) seems to have been the first author of a printed book to discuss the matter, and various 16th century writers had much to say about it.[4]

Few writers before the 17th century made any attempt at explaining the process, although Trenchant (1566) devoted some attention to it, using the illustration of a square cut into smaller squares.[5]

[1] "¶Se 3 ducati e vn quarto de ducato . . . guadagnano 5 fiorini e vn terzo," much as we say "a dollar and a half."

[2] ". . . dañ ich hab die sechs nur ein halbs mal haben wöllen" (1534 ed., fol. C iiij, *v.*).

[3] This is given by al-Karkhî (*c.* 1020).

[4] Among them was Ramus (1569). In the 1586 edition (p. 73) his commentator, Schoner, gives a whole page to it. Cataneo (1546) also devotes a page to it, seeking particularly to combat Borghi and Pacioli. (See 1567 edition, fol. 21, *v.*) Even Tartaglia did not see the point of the controversy (see 1592 ed., I, fol. 187, *r.*).

[5] Cardinael's *School Boecken* (1650; 1674 ed.) goes into the matter more fully, using several diagrams.

Division of Fractions. Naturally the most difficult operation was division. Multiplication by the inverted divisor is so simple that we hardly realize that it has come into general use only recently, although it was known in the early Middle Ages by both the Hindus and the Arabs.[1] Influenced by the notion that only fractions could deal with fractions, medieval writers often substituted for the division of a fraction by an integer the process of multiplying by the reciprocal of the integer;[2] that is, $\frac{2}{3} \div 4 = \frac{2}{3} \times \frac{1}{4}$.

The early printed books gave two leading methods. The first of these reduced the fractions to a common denominator and took the quotient of the numerators,[3] as in the case of

$$\frac{2}{3} \div \frac{3}{4} = \frac{8}{12} \div \frac{9}{12} = \frac{8}{9}.$$

The second method is one of cross multiplication. Thus, in the case of $\frac{2}{3} \div \frac{3}{4}$ we have

$$\begin{array}{ccc} 2 & & 3 \\ & \diagdown\!\diagup & \\ 3 & & 4 \end{array} = \frac{8}{9},$$

which involves the same operations that enter with the inverted divisor. This was the favorite method in the early printed books,[4] and the name "cross multiplication" or its equivalent[5] was common, the divisor being usually placed on the left, but sometimes on the right.[6] One writer expresses the opinion that the divisor was placed at the left because the process may have come from the Hebrews, who write toward the left.[7]

[1] Brahmagupta (*c.* 628) and Bhāskara (*c.* 1150) both gave it (Colebrooke translation, pp. 17, 278), and al-Ḥaṣṣâr (*c.* 1175?) recognized it, at least with integral dividends (*Bibl. Math.*, II (3), p. 36).

[2] Thus Rollandus (*c.* 1424).

[3] *E.g.*, Chuquet (1484, fol. 16, whose manuscript was so extensively appropriated by De la Roche and in part printed in 1520), Trenchant (1566), and Ramus (1555).


[4] *E.g.*, Widman (1489): "Nū wiltu teilē $\frac{9}{13}$ in $\frac{5}{6}$ sprich 6 mal 9 ist 54 dy setz für den zeler vnd sprich darnach 5 mal 13 ist 65 die setz für den nēner also $\frac{54}{65}$" (1508 ed., fol. 30, *v.*).

[5] Thus Hodder (1672 ed.) says "multiply cross wise"; Riese (1522), "so multiplicir im creutz"; Peletier (1549), "multiplier en croix"; Pagani (1591), "moltiplica in croce."

[6] Thus Hudalrich Regius (1536), Pagani (1591), Mots (1640), and others.

[7] Wentsel, 1599, p. 88.

The idea would have been more reasonable, so far as immediate origin is concerned, if he had spoken of the Arabs.



EARLY DIVISION OF FRACTIONS

From an anonymous Italian MS. of 1545 in Mr. Plimpton's library

The Inverted Divisor. As already said, the method of multiplying by the inverted divisor was known to certain Hindu and Arab writers. It seems, however, to have dropped out of

sight for three or four hundred years, reappearing in Stifel's works in 1544.[1] It was not at once accepted, only a few of the 16th century writers making any use of it,[2] but in the 17th century it became fairly common.

Before the inverted divisor came into general use there were several special rules that met with some favor. One of these, given by Gemma Frisius (1540), may be expressed in modern symbols thus:

$$\frac{a}{b} \div \frac{ka}{c} = \frac{c}{kb},$$

as in

$$\frac{3}{5} \div \frac{12}{13} = \frac{3}{5} \div \frac{4 \times 3}{13} = \frac{13}{4 \times 5} = \frac{13}{20}.$$

2. Sexagesimal Fractions

Nature of Sexagesimals. For scientific purposes the medieval writers usually followed the late Alexandrian astronomers in the use of fractions written on the scale of sixty.[3] This custom has continued until now in the measures of time, angles, and arcs, as when we write 2 hr. 20 min. 45 sec., that is, $(2 + \frac{20}{60} + \frac{45}{3600})$ hours, instead of $(2 + \frac{1}{3} + \frac{1}{80})$ hours. The measure of time meets a popular need, and so the sexagesimal fraction gives no present evidence of being abandoned for this purpose, but for circular measure it is losing its hold as decimals become better known, and seems destined soon to disappear.

In the Middle Ages the scientific workers carried the sexagesimal divisions still farther than the Greeks, as if we were to write 2 10′ 30″ 45‴ 5^{iv} 7^{v}, meaning thereby

$$2 + \frac{10}{60} + \frac{30}{60^2} + \frac{45}{60^3} + \frac{5}{60^4} + \frac{7}{60^5}.$$

[1] "Ego Diuisionis regulam reduco ad regulam Multiplicationis Minutiarum, hoc modo: Diuisoris terminos commuto," etc. *Arithmetica Integra*, 1544, fol. 6, *r*.

"Thu im also. Den Teyler . . . kere vmb/also ausz dem Zeler werde der nenner/vnd ausz dem nenner der Zeler. So steht denn das exemplum mit vmbgekereten Teyler also" (*Deutsche Arithmetica*, 1545, fol. 13, *v*.).

[2] Among them were Thierfelder (1587) and Clavius (1583). The latter says: "ac si termini diuisoris commutentur, & regula multiplicationis seruetur" (1585 ed., p. 118, and similarly the Italian edition of 1586, p. 106).

[3] Latin *sexagesimus* or *sexagensumus*, sixtieth, from *sexaginta*, sixty.

Thus Sibṭ al-Mâridînî,[1] an astronomer at the mosque of al-Azhar in Cairo in the middle of the 15th century,[2] gave $45^\circ\ 50' \div 1^\circ\ 25' = 33^\circ\ 45'\ 52''\ 56'''\ 28^{iv}\ 14^{v}\ 7^{vi}\ 3^{vii}\ 31^{viii}\ 45^{ix}\ 52^{x} \cdots$, and similar cases occur in many medieval works.

Names of Sexagesimals. Sexagesimals were usually known as physical fractions in the Middle Ages.[3] The name may possibly have come from their use in physics, this word (more frequently "physic"), as applied to natural philosophy, not being so recent as is sometimes thought. On the other hand, it may come from the fact that the denominators were understood to proceed in the natural[4] order of the powers of 60, somewhat as we speak of "natural numbers" at present, this being an opinion expressed in the 16th century.[5]

They were also called astronomical fractions,[6] the reason being quite apparent.[7]

Origin of Sexagesimals. There is a common idea that sexagesimal fractions came from Babylon,—an idea which arose from the fact that 60 plays an important part in the number

[1]Mohammed ibn Mohammed ibn Aḥmed, Abû 'Abdallâh, Bedr ed-dîn al-Miṣri, born in 1423, died in 1494/95. He wrote a number of works on arithmetic and astronomy.

[2]He gives the result only to 31^{viii}, the fraction then repeating,—an interesting case of a circulating sexagesimal. See Carra de Vaux, "Sur l'histoire de l'arithmétique arabe," *Bibl. Math.*, XIII (2), 33. The above symbols are, of course, modern. The problem is substantially that of $45\frac{5}{6} \div 1\frac{5}{12} = 33\frac{3}{4} +$.

[3]Thus we find in the MSS. such expressions as "Modum representationis minuciarum vulgarium et physicarum" (anonymous MS. of 1466) and "Minucie duplices sunt scilicet phisice et vulgares" (anonymous MS. of 15th century). In the early printed books they are called "fractiones phisice" (Ciruelo, 1495), "fraciones fisicas" (Texeda, 1546), "Minucciamenti Fisici" (Italian edition of Gemma Frisius, 1567), and by other similar names. [4]Φύσις (*phy'sis*, nature).

[5]Thus Trenchant (1566): "S'apele phisic, c'est à dire, naturel: pour ce que ses denominateurs, & caracteres, sont selon l'ordre naturel du nombre commençant à l'vnité (1578 ed., p. 19).

[6]"De Fractionibus Astronomicis, siue de minutiis Physicis," as Gemma Frisius (1540) says in his Latin editions, the Italian having "Rotti Astronomici." Trenchant has "Du nombre phisic, ou fractions astronomiques." Peletier (Peletarius) in his notes on Gemma Frisius (1563 ed.) speaks of "Fractiones Astronomicae, quas vulgò Physicis vocât." The name was used by Abraham ben Ezra (*c.* 1140) and probably by the late Greek writers.

[7]As Peletier (1549) says, because they "seruent aux supputations des mouuements celestes."

system of that country. The assertion of this origin was first made, so far as we know, by Achilles Tatius, an Alexandrian rhetorician of the 5th or 6th century. It has also been assumed that the Babylonians divided the circle into 360 equal parts, because of the early notion that a year consisted of 360 days, and because their scientists knew that the radius employed in stepping around a circle divided it into six equal arcs, thus making 60 a mystic number. This reason may possibly be valid, but there is no authority for asserting that it is historical. The Babylonians divided the circle into 8, 12, 120, 240, and 480 equal parts, but not into 360 such parts.[1] Thus in a tablet from the palace of Sennacherib (*c.* 700 B.C.) now in the British Museum the division into 480 parts is given. It is true that six-spoked wheels are found represented on the Babylonian monuments, but no more frequently than the eight-spoked wheels, and the six-spoked type is more common in Egypt where the number 60 was not used to any great extent. It would seem, therefore, that the number 60 was not derived from the division of the circle into six equal arcs.

It is true, however, that the Babylonians wrote the equivalent of 11 for $60 + 1$, 111 for $60^2 + 60 + 1$, and 44 26 for $44 \times 60 + 26$, although there is no reason for believing that this is a proof of their use of sexagesimal fractions. In a certain tablet of *c.* 2000 B.C., for example, the equivalent of the square of 44 26 40 is given as 32 55 18 31 6 4. This may be interpreted to mean the square of either $44 \times 60^2 + 26 \times 60 + 40$ or $44 + \frac{26}{60} + \frac{40}{60^2}$. In the latter case we have sexagesimal fractions; in the former, numbers written on the scale of sixty, an interpretation more in harmony with the system of compound num-

[1] On this entire discussion see A. H. Sayce and R. H. M. Bosanquet, "The Babylonian Astronomy," in *Monthly Notices of the Royal Astron. Society*, XL, 108; E. Hoppe, *Archiv der Math.*, XV (3), 304; E. Löffler, *ibid.*, XVII (3), 135; and Hochheim, *Kâfî fil Hisâb*, p. 23. The claim that the Chinese used a sexagesimal system in the third millennium B.C. (Vol. I, p. 24) is not supported by sufficient evidence to be considered at present. It is very improbable that it involved anything more than a recognition of 60 as a convenient unit for subdivision. On the Greek development of sexagesimals see Heath, *History*, I, 44.

bers used by all ancient peoples. Similarly, we find the case of $1 \div 81$, but whether this is to be interpreted as having the dividend 60 or some power of 60 is uncertain. In any case we have no evidence of any such general use of sexagesimal fractions as is found among the Greek astronomers.[1]

The division of the circle into 360 parts as practiced by such Greek astronomers as Ptolemy (*c.* 150) was probably the outgrowth rather than the origin of the sexagesimal system. The Babylonians counted decimally by preference, although the base of 60 played a considerable part in their system. They counted decimally to 60, that is, to a *soss*; then by *sosses* and the number over to the *ner*, which was 10 *sosses*, or 600; then by *ners*, *sosses*, and the number over to the *saru*, which was 6 *ners*, or 3600; but they never counted 60, 360, 3600, so that 360 was not a natural step in their sexagesimal system.[2]

Greek Use of Sexagesimals. We do not know why the Greek astronomers should have developed a scale of 60 in such a complete form, although we can readily surmise the cause. There seems to be no reason to doubt that the number 60 was suggested to them from Babylon, but the system of sexagesimal fractions, as we know and use it, was, so far as now appears, their own invention. Ptolemy used these fractions to represent his chords in terms of a radius 60;[3] that is, the chord of 24° would then be 24.9494, or, in sexagesimals, 24 56′ 58″. It seems clear, however, that the Greeks needed for their astronomical work a better type of fraction than the unit type of the Egyptians; that their habit of using such submultiples, as in feet and inches, naturally led them to a similar usage in fractions, as would be the case with degrees and minutes; and that the 60 of Babylon was a convenient radix, since it has as factors 2, 3, 4, 5, 6, 10, 12, 15, 20, and 30, and so permits of the ready use of halves, thirds, fourths, fifths, sixths, tenths, twelfths, and so

[1] For arguments in favor of the fraction interpretation see F. Cajori, "Sexagesimal Fractions among the Babylonians," *Amer. Math. Month.*, XXIX, 8. See also Heath, *History*, I, 29. [2] Hilprecht, *Tablets*.

[3] For a discussion of this point see A. Schülke, "Zur Dezimalteilung des Winkels," *Zeitsch. für math. und naturw. Unterr.*, XXVII, 339; Heath, *History*, I, 45.

on. The Greeks may thus have been led to divide the radius into 60 equal parts and the diameter into 120 of these parts. Since the common value of π was 3 in ancient times, the circumference was naturally taken as 3×120, or 360.

Such was the influence of the Greek scholars that all the medieval astronomers, Christian, Jewish, and Mohammedan, used the sexagesimal system;[1] but some of the mathematical writers referred the system to India instead of Greece, influenced therein by the belief that our numerals came from the Hindus.[2]

Terms Used. When the Greeks decided to take $\frac{1}{360}$ of a circle as a unit of arc measure, they called this unit a degree.[3] They called $\frac{1}{60}$ of a degree a first part,[4] $\frac{1}{3600}$ a second part,[5] and so on.

Multiplication involving Sexagesimals. The operations of addition and subtraction with sexagesimals involved no difficulties, but multiplication and division were not so simple. It is meaningless to us to multiply $4^\circ\ 7'\ 38''$ by $5^\circ\ 6'\ 29''$, or even $4\ 7'\ 38''$ by $5\ 6'\ 29''$, but to the medieval scientist it meant

[1] As a noteworthy illustration, see the *Libros del saber de Astronomia del Rey Alfonso X*, Madrid, 1863. The Alfonsine astronomical tables date from *c.* 1254, but for argument as to a later date, see A. Wegener, "Die Astronomischen Werke Alfons X," *Bibl. Math.*, VI (3), 138.

[2] Thus Johannes Hispalensis (*c.* 1140): "placuit tamen Indis, denominationem suarum fractionum facere a sexaginta. Diuiserunt enim gradum unum in sexaginta partes, quas uocauerunt minuta" (B. Boncompagni, *Trattati*, II, 49). He may have had his idea from al-Khowârizmî (*c.* 825): "Set indi posuerunt exitum partium suarum ex sexaginta: diuiserunt enim unum in .LX. partes, quas nominauerunt minuta" (from a Cambridge MS. of the *Algoritmi de Numero Indorum*, in the *Trattati*, I, 17).

[3] Μοῖρα (*moi'ra*); medieval Latin, *de* + *gradus* (step). The Arabs translated μοῖρα by *daraja* (ladder, scale, step), which led G. H. F. Nesselmann (*Die Algebra der Griechen*, p. 137 (Berlin, 1842), hereafter referred to as Nesselmann, *Alg. Griechen*) to think that this word was the original form of the word "degree." It may have influenced the final form.

[4] Πρῶτα ἑξηκοστά (*pro'ta hexekosta'*); Latin, *pars minuta prima* (first small or fractional part). From this came our "minute." The Greeks also used λεπτά (*lepta'*, minute, the adjective). In the 12th century Walcherus (see Volume I, page 205) spoke of the minutes as *puncta*, and the same term is so used in an algorism of *c.* 1200. See L. C. Karpinski, "Two Twelfth Century Algorisms," *Isis*, III, 396.

[5] Δεύτερα ἑξηκοστά (*deu'tera hexekosta'*); Latin, *pars minuta secunda*, from which our "second." See also Wertheim, *Elia Misrachi*, p. 19 n.

simply the finding of $(4 + \frac{7}{60} + \frac{38}{3600}) \times (5 + \frac{6}{60} + \frac{29}{3600})$. In the operation there is, for example, $7' \times 6' = 42''$, which means simply that $\frac{7}{60} \times \frac{6}{60} = \frac{42}{3600}$.[1] In such work it became convenient to have multiplication and division tables, and these are found in various medieval manuscripts.[2] Some idea of the difficulty of operating with these fractions may be inferred from a problem in the work of Maximus Planudes (*c.* 1340).[3] His multiplication of 14° 23′ by 8° 16′, giving the product 3 signs[4] 28° 54′ 8″, is here shown.

3	28	54	8
	14	23	
	8	16	

Division involving Sexagesimals. In division Maximus Planudes reduced all the terms to the same denomination. For example, the operation

$$\frac{3^\circ\ 23'\ 54''}{2^\circ\ 34'\ 24''} = 1^\circ\ 19'\ 14'',$$

or

$$\frac{3 + \frac{23}{60} + \frac{54}{3600}}{2 + \frac{34}{60} + \frac{24}{3600}} = 1 + \frac{19}{60} + \frac{14}{3600},$$

is worked out as illustrated here.[5]

3	23	54
2	34	24
	12234	1
	9264	
	2970	
	178200	19
	9264	
	2184	
	131040	14
	9264	

The finding of roots by the aid of sexagesimals appears in the works of

[1] Thus in the translation of al-Khowârizmî (*c.* 825) attributed to Adelard of Bath (*c.* 1120): "Sex minuta multiplicata in VII. minuciis, erunt .XLII. secunda" (Boncompagni, *Trattati*, I, 18).

[2] In the adaptation of the *Liber Algorismi* by Johannes Hispalensis (*c.* 1140) the multiplication table is given up to *nona* times *nona*, that is, up to $\frac{1}{60^9} \times \frac{1}{60^9} = \frac{1}{60^{18}}$ (see Boncompagni, *Trattati*, II, 103). The printed arithmetics occasionally gave such tables; *e.g.*, those of Cardan (1539, cap. 38), Fine (1530; 1555 ed., fol. 38, *r.*), Trenchant (1566), and Peletier (1549). Schoner, in his *De logistica sexagenaria* (1569; 1586 ed., p. 370), calls it the "abacus logisticus," and a table of products up to 60 × 60, for use with sexagesimals, is called by Fine (1530) a "tabvla proportionalis." Division tables are also given by various writers; *e.g.*, Fine (1530) and Trenchant (1566).

[3] Wäschke, *Planudes*, p. 34.

[4] The "sign" was 30°, and the 12 signs of the zodiac gave 360°.

[5] He says that the division may be continued farther.

Theon of Alexandria[1] (*c.* 390) and Maximus Planudes[2] (*c.* 1340), and in several of the 16th century arithmetics.[3] Its nature may be inferred from the work in division.

Symbols. The symbols (° ′ ″) are modern. In medieval and Renaissance times there were several methods used for designating the sexagesimal orders. Thus in a manuscript of Leonardo of Cremona[4] (15th century) we have

$$.5.\ \overset{\text{aj}''}{19}\ .\ \overset{2''}{57}\ .\ \overset{3''}{38} \quad \text{for} \quad 5°\ 19'\ 57''\ 38''',$$

and

$$\overset{\text{aj}}{.46.}\ \overset{2}{39}\ .\ \overset{3}{12}\ .\ \overset{4}{36} \quad \text{for} \quad 46'\ 39''\ 12'''\ 36^{\text{iv}}.$$

Gemma Frisius (1540) wrote[5]

$$\begin{array}{cccccc} \text{S.} & \breve{\text{g}}. & \hat{\text{m}}. & \check{2}. & \hat{3}. & \check{4} \\ 1. & 16. & 25. & 17. & 21. & 27 \end{array}$$

for $1^{\text{s}}\ 16°\ 25'\ 17''\ 21'''\ 27^{\text{iv}}$.

Peletier (1571 edition of Gemma Frisius) used m˜ or 1˜ for minutes, 2˜ for seconds, 3˜ for thirds, and so on. Jean de Lineriis (*c.* 1340)[6] used the symbols s̈, g̈, m̈, 2̈, 3̈, 4̈, and these, with slight modifications, are the ones most commonly seen up to the close of the 16th century. About that time there came into use such forms as

$$\begin{array}{ccccccc} \text{IIae} & \text{Iae} & \circ & \text{I} & \text{II} & \text{III} \\ 3 & 15. & 7. & 50. & 34. & 23. \end{array}$$

[1] The process is given in Heath, *History*, I, 60, and in Gow, *Greek Math.*, p. 55.

[2] Heath, *History*, II, 547, where the date of Planudes is given a little earlier.

[3] *E.g.*, the Peletier (Peletarius) revision (1545) of Gemma Frisius.

[4] In Mr. Plimpton's library; *Rara Arithmetica*, p. 474. The title is *Artis metrice pratice compilatio.*

[5] "Circulus 12 Signis constat: Signum, 30 Gradibus." The relation to the zodiac is apparent.

[6] In the *Algorismvs de Minutijs* appended to Beldamandi's work (1483 ed.).

in which $\frac{0}{7}$ stands for units,[1] and in which the symbols are evidently the forerunners of the ones now in common use.[2]

One curious example of symbolism is seen in the multiplication table given by Fine (1530) for use in sexagesimal computation, the product 8×42 being given as 5.36, that is, $5 \times 60 + 36$, the period being essentially a sexagesimal point.

3. Decimal Fractions

Need for Decimal Fractions. Before the beginning of printing, operations with common fractions having large terms are not frequently found. In mercantile affairs they were not needed, and in astronomical work the sexagesimal fraction served the purpose fairly well. The elaborate Rollandus manuscript of *c.* 1424 contains the addition of no common fractions more difficult than $\frac{2}{2}\frac{0}{2}\frac{7}{0}$ and $\frac{1}{2}\frac{9}{8}\frac{7}{0}$, and the work in multiplication involves no fraction more elaborate than $\frac{2}{3}\frac{9}{6}$. There are exceptions to the general rule,[3] but they are not numerous. In the recording of results in division, however, elaborate common fractions were frequently used.[4] By the advent of printing, writers were led into various excesses. Widman (1489), for example, used in business computations fractions far beyond any commercial needs, his successors were even more reckless,[5] and the theorists naturally went still farther.[6]

[1] This example is from Schoner, *De logistica sexagenaria*, 1569; 1586 ed., p. 366. Of the 7 he says, "qui & partium numerus dicitur, circulus," and he speaks of "7 unitates."

[2] Peletier (1549) remarks: "Les Degrés dont seront au milieu de la numeration Astronomique: & seront representés par °: les Minutes par ′ les Secondes par ″: . . . Et ansi des autres" (*L'Arithmetiqve*, p. 107 (1607 ed.)).

[3] *E.g.*, in a Dutch MS. of the 15th century (Boncompagni sale cat., No. 477) the square root of $2520\frac{24267352095}{319795477401 6}$ is required.

[4] *E.g.*, in the *Svmme Arismetice* of Stephano di Baptista delli Stephani da Mercatello (MS. of c. 1522), a pupil of Pacioli's, there are results like $34\frac{2}{5}\frac{9}{5}\frac{8}{5}\frac{7}{9}\frac{2}{8}$ (fol. 74, *r.*).

[5] Thus Widman uses $\frac{3}{6}\frac{6}{9}\frac{5}{1}\frac{1}{8}\frac{5}{5}\frac{7}{7}$, Trenchant (1566) has $1980\frac{1}{1}\frac{7}{0}\frac{3}{1}\frac{9}{0}\frac{2}{2}\frac{5}{0}\frac{1}{2}\frac{4}{0}\frac{1}{1}\frac{1}{6}$ (1578 ed., p. 286), and Wentsel (1599) has several fractions as difficult as $\frac{48}{92}\frac{10}{15}\frac{79}{214}$; and all of these were commercial writers.

[6] As when Scheubel, in his *De nvmeris, tractatus quintus* (1545), gives $37\frac{2}{8}\frac{2}{4}\frac{2}{7}\frac{2}{1}\frac{7}{2}\frac{9}{4}\frac{7}{2}\frac{1}{1}\frac{4}{7}\frac{2}{2}\frac{9}{1}\frac{0}{7}$, and Coutereels (Eversdyck's edition of 1658) gives a result like $2\frac{1}{3}\frac{1}{3}\frac{7}{7}\frac{2}{5}\frac{4}{6}\frac{9}{6}\frac{1}{8}\frac{1}{7}\frac{2}{5}\frac{2}{4}\frac{4}{2}$ days.

Forerunners in the Invention. As usual in the case of an important invention there were various scholars who had some intuition of the need for such a device as the decimal fraction long before it was finally brought to light. Such a man was Joannes de Muris, or Jean de Meurs, who wrote early in the 14th century.[1] The most interesting of the early influences tending to the invention, however, was a certain rule for the extraction of $\sqrt[n]{a}$, expressed in modern symbols by $\frac{\sqrt{a \cdot 10^{kn}}}{10^k}$. In particular, $\sqrt{3} = \frac{\sqrt{30000}}{100}$, or $\frac{\sqrt{3000000}}{1000}$, the actual process of extracting the root being quite like our present one with decimals. It was known to the Hindus, to the Arabs, and to Johannes Hispalensis (*c.* 1140), and is found in the works of Johann von Gmünden (*c.* 1430), Peurbach (*c.* 1460), and their successors until the close of the 16th century.[2] The most interesting step from this rule in the direction of the decimal fraction appears in certain tables of square roots, in connection with which the statement is made that, the numbers having been multiplied by 1,000,000, the roots are 1000 times too large. Such a table, from Adam Riese's *Rechnung auff der Linien vnd Federn* (Erfurt, 1522), is reproduced on page 237. The same plan is given by such later writers as Trenchant (1566) and Bartjens (1633). Even after the decimal fraction was well known, the analogous plan of using a radius of 10,000,000, in order to express the trigonometric functions as whole numbers, remained in use for more than two centuries. It even extended to the reckoning of interest "to the Radius 100,000," as Thomas Willsford says in his appendix to the 1662 edition of Recorde's *Grovnd of Arts*,[3] so as to avoid decimals.

[1] L. C. Karpinski, *Science* (N. Y.), XLV, 663.

[2] Buckley, for example, an English arithmetician, who died *c.* 1570, gave the rule in Latin verse as follows:

> Quadrato numero senas præfigito ciphras
> Producti quadri radix per mille secetur.
> Integra dat Quotiens, et pars ita recta manebit
> Radici ut veræ, ne pars millesima desit.
>
> *Arithmetica memorativa. c.* 1550

[3] As spelled in this edition.

ſo: Den erſten Punct ſetz 1. vnd ſetze dafür drey nulla / Ziehe dañ Radicem quadratam daruon ſo kommen 1000. Dann preponir dem anderen Puncten / das iſt der Ziffern 2. auch ſechs 0 / vnd ziehe Radicem quadratã dauon / ſo komen 414.

Den dritten Punct mach auch alſo. Setz 3. vñ darnach ſechs 0. Extrahir dann Radicem quadratam dauon / kommen 832. Alſo thũ mit allen Puncten / ſo machſtu die Tafel ſelber. Es iſt aber groß mühe vnd verdroſſen arbeyt / Darumb hab ich dir hie ein Tafel außgezogen / die gehet biß vff 240. Punct der tieffe / der mañ gnũg hat vff groß oder kleyne vaß.

Tabula Radicum quadratarum.

1	1	1000	17	123	33	747
	2	414	18	242	34	833
	3	732	19	358	35	917
2	4	1000	20	472	6	36	1000
	5	234	21	584	37	82
	6	449	22	692	38	163
	7	645	23	767	39	244
	8	828	24	900	40	324
3	9	1000	5	25	1000	41	403
	10	162	26	98	42	481
	11	316	27	195	43	558
	12	446	28	290	44	634
	13	606	29	384	45	709
	14	741	30	477	46	783
	15	873	31	567	47	856
4	16	1000	32	659	48	928

EARLY STEPS TOWARD DECIMALS (1522)

From Adam Riese's arithmetic, showing a table of square roots in which the figures of the decimal fractions appear, but without any form of decimal point. From a later edition of the work

Another influence leading to the invention of the decimal fraction was the rule for dividing numbers of the form $a \cdot 10^n$, attributed by Cardan (1539) to Regiomontanus. This appears in several manuscripts of the 15th century,[1] as in the case of $470 \div 10 = 47$ and $503 \div 10 = 50\frac{3}{10}$. Borghi (1484) elaborates this rule, but it appears in its most interesting form in the rare arithmetic of Pellos (1492), who unwittingly made use of the decimal point for the first time in a printed work (p. 239). The use of the dot before and after integers had been common in the medieval manuscripts, as in the case of Chuquet's work already mentioned, but its use to separate the integer from what is practically a decimal fraction is first seen here. Later writers commonly used a bar for this purpose, as was the case with Rudolff (1530; see page 241), Cardan (1539), Cataneo (1546), and various other writers. Even as late as the 1816 edition of Pike's *Arithmetick* (New York, 1816) 46,464 is divided by 7000 thus:

$$7|000)\,46|464\;(6\tfrac{4}{7}\tfrac{4}{0}\tfrac{6}{0}\tfrac{4}{0}$$
$$\underline{42|}$$
$$4|464$$

Pellos, however, did not recognize the significance of the decimal point, as is evident from the facsimile on page 239, and no more did Cardan appreciate the significance of the bar that he used for the same purpose.[2]

The initial steps in the invention of the decimal fraction were not confined to the West, however; indeed, the credit for first recognizing the principle of this type of fraction may well be given to al-Kashî,[3] the assistant of the prince astronomer Ulugh Beg and the first director of the latter's observatory at Samarkand. In his *al-Risâli al-moḥîṭîje* (*Treatise on the circum-*

[1] Thus Chuquet (1484): "Comme qui vouldroit partir .470. par .10. fault oster .0. qui est la p̄me' figure de .470. et demeurent .47. et tant monte la part. Ou que vouldroit partir 503. par .10. fault oster .3. et les mettre dessus .10. et lon aura .50. $\frac{3}{10}$. pour quotiens." Fol. 8, *v*.

[2] See his *Practica* (1539), cap. 38.

[3] See Volume I, page 289, n. 5. He died *c.* 1436, or possibly, as some writers assert, *c.* 1424.

℄ Partir per 2 ol

7 9 6 5 4 8 3 9 . 7

quocient 3 9 8 2 7 4 1 9 $\frac{17}{20}$

℄ Partir per 3 0

5 8 3 6 0 4 . 3

quocient 1 9 4 5 6 4 $\frac{23}{30}$

℄ Partir per 7 0

9 5 3 7 9 1 . 9

quocient 1 3 6 2 5 5 $\frac{69}{70}$

℄ Partir per 1 0 0

6 9 7 6 5 . 8 7

quocient

℄ Partir per 4 0 0l

7 8 9 6 5 . 7 3

quocient 1 9 7 4 1 $\frac{173}{400}$

℄ Partir per 3 0 0 0

8 7 6 5 8 . 7 9 1

quocient 2 9 2 1 9 $\frac{1791}{3000}$

FROM THE PELLOS ARITHMETIC (1492)

ference) he not only gives the value of π to a higher degree of accuracy than any of his predecessors, but he writes it (using Arabic characters) as follows:

sah-hah

3 1415926535898732,

the word *sah-hah* meaning complete, correct, integral.[1] We have, therefore, a fraction which we may express as follows:

Integer

3 14159 · · ·,

the part at the right being the decimal. Manifestly it is, therefore, a clear case of a decimal fraction, and it seems to be earlier than any similar one to be found in Europe.

The Invention. The first man who gave evidence of having fully comprehended the significance of all this preliminary work seems to have been Christoff Rudolff, whose *Exempel-Büchlin* appeared at Augsburg in 1530. In this work he solved an example in compound interest, and used the bar[2] precisely as we should use a decimal point today (see page 241). If any particular individual were to be named as having the best reason to be called the inventor of decimal fractions, Rudolff would seem to be the man, because he apparently knew how to operate with these forms as well as merely to write them, as various predecessors had done. His work, however, was not appreciated, and apparently was not understood, and it was not until 1585 that a book upon the subject appeared.

The first to show by a special treatise that he understood the significance of the decimal fraction was Stevin, who published a work[3] upon the subject in Flemish, followed in the same year

[1] The modern Turkish form is *sahih*. I am indebted for these facts to Professor Salih Mourad of Constantinople.

[2] On the general question of notation see Gravelaar, "De Notatie der decimale Breuken," *Nieuw Archief voor Wiskunde*, IV (2).

[3] *De Thiende*. A copy of this rare pamphlet was fortunately saved at the time of the destruction of the Louvain library, having been borrowed a few days before by the Reverend H. Bosmans, S. J. See the *Revue des Questions Scientifiques*, January, 1920. There was an English translation by Robert Norton, London, 1608.

3>5. 18>5.
fl. 393|>5 hauptgůt vñ gewin des erstē jars.
19683>5
413|43>5 Andern
206>18>5
434|10933>5 Dritten
21>05468>5
455|8148843>5 Vierdten
22>90>4218>5
478|6055585933>5 Fünfften
23 9302>92968>5
502|535865234 3>5 Sechsten
25 126>93261>18>5
527|66265849609 3>5 Sibenden
263831329248046 8>5
554|045>9142080843>5 Achteten
2>>022895>104492 18>5
581|>48080991943359 3>5 Neündtē
29 08>4040905 9>16>968>5
fl.610|835485041540523433>5 Zehēdē
ß 6|68>88033232421 8>5000
℔ 20|6164099 69>2656250000.
>2 Die 120 fl tragē 2 jar p hauptgůt zins vnd zinszins 132 fl 2ß 12 ₰. Bringt zinß vñ zinszins 12 fl 2 ß 12 ₰. Darnach die 250 fl tragē 3 jar hauptg. zins vñ zinszinß 289 fl 3ß > ₰ ½. Vnd ist halber zins des vierdē jars > fl 1ß 26. ₰ 7/16

℔ ij

EARLY APPROACH TO DECIMAL FRACTIONS

From the 1540 edition of Rudolff's *Exempel-Büchlin* (1530), showing the use of decimal fractions in compound interest

(1585) by a French translation. This work, entitled in French *La Disme*, set forth the method by which all business calculations involving fractions can be performed as readily as if they involved only integers.[1] Stevin even went so far as to say that the government should adopt and enforce the use of the decimal system, thus anticipating the modern metric system.[2] He was the first to lay down definite rules for operating with decimal fractions, and his treatment of the subject left little further to be done except to improve the symbolism. Some idea of his treatment of the subject and of his symbols may be obtained from the facsimile shown on page 243.

The Symbolism. The decimal fraction had now reached the stage in its progress when the symbolism had to be settled. As already stated, Pellos (1492) had used a period to separate the decimal from the integral part, but he had not comprehended the nature of the fraction. This, however, was hardly more strange than that as good a computer as Vlacq[3] (1628) should use decimals in his calculations and tables and yet give a result in the form $1295\frac{65032}{100000}$. Several writers had used the bar to mark off the decimal part, and Rudolff had probably grasped the significance of the new fraction. Stevin had fully comprehended and clearly expounded the theory, but his symbols were not adapted to use. The improvement in the symbolism was due largely to Bürgi, Kepler, and Beyer, and to the English followers of Napier.

Jobst Bürgi (1552–1632) dropped the plan used by Regiomontanus—that of taking 10,000,000 as the *sinus totus* in trigonometry—and took 1 instead, the functions therefore becoming decimal fractions. He was not clear as to the best method of representing these fractions, however, and in his manuscript of 1592 he used both a period and a comma for the

[1] ". . . facilement expedier par nombres entiers sans rompuz toutes comptes se rencontrans aux affaires des Hommes."

[2] Adriaen van Roomen (1609) tells us that Bishop Ernst of Bavaria had similar ideas as to measures.

[3] *Arithmetica Logarithmica*, pp. 35 *et passim* (Gouda, 1628), evidently thinking that the decimal form of the result would not be understood by most readers.

SECONDE PARTIE DE LA DISME DE L'OPERATION.

PROPOSITION I, DE L'ADDITION.

Estant donnez nombres de Disme à ajouſter : Trouver leur ſomme :

Explication du donné. Il y a trois ordres de nombres de Diſme, deſquels le premier 27 ⓪8①4②7③, le deuxieſme 37 ⓪8①7②5③, le troiſieſme 875 ⓪7①8②2③.

Explication du requis. Il nous faut trouver leur ſomme. *Conſtruction.* On mettra les nombres donnez en ordre comme ci joignant, les aiouſtant ſelon la vulgaire maniere d'aiouſter nombres entiers, en ceſte ſorte:

⓪	①	②	③
	2	7	8	4	7
	3	7	6	7	5
8	7	5	7	8	2
9	4	1	3	0	4

Donne ſomme (par le 1 probleme de l'Arithmetique) 941304, qui ſont (ce que demonſtrent les ſignes deſſus les nombres) 941 ⓪3①0②4③. Ie di, que les meſmes ſont la ſomme requiſe. *Demonſtration.* Les 27 ⓪8①4②7③ donnez, ſont (par la 3e definition) 27 $\frac{8}{10}$, $\frac{4}{100}$, $\frac{7}{1000}$, enſemble 27 $\frac{847}{1000}$, & par meſme raiſon les 37 ⓪6①7②5③ vallent 37 $\frac{675}{1000}$, & les 875 ⓪7①8②4③ feront 875 $\frac{782}{1000}$, leſquels trois nombres, comme 27 $\frac{847}{1000}$, 37 $\frac{675}{1000}$, 875 $\frac{782}{1000}$, ſont enſemble (par le 10e probleme de l'Arith.) 941 $\frac{304}{1000}$, mais autant vaut auſſi la ſomme 941 ⓪3①0②4③, c'eſt

A PAGE FROM STEVIN'S WORK, 1634 EDITION

From the first work devoted to decimal fractions. The first edition was published at Leyden, 1585

decimal point,[1] and also wrote $14\overset{\circ}{1}4$ for 141.4. In his use of these fractions he was followed by Prætorius, in a manuscript of 1599.[2]

In his tables of 1612[3] Pitiscus assumed the radius to be 100,000 and gave sin 10″ as 4.85. Since this sine for the radius 1 is 0.00004848, the point after the 4 is possibly intended as a decimal point. Occasionally he used several points, as when he gave sin 89° 59′ 30″ as 99999. 99894. 23. In his trigonometry, of which the tables are a part, he used a vertical line to mark off the decimal.[4] In the 1600 edition both the point and the vertical line are used for other purposes, the former to separate sexagesimals[5] and the latter to separate (as above) a large number into periods, usually of five figures each.

It is unquestionably true that the invention of logarithms had more to do with the use of decimal fractions than any other single influence. When Napier published his tables in 1614 he made no explicit use of decimal fractions, the sine and the logarithm each being a line of so many units. In the 1616 translation of this work, however, the translator, Edward Wright, made use of the decimal point. One line will serve to show the appearance of the table:

Deg. o. Min.	Sines.	Logarith.	+|− Differen	Logarith.	Sines	
30	8726	4741385	4741347	38 · 1	999961 · 9	30

In his *Rabdologiae . . . Libri Dvo* of 1617 Napier made some observations upon the subject and wrote both 1993,273 and 1993, 2′ 7″ 3‴ for the number which we now, in America, write 1993.273. In the Leyden edition of this work (1626) the

[1] Cantor, *Geschichte*, II (2), 617. [2] *Ibid.*, p. 619.

[3] *Canon Triangulorum Emendatissimus et ad usum accommodatissimus. Pertinens ad Trigonometriam Bartholomaei Pitisci . . .*, Frankfort, 1612.

[4] "Deinde pro latere AC nuper invento 13|00024 assumo 13 fractione scilicet $13\frac{24}{100000}\cdots$." *Trig. Problematum Geod. Liber Unus*, p. 12.

[5] As in this subtraction (p. 67):

70°. 0′
46. 8
23. 52′

Stevin notation is used, by which the above number would appear as 1993, ⓪2,①7,②3.[1] In any case it is evident that Napier understood something about the decimal fraction, that he did not invent our modern symbolism, and that the practical use of logarithms soon made a knowledge of decimals essential.

In 1616 Kepler wrote a work on mensuration[2] in which he distinctly took up the decimal fraction, using both a decimal point (comma) and the parentheses to separate the fractional part.[3] He stated it as his opinion that these fractions were due to Bürgi,[4] although it seems strange that he was not familiar with the work of Stevin. In his edition of Tycho Brahe's *Tabulae Rudolphinae*, published at Ulm in 1627, he uses (p. 25) the period for a decimal point, thus: "29.032 valet $29\frac{32}{1000}$."

In the year 1616 Johann Hartmann Beyer (1563–1625) wrote to Kepler concerning his work, and in the letter he used both the decimal comma and the sexagesimal symbolism for the decimal, writing 314, 1′ 5″ 9‴ 2⁗ 6⁗′ 5⁗″ + for 314.15926 +. Beyer had before this (1603) published a work on these fractions, *Logistica decimalis*, and on this account had laid claim to their invention, although he had long been preceded by Stevin. Adriaen Metius (1571–1635) took about the same step in symbolism when he wrote both 47852°:8′0″4‴ and 47852/8′0″4‴ for 47852.804. He also spoke[5] of "$\frac{4814841}{10000000}$ ofte 4′8″1‴4⁗8⁗′1⁗″."

[1] C. G. Knott, *Napier Memorial Volume*, pp. 77, 182, 188, 190, 191. Edinburgh, 1914.

[2] *Ausszug auss der uralten Messe-Kunst Archimedis*. It appears in Volume V of the Frisch edition of Kepler's works, 1864.

[3] "Fürs ander, weil ich kurtze Zahlen brauche, derohalben es offt Brüche geben wirdt, so mercke, dass alle Ziffer, welche nach dem Zeichen (,) folgen, die gehören zu dem Bruch, als der zehler, der nenner dazu wirt nicht gesetzt." He then gives an example in interest:

3,65
6 mal
facit 21 (90)

[4] "Dise Art der Bruchrechnung ist von Jost Bürgen zu der sinusrechnung erdacht."

[5] *Opera Omnia*, 1633, pp. 19, 31, 49, 50. When De Morgan (*Arithmetical Books*, p. 41) said of a 1640 edition of Metius that "sexagesimal fractions are taught, but not decimal ones," he may have confused the symbols.

There are numerous examples of writers of the same period who used these awkward symbols. Girard, the editor of Stevin, whose first edition of the latter's works appeared in 1625, did much to make known the works of his master, but he apparently added nothing to the theory or the symbolism. Even as late as 1655 we find the period used to separate an integer and common fraction,[1] as in the case of $198.\frac{1}{2}$, and in the 1685 edition of Casati's work[2] we have 0.00438 represented by

$$\frac{438}{100000}$$

In 1657 Frans van Schooten[3] used the symbol 17579625 . . . ③ for 17579.625. It had the advantage that, in finding the product of two decimals, the indices in the circles need only be added in order to determine the proper index in the result.

The use of smaller type for the decimal part was not uncommon,[4] and it is still seen on the continent of Europe. As to the development of these fractions in England, Professor Cajori has suggested that Oughtred's (1631) use of the symbol $0\,|\,\overline{56}$ for 0.56 was one of the causes for delay in the general adoption of the decimal point.

It should also be said that the symbolism is by no means settled even yet. In England $23\frac{45}{100}$ is written[5] $23\cdot45$, in the United States it appears as 23.45, and on the Continent such forms as 23,45 and 23_{45} are common. Indeed, in America we commonly write $\$23.\overline{45}$ or $\$23\frac{45}{100}$ instead of \$23.45, to avoid forgery.

[1] B. Capra, *Vsvs et fabrica circini* . . ., p. 25 (Bologna, 1655). The first edition, however, seems to have been 1607.

[2] P. Casati, *Fabrica et vso Del Compasso di Proportione*, p. 123. Bologna, 1685. In the first edition (1664), however, he writes such a number (p. 86) as a common fraction, with the bar between the terms.

[3] *Exercitationum Mathematicarum Libri quinque*, liber primus, p. 33. Leyden, 1657.

[4] *E.g.*, in some editions of Vieta's tables; also in R. Butler, *The Scale of Interest* (London, 1630), where 1_{125} is used for 1.125.

[5] Not always, however. In a MS. at the Woolwich Academy, of date 1736, the decimal point is always a comma. Hodder wrote a *Decimal Arithmetick* in 1668, in which he used both the comma and the dot.

Summary. The historical steps in the invention of the decimal fraction may be summed up as follows: Pellos (1492) used a decimal point where others had used a bar, but the idea of the decimal fraction was not developed by him. Rudolff (1530) worked intelligently with decimal fractions, using a bar for the separatrix, but he did not write upon the theory. Stevin (1585) wrote upon the theory but had a poor symbolism. About 1600, several writers attempted to improve the symbolism, and Bürgi, in 1592, actually used a comma for the decimal point, without the common sexagesimal marks, and comprehended the nature and advantages of these fractions. Napier knew something of the theory of decimals and rendered their use essential, but did not himself contribute to the symbolism.[1] In the mere writing of the decimal fraction, at least, all these efforts had been anticipated by al-Kashî (*c.* 1430), whose symbolism was quite as good as that of any European writer for the next century and a half.

It is thus difficult to pick out the actual inventor, although Rudolff and Stevin are entitled to the most credit for bringing the new system to the attention of the world. It should be added that these fractions were mentioned by Richard Witt in his *Arithmeticall Questions* in 1613, and that Henry Lyte (1619) wrote *The Art of Tens, or Decimall Arithmeticke*,—a work which did for England what the work of Stevin had done for the Continent.

Percentage. Long before the decimal fraction was invented the need for it was felt in computations by tenths, twentieths, and hundredths, and this need gave rise to a peculiar notation which took the place of the decimal forms and which has persisted to the present time in the symbol %.

The computations of the Romans that led up to the subject of percentage may be illustrated by the *vicesima libertatis*, a tax of $\frac{1}{20}$ on every manumitted slave; by the *centesima rerum venalium*, a tax of $\frac{1}{100}$ levied under Augustus on goods sold at

[1] How imperfect his knowledge was may be seen by examining his *De arte logistica*, pp. 60, 65, 75, *et passim*.

auction; and by the *quinta et vicesima mancipiorum*, a tax of $\frac{1}{25}$ on every slave sold.[1] Without recognizing per cents as such, the Romans thus made use of fractions which easily reduce to hundredths.

In the Middle Ages, both in the East[2] and in the West, there was a gradual recognition of larger denominations of money than the ancients had commonly known, and this led to the use of 100 as a base in computation. In the Italian manuscripts of the 15th century it is common to find examples involving such expressions as 20 ℔ 100, x℔ cento, and vi ℔ c°, for our 20%, 10%, and 6%.

When commercial arithmetics began to be printed, this custom was well established, and so in Chiarino's work of 1481 there are numerous expressions like "xx. per .c." for 20%, and "viii in x percēto" for 8 to 10%. Borghi (1484) and Pellos (1492) made less use of per cents than one would expect of such commercial authorities, although each recognized their value.[3] The demand was growing, however, and Pacioli (1494), familiar with the large commerce of the *giudecca* at Venice, had much to say of it.[4] Beginning early in the 16th century the commercial arithmetics made considerable use of per cents in connection with interest and with profit and loss, sometimes in relation to the Rule of Three,[5] so popular with merchants of that period, but more frequently in relation to isolated problems.[6]

[1] Harper's *Dict. Class. Lit.*, p. 1634.

[2] *E.g.*, Bhāskara (*c.* 1150) uses per cents in the interest problems in the *Lilāvati*. See Taylor, *Lilawati*, p. 47; Colebrooke, *Lilāvati*, p. 39.


[3] *E.g.*, Borghi: ". . . guadagno a rason de .20. per cento"; and Pellos: ".12. ℔ .100.," "p .3. ans a rason de .16. ℔ cent."

[4] His printed forms include "℔ cēto" (fol. 65), ".10. ℔ cento" (fol. 66), and "per cēto" (fol. 66).

[5] Thus Ortega (1512; 1515 ed.) has a chapter on "Regvla de tre de centenare" (fol. 51).

[6] Thus Wålckl (1536): "Itē einer leihet einem 200 fl. 3 iar vnd eines ieden iars nimbt er 10fl vō 100 ist die frag wieuil die 3 iar thůt gwiñ vnd gwiñsgwinn" (fol. B 7). So, also, Rudolff (1530; 1540 ed.): "Weñ man vom hundert zu jårlichem zins geben sol 5 flo . . ." (Ex. 71 in the *Exempel-Büchlin*). He shows that the Italian "pro cento" was not yet well known in Germany, for he says (Ex. 156): ". . . vnd wieuil pro cento (verstee an Hundert floren)."

In America at present the expression 6% is identical in meaning with 0.06, per cent having come to signify merely hundredths. This was not the original meaning, nor does it conform to the present usage in England and certain other countries, where expressions like "£6 per cent" are in common use. This usage is historically correct, the 15th and 16th century writers, with whom percentage begins in any large way, having always employed it.[1]



EARLY PER CENT SIGN

From an Italian MS. of *c.* 1490. Notice also the old symbol for pounds, which may have suggested the dollar sign


Chief Use for Per Cents. The chief use for per cents in the 16th century was in relation to the computation of interest, and by the beginning of the 17th century the rate was usually quoted in hundredths.[2] It also appears in computing profit and loss, at first indirectly, as in the following addition to Recorde by John Mellis (1594): "If one yard cost 6s——8 pence:

[1] Thus Sfortunati (1534; 1545 ed.) uses "libre .30. per 100"; Riese (1522) uses "10 lb. von 100," "10 fl zum 100," "10 fl am 100," and other similar forms; and Albert (1534) has "10 fl mit 100 fl."

[2] *E.g.*, Trenchant (1566) has "à raison de 12 pour 100" with a 12% interest table; Petri (1567) speaks of "8 ten hondert" and "12 ten 100"; Raets (1580) gives the rate as "15 ten hondert," and Wentsel (1599) as "10. ten 100 tsjaers,"—all of which shows the high rates of interest prevailing.

and the same is sold againe for 8 s̄——6 pence: the question is, what is gayned in 100 pounds laying out on such commoditie." Many books, however, stated the problems substantially as at present.

The Per Cent Sign. In its primitive form the per cent sign (%) is found in the 15th century manuscripts on commercial arithmetic, where it appears as "per c̊" or "p c̊," a contraction



THE PER CENT SIGN IN THE 17TH CENTURY

From an anonymous Italian MS. of 1684

for "per cento."[1] As early as the middle of the 17th century it had developed into the form "per $\frac{0}{0}$," after which the "per" finally dropped out. The solidus form (%) is modern.

Permillage. It is natural to expect that percentage will develop into permillage, and indeed this has not only begun, but it has historic sanction. Bonds are quoted in New York "per M," and so in various other commercial lines. This was already common in the 16th century.[2] At present, indeed, the symbol $\frac{0}{00}$ is used in certain parts of the world, notably by German merchants, to mean per mill, a curious analogue to % developed without regard to the historic meaning of the latter symbol.

[1] *Rara Arithmetica*, pp. 439, 441, 458, with facsimiles.

[2] Thus Cardan (1539) says that "tara cōputāda est ad 100. vel ad 1000." *Arithmetica*, 1539, capp. 57, 59.

4. Surd Numbers

Ancient Idea of Irrationals. Proclus (*c.* 460) tells us that the Pythagoreans discovered the incommensurability of the diagonal and the side of a square,[1] which is only a geometric view of the irrationality of $\sqrt{2}$. Proclus also states that they were led to study the subject of commensurability through their work with numbers. Plato says[2] that Theodorus of Cyrene (*c.* 425 B.C.) discovered that "oblong numbers, 3, 5, 6, 7, $\cdots$, are composed of unequal sides." He also states that "Theodorus was writing out for us something about roots, such as the roots of three or five, showing that they are incommensurable by the unit: he selected other numbers up to seventeen—there he stopped."[3]

With respect to other writers on incommensurable lines, Diogenes Laertius (2d century) tells us that Democritus (*c.* 400 B.C.) composed a treatise upon the subject.[4]

Summary of Greek Ideas on Irrationals. Summarizing the work of the Greeks, there seems to be good reason for believing that the immediate followers of Pythagoras knew and demonstrated the incommensurability of the diagonal and the side of a square, but that they looked upon this case of irrationality as a peculiarity of the square. Theodorus seems to have carried the investigation farther, recognizing that irrationality of square roots was not confined to $\sqrt{2}$. Theætetus (*c.* 375 B.C.) appears to have laid the foundations for a general theory of quadratic irrationals[5] and to have established their leading

[1] Heath's *Euclid*, Vol. III, p. 1 seq., to the notes of which the reader is referred. See also Cantor, *Beiträge*, p. 108. The proof is given in Euclid's *Elements*, numbered X, 117 in early editions, but is now relegated to an appendix. See also H. Vogt, "Die Entdeckungsgeschichte des Irrationalen nach Plato . . .," *Bibl. Math.*, X (3), 97; Heath, *History*, I, 65, 90, 154.

[2] *Theætetus*, 147 D; Jowett translation, IV, 123; Heath, *History*, I, 203.

[3] It should be observed that the method of proof for $\sqrt{3}$ is quite different from that for $\sqrt{2}$, and so for other surds. See Heath, *Euclid*, Vol. III, p. 2, and *History*, I, 155.

[4] Περὶ ἀλόγων γραμμῶν καὶ νασταῖν β′. See F. Hultsch, *Neue Jahrbücher für Philologie und Pädagogik*, CXXIII, 578; Heath, *History*, I, 156.

[5] Heath, *History*, I, 209.

properties. Euclid (*c.* 300 B.C.) took the final important step due to the Greek geometers, classifying square roots and introducing the idea of biquadratic irrationals.[1]

This discovery, then, was the second noteworthy step in the creation of types of artificial numbers. The Greeks showed that all magnitudes are either rational (*ῥητά, rheta'*) or irrational (*ἄλογα, a'loga*), their idea of an irrational number being such a number as cannot be expressed as the ratio of two integers.

The geometric treatment of incommensurables naturally led to the arithmetic and algebraic treatment of irrationals, the subject of the present discussion.[2]

The Name "Surd." Al-Khowârizmî (*c.* 825) spoke of rational numbers as "audible" and of surds as "inaudible,"[3] and it is the latter that gave rise to the word "surd" (deaf, mute). So far as now known, the European use of this word begins with Gherardo of Cremona (*c.* 1150).[4] The term was also used by Fibonacci (1202), but to represent a number that has no root.[5] The Arabs and Hebrews often called surds "nonexpressible numbers,"[6]—a name which may have suggested the "inexplicable sides" of the Renaissance writers.[7] It is simply a translation from the Euclidean term *ἄλογος* (*a'logos*, without ratio, irrational, incommensurable).

As to what constitutes a surd, however, there has never been a general agreement. It is admitted that a number like $\sqrt{2}$

[1] Vogt, *loc. cit.* On Professor Zeuthen's discussion of Vogt's conclusions, see H. Bosmans, in the *Revue des Questions Scientifiques*, July, 1911. See also Heath, *History*, I, 402.

[2] On the history of transcendental numbers see the statement on page 268.

[3] Rosen ed., p. 192. [4] *Bibl. Math.*, I (3), 516.

[5] "Nam quidem numeri habent radices, et uocatur [*sic*] quadrati; et quidam non; quorum radices, que surde dicuntur, cum inpossibile sit eas in numeris inuenire . . . " (*Liber Abaci*, p. 353). By "root" he refers here, as usual, to square root.

[6] *E.g.*, al-Karkhî (*c.* 1020); see Hochheim, *Kâfî fil Hisâb*, II, 12.

[7] *E.g.*, Schoner, in his *De numeris figuratis* (1569; 1586 ed., p. 213), says: "Explicabile latus est, cujus ad 1. ratio explicari potest. Ut latus 4 est 2, & dicitur explicabile. . . . Inexplicabile latus contra est, cujus ad 1. ratio explicari non potest. Ut 3. . . ."

Stevin (1585) speaks of "nombres, comme $\sqrt{8}$, & semblables, qu'ils appellent absurds, irrationels, irreguliers, inexplicables, sourds, &c" (1634 ed., p. 9).

is a surd, but there have been prominent writers who have not included $\sqrt{6}$, since $\sqrt{6} = \sqrt{2} \times \sqrt{3}$;[1] and $\sqrt{2+\sqrt{3}}$ is commonly excluded.[2]

Approximate Values. An interest in the irrational showed itself strongly among the ancients. Here was a mystery to be fathomed, and from the time of Pythagoras to that of Weierstrass the nature of irrationals and the ability to work with them occupied the attention of a considerable part of the mathematical world. Among the noteworthy efforts was the one which sought to find an approximate value for an expression like $\sqrt{2}$. As already said (p. 144), the Greeks found the square root of a number in much the same way as that which is commonly taught in school today, but their ignorance of the decimal fraction made the process of approximation very difficult in the case of surds. For this reason the ancient and medieval writers resorted to various rules which can best be appreciated by first considering the principle involved.

Let a be an approximation to $\sqrt{A}$ by defect. Then A/a must be an approximation by excess, and the arithmetic mean, $a_1 = \frac{1}{2}\left(a + \frac{A}{a}\right)$, is an approximation of the second order by excess,[3] and the harmonic mean, A/a_1, is an approximation of the second order by defect. This process may evidently be carried on indefinitely. If $A = a^2 + r$, we have in particular,

$$a_1 = a + \frac{r}{2a}, \qquad \frac{A}{a_1} = a + \frac{r}{2a + \frac{r}{a}},$$

[1] *E.g.*, Behâ Eddîn (*c.* 1600), al-Karkhî (*c.* 1020), and other Arab writers included only non-squares not divisible by the digits 2, 3, . . . , 9. See Hochheim, *Kâfî fil Hisâb*, II, 13 n.

[2] G. Chrystal, *Algebra*, 2d ed., I, 203 (Edinburgh, 1889): ". . . a surd number is the incommensurable root of a commensurable number. . . . For example . . . $\sqrt{e}$ is not a surd. . . . Neither is $\sqrt{(\sqrt{2}+1)}$."

[3] For $\frac{x^2}{x-n} = x + n + \frac{n^2}{x-n}$, so that if we divide by a number that is n less than the square root, we shall have a result that is more than n in excess of the square root.

and so on. Recognizing that

$$a_1 = \tfrac{1}{2}\left(a + \frac{A}{a}\right) = a + \frac{r}{2a}$$

is an approximation by excess, the Arabs took

$$a_1' = a + \frac{r}{2a+1}$$

as an approximation by defect, but this rule is not found among the Greeks.[1]

Of the various rules for approximation to $\sqrt{A}$, the one most commonly used in the past may be expressed as

$$\sqrt{A} = \sqrt{a^2 + r} = a + \frac{r}{2a},$$

as in

$$\sqrt{10} = \sqrt{9+1} = 3 + \frac{1}{2\times 3} = 3\tfrac{1}{6},$$

an approximation by excess. The corresponding approximation by defect is:

$$\sqrt{A} = \sqrt{a^2 + r} = a + \frac{r}{2a+1},$$

as in

$$\sqrt{10} = \sqrt{9+1} = 3 + \tfrac{1}{7} = 3\tfrac{1}{7},$$

which probably explains why $\sqrt{10}$ was so often used for π by early writers. Of these approximations, the one by excess is found in the works of Heron (*c.* 50?).[2] The medieval writers used both of these approximations,[3] often with variations. For

[1] P. Tannery, "L'extraction des racines carrées d'après Nicolas Chuquet," *Bibl. Math.*, I (2), 17; "Du rôle de la musique grecque dans le développement de la mathématique pure," *ibid.*, III (3), 171.

[2] A fact noted by Clavius, *Epitome*, 1583; 1585 ed., p. 318. It should occasionally be repeated that, as stated in Volume I, page 125, this date is uncertain. Heron may have lived as late as the 3d century.

[3] *E.g.*, al-Ḥaṣṣâr (*c.* 1175?) made use of both the one by defect and the one by excess, together with $a + \frac{r+1}{2a+2}$ and $a + \frac{r}{2a} - \frac{\left(\frac{r}{2a}\right)^2}{2\left(a + \frac{r}{2a}\right)}$ for closer work.

For other cases see Hochheim, *Kâfî fîl Hisâb*, II, 14; Wertheim, *Elia Misrachi*, p. 21; B. Boncompagni, *Atti Pontif.*, XII, 402.

example, Rhabdas (*c.* 1341), following an Arabic method,[1] obtained a first approximation to $\sqrt{10}$ by using a rule equivalent to

$$\sqrt{A} = a + \frac{A - a^2}{2a},$$

giving

$$\sqrt{10} = 3 + \frac{10 - 9}{6} = 3\tfrac{1}{6} = \frac{19}{6}.$$

Then, since $10 \div \frac{19}{6} = 3\frac{3}{19}$, he takes the mean of $3\frac{1}{6}$ and $3\frac{3}{19}$, which is $3\frac{3}{22}\frac{7}{8}$. This he shows to be a close approximation, since $(3\frac{3}{22}\frac{7}{8})^2 = 10\frac{1}{5}\frac{1}{9}\frac{1}{8}\frac{1}{4}$.

A somewhat different method, also involving averages, is given by Chuquet (1484). Let two approximate values of $\sqrt{A}$ be $\frac{p_0}{q_0}$ and $\frac{p_1}{q_1}$, the first being too great and the second being too small, and let $p' = p_0 + p_1$ and $q' = q_0 + q_1$. Then p'/q' is a new approximation intermediate in value, and whether it is by excess or defect is found by squaring. In the same way an approximation is found between p'/q' and one of the others, and so on. This rule was employed by several later writers.[2] De la Roche (1520), who plagiarized Chuquet, asserted that any study of "imperfect roots" was useless, although custom required it.[3] Such approximations are common in the works of the 16th century, together with similar rules for cube root.[4] With all this there naturally developed many evidences of ignorance, as when Peter Halliman (1688) gave substantially the rule

$$\sqrt[n]{a^n + r} = a + \frac{r}{(a+1)^n - r^n},$$

[1] P. Tannery, *Notices et extraits des manuscrits de la Bibl. nat.*, XXXII, 185.

[2] *E.g.*, Ortega (1512); substantially by Clavius (*Epitome*, 1583; 1585 ed., p. 318, where he gives $\sqrt{20} = 4\frac{5}{1}\frac{4}{5}\frac{7}{9}\frac{3}{2}$, approximately); and substantially by Metius when he found the value of π by interpolating between $3\frac{1}{2}\frac{7}{0}$ and $3\frac{1}{0}\frac{5}{6}$, the result being the Chinese value, $3\frac{5}{1}\frac{5}{3}$. See P. Tannery, *Bibl. Math.*, I (2), 17.

[3] The study of "racines imparfaites" was "labeur sans vtile," but "pour la perfection de ce liure" he gave a method "par la regle de mediation entre le plus et le moins,"—an elementary method of interpolation. This is described by Treutlein in the *Abhandlungen*, I, 66.

[4] Thus Stevin gives substantially the rule that

$$\sqrt[3]{A} = \sqrt[3]{a^3 + r} = a + \frac{r}{3a(a+1) + 1}.$$

Œuvres, 1634 ed., p. 30.

celebrating his discovery by the doggerel verse,

Now Logarithms lowre your sail,
And Algebra give place,
For here is found, that ne'er doth fail,
A nearer way, to your disgrace.[1]

It should also be understood that such rules for roots are ancient. For example, Heron (*c.* 50?) gives what may possibly be the equivalent of the formula

$$\sqrt[3]{n} = b + \frac{a(n-b^3)}{n+a(n-b^3)},$$

where $a > \sqrt[3]{n} > b$, and $a - b = 1$. By means of this rule we should find that $\sqrt[3]{109} = 4.7785$ instead of 4.7769.[2]

Criteria for Squares. In order to determine whether or not $\sqrt{a}$ is a surd, those who were interested in number theory developed from time to time criteria for ascertaining whether a number is a square. Such criteria are found in various ancient and medieval works, both Arab[3] and European. A Munich manuscript[4] of the 15th century, for example, states that if a square ends in an even number, it is divisible by 4;[5] that if it ends in zeros, it ends in an even number of zeros;[6] that it cannot end in 2, 3, 7, or 8;[7] and that every square is of the form $3n$ or $3n+1$.[8] Such rules, often extended, are found in various works of the classical and Renaissance periods.[9]

[1] From *The Square and Cube Root compleated and made easie* (London, 1688), quoted by A. De Morgan, *Arithmetical Books*, p. 52.

[2] J. G. Smyly, "Heron's formula for cube root," *Hermathena*, XLII, 64, correcting M. Curtze, *Zeitschrift*, Hl. Abt. (1897), p. 119, and referring to Heron's *Metrica*, III, 20. See also the interpretations in Heath, *History*, II, 341.

[3] *E.g.*, al-Karkhî (*c.* 1020) and al-Qalasâdî (*c.* 1475). See Hochheim, *Kâfî fil Hisâb*, II, 13.

[4] No. 14,908, described by Curtze in *Bibl. Math.*, IX (2), 38.

[5] "Omnis quadratus, cuius prima est par, est per 4 divisibilis."

[6] "Omnis quadratus in primis locis habet parem numerum ciffrarum."

[7] "Nullus quadratus recipit in primo loco 2, 3, 7 vel 8, sed alios bene." This is a very old rule.

[8] "Omnis quadratus est simpliciter vel subtracte unitate per 3 divisibilis."

[9] Thus Buteo (1559) adds that a square number cannot end in 5 unless it ends in 25.

Surds in Algebra. The placing of the study of surds in the books on algebra is a tradition which began with the Renaissance. The books on logistic, used in commercial schools, had no need for the subject; it properly belonged in the books on the theory of numbers, the ancient *arithmetica*. Since, however, algebra took over a considerable part of the latter in the Renaissance period, surds found a place in this science. Furthermore, since these forms are needed in connection with irrational equations, they were usually considered before that topic in the study of algebra. In the 15th century, however, they are often found in the theoretical arithmetics.[1]

5. Negative Numbers

Early Use of Negative Numbers. No trace of the recognition of negative numbers, as distinct from simple subtrahends, has yet been found in the writings of the ancient Egyptians, Babylonians, Hindus, Chinese, or Greeks. Nevertheless the law of signs was established, with the aid of such operations as $(10-4)\cdot(8-2)$, and was known long before the negative number was considered by itself.

The Chinese made use of such numbers as subtrahends at a very early date. They indicated positive coefficients by red computing rods, and negative ones by black, and this color scheme is also found in their written works.[2] The negative number is mentioned, at least as a subtrahend, in the *K'iu-ch'ang Suan-shu* (*c.* 200 B.C.),[3] and in various later works, but the law of signs is not known to have been definitely stated in any Chinese mathematical treatise before 1299, when Chu Shï-kié gave it in his elementary algebra, the *Suan-hio-ki-möng* (*Introduction to Mathematical Studies*).

[1] Thus the Rollandus MS. (*c.* 1424) has surds in the arithmetic just before algebra is begun, and similarly in Pacioli's *Sūma* (1494). On the modern problem consult the *Encyklopädie der mathematischen Wissenschaften*, I, 49 (Leipzig, 1898–); hereafter referred to as *Encyklopädie*.

[2] Mikami, *China*, pp. 18, 20, 21, 27, 89, *et passim*; Cantor, *Geschichte*, I (2), 642.

[3] See Volume I, page 31, for discussion as to earlier date for the original work. It may have been written before 1000 B.C.

The first mention of these numbers in an occidental work is in the *Arithmetica* of Diophantus (*c.* 275),[1] where the equation $4x + 20 = 4$ is spoken of as absurd (ἄτοπος), since it would give $x = -4$. Of the negative number in the abstract, Diophantus had apparently no conception. On the other hand, the Greeks knew the geometric equivalent of $(a - b)^2$ and of $(a+b)\ (a-b)$; and hence, without recognizing negative numbers, they knew the results of the operations $(-b)\cdot(-b)$ and $(+b)\cdot(-b)$.

In India the negative number is first definitely mentioned in the works of Brahmagupta (*c.* 628). He speaks of "negative and affirmative quantities,"[2] using them always as subtrahends but giving the usual rules of signs. The next writer to treat of these rules is Mahāvīra (*c.* 850), and after that time they are found in all Hindu works on the subject.

The Arabs contributed nothing new to the theory, but al-Khowârizmî (*c.* 825) states the usual rules,[3] and the same is true of his successors.

When Fibonacci wrote his *Liber Abaci* (1202) he followed the Arab custom of paying no attention to negative numbers, but in his *Flos* (*c.* 1225) he interpreted a negative root in a financial problem to mean a loss instead of a gain.[4] Little further was done with the subject by medieval writers, but as we approach the Renaissance period we find the negative number as such receiving more and more recognition. For example, among the problems set by Chuquet (1484) is one[5] which leads to an equation with roots "$\tilde{m}.7.\frac{3}{11}$" and "$27.\frac{3}{11}$," that is, $-7\frac{3}{11}$ and $27\frac{3}{11}$.

Modern Usage. The first of the 16th century writers to give noteworthy treatment to the negative number was Cardan. In

[1] Nesselmann, *Alg. Griechen*, p. 311; Heath, *Diophantus*, 2d. ed., pp. 52, 200 (Cambridge, 1910); Cantor, *Geschichte*, I (2), 441.

[2] Colebrooke translation, pp. 325, 339.

[3] Rosen translation, p. 26.

[4] "Hanc quidem quaestionem insolubilem esse monstrabo, nisi concedatur, primum hominem habere debitum." *Scritti*, II, 238.

[5] Boncompagni's *Bullettino*, XIV, 419, Ex. xiv. Chuquet adds, "Ainsi ce calcule est vray que aulcuns tiennent Impo.ble"

his *Ars Magna* (1545) he recognized negative roots of equations and gave a clear statement of the simple laws of negative numbers.[1]

Stifel (1544) distinctly mentioned negative numbers as less than zero,[2] and showed some knowledge of their use. By this time the rules of operation with numbers involving negative signs were well understood, even though the precise nature of the negative number was not always clear. Thus Bombelli (1572) gave these rules and applied them intelligently to such cases as $(+15)+(-20)=-5$.[3] It was due to the influence of men like Vieta, Harriot, Fermat, Descartes, and Hudde, however, that the negative number came to be fully recognized and understood. The idea of allowing a letter, with no sign prefixed, to represent either a positive or a negative number seems due to Hudde (1659).

Names and Symbols for Negative Numbers. The Hindu writers who mentioned negative numbers, or numbers used as subtrahends, placed a dot or a small circle over or beside each, as stated on page 396. The names used were the equivalent of our word "negative." The early European usage has already been mentioned, but it remains to speak of the establishing of our modern terminology.

As already stated, the Chinese wrote positive numbers in red and negative numbers in black, and so indicated them by their stick symbolism. They also had another method for indicating negative coefficients, one that may have been due to Li Yeh (1259). This consisted in drawing a diagonal stroke through the right-hand digit figure of a negative number, as in the case of 𝍠 for $-10{,}724$, and of 𝍠 for $-10{,}200$.[4]

In the 15th century the names "positive" and "affirmative" were used to indicate positive numbers, as also "privative"

[1] Thus on fol. 3, *v.*, speaking of squares, he says: "At uero quòd tam ex 3, quàm ex m: 3, fit 9, quoniam minus in minus ductū ꝑducit plus."

[2] "Finguntur numeri infra o, id est, infra nihil." *Arithmetica Integra*, fol. 249, *r.*

[3] "Più via più fà più. Meno via meno fà più" etc. (p. 70). Also: "E p.15 con m.20 fà m.5. perche se io mi trouassi scudi 15, e ne fossi debitore 20, pagati li 15 restarei debitore 5" (p. 72).

[4] Mikami, *China*, p. 82.

and "negative" for negative numbers,[1]—a usage followed by Scheubel (1551).[2]

Cardan (1545) spoke of "minus in minus" as being plus, but in general he called positive numbers *numeri ueri*[3] and negative numbers *numeri ficti*.[4] His symbol for a negative number is simply m:, as in the case of m:3 for -3.[5]

Stifel (1544) called these numbers "absurd" and wrote $0-3$ as an illustration.[6]

Tartaglia (1556) spoke of a negative number as "the term called minus,"[7] laying down the usual rules.[8]

Bombelli (1572) used the word "minus" (*meno*) as we do in such rules as "minus times minus gives plus," his symbol for -5 being m.5. Unlike Cardan, he had a definite sign for $+5$ also, writing it p.5.

Tycho Brahe, the astronomer (1598), spoke of the negative number as "privative" and indicated it by the minus sign.[9]

Napier (*c.* 1600) used the adjectives *abundantes* and *defectivi* to designate positive and negative numbers, Sturm (1707) spoke of *Sache* and *Mangel*, and various other names and symbols have been suggested.[10]

[1] H. E. Wappler, *Zur Geschichte der deutschen Algebra im XV. Jahrhundert*, Prog., p. 31 (Zwickau, 1887). [2] Tropfke, *Geschichte*, II (2), 79.

[3] Or *ueri numeri*. He used both forms, as on fol. 3, *v.*, of the *Ars Magna*.

[4] So in speaking of the roots of an equation he says "una semper est rei uera aestimatio, altera ei aequalis, ficta."

[5] So he gives the roots of $x^2 = 16$ thus: "res est 4, uel m: 4."

[6] "Finguntur numeri minores nihilo ut sunt $0-3$" (*Arithmetica Integra*, fol. 48, *r.*). Later (fol. 249, *v.*) he speaks of zero as "quod mediat inter numeros veros et numeros absurdos."

[7] ". . . il termine chiamato men" (*General Trattato*, II, fol. 83, *r.*).

[8] *E.g.*, "Terzo regola, a multiplicare men fia men fa sempre piu" (*ibid.*, fol. 85, *v.*). His illustration is as follows:

a multiplicar	9 m̃ 2
per	8 m̃ 3
fa	72 men 43 piu 6
che sara	35 a ponto

[9] ". . . quod eorum alii positivi sunt alii privativi; positivi ii quibus vel nullum signum est additum vel praefigi debet hoc+ ; privativi verò qui praefixum habere debent signum hoc —" (*Tabulae Rudolphinae*, p. 9 (Ulm, 1627)).

[10] On the later treatment of the negative number see Cantor, *Geschichte*, IV, 79–88.

6. Complex Numbers

Early Steps in Complex Numbers. The first trace of the square root of a negative number to be found in extant works is in the *Stereometria* of Heron of Alexandria (*c.* 50?), where $\sqrt{81-144}$ is taken to be $\sqrt{144-81}$, or $8-\frac{1}{16}$. The problem involved is impossible of solution, and this step should have been left $\sqrt{-63}$ or $3\sqrt{-7}$,[1] but whether the error is due to Heron or to some copyist is uncertain.

The next known recognition of the difficulty is found in the *Arithmetica* of Diophantus[2] (*c.* 275). In attempting to compute the sides of a right-angled triangle of perimeter 12 and area 7, Diophantus found it necessary to solve the equation $336x^2+24=172x$. He stated that the equation cannot be solved unless the square of half the coefficient of x diminished by 24×336 is a square, not otherwise seeming to notice that this equation has complex roots.

Mahāvīra (*c.* 850) was the first to state the difficulty clearly, saying, in his treatment of negative numbers, that, "as in the nature of things a negative [quantity] is not a square [quantity], it has therefore no square root."[3]

Bhāskara (*c.* 1150) used about the same language in his *Bija Gaṇita*:

> The square of an affirmative or of a negative quantity is affirmative; and the square root of an affirmative quantity is two-fold, positive and negative. There is no square-root of a negative quantity: for it is not a square.[4]

The Jewish scholar Abraham bar Chiia (*c.* 1120) set forth the same difficulty in discussing the equations $xy=48$ and $x+y=14$.[5]

[1] On this general topic see W. W. Beman, "A Chapter in the History of Mathematics," vice-presidential address in Section A, *Proc. of the American Assoc. for the Adv. of Sci.*, 1897; E. Study, in the *Encyklopädie der Math. Wissensch.*, I, A, 4 (Leipzig, 1898); H. Hankel, *Vorlesungen über die complexen Zahlen*, Leipzig, 1867; G. Loria, *Scientia*, XXI, 101; F. Cajori, *Amer. Math. Month.*, XIX, 167. On Heron, see the Schmidt edition (Leipzig, 1914), V, 35.

[2] Heath, *Diophantus*, 2d ed., p. 244 (Cambridge, 1910).

[3] *Gaṇita-Sāra-Saṅgraha*, p. 7.

[4] Colebrooke's translation, p. 135.

[5] *Abhandlungen*, XII, 46.

The Arabs and Persians seem to have paid no special attention to the subject, and the next step was taken in Italy and after the invention of printing.

Early European Efforts. Pacioli (1494) stated in his *Sūma* that the quadratic equation $x^2 + c = bx$ cannot be solved unless $\frac{1}{4}b^2 \geqq c$,[1] so that he recognized the impossibility of finding the value of $\sqrt{-a}$. About the same time Chuquet (1484) seems to have found that $\sqrt{-a}$ represents an impossible case.[2]

Cardan (1545) spoke of the equation $x^4 + 12 = 6x^2$ as being impossible,[3] referring to the roots of such equations as *ficta* or *per m̃.* He was the first to use the square root of a negative number in computation, the problem being to divide 10 into two parts whose product is 40.[4] He found the number to be $5 + \sqrt{-15}$ and $5 - \sqrt{-15}$, spoke of the solution "by the minus root,"[5] and proved by multiplication that his results were correct.

The next to attack the problem was Bombelli (1572). In his algebra[6] he speaks of such quantities as $+\sqrt{-a}$ and $-\sqrt{-a}$, but he made no advance upon Cardan's theory.

Stevin (1585) noted the difficulty of working with imaginaries, but could only remark that the subject was not yet mastered.[7] Girard (1629) found it necessary to recognize complex roots in order to establish the law as to the number of roots of an equation.[8]

[1] Fol. 147, *r.* *E.g.*, $x^2 + 7 = x + 5$: "Dico q̄sto essere impossibile." As in all such cases, the symbolism here shown is modern.

[2] Boncompagni's *Bullettino*, XIV, 444.

[3] "Quòd si caruerit estimatiõe uera, carebit etiam ea, quę est per m: uelut 1 q̄d^{ti} q̄d^{m} p: 12, aeq̄tur 6 q̄d^{tis}, quia non potest aequationẽ ueram habere, carebit etiam ficta, sic eã uocamus eam, quae debiti est seu minoris" (*Ars Magna*, fol. 3, *v.*).

[4] *Opera*, IV, 287. Lyons, 1663.

[5] "Per radicem m̃."

[6] *L'Algebra*, p. 294 seq. Bologna, 1572.

[7] *Œuvres*, 1634 ed., pp. 71, 72.

[8] "On pourroit dire à quoy sert ces solutions qui sont impossibles, je respond pour trois choises, pour la certitude de la reigle generale, et qu'il ny a point d'autre solutions, et pour son utilité" (*Invention nouvelle en l'algebre*, fol. F 1 (Amsterdam, 1629)). The solution of "1 (2) est esgale à 6 (1) − 25," that is, $x^2 = 6x - 25$, which gives $x = 3 \pm \sqrt{-16}$, he calls "inexplicable" (p. 114). He places in the same category (p. 130) numbers like $\sqrt{-3}$. From this point on, the reader may profitably consult the *Encyklopädie*, I, 148.

Approach to a Graphic Representation. Wallis (1673) seems to have been the first to have any idea of the graphic representation of these quantities. He stated[1] that the square root of a negative number was thought to imply the impossible, but that the same might also be said of a negative number, although we can easily explain the latter in a physical application:

These *Imaginary* Quantities (as they are commonly called), arising from the *Supposed* Root of a Negative Square (when they happen,) are reputed to imply that the Case proposed is Impossible.

And so indeed it is, as to the first and strict notion of what is proposed. For it is not possible that any Number (Negative or Affirmative) Multiplied into itself can produce (for instance) -4. Since that Like Signs (whether $+$ or $-$) will produce $+$; and therefore not -4.

But it is also Impossible that any Quantity (though not a Supposed Square) can be *Negative*. Since that it is not possible that any *Magnitude* can be *Less than Nothing* or any *Number Fewer than None*.

Yet is not that Supposition (of Negative Quantities,) either Unuseful or Absurd; when rightly understood. And though, as to the bare Algebraick Notation, it import a Quantity less than nothing: Yet, when it comes to a Physical Application, it denotes as Real a Quantity as if the Sign were $+$; but to be interpreted in a contrary sense.

Having shown that we may have negative lines, he asserts that we may also have negative areas and that a negative square must have a side, thus:

Now what is admitted in Lines must, on the same Reason, be allowed in Plains also...

But now (supposing this Negative Plain, -1600 Perches, to be in the form of a Square;) must not this Supposed Square be supposed to have a Side? And if so, what shall this Side be?

We cannot say it is 40, nor that it is -40 ...

But thus rather that it is $\sqrt{-1600}$, or ... $10\sqrt{-16}$, or $20\sqrt{-4}$, or $40\sqrt{-1}$.

[1] *Algebra*, cap. LXVI; Vol. II, p. 286, of the Latin edition; but for his 1673 statement, differing somewhat from that in his algebra, see Cajori in *Amer. Math. Month.*, XIX, 167. See also G. Eneström, *Bibl. Math.*, VII (3), 263.

Where $\surd$ implies a Mean Proportional between a Positive and a Negative Quantity. For like as $\surd bc$ signifies a Mean Proportional between $+b$ and $+c$; or between $-b$ and $-c$; $\cdots$ So doth $\surd -bc$ signify a Mean Proportional between $+b$ and $-c$, or between $-b$ and $+c$.

He therefore reached the position where he would be supposed to draw a line perpendicular to the real axis and say that this might be taken as an imaginary axis, but although he touched lightly upon this possibility, he did nothing of consequence with the idea.

Leibniz on Complex Numbers. Leibniz was the next to take up the study of imaginaries. He showed (1676) that $\sqrt{1+\sqrt{-3}}+\sqrt{1-\sqrt{-3}}=\sqrt{6}$, and (1702) that x^4+a^4 is equal to $\left(x+a\sqrt{-\sqrt{-1}}\right)\left(x-a\sqrt{-\sqrt{-1}}\right)\left(x+a\sqrt{\sqrt{-1}}\right)\left(x-a\sqrt{\sqrt{-1}}\right)$.[1] He was much impressed by the possibilities of the imaginary, but he seems never to have grasped the idea of its graphic representation.[2]

Modern Analytic Treatment. In 1702 Jean Bernoulli brought the imaginary to the aid of higher analysis by showing the relation between the $\tan^{-1}x$ and the logarithm of an imaginary number.[3] Newton's work with imaginaries (1685) was confined to the question of the number of roots of an equation,[4] a subject that was continued by Maclaurin[5] and other English algebraists.

[1] *Werke*, Gerhardt ed. (Berlin, 1850), II (3), 12; (Halle, 1858), V (3), 218, 360. See Tropfke, *Geschichte*, I (1), 171.

[2] "Itaque elegans et mirabile effugium reperit in illo Analyseos miraculo, idealis mundi monstro, pene inter Ens et non-Ens Amphibio, quod radicem imaginariam appellamus" (*Werke*, V, 357).

"Ex irrationalibus oriuntur quantitates impossibiles seu imaginariae, quarum mira est natura, et tamen non contemnenda utilitas; etsi enim ipsae per se aliquid impossibile significent, tamen non tantum ostendunt fontem impossibilitatis, et quomodo quaestio corrigi potuerit, ne esset impossibilis, sed etiam interventu ipsarum exprimi possunt quantitates reales" (*ibid.*, VII, 69).

[3] *Opera*, I, 393 (Lausanne, 1472); Tropfke, *Geschichte*, II (2), 83; Cantor, *Geschichte*, III, 348.

[4] *Arithmetica Universalis*, p. 242 (Cambridge, 1707), the imaginaries being called "radices impossibiles."

[5] *Phil. Trans.*, XXIV (1726), 104; XXXVI, 59; *Algebra*, 1748 ed., p. 275.

The first important step in the new theory to be taken by a British mathematician was made by Cotes (*c.* 1710) when he stated[1] that $\log(\cos\phi + i\sin\phi) = i\phi$, corollaries to which are the important formula

$$e^{\phi i} = \cos\phi + i\sin\phi,$$

which bears Euler's name, and the well-known relation

$$(\cos\phi + i\sin\phi)^n = \cos n\phi + i\sin n\phi,$$

suggested by De Moivre in 1730 but possibly known by him as early as 1707.[2] Euler (1743, 1748) was the first to prove that this relation holds for all values of n, and also that

$$\sin\phi = \frac{e^{\phi i} - e^{-\phi i}}{2i}$$

and

$$\cos\phi = \frac{e^{\phi i} + e^{-\phi i}}{2}.$$

Graphic Representation. Although some approach to the graphic representation of the complex number had been made by Wallis, and although the goal had been more nearly attained by H. Kühn, of Danzig, it was a Norwegian surveyor, Caspar Wessel,[3] who first gave the modern geometric theory. In 1797 he read a paper on the subject before the Royal Academy of Denmark. This was printed in 1798 and appeared in the memoirs of the Academy in 1799.[4] In this he says:

Let us designate by $+1$ the positive rectilinear unit, by $+\epsilon$ another unit perpendicular to the first and having the same origin; then the angle of direction of $+1$ will be equal to 0°, that of -1 to 180°, that of ϵ to 90°, and that of $-\epsilon$ to $-90°$ or to 270°.

[1] *Harmonia mensurarum* (posthumous), p. 28 (Cambridge, 1722): "Si quadrantis circuli quilibet arcus, radio *CE* descriptus, sinum habeat *CX*, sinumque complementi ad quadrantem *XE:* sumendo radium *CE* pro Modulo, arcus erit rationis inter $EX + XC\sqrt{-1}$ & *CE* mensura ducta in $\sqrt{-1}$." See also *Bibl. Math.*, II (3), 442.

[2] *Bibl. Math.*, II (3), 97–102.

[3] Born at Jonsrud, June 8, 1745; died 1818.

[4] See the French translation, *Essai sur la représentation analytique de la direction,* Copenhagen, 1897.

The plan was therefore the same as the one now used and until recently attributed to various other writers, including Henri Dominique Truel[1] (1786), A. Q. Buée[2] (1805), J. R. Argand[3] (1806), Gauss,[4] J. F. Français[5] (1813), and John Warren.[6] W. J. G. Karsten (1732–1787), at one time professor at Halle, gave a method (1768) of representing imaginary logarithms.[7]

Gauss took as his four units the roots of the equation $x^4 - 1 = 0$. Eisenstein[8] developed a theory based on the roots of the equation $x^3 - 1 = 0$, and Kummer[9] based his on the roots of the equation $x^n - 1 = 0$.[10]

Terms and Symbols. As already mentioned, Cardan (1545) spoke of a solution like $5 + \sqrt{-15}$ as "per radicem m̃," or "sophistic quantities," and Bombelli called numbers like $+\sqrt{-n}$ and $-\sqrt{-n}$ *piu di meno* and *meno di meno*,[11] abbreviated to *p. di m.* and *m. di m.* Descartes (1637) contributed the terms "real" and "imaginary."[12]

Most of the 17th and 18th century writers spoke of $a + b\sqrt{-1}$ as an imaginary quantity.[13] Gauss (1832) saw the desirability of having different names for $a\sqrt{-1}$ and $a + b\sqrt{-1}$,

[1] Cauchy mentions this fact, but nothing is known of the man. His MSS. are lost.

[2] In a paper *Sur les Quantités Imaginaires*, read before the Royal Society, London, 1805. See the *Philosophical Transactions*, London, 1806.

[3] *Essai sur une manière de représenter les quantités imaginaires dans les constructions géométriques*, Paris, 1806; 2d ed., Paris, 1874.

[4] He refers to the subject in his *Demonstratio nova* (1798), but does nothing with it. In his *Theoria residuorum biquadraticorum, Commentatio secunda* (1831) he presents the theory in its present form, evidently ignorant of Wessel's work.

[5] See Gergonne's *Annales*, IV, 61.

[6] *A Treatise on the Geometrical Representation of the Square Roots of Negative Quantities*, Cambridge, 1828. The general plan is that of Wessel, but the treatment of the subject is very abstract.

[7] F. Cajori, *Amer. Math. Month.*, XX, 76.

[8] On Gauss's estimate of him see Volume I, page 509.

[9] See Volume I, pages 507, 508.

[10] Tropfke, *Geschichte*, II (2), 88.

[11] "Plus of minus" and "minus of minus." *L'Algebra*, p. 294 seq. (Bologna, 1572).

[12] "Au reste tant les vrayes racines que les fausses ne sont pas toujours réelles, mais quelquefois seulement imaginaires." *La Geometrie*, 1705 ed., p. 117.

[13] Thus d'Alembert (1746): "Une fonction quelconque de tant et de telles grandeurs imaginaires, qu'on voudra, peut toujours être supposée égale à $p + q\sqrt{-1}$" (*Hist. de l'Acad. d. Berlin*, II, 195).

and so he gave to the latter the name "complex number."[1] The use of i for $\sqrt{-1}$ is due to Euler (1748).[2] Cauchy[3] (1821) suggested the name "conjugates" (*conjuguées*) for $a+bi$ and $a-bi$ and the name "modulus" for $\sqrt{a^2+b^2}$. Weierstrass called the latter the "absolute value" of the complex number and represented it by $|a+bi|$. Gauss had already given to a^2+b^2 the name of "norm."[4]

Quaternions and Ausdehnungslehre. The development of complex numbers, with their graphic representation in a plane, naturally led to the consideration of numbers of this type that might be graphically represented in a space of three dimensions. Argand (1806) attempted to take this step but found himself unable to do so, and Servois (1813) also made the attempt and failed.

In 1843 Sir William Rowan Hamilton[5] discovered the principle of quaternions, and presented his first paper on the subject before the Royal Irish Academy. His first complete treatment was set forth in his *Lectures on Quaternions* (1853). His discovery necessitated the withdrawal of the commutative law of multiplication, the adherence to which had proved to be a bar to earlier progress in this field.

The most active of the British scholars who first recognized the power of quaternions was Peter Guthrie Tait. Becoming acquainted with Hamilton soon after the latter's *Lectures on Quaternions* appeared, Tait began with him a correspondence that was carried on until his death.[6] Tait had been a classmate

[1] "Tales numeros vocabimus numeros integros complexos" (*Werke*, II, 102 (Göttingen, 1876)).

[2] ". . . formulam $\sqrt{-1}$ littera i in posterum designabo, ita vt sit $ii = -1$" (*Institutionum calculi integralis volumen IV*, 184 (Petrograd, 1794)). In his *Introductio in Analysin Infinitorum* (Lausanne, 1748), he first used the symbol: "Cum enim numerorum negativorum Logarithmī sint imaginarii . . . erit $l.-n$ quantitas imaginaria, quae sit $= i$." See also W. W. Beman in *Bulletin of the Amer. Math. Soc.*, IV, 274, 551.

[3] *Cours d'Analyse algébrique*, p. 180 (Paris, 1821).

[4] Tropfke, *Geschichte*, II (2), 90. For other attempts at explaining the imaginary number see Cantor, *Geschichte*, IV, 88–91, 303–318, 573, 712–715, 729–731.

[5] A. Macfarlane, *Ten Brit. Math.*, p. 43 (New York, 1916). See also P. G. Tait's article on "Quaternions" in the *Encyc. Britan.*, 9th ed., XX, 160.

[6] A. Macfarlane, "Peter Guthrie Tait," *Physical Review*, XV, 51.

of Clerk Maxwell's at the Edinburgh Academy and, like him, was deeply interested in physical studies. Partly as a result of this early training he soon began to apply the theory of quaternions to problems in this field, his results appearing in the *Messenger of Mathematics* and the *Quarterly Journal of Mathematics*. Recognizing Hamilton's wish, if not at his request, he delayed the publication of his work on the theory until after the former's *Elements* appeared. His own *Elementary Treatise on Quaternions* was therefore not issued until 1866, after which he continued to write upon the subject until his death. In 1873 he published with Professor Kelland a work entitled *An Introduction to Quaternions*, which did much to make the subject known to physicists. The theory has not, however, been as favorably received by scientists as had been anticipated by its advocates. It should be added that Gauss, about the year 1820, gave some attention to the subject but without developing any theory of importance.[1]

At about the same time that Hamilton published his discovery of quaternions Hermann Günther Grassmann published his great work, *Die lineale Ausdehnungslehre* (1844), although he seems to have developed the theory as early as 1840.[2]

7. Transcendental Numbers

Transcendental Numbers Considered Elsewhere. Among the artificial numbers should, of course, be included not only such types as surds but also all such nonalgebraic types as are found in connection with the trigonometric functions, logarithms, the study of the circle, and the theory of transcendental numbers in general. These may, however, be more conveniently considered in connection with algebra, geometry, and trigonometry, as they are commonly found in the teaching of these subjects.

[1] Tropfke, *Geschichte*, II (2), 88.

[2] V. Schlegel, "Die Grassmann'sche Ausdehnungslehre," Schlömilch's *Zeitschrift*, XLI. For A. Macfarlane's digest of the views of various writers see *Proceedings of the American Assoc. for the Adv. of Sci.*, 1891. See also E. Jahnke, in *L'Enseignement Mathématique*, XI (1909), 417; F. Engel, in Grassmann's *Gesammelte math. und physikal. Werke*, III (Leipzig, 1911).

TOPICS FOR DISCUSSION

1. The sequence of development of artificial numbers, with the causes leading to the successive steps.

2. General nature of compound numbers at various periods in their development.

3. Nature of fractions among the Egyptians, and the reasons for the persistence of the unit fraction.

4. Greek symbolism for fractions compared with that which the Romans used.

5. Origin and development of our common fractions.

6. The etymology of terms used in fractions, with the change in these terms from time to time.

7. The sequence of operations with fractions in the early printed works on arithmetic.

8. The development of methods of performing the operations with common fractions.

9. The origin, development, symbolism, and present status of sexagesimal fractions.

10. The origin and development of decimal fractions, including the question of symbolism.

11. The human needs that led to the development of the various types of fraction.

12. The origin and development of the idea of per cent, including the question of symbolism.

13. The reason for the interest of the Greek mathematicians in incommensurable numbers.

14. The development of surd numbers, particularly among the Greek and Arab writers.

15. History of the various methods of approximating the value of a surd number.

16. The origin and development of negative numbers, including the question of symbolism.

17. The origin and development of the idea of complex numbers, including the question of symbolism.

18. The origin and development of the graphic representation of complex numbers.


19. The origin and development of the idea of complex exponents in algebra.

CHAPTER V

GEOMETRY

1. General Progress of Elementary Geometry

Intuitive Geometry. All early geometry was intuitive in its nature; that is, it sought facts relating to mensuration without attempting to demonstrate these facts by any process of deductive reasoning. The prehistoric geometry sought merely agreeable forms, as in the plaiting of symmetric figures in a mat. The next stage was that of the mensuration of rectangles and triangles, and geometry was in this stage when the Ahmes Papyrus (*c.* 1550 B.C.)[1] was written. In this work the area of an isosceles triangle of base b and sides s is given as $\frac{1}{2}bs$. For the area of a circle of diameter d Ahmes used a rule which may be expressed in modern symbols as $(d-\frac{1}{9}d)^2$, which shows that he took 3.1605– as the value of π,—a value based on experiment.[2]



In Babylonia the same conditions existed. The tablets which have come down to us contain a few cases of mensuration,[3] but the rules are based merely on experiment.

[1] It should be recalled that Professor Peet (*Rhind Papyrus*, p. 3) puts this date as probably before 1580 B.C.

[2] On the general history of the development of geometry see G. Loria, *Il Passato ed il Presente delle Principali Teorie Geometriche*, Turin, 3d ed., 1907; hereafter referred to as Loria, *Passato-Presente Geom.* This work first appeared in the Turin *Memorie della R. Accad.*, XXXVIII (2), and was translated into German by F. Schütte, Leipzig, 1888. See also the *Encyklopädie*, Vol. III; R. Klimpert, *Geschichte der Geometrie*, Stuttgart, 1888; E. F. August, *Zur Kenntniss der geometrischen Methode der Alten*, Berlin, 1843; and the various general histories of mathematics. On the early Egyptian geometry see E. Weyr, *Ueber die Geometrie der alten Aegypter*, Vienna, 1884; Eisenlohr, *Ahmes Papyrus*. On a papyrus which may be slightly earlier than that of Ahmes, see page 293.

[3] Hilprecht, *Tablets*.

The native mathematics of China was also of this type. The *Nine Sections*, written perhaps *c.* 1100 B.C., contains statements which show that the author knew the relations of the sides of certain right-angled triangles,[1] but there is no evidence of any proof of such relations.

In the later Chinese mathematics there are many ingenious examples involving mensuration, but nowhere does there appear any further idea of geometric demonstration, as we understand the term, than is found in the earliest works.

In India the same conditions existed, the native geometry giving us no evidence of any approach to a sequence of deductive proofs. There was a large amount of mensuration,[2] and considerable ability was shown in the formulation of rules, but the basis of the work was wholly empirical.

The Romans were interested in mathematics only for its immediately practical value. The measurement of land, the laying out of cities, and the engineering of warfare appealed to them, but for demonstrative geometry they had no use. Indeed, it may be said that, outside of those lands which were affected by the Greek influence, the ancient world knew geometry only on its intuitive side. Demonstrative geometry was Greek in its origin, and in the Greek civilization it received its only encouragement for more than a thousand years.

Demonstrative Geometry. The idea of demonstrating the truth of a proposition which had been discovered intuitively appears first in the teachings of Thales (*c.* 600 B.C.). It is probable that this pioneer knew and proved about six theorems,[3] each of which would have been perfectly obvious to anyone without any demonstration whatever. The contribution of Thales did not lie in the discovery of the theorems, but in their proofs. These proofs are lost, but without them his work in geometry would have attracted no attention, either among his contemporaries or in the history of thought.

[1] See Volume I, pages 30, 33.

[2] *E.g.*, consider Bhāskara's *Lilāvati*, with special sections on ponds, walls, timber, heaps, shadows, and excavations.

[3] See Volume I, page 67; Heath, *History*, I, 130.

From the time of Thales until the decay of their ancient civilization demonstrative geometry was the central feature of the mathematics of the Greeks. The history of the general progress of the science has been sufficiently outlined elsewhere in this work.[1]

The Arabs recognized the Greek culture more completely than any other people until the period of the awakening of Western Europe. They translated the Greek classics in geometry as they did also in philosophy and natural science, but they never made any additions of real significance to the works of Euclid and Apollonius.

It was chiefly through the paraphrase of Boethius (*c.* 510) that Euclid's *Elements* (*c.* 300 B.C.) was known in the Dark Ages of Europe. The study of geometry received some encouragement at the hands of Gerbert (*c.* 1000), Fibonacci (1220), and a few other medieval scholars, but no progress was made in the advance of the great discipline which had been so nearly perfected in Alexandria more than a thousand years earlier.

With the invention of European printing the work of Euclid became widely known, the first printed edition appearing in 1482.[2] Little by little new propositions began to be suggested, but the invention of analytic geometry early in the 17th century took away, for a considerable period, much of the interest in improving upon the ancient theory.

The next advance in the pure field was made in the 17th century, when Desargues[3] (1639) published a work which treated of certain phases of projective geometry. The new analytic treatment of the subject, however, was so novel and powerful as to take the attention of mathematicians from the work of Desargues, and it was not until the 19th century that pure geometry again began to make great progress.

The Greek theory of conics has already been considered sufficiently for our purposes (Vol. I, Chap. IV). The analytic and modern synthetic geometries are considered later.

[1] See Volume I, pages 59–146; Heath, *History*, I, chap. iv.

[2] Venice, Erhard Ratdolt; the Campanus translation. See Volume I, page 251.

[3] See Volume I, page 383.

2. Name for Geometry

Reason for Uniformity. When we consider that our elementary geometry is essentially the *Elements* of Euclid, and that the subject never flourished in ancient times outside the Greek sphere of influence, it is apparent that the Greek name would be the one generally used to designate the science. It is derived from the words for "earth" and "measure"[1] and therefore was originally, as it is in some languages today, synonymous with the English word "surveying." Since the latter science was well developed in Egypt before the Greeks founded Alexandria, the name is probably a translation of an Egyptian term. It was in use in the time of Plato and Aristotle, and doubtless goes back at least to Thales.

Euclid did not call his treatise a geometry, probably because the term still related to land measure, but spoke of it merely as the *Elements*.[2] Indeed, he did not employ the word "geometry" at all, although it was in common use among Greek writers.[3]

When *Euclid* was translated into Latin in the 12th century, the Greek title was changed to the Latin form *Elementa*,[4] but the word "geometry" is often found in the title-page, first page, or last page of the early printed editions.[5]

There have been, as would naturally be expected, various fanciful names for textbooks on geometry. In the 16th century such names were common in all branches of learning. Among the best-known of these titles is the one seen in Robert Recorde's *The pathway to Knowledg* (London, 1551 and 1574).

[1] Γῆ (*gê*), earth, and μετρεῖν (*metrein'*), to measure.

[2] In Greek, στοιχεία (*stoichei'a*). So in the *editio princeps* of the Greek text (Basel, 1533) the title appears as ΕΥΚΛΕΙΔΟΥ ΣΤΟΙΧΕΙΩΝ ΒΙΒΛΙΑ ΙΕ.

[3] Thus Plato (*Theætetus*, 173 E; *Meno*, 76 A; *Republic*, 546 C, 511 D), Xenophon (*Symposium*, 6, 8, etc.), and Herodotus (II, 109) use the word in some of its forms, but always to indicate surveying.

[4] So in the *editio princeps* (1482) the first page begins: "Preclarissimus liber elementorum Euclidis perspicacissimi: in artem geometrie incipit quâ foelicissime." The colophon also has the name *geometria*.

[5] *E.g.*, the first English edition (London, 1570) has the title *The Elements of Geometrie of the most auncient Philosopher Evclide of Megara.*

3. Technical Terms of Euclidean Geometry

Point. The history of a few typical terms of elementary geometry will now be considered.[1] The Pythagoreans defined a point as "a monad having position,"[2] and this definition was adopted by Aristotle (*c.* 340 B.C.). Plato (*c.* 380 B.C.) called a point "the beginning of a line,"[3] and Simplicius (6th century) called it "the beginning of magnitudes and that from which they grow," adding that it is "the only thing which, having position, is not divisible." Euclid (*c.* 300 B.C.) gave the definition: "A point is that which has no part." Heron (*c.* 50?) used the same words, but added "or a limit without dimension or a limit of a line." When Capella (*c.* 460) translated the definition into Latin, he made it read, "A point is that of which a part is nothing,"[4] which is a different matter.

Modern writers usually resort to analogy and give only a quasi definition, or else they make use of the idea of limit.

Line. The Platonists defined a line as length without breadth, and Euclid[5] did the same. Aristotle objected to such a negative definition, although Proclus (*c.* 460) observes that it is positive to the extent that it affirms that a line has length. An unknown Greek writer[6] defined it as "magnitude extended one way,"[7] a phrase not unlike one used by Aristotle. The latter defined it as a magnitude "divisible in one way only,"[8] in contrast to a surface, which is divisible in two ways, and to a solid, which is divisible in three ways. Proclus suggested defining a line as the "flux of a point,"[9] an idea also going back to

[1] For further discussion see J. H. T. Müller, *Beiträge zur Terminologie der griechischen Mathematiker*, 1860; Heath, *Euclid*, Vol. I, p. 155; H. G. Zeuthen, "Sur les définitions d'Euclide," *Scientia*, XXIV, 257, on the general nature of Euclid's definitions. See also Heath, *History*, on all such details.

[2] Μονὰς προσλαβοῦσα θέσιν (monad with position added).

[3] Ἀρχὴ γραμμῆς. On this and other definitions consult Heath, *Euclid*, Vol. I, p. 155; H. Schotten, *Inhalt und Methode des planimetrischen Unterrichts*, Vol. I (Leipzig, 1890); Vol. II (Leipzig, 1893); hereafter referred to as Schotten, *Inhalt*.

[4] "Punctum est cuius pars nihil est."

[5] Γραμμὴ δὲ μῆκος ἀπλατές.

[6] Alluded to by al-Nairîzî (*c.* 910) as one Heromides or Herundes.

[7] Μέγεθος ἐφ' ἓν διάστατον. [8] Μοναχῇ διαιρετόν. [9] Ῥύσις σημείου.

Aristotle, who remarked that "a line by its motion produces a surface, and a point by its motion a line." This occasionally appears as "A line is the path of a moving point."

Straight Line. It is evident that certain terms are so elementary that no simpler terms exist by which to define them. This is true of "point" and "line," but it is more evidently true of terms like "straight line" and "angle." Plato defined a straight line as "that of which the middle covers the ends," that is, relatively to an eye placed at either end and looking along the line. Euclid endeavored to give up the appeal to sight and defined it as "a line which lies evenly with the points on itself." Proclus explains that Euclid "shows by means of this that the straight line alone [of all lines] occupies a distance equal to that between the points on it," adding that the distance between two points on a circumference or any other line, and measured on this line, is greater than the interval between them. Archimedes (*c.* 225 B.C.) stated this idea more tersely by saying that "of all lines having the same extremities the straight line is the shortest," which is the source of the definition often found in textbooks, "a straight line is the shortest distance between two points," although "line" and "distance" are two radically different concepts. "The shortest path between two points" is an expression that is less objectionable, but it merely shifts the difficulty.

Heron (*c.* 50?) defined a straight line as "a line stretched to the utmost toward the ends," and Proclus adopted this phrase with the exception of "toward the ends." It is evidently objectionable, however, because it appeals to the eye and relates to a physical object. Heron also suggested the idea that "all its parts fit on all [other parts] in all ways," a definition substantially adopted by Proclus. Still another definition due to Heron is "that line which, when its ends remain fixed, itself remains fixed when it is, as it were, turned round in the same plane." This too was used with slight change by Proclus, and it appears in various modern works as "that which does not change its position when it is turned about its extremities (or any two points in it) as poles."

Surface. The Pythagoreans used a word[1] meaning "skin" or "color" to designate a surface. Aristotle, like Plato, used other words,[2] and spoke of a surface as extended or continuous or divisible in two ways, and as the extremity or the section of a solid. Aristotle recognized as common the idea that a line by its motion produces a surface.[3] Euclid defined a surface as "that which has only length and breadth."

Plane Surface. The same difficulties that the ancients had in defining a straight line were met when they attempted to define a plane. Euclid stated that "a plane surface is a surface which lies evenly with the straight lines on itself." Heron (*c.* 50?) added that it is "the surface which is stretched to its utmost," this being analogous to his definition of a straight line. He also defined it as "a surface all the parts of which have the property of fitting on" [each other], and as "such that if a straight line passes through two points on it, the line coincides wholly with it at every spot, all ways." Proclus (*c.* 460), adopting an assumption stated by Archimedes, defined it as "the least surface among all those which have the same extremities," and also used a modification of Euclid's definition, "a surface such that a straight line fits on all parts of it," or "such that the straight line fits on it all ways." There was no material improvement on these statements until the 18th century, when Robert Simson (1758) suggested the definition that "a plane superficies is that in which any two points being taken, the straight line between them lies wholly in that superficies,"[4] a statement which Gauss (*c.* 1800) characterized as redundant. Fourier (*c.* 1810) gave the definition that a plane is formed by the aggregate of all the straight lines which, passing through one point on a straight line in space, are perpendicular to that

[1] Χροιά (*chroia'*).

[2] Ἐπιφάνεια (*epipha'neia*) and ἐπίπεδον (*epi'pedon*). From the former, a word meaning "appearance," we have our word "epiphany." The latter word, meaning a plane surface, occurs in our word "parallelepiped." Later Greek writers also used ἐπιφάνεια to indicate any kind of surface, and Plato used ἐπίπεδον in the same way.

[3] On the different kinds of lines and surfaces, consult Heath, *Euclid.*

[4] Compare one of Heron's definitions above.

straight line. This is, of course, merely putting into another form a well-known theorem of Euclid.[1] Crelle (1834) suggested that a plane is the surface containing throughout their entire lengths all the straight lines passing through a fixed point and also intersecting a straight line in space.[2]

Angle. Euclid's definitions of an angle are as follows:

A plane angle is the inclination to one another of two lines in a plane which meet one another and do not lie in a straight line.

And when the lines containing the angle are straight, the angle is called rectilineal.

This excludes the zero angle, straight angle, and in general the angle $n\pi$, and defines angle by the substitution of the idea of inclination,—in modern form, the difference in direction.

Even less satisfactory is the definition of Apollonius (*c.* 225 B.C.) which asserts that an angle is "a contracting of a surface or a solid at one point under a broken line or surface." Plutarch (1st century) and various other writers defined it as "the first distance under the point," which Heath[3] interprets as "an attempt (though partial and imperfect) to get at the rate of divergence between the lines at their point of meeting." Perhaps this idea was also in the mind of Carpus of Antioch (1st century) when he said that the angle is "a quantity, namely a distance between the lines or surfaces containing it."

Later writers often return to the qualitative idea of Aristotle, as in the definition that an angle is a figure formed by two lines which meet. This was refined by Professor Hilbert of Göttingen[4] as follows:

Let α be any arbitrary plane and h, k any two distinct half-rays lying in α and emanating from the point O so as to form a part of two different straight lines. We shall call the system formed by these two half-rays h, k an *angle*.

[1] *Elements*, XI, 5.

[2] For further consideration of modern definitions see Heath, *Euclid*, Vol. I, p. 174, and Schotten, *Inhalt*, II.

[3] Heath, *Euclid*, Vol. I, p. 177.

[4] *Foundations of Geometry*, translated by E. J. Townsend, p. 13 (Chicago, 1902).

Circle. The ancient writers defined a circle substantially as Euclid did:

A circle is a plane figure contained by one line such that all the straight lines falling upon it from one point among those lying within the figure are equal to one another;

And the point is called the center of the circle.

Euclid had already defined a figure as "that which is contained by any boundary or boundaries," so that a circle is, in his view, the portion of a plane included in the bounding line. This bounding line Euclid usually calls the periphery (*περιφέρεια*), a word translated into the Latin as *circumferentia*, whence our "circumference." Euclid is not consistent however, for he speaks of a circle as not cutting a circle in more than two points,[1] the word "circle" here referring to the bounding line.

This uncertain use of the term has been maintained until recent times. The influence of analytic geometry has led to defining the circle as a line, but there is still no uniformity in the matter.

Diameter and Radius. Euclid used the word "diameter"[2] in relation to the line bisecting a circle and also to mean the diagonal[3] of a square, the latter term being also found in the works of Heron.

The term "radius" was not used by Euclid, the term "distance" being thought sufficient. Boethius (*c.* 510) seems to have been the first to use the equivalent of our "semidiameter."[4] A similar use also appears in India, in the writings of Āryabhaṭa[5] (*c.* 510). Ramus[6] (1569) used the term, saying: "Radius est recta a centro ad perimetrum."

[1] *Elements*, III, 10, where the Greek κύκλος (*ky'klos*, circle) is used.

[2] Διά (*dia'*, through) + μετρεῖν (*metrein'*, to measure).

[3] Διαγώνιος, from διά (*dia'*, through) + γωνία (*goni'a*, angle).

[4] Thus in the *Ars geometriae*, ed. Friedlein, p. 424: "Conscribitur age emicyclus XXVIII in basi et in semidiametro XIIII pedes habens" (MS. of the 11th century).

[5] L. Rodet, "Leçons de Calcul d'Âryabhaṭa," *Journ. Asiatique*, XIII (7), 398; reprint (Paris, 1879), p. 10: "The chord of the sixth part of the circumference is equal to a semidiameter."

[6] On the general question see *Bibl. Math.*, II (3), 361, and P. Ramus, *Scholarvm Mathematicarvm, Libri vnvs et triginta*, p. 155 (Basel, 1569).

From India it seems to have passed over to Arabia and thence to Europe. So Plato of Tivoli (*c.* 1120) used *mediatas diametri* and *dimidium diametri*, Fibonacci (1220) used *semidyameter*,[1] and Jordanus Nemorarius (*c.* 1225) preferred the form *semidiameter*.[2]

The early printed books,—such as those of Maurolico (1558), Tartaglia (1560), and Pedro Nunes (1564),—commonly used the word *semidiameter*.[3]

The word "radius" as used in this sense is modern. It appears, as above stated, in the *Scholarvm Mathematicarvm, Libri vnvs et triginta* of Ramus (1569), and a little later was used by Thomas Fincke[4] (1583) in his *Geometria Rotundi*. It was then adopted by Vieta[5] (*c.* 1590), and after that time it became common.

Parallel Lines. The word "parallel"[6] means "alongside one another." Euclid defined parallel straight lines as "straight lines which, being in the same plane and being produced indefinitely in both directions, do not meet one another in either direction." Rather less satisfactory is the definition of Poseidonius (*c.* 100 B.C.) as those lines "which, in one plane, neither converge nor diverge, but have all the perpendiculars equal which are drawn from the points of one line to the other." This definition is substantially that ascribed to Simplicius (6th century), that two straight lines are parallel "if, when they are produced indefinitely both ways, the distance between them, or the perpendicular drawn from either of them to the other, is always equal and not different." The direction theory, one of the least satisfactory of all, is due to Leibniz.[7]

[1] *Scritti*, II, 85. See also *dimidium dyametri* on page 86.

[2] See his "De Triangulis" in the *Mitteilungen des Coppernicus-Vereins . . . zu Thorn*, VI (1887).

[3] Tropfke, *Geschichte*, IV (2), 108.

[4] Variously spelled. A Danish mathematician (1561–1646). See Volume I, page 348.

[5] "Posito X radio seu semidiametro circuli." See Tropfke, *Geschichte*, IV (2), 108.

[6] Παράλληλος (*paral'lelos*).

[7] For further discussion, including the various bases for a definition, see Heath, *Euclid*, Vol. I, p. 192; Schotten, *Inhalt*, II, 188.

4. Axioms and Postulates

Distinction between Axioms and Postulates. The Greek writers recognized the existence of first principles "the truth of which," as Aristotle affirmed, "it is not possible to prove." These, he stated, were of two kinds: (1) those which are common to all sciences, and for which the name "axiom" was used by the Stoic philosophers and by Aristotle himself; (2) those which relate to the particular science, and to which the name "postulate" was given by later writers.[1] The distinction was not completely recognized, even by Euclid, for his fourth axiom (see page 281) is rather a geometric postulate or a definition.

Aristotle had other names for axioms, speaking of them as "the common [things]"[2] or "common opinions."[3]

Proclus (*c.* 460) states that Geminus (*c.* 77 B.C.) taught that axioms and postulates "differ from one another in the same way as theorems are also distinguished from problems,"—an opinion which is quite at variance with that of Aristotle.

Euclid seems to have used the term "common notion" to designate an axiom, although he may have used the term "axiom" also.[4]

The word "postulate" is from the Latin *postulare*, a verb meaning "to demand." The master demanded of his pupils that they agree to certain statements upon which he could build. It appears in the early Latin translations of Euclid[5] and was commonly used by the medieval Latin writers.

As to the number of these assumptions, Aristotle set forth the opinion which has been generally followed ever since, that "other things being equal that proof is the better which proceeds from the fewer postulates or hypotheses or propositions."

[1] Heath, *Euclid*, Vol. I, p. 117. On the general question of foundation principles in geometry, see the *Encyklopädie*, II, 1.

[2] Τὰ κοινά (*ta koina'*). [3] Κοιναὶ δόξαι (*koinai' dox'ai*).

[4] But not in his extant writings. On the doubt that has been raised as to his giving a list of axioms at all, see Heath, *Euclid*, Vol. I, p. 221.

[5] "Postulata. I. Postuletur, ut a quouis puncto ad quoduis punctum recta linea ducatur." The Greek word for postulates used by Euclid was αἰτήματα (*aite'mata*) (*Euclid*, ed. Heiberg, I, 8, 9).

Axioms. Euclid laid down certain axioms, or "common notions," probably five in number, as follows:

1. Things which are equal to the same thing are equal to one another.
2. If equals be added to equals, the wholes are equal.
3. If equals be subtracted from equals, the remainders are equal.
4. Things which coincide with one another are equal to one another.[1]
5. The whole is greater than the part.

The axioms of inequality, of doubling, and of halving may have been given by Euclid, but we are not certain.[2]

It will be observed that Euclid built up his geometry on a smaller number of axioms than many subsequent writers have thought to be necessary.

Postulates. Euclid does not use a noun equivalent to the Latin *postulatum*, but says:

Let the following be postulated:

1. To draw a straight line from any point to any point.
2. To produce a finite straight line continuously in a straight line.
3. To describe a circle with any center and [any] distance.
4. That all right angles are equal to one another.
5. That, if a straight line falling on two straight lines makes the interior angles on the same side less than two right angles, the two straight lines, if produced indefinitely, meet on that side on which are the angles less than the two right angles.[3]

Considerable criticism of these postulates developed among the later Greeks. Zeno of Sidon (1st century B.C.) asserted that it was necessary to postulate that two [distinct] straight lines cannot have a segment in common. If this is not done, he claimed, one or more of the proofs in Book I are fallacious.


Others asserted that postulates 4 and 5 are theorems capable of proof. Proclus (*c.* 460) attempted a proof of postulate 4,

[1] Essentially a postulate or a definition.


[2] For the evidence *pro* and *contra*, see Heath, *Euclid*, Vol. I, p. 223. Heiberg's edition of Euclid (I, 10, 11) numbers these 1, 2, 3, 7, 8, giving in Greek the doubtful axioms.

[3] Heath, *Euclid*, Vol. I, pp. 154, 195.


but it was fallacious. He also claimed that the converse is not necessarily true which asserts that an angle which is equal to a right angle is also a right angle, for he said that, in the figure given, $\angle ABC = \angle XBY$ and yet $\angle XBY$ is not a right angle. Saccheri[1] (1733) gave a proof of the postulate, but he assumed other statements equally fundamental upon which to base his argument. Modern writers often adduce a simple proof based upon the postulate of the equality of straight angles, but this simply substitutes one postulate for another.



Postulate 5, the "Postulate of Parallels," has been frequently attacked.[2] Ptolemy attempted to prove it, one of his arguments being that if $a + b$ is a straight angle, then $c + d$ must be a straight angle. Hence if the lines meet at P they also must meet at Q, and in that case the two straight lines inclose space. Proclus gave a more seductive argument relating to the meeting of lines in general, thus:



Draw the lines AK and CL so that $\angle A + \angle C <$ two right angles.



Bisect AC at E and lay off $AF = AE$ and $CG = EC$.

Then AF and CG cannot meet on FG, as at H; for if they did we should have $AH = AE$ and $CH = CE$, and so the sum of two sides of a triangle would be equal to the third side.

Bisect FG at H, make $FK = FH = HG = GL$.

[1] *Euclides ab omni naevo vindicatus*, p. x (Milan, 1733).

[2] Heath, *Euclid*, Vol. I, p. 202; Engel and Stäckel, *Die Theorie der Parallellinien*, Leipzig, 1895; G. B. Halsted, *Saccheri's Euclides Vindicatus* (translation), p. 7 (Chicago, 1920); G. Boccardini, *L'Euclide emendato del P. Gerolamo Saccheri*, Milan, 1904 (incomplete translation); R. Bonola, "Sulla teoria delle parallele e sulle geometrie non-euclide," in F. Enriques, *Questioni riguardanti le matematiche elementari*, p. 248 (Bologna, 1912), with an English translation, from the German edition, by H. S. Carslaw, Chicago, 1912. On the modern theory and treatment of postulates see C. J. Keyser, *Mathematical Philosophy*, Lecture II (New York, 1922).

Then FK and GL cannot meet on KL, for the same reason, and so on however far we go.

Hence the lines described in the postulate cannot meet at all, even though $\angle A + \angle C < 180°$.[1]

Further attempts at a proof of Postulate 5 were made by al-Ṭûsî (*c.* 1200), Wallis (*c.* 1660), Saccheri (1733), Lambert (*c.* 1766), Legendre (1794 and later), and many others.[2]

As an alternative postulate Proclus stated in substance, and Playfair (1795) made well known to the modern world, the following:

> Through a given point only one parallel can be drawn to a given straight line; [or]
>
> Two straight lines which intersect one another cannot both be parallel to one and the same straight line.

Playfair's form of the postulate was practically given, however, somewhat earlier than 1795. Joseph Fenn, in his edition of Euclid's *Elements*, published at Dublin in 1769,[3] stated it as follows: "A straight line which cuts one of two parallel lines will necessarily cut the other, provided this cutting line is sufficiently produced." Substantially the same assumption was also given by William Ludlam[4] in 1785, and, indeed, was given by Proclus (*c.* 460), as asserted above, in a note to Euclid, I, 31.

It has been observed by various writers that Euclid tacitly assumed other postulates, such as one relating to the intersection of plane figures and one which asserts that space is homogeneous or that a figure may be transposed without deformation.

[1] For the rest of his treatment of Postulate 5, see Heath, *Euclid*, Vol. I, p. 207.

[2] On the general question of the validity of the postulate, see page 335.

[3] *First Volume of the Instructions given in the Drawing School established by the Dublin-Society . . . under the direction of Joseph Fenn, heretofore Professor of Philosophy in the University of Nants*, Dublin, 1769. F. Cajori, "On the history of Playfair's parallel-postulate," *School Science and Mathematics*, XVIII, 778.


[4] Born *c.* 1718; died in Leicestershire, March 16, 1788. He wrote various works on astronomy. His *Rudiments of Mathematics* first appeared in 1785.

5. Typical Propositions of Plane Geometry

Pons Asinorum. Most of the basic theorems of elementary plane geometry are found in Euclid's *Elements*. Of these a relatively small number have any interesting history, and only a few typical ones need be considered, the first being Proposition 5 of Book I. As given by Euclid, this reads as follows:

In isosceles triangles the angles at the base are equal to one another, and, if the equal straight lines be produced further, the angles under the base will be equal to one another.

Proclus states that Thales (*c.* 600 B.C.) was the first to prove this proposition. At any rate it was well known to Aristotle (*c.* 340 B.C.), who discusses one of the proofs then possibly current. Proclus (*c.* 460) says that Pappus (*c.* 300) proved the theorem without using any auxiliary lines, simply taking the triangle up, turning it over, and laying it down upon itself. The question as to how he could lay the triangle itself down upon itself has caused a change in the phraseology on the part of modern writers.



The proposition represented substantially the limit of instruction in many courses in the Middle Ages. It formed a bridge across which fools could not hope to pass, and was therefore known as the *pons asinorum*, or bridge of fools.[1] It has also been suggested that the figure given by Euclid resembles the simplest form of a truss bridge, one that even a fool could make. The name seems to be medieval.

The proposition was also called *elefuga*, a term which Roger Bacon (*c.* 1250) explains as meaning the flight of the miserable ones, because at this point they usually abandoned geometry.[2]


[1] The term is sometimes applied to the Pythagorean Theorem.


[2] "Sic est hic quod isti qui ignorant alicujus scientiae, ut sit geometriae, nisi sint pueri qui coguntur per virgam, resiliunt et tepescunt, ut vix volunt tres vel quatuor propositiones scire. Unde ex hoc accidit quod quinta propositio geometriae Euclidis dicitur *Elefuga*, id est, fuga miserorum; *elegia* enim Graece dicitur, Latine *miseria*; et *elegi* sunt *miseri*." *Opus Tertium*, cap. vi.

Congruence Theorems. The second of the usual congruence theorems relates to the case of two angles and the included side of a triangle. Proclus (*c.* 460) says of this:

> Eudemus (*c.* 335 B.C.) in his geometrical history refers this theorem to Thales (*c.* 600 B.C.). For he says that, in the method by which they say that Thales proved the distance of ships in the sea, it was necessary to make use of this theorem.

How Thales could have used this theorem for the purpose is purely a matter of conjecture. He might have stood at *T*, the top of a cliff *TF*, and sighted to the ship *S*, using two hinged rods to hold the angle *STF*. He could then have turned and sighted along the same rod to a point *P* along the shore. If he kept the angle constant, he would then merely have to measure *FP* to find the unknown distance *FS*. Since in those days the ships were small and remained near the shore in good weather, this plan would have been quite





EARLY METHODS OF MEASURING DISTANCES

From Belli's *Libro del Misvrar con la vista*, Venice, 1569, but representing essentially the method probably used by Thales

feasible. Thales probably had some simple instrument like the astrolabe by which he could measure angles when observing the

stars, and he could have used this. We shall presently see that such instruments, in primitive form, were known to the Babylonians before his time.

Euclid stated the proposition in a more complicated form than the one now in use, but his proof had the advantage that it did not employ superposition. The latter form of proof is given by al-Nairîzî (*c.* 910), who ascribes it to some unknown predecessor.

Renaissance writers often used the theorem in practical mensuration, as Thales is thought to have done. The illustration on page 285, from Belli's work[1] of 1569, shows two methods of using it, and there is a story that one of Napoleon's engineers gained imperial favor by quickly applying it on an occasion when the army was held up by a river.

Areas. The sources of such propositions as those relating to the area of the triangle, the rectangle, the trapezium (trapezoid), and other rectilinear figures are, of course, unknown. The theorems relating to these areas are found in Euclid and apparently were common property long before his time. It is interesting to know, however, that the Egyptian surveyors, even after the time of Euclid, were in the habit of finding the area of a field by taking the product of the half-sums of the opposite sides. This is correct in the case of a rectangle, but in the case of a general convex quadrilateral it gives a result that is too large. The error was corrected to a certain extent by omitting in the calculation all fractions less than $\frac{1}{64}$ of the large unit of length.[2]

The rule for the area of an inscribed convex quadrilateral, expressed by the formula

$$A = \sqrt{(s-a)(s-b)(s-c)(s-d)},$$

was given by both Brahmagupta (*c.* 628) and Mahāvīra (*c.* 850), but without the limitation that it holds only for an

[1] S. Belli, *Libro del Misvrar con la vista*, Venice, 1569.

[2] This unit was the σχοῖνος (*schoi'nos*), equal to 100 cubits. Our information comes from a papyrus in the British Museum. See H. Maspero, *Les Finances de l'Égypte sous les Lagides*, p. 135 (Paris, 1905).

inscribed figure. If $d = 0$, the figure becomes a triangle and the formula reduces to

$$A = \sqrt{s(s-a)(s-b)(s-c)},$$

the rule being given by Heron (*c.* 50?), but being ascribed on Arabic authority to Archimedes. Both Brahmagupta and Mahāvīra[1] gave the equivalents of the following formulas, without limitations, for the lengths of the diagonals of a quadrilateral:

$$m = \sqrt{\frac{(ac+bd)(ab+cd)}{ad+bc}},$$

$$n = \sqrt{\frac{(ac+bd)(ad+bc)}{ab+cd}}.$$

The first of these formulas was rediscovered by W. Snell, who gave it in his edition (1619) of Van Ceulen's works.

Among the other interesting formulas related to those concerning areas is one discovered by Lhuilier and published in 1782. It gives the radius of the circle circumscribing a quadrilateral and reduces to the following:[2]

$$r = \tfrac{1}{4}\sqrt{\frac{(ab+cd)(ac+bd)(ad+bc)}{(s-a)(s-b)(s-c)(s-d)}}.$$

Angle Sum. The fact that the sum of the angles of a triangle is equal to two right angles has long been recognized as one of the most important propositions of plane geometry. Eutocius (*c.* 560) tells us that Geminus (*c.* 77 B.C.) stated that "the ancients investigated the theorem of the two right angles in each individual species of triangle,—first in the equilateral, again in the isosceles, and afterwards in the scalene triangle." Proclus (*c.* 460) says that Eudemus (*c.* 335 B.C.) ascribed the theorem to the Pythagoreans.

There is also a possibility that Thales knew this property of the triangle, for Diogenes Laertius (2d century) quotes Pamphilius (1st century) as saying that he was the first to inscribe

[1] See Volume I, page 163.

[2] I am indebted to Professor R. C. Archibald for this information, as for many other valuable suggestions.

a right-angled triangle in a circle, the proof of which solution requires this proposition, at least in a special case. The theorem was certainly well known before Euclid, for Aristotle refers to it several times.

Pythagorean Theorem. The relation of the sides of a triangle when these sides are 3, 4, and 5 (that is, $3^2 + 4^2 = 5^2$) was well known long before the time of Pythagoras. We find in the *Nine Sections* of the Chinese, perhaps written before 1100 B.C., this statement: "Square the first side and the second side and add them together; then the square root is the hypotenuse."

The Egyptians also knew the numerical relation for special cases, for a papyrus of the 12th dynasty (*c.* 2000 B.C.), discovered at Kahun, refers to four of these relations, one being $1^2 + (\frac{3}{4})^2 = (1\frac{1}{4})^2$. It was among these people that we first hear of the "rope stretchers,"[1] those surveyors who, it is usually thought, were able by the aid of this property to stretch a rope so as to draw a line perpendicular to another line, a method still in use at the present time.

Pythagorean Numbers in India. The Hindus knew the property long before the beginning of the Christian era, for it is mentioned in the *Śulvasūtras*,[2] the sacred poems of the Brahmans. The *Śulvasūtra* of Āpastamba gives rules for constructing right angles by stretching cords of the following lengths: 3, 4, 5; 12, 16, 20; 15, 20, 25; 5, 12, 13; 15, 36, 39; 8, 15, 17; and 12, 35, 37. Although the date of these writings is uncertain,[3] it is evident that the relations were known rather early in India.[4]

Did Pythagoras prove the Theorem? The proof of the proposition is attributed to Pythagoras (*c.* 540 B.C.) by various writers, including Proclus (*c.* 460), Plutarch (1st century), Cicero

[1] Ἀρπεδονάπται (*harpedonap'tae*) (Vol. I, p. 81). See Peet, *Rhind Papyrus*, p. 32.

[2] Curiously the word is sometimes interpreted to mean rope-stretching.


[3] Perhaps the 4th or 5th century B.C.

[4] A. Bürk, "Das Āpastamba-Śulba-Sūtra," in the *Zeitschrift der deutschen morgenländischen Gesellschaft*, LV, 543, and LVI, 327; G. Thibaut, *Journal of the Royal Asiatic Society of Bengal*, XLIV, reprint 1875, and his articles in *The Pandit*, Benares, 1875/6 and 1880; Heath, *Euclid*, Vol. I, p. 360.

(*c.* 50 B.C.), Diogenes Laertius (2d century), and Athenæus (*c.* 300). No one of these lived within, say, five centuries of Pythagoras, so that we have only a weak tradition on which to rest the general belief that Pythagoras was the first to prove the theorem.[1] It would seem as if such an important piece of history would have some mention in the works of a man like Aristotle; but, on the other hand, it is difficult to see how such a tradition should be so generally received unless it were well founded. Not only are we not positive that the proof is due to Pythagoras at all, but we are still more in doubt as to the line of demonstration that he may have followed.

Hundreds of proofs have been suggested for the proposition, but only two are significant enough to be mentioned at this time. Of these the first is the one in the *Elements*, a proof which Proclus tells us was due to Euclid himself. Although Schopenhauer, the philosopher, calls it "a proof walking on stilts" and "a mousetrap proof," it has stood the test of time better than any other.

The second noteworthy proof is that of Pappus (*c.* 300). In this figure we have any triangle ABC, with CM and CN any parallelograms on AC and BC, and with QR equal to PC. Then $AT = CM + CN$, a relation that reduces to the Pythagorean Theorem when ABC is a right-angled triangle and when the parallelograms are squares.[2]



The Pythagorean Theorem is not uncommonly called the *pons asinorum* by modern French writers. The Arabs called it the "Figure of the Bride," possibly because it represents two joined in one. It is also called the "Bride's Chair," possibly because the Euclid figure is not

[1] For arguments against this belief see H. Vogt, *Bibl. Math.*, VII (3), 6, and IX (3), 15; G. Junge, *Wann haben die Griechen das Irrationale entdeckt*, Halle, 1907.

[2] *Mathematicae Collectiones*, ed. Commandinus, Bologna, 1660, liber quartus, p. 57; Hultsch ed., IV, 177.

unlike the chair which a slave carries on his back and in which the Eastern bride is sometimes transported to the ceremony.[1]

Of the rules for forming rational right-angled triangles the following related ones are among the most important:[2]

$$n^2+\left(\frac{n^2-1}{2}\right)^2=\left(\frac{n^2+1}{2}\right)^2 \quad \text{Pythagoras } (c.\ 540 \text{ B.C.})$$

$$(2n)^2+(n^2-1)^2=(n^2+1)^2 \quad \text{Plato } (c.\ 380 \text{ B.C.})$$

$$(2n+1)^2+\left[\frac{(2n+1)^2-1}{2}\right]^2=\left[\frac{(2n+1)^2-1}{2}+1\right]^2$$

Proclus ($c.$ 460)

Recent Geometry of the Triangle. In the 19th century the geometry of the triangle made noteworthy progress. Crelle (1816) made various discoveries in this field, Feuerbach (1822) soon after found the properties of the nine-point circle, and Steiner set forth some of the properties of the triangle, but it was many years before the subject attracted much attention. Lemoine[3] (1873) was the first to take up the subject in a systematic way and to contribute extensively to its development. His theory of "transformation continue" and his "géométrographie" should also be mentioned. Brocard's contributions to the geometry of the triangle began in 1877, and certain critical points of the triangle bear his name.

The Pentagon and Decagon. The tenth proposition of Book IV of Euclid is the problem: "To construct an isosceles triangle having each of the angles at the base double the remaining one." This makes the vertical angle 36° and each of the others 72°, and therefore permits of the construction of

[1] The Greeks are said to have called it the "theorem of the married women," and Bhāskara to have spoken of it as the "chaise of the little married women." E. Lucas, *Récréations Mathématiques*, II, 130.

[2] H. A. Naber, *Das Theorem des Pythagoras*, Haarlem, 1908; Heath, *Euclid*, Vol. I, p. 350; the names are those of the authors of rules approximately represented by these formulas. The assertion as to Pythagoras is open to doubt.

[3] D. E. Smith, "Émile-Michel-Hyacinthe Lemoine," *Amer. Math. Month.*, III, 29; J. S. Mackay, various articles on modern geometry in the *Proceedings of the Edinburgh Mathematical Society*; E. Vigarié, "La bibliographie de la géométrie du triangle," *Mathesis*, XVI, suppl., p. 14.

a regular decagon and of a regular pentagon. The problem seems to have been known to the Pythagoreans, for Proclus (*c.* 460) tells us that they discovered "the construction of the cosmic figures,"—a statement anticipated by Philolaus (*c.* 425 B.C.) and Iamblichus (*c.* 325), and this construction requires the use of the problem. Lucian (2d century), the scholiast to the *Clouds* of Aristophanes, tells us that the pentagram, the star pentagon, was the badge of the Pythagorean brotherhood, and the construction of such a figure depends upon this proposition.[1]

The solution is related to that of the division of a line in extreme and mean ratio.[2] This was referred to by Proclus when he said that Eudoxus (*c.* 370 B.C.) "greatly added to the number of the theorems which Plato originated regarding the *section*." This is the first trace that we have of this name for such a cutting of the line.

In comparatively modern times the section appears first as "divine proportion,"[3] and then, in the 19th century,[4] as the "golden section."

6. Typical Propositions of Solid Geometry

Prism. Since the Greeks were so much more interested in the logic of geometry than in its applications to mensuration, and since they found a sufficient field for their activities in the work with plane figures, they did not develop the science of solid geometry to any great extent, as witness the *Elements* of Euclid. This is one reason why even the technical terms were not so completely standardized as those of plane geometry. Of the special types of prism the right parallelepiped is naturally

[1] Heath, *Euclid*, Vol. II, p. 97. [2] Euclid, II, 11, and VI, 30.

[3] So Pacioli gave to his work of 1509 the title *De diuina proportione*. Ramus (*Scholarvm Mathematicarvm, Libri vnvs et triginta*, Basel, 1569; *ibid.*, 1578; Frankfort, 1599, p. 191) referred to it in these words: "Christianis quibusdam divina quaedam proportio hic animadversa est . . ." ; and Kepler (Frisch ed. of his *Opera*, I, 377 (Frankfort, 1858)) spoke of it in the following terms: "Inter continuas proportiones unum singulare genus est proportionis divinae . . . ."

[4] The term seems to have come into general use in the 19th century. It is found in the *Archiv der Math. und Physik* (IV, 15–22) as early as 1844.

the most important. The word "parallelepiped"[1] means parallel surfaces. Although it is a word that would naturally be used by Greek writers, it is not found before the time of Euclid. It appears in the *Elements* (XI, 25) without definition, in the form of "parallelepipidal solid," the meaning being left to be inferred from that of the word "parallelogrammic" as given in Book I.

We have as yet no generally accepted name for a rectangular solid or right parallelepiped, nor had the Greeks. The word "cuboid" is significant and in modern times has had some sanction.

Euclid used *cubus*[2] for cube, and Heron (*c.* 50?) did the same. Heron also used "hexahedron"[3] for this purpose and then applied *cubus* to any right parallelepiped.

Although the Greeks knew that the diagonal of a right parallelepiped of edges a, b, c was $\sqrt{a^2+b^2+c^2}$, strangely enough the statement is not found in any of their works. It first appears, so far as now known, in the *Practica geometriae* of Fibonacci (1220).[4]

The word "prism" is Greek.[5] Euclid defines and treats it as we do at the present time.[6]

Pyramid. The Greeks probably obtained the word "pyramid"[7] from the Egpytian. It appears, for example, in the Ahmes Papyrus (*c.* 1550 B.C.). Because of the pyramidal form of a flame the word was thought by medieval and Renaissance writers to come from the Greek word for fire,[8] and so a pyramid was occasionally called a "fire-shaped body."[9]

[1] From παράλληλος (*paral'lelos*, parallel) + ἐπίπεδον (*epi'pedon*, plane surface).

[2] Κύβος, Latin *cu'bus*. See *Elements*, XI, def. 25.

[3] Ἑξάεδρον, from ἕξ (*hex*, six) + ἕδρα (*hed'ra*, seat).

[4] ". . . ut in Solido .aei. cuius dyameter sit linea .tb." (*Scritti*, II, 163).

[5] Πρίσμα (*pris'ma*), from πρίζειν (*pri'zein*), to saw; hence something sawed off.

[6] *Elements*, XI, def. 13.

[7] Πυραμίς (*pyramis'*), pl. πυραμίδες (*pyrami'des*), perhaps from the Egyptian *piromi*, but also thought to come from πυρός (*pyros'*, grain), as if a granary. On the uncertainty of the origin see Peet, *Rhind Papyrus*, p. 98.

[8] Πῦρ (*pyr*), as in "pyrotechnic."

[9] Thus the 16th century writer W. Schmid (*Das erste Buch der Geometrie*, Nürnberg, 1539) speaks of the "feuerförmige Cörper."

Euclid's treatment of the pyramid has remained substantially unchanged, except as to the proposition relating to the equivalence of pyramids of the same height and of equivalent bases. Cavalieri (1635) applied to this proposition his method of indivisibles, and Legendre[1] (1794) gave a simple proof that is now in common use. Āryabhaṭa (*c.* 510) gave the volume as half the product of the base and height, or at least it so appears in the extant manuscripts.[2]

Frustum of a Pyramid. The method of finding the volume of the frustum of a pyramid is found in Heron's *Stereometry*,[3] his rule reducing to the modern form of $V=\frac{1}{3}h(b+b'+\sqrt{bb'})$; but the actual rule first appears in Fibonacci's *Practica geometriae* (1220),[4] unless we accept a gloss upon an Arabic manuscript of the 12th century as evidence that it was known at that time.[5] The method itself was probably known to the Egyptians, at least for a special case, long before the time of Heron; for in a hieratic papyrus, apparently a little earlier than that of Ahmes, there is a statement which seems to show familiarity with the method for the case of a square pyramid.[6]

Brahmagupta (*c.* 628) also gave a rule[7] for the volume of a frustum of a pyramid with square base of sides s_1 and s_2, substantially as follows: $V=\frac{1}{3}h(s_1^2+s_2^2+s_1s_2)$.

[1] *Elémens*, 1st ed., VI, 17 (Paris, 1794). The order differs slightly in the different editions.

[2] Rodet, "Leçons de Calcul d'Âryabhaṭa," *Journal Asiatique*, XIII (7), 393; reprint, pp. 9, 10, 20.

[3] I, capp. 33, 34. See Tropfke, *Geschichte*, II (1), 383, with reference to the MS. of the Μετρικά (*metrika'*) discovered recently and published by Schöne, Leipzig, 1903. In his *Stereometry* he considers the pyramid with a square base; in the *Metrica*, with a triangular base.

[4] *Scritti*, II, 177.

[5] See the *Sitzungsberichte der physikalisch-medizinischen Societät zu Erlangen*, 50.–51. Band, p. 270, hereafter referred to as *Erlangen Sitzungsberichte*.

[6] This was first published by B. A. Touraeff (Turajev) in 1917, my attention being called to the fact by Professor R. C. Archibald. The manuscript is now in Moscow. See *Ancient Egypt*, Part III, p. 100 (London, 1917).

[7] Colebrooke ed., p. 312. On Āryabhaṭa see Rodet, *loc. cit.*, pp. 9, 10. On Bhāskara, see Colebrooke ed., p. 97. On Mahāvīra, see his work, p. 259. On the later treatment of the tetrahedron see Tropfke, *Geschichte*, II (1), 385. For the cylinder and cone, which are closely related to the prism, pyramid, and circle, see *ibid.*, p. 387.

Frustum of a Cone. The late Greeks knew how to find the volume of the frustum of a cone, deriving it from the rule that Heron used for the frustum of a pyramid, and thereafter the same method appeared in various mathematical treatises. Heron, however, used an approximation method which is probably of Egyptian origin, namely, that of taking the product of the altitude and the area of the circle midway between the bases.[1] An interesting example of the use of this approximation method has been found among the Greek papyri on arithmetic, being probably the work of a schoolboy of about the 4th century. The problems in this papyrus resemble those found in the Akhmîm Papyrus, which was written somewhat later. The first one of these problems relates to finding the contents of a circular pit of which the circumference is 20 cubits at the top and 12 cubits at the bottom, and of which the altitude is 12 cubits. The writer makes an error in his methods as well as his calculations, but endeavors to use Heron's approximation.[2]

Sphere. The word "sphere" comes to us from the Greek through the Latin.[3] The pure geometers of Greece had little interest in its measurement, although Archimedes tells us that "earlier geometers . . . have shown . . . that spheres are to one another in the triplicate ratio of their diameters."[4]

Archimedes also states that the volume of any sphere is four times that of the cone with base equal to a great circle of the sphere and with height equal to the radius of the sphere,—a statement that amounts to saying that $V=\frac{4}{3}\pi r^3$. He also stated that a cylinder with base equal to a great circle of the sphere and with height equal to the diameter of the sphere is equal to $1\frac{1}{2}$ times the sphere,—a statement that amounts to the same thing.[5]

[1] *Stereometrie*, ed. Hultsch, p. 157.

[2] J. G. Smyly, "Some Examples of Greek Arithmetic," *Hermathena*, XLII (1920), 105.

[3] Σφαῖρα (*sphai'ra*), Latin *sphaera*.

[4] Allman, *Greek Geom.*, p. 96; Archimedes, ed. Heiberg, II, 265; Heath, *Archimedes*, 234.

[5] Heiberg ed., Vol. I, *De sphaera et cylindro*.

Polyhedrons. The word "polyhedron"[1] is not found in the *Elements* of Euclid; he uses "solid," "octahedron," and "dodecahedron," but does not mention the general solid bounded by planes. The chief interest of the Greeks in figures of this type related to the five regular polyhedrons. It seems probable that Pythagoras (*c.* 540 B.C.)[2] brought his knowledge of the cube, tetrahedron, and octahedron from Egypt, but the icosahedron and the dodecahedron seem to have been developed in his own school. The Pythagoreans assigned the tetrahedron to fire, the octahedron to air, the icosahedron to water, the cube to earth, and the dodecahedron, apparently the last one discovered, to the universe. They seem to have known that all five polyhedrons can be inscribed in a sphere. They passed the study of these solids on to the school of Plato (*c.* 380 B.C.), where they attracted so much attention as to be known to later writers as "Platonic bodies" or "cosmic figures." It is not probable, however, that the early Pythagoreans actually constructed the figures in the sense that Euclid (*c.* 300 B.C.)[3] and Pappus (*c.* 300)[4] constructed them.

We have specimens extant of icosahedral dice that date from about the Ptolemaic period in Egypt.[5] There are also a number of interesting ancient Celtic bronze models of the regular dodecahedron still extant in various museums. There was probably some mystic or religious significance attached to these forms. Since a stone dodecahedron found in northern Italy dates back to a prehistoric period, it is possible that the Celtic people received their idea from the region south of the Alps,[6] and it is also possible that this form was already known in Italy when the Pythagoreans began their teaching in Crotona.

[1] From πολύς (*polys'*, many) + ἕδρα (*hed'ra*, seat).

[2] Heath, *Euclid*, Vol. III, p. 525.

[3] *Elements*, XIII, props. 13–17. See Heath, *Euclid*, Vol. III, pp. 467–503, with notes on the solutions suggested by Pappus.

[4] Pappus, ed. Commandino, p. 45 seq. (Bologna, 1660); ed. Hultsch, III, 142 seq.

[5] See the illustration on page 49.

[6] F. Lindemann, "Zur Geschichte der Polyeder und der Zahlzeichen," *Sitzungsberichte der math.-physik. Classe der K. Bayerischen Akad. der Wissensch. zu München*, Munich, XXVI, 625.

The five regular polyhedrons attracted attention in the Middle Ages chiefly on the part of astrologers. At the close of this period, however, they were carefully studied by various mathematicians. Prominent among the latter was Pietro Franceschi, whose work *De corporibus regularibus* (*c.* 1475) was the first to treat the subject with any degree of thoroughness. Following the custom of the time, Pacioli (1509) made free use of the works of his contemporaries, and as part of his literary plunder he took considerable material from this work and embodied it in his *De diuina proportione.*[1]

Albrecht Dürer, the Nürnberg artist, showed how to construct the figures from a net[2] in the way commonly set forth in modern works. The subject of stellar polyhedrons begins with Kepler[3] (1619) and has attracted considerable attention since his time.

Polyhedron Theorem. Among the most interesting of the modern formulas relating to a polyhedron is the one connecting the faces, vertices, and edges. This formula, often known as Euler's Theorem, may be stated as $f+v=e+2$. It was possibly known to Archimedes (*c.* 225 B.C.),[4] but not until the 17th century was it put into writing in a form still extant. Descartes (*c.* 1635) was the first to state it.[5] Euler seems to have come upon it independently. He announced it in Petrograd in 1752, but with simply an inductive proof. General proofs have been given by various writers.

Pappus-Guldin Theorem. Pappus of Alexandria (*c.* 300) stated, in substance, the basis of the theorem that the volume of a figure formed by the revolution of a plane figure about an axis is equal to the area of the figure multiplied by the length of the line generated by the center of gravity.[6] He also con-

[1] Venice, 1509.

[2] *Underweyssung der messung mit dem zirckel uñ richtscheyt, in Linien, ebnen unnd gantzen corporen*, Nürnberg, 1525.

[3] Frisch edition of his works, V, 126, where he speaks of such a figure as the "stella pentagonica" and carries the discussion "in solido."

[4] Tropfke, *Geschichte*, II (1), 398.

[5] But the fact was not made known until his *Œuvres inédites* appeared in 1860, where it appears on page 218.

[6] Hultsch edition of his works, p. 682.

sidered the case in which there was not a complete revolution. The proposition was soon forgotten and so remained until revived by Kepler (1615),[1] who extended the theory to include the study of the revolution of various plane figures. He gave special attention to the torus formed by the revolution of a circle or an ellipse. When the axis was tangent to the circle or the ellipse, he called the resulting solid an *annulus strictus*. Although he treats of only special cases, he doubtless knew the general theorem.

Various writers made use of the general principle in the 17th century, but it was brought into special prominence by Habakuk Guldin[2] (1577–1643), a Swiss scholar. It appeared in Book II of his *Centrobaryca* (1641). He added nothing to the theory except as he stated the general proposition.

In 1695 Leibniz suggested that the proposition could be extended to include the case of a plane revolving about an axis on any other path than a circle, provided it is always perpendicular to this path,[3] an idea that was considered also by Euler in 1778.

7. The Three Famous Problems

Nature of the Problems. The Greeks very early found themselves confronted by three problems which they could not solve, at least by the use of the unmarked ruler and the compasses alone.[4]

The first was the trisection of any angle. The trisection of the right angle was found to be simple, but the trisection of any arbitrary angle whatever attracted the attention and baffled the efforts of many of their mathematicians. To this problem may be added the related ones of dividing any given angle into any required number of equal parts and of inscribing in a circle a regular polygon of a given number of sides.

[1] In his *Stereometria Doliorum*. See the *Opera Omnia*, ed. Frisch, IV, 551–670.

[2] Known as Paul Guldin after he entered the Catholic Church.

[3] *Acta Eruditorum*, 1695, p. 493.

[4] See any of the histories of Greek mathematics. A good summary is given in H. G. Zeuthen, *Histoire des Mathématiques dans l'Antiquité et le Moyen Age*, French translation by J. Mascart, p. 57 (Paris, 1902).

The second problem was the quadrature of the circle, that is, the finding of a square whose area is the same as that of a given circle. The solution would be simple if we could find a straight line that is equal in length to the circumference of the circle; that is, if we could rectify the circumference. This is easily accomplished by rolling the circle along a straight line, but such a proceeding makes use of an instrument other than the ruler and compasses, namely, of a cylinder with a marked surface.

The third problem was the duplication of a cube,[1] that is, the finding of an edge of the cube whose volume is twice the volume of a given cube. This was known as the Delian Problem, one story of its origin being that the Athenians appealed to the oracle at Delos to know how to stay the plague which visited their city in 430 B.C. It is said that the oracle replied that they must double in size the altar of Apollo. This altar being a cube, the problem was that of its duplication. Since problems about the size and shape of altars appear in the early Hindu literature, it is not improbable that this one may have found its way, perhaps through Pythagoras, from the East. It was already familiar to the Greeks in the 5th century B.C., for we are told by Eratosthenes[2] that Euripides (*c.* 485–406 B.C.) refers to it in one of his tragedies which is no longer extant.[3]

Trisection Problem: the Conchoid. There are various ways of trisecting any plane angle, but it will suffice at this time to give only a single one. Probably the best known of the Greek attempts is the one made by Nicomedes (*c.* 180 B.C.). He used a curve known as the conchoid.[4] We take a fixed point O which


[1] N. T. Reimer, *Historia problematis de cubi duplicatione* (Göttingen, 1798); C. H. Biering, *Historia problematis cubi duplicandi* (Copenhagen, 1844); Archimedes, *Opera*, ed. Heiberg, III, 102; A. Sturm, *Das Delische Problem* (Linz, 3 parts, 1895, 1896, 1897), a critical historical study with extensive bibliography.

[2] See Archimedes, *Opera*, ed. Heiberg, III, 102.

[3] Too small hast thou designed the royal tomb.
Double it; but preserve the cubic form.


[4] Thus Proclus: "Nicomedes trisected every rectilineal angle by means of the conchoidal lines, the inventor of whose particular nature he is, and the origin, construction, and properties of which he has explained. Others have solved the same problem by means of the quadratrices of Hippias and Nicomedes

is d distant from a fixed line AB, and we draw OX parallel to AB and OY perpendicular to OX. We then take any line OA through O, and on OA produced lay off $AP = AP' = k$, a constant. Then the locus of points P and P' is a conchoid. According as $k \gtrless d$ we have O a node, a cusp, or a conjugate point. The equation of the curve is $(x^2+y^2)(x-d)^2 - k^2x^2 = 0$.



In order to trisect a given angle we proceed as follows:

Let YOA be the angle to be trisected. From A construct AB perpendicular to OY. From O as pole, with AB as a fixed straight line and 2 AO as a



TRAMMEL FOR CONSTRUCTING THE CONCHOID

From Bettini's *Apiaria Universae Philosophiae Mathematicae*, Bologna, 1641

constant distance, describe a conchoid to meet OA produced at P and to cut OY at Q. At A construct a perpendicular to AB

. . .; others, again, starting from the spirals of Archimedes" (Proclus, ed. Friedlein, p. 272 (translation by Allman); see also Gow, *Greek Geom.*, p. 266; Heath, *History*, I, 235, 238).

meeting the curve at T. Draw OT and let it cut AB at N. Let M be the mid-point of NT.

Then $MT = MN = MA$.


But $NT = 2\,OA$ by construction of the conchoid.

Hence $MA = OA$.

Hence $\angle AOM = \angle AMO = 2\angle ATM = 2\angle TOQ$.


That is, $\angle AOM = \frac{2}{3}\angle YOA$, and $\angle TOQ = \frac{1}{3}\angle YOA$.

The Quadratrix. A Greek geometer named Hippias, probably Hippias of Elis[1] (*c.* 425 B.C.), invented a curve which he used in the trisection of an angle. In this figure, X is any point on the quadrant AC. As the radius OX revolves at a uniform rate from the position OC to the position OA, the line MN moves at a uniform rate from the position CB to the position OA, always remaining parallel to OA. Then the locus of P, the intersection of OX and MN, is a curve CQ. Manifestly, when OX is one nth of the way from OC around to OA, MN is one nth of the way from CB down to OA. If, therefore, we make $CM = \frac{1}{3}CO$, MN will cut CQ at a point P' such that OP' will trisect the right angle. In the same way, by trisecting OM we can find a point P' on CQ such that OP' will trisect angle AOX, and so for any other angle. The method evidently applies to the multisection as well as to the trisection of an angle.



Other Methods of Trisection. The next prominent investigation of the problem is one that is attributed to Archimedes[2] (*c.* 225 B.C.), although probably not due to him in the form that has come down to us. The plan is as follows:

Produce any chord AB of a circle until the part produced, BC, is equal to r, the radius. Join C to the center O and produce CO to the circle at D.




[1] But this has been questioned. For the arguments, see Allman, *Greek Geom.*, p. 93. See also Volume I, page 82, n., and Heath, *History*, I, 225.

[2] See Heath, *Archimedes*; Allman, *Greek Geom.*, p. 90, with references.

Then arc AD is three times arc EB; that is, $\angle EOB$ is $\frac{1}{3}\angle AOD$. This, however, is manifestly no solution of the problem.

Vieta (*c.* 1590) was led by this to suggest the following:

Let $\angle AOB$ be the angle to be trisected. Describe any circle with center O. Suppose the problem solved and that $\angle AOP = \frac{1}{3}\angle AOB$. Through B suppose BRQ drawn parallel to PO. Then $\angle RBO = \angle ORB = 2\angle AOP = 2\angle Q$. But $\angle ORB = \angle Q + \angle ROQ$, and hence $\angle Q = \angle ROQ$, and so $OR = QR$. The problem is therefore reduced to the following:



From B draw BRQ so that the part RQ, intercepted between the circle and the diameter AS produced, shall be equal to the radius, a construction involving the use of a marked straightedge.

Regular Polygons. If we can trisect an angle of 360° we can inscribe a regular polygon of three sides in a circle, and similarly for the inscription of other regular polygons. The trisection problem therefore naturally suggests the larger problem of the inscription of a polygon of any given number of sides. It was for a long time believed that the Greeks had exhausted all the possibilities in this line. In 1796, however, Gauss showed that it was possible, by the use of the straightedge and compasses alone, to inscribe a polygon of 17 sides. He even extended the solution to include polygons of 257 and 65,537 sides. The general proposition, as it now appears, is as follows: A regular polygon of p sides, where p is a prime number greater than 2, can be constructed by ruler and compasses if and only if p is of the form $2^{2^t} + 1$. For $t = 0, 1, 2, 3$, and 4 the values of p are respectively 3, 5, 17, 257, and 65,537; but if $t = 5$, $p = 641 \times 6700417$ and hence is not prime.[1] Gauss has left the following interesting record of the discovery:

The day was March 29, 1796, and chance had nothing to do with it. Before this, indeed during the winter of 1796 (my first semester in

[1]For the theory see J. W. A. Young, *Monographs on Modern Mathematics* (New York, 1911), article by L. E. Dickson, p. 378, with references.

Göttingen), I had already discovered everything related to the separation of the roots of the equation

$$\frac{x^p - 1}{x - 1} = 0$$

into two groups. . . . After intensive consideration of the relation of all the roots to one another on arithmetical grounds, I succeeded, during a holiday in Braunschweig, on the morning of the day alluded to (before I had got out of bed), in viewing the relation in the clearest way, so that I could immediately make special application to the 17-side and to the numerical verification. . . . I announced this discovery in the *Literaturzeitung* of Jena, where my advertisement was published in May or June, 1796.[1]

Squaring the Circle. The second famous problem of antiquity was that of squaring the circle. The first attempts were of course empirical. They were made long before the scientific period of the Greek civilization, and they naturally resulted in rude approximations.

The first definite trace that we have of an approximate value of π is in the Ahmes Papyrus[2] (*c.* 1550 B.C.). There is given in that work a problem requiring the finding of the area of a circle, the method, expressed in modern symbols, being as follows:

$$A = (d - \tfrac{1}{9}\, d)^2,$$

which amounts to saying that $\pi = 3.1605 -$, a result apparently arrived at empirically, as already stated on page 270.

It is probable that 3 is a much older value of π, although we have no extant literature to prove this fact. We find such a value in early Chinese works,[3] in the Bible,[4] in the Talmud,[5]

[1] R. C. Archibald, "Gauss and the Regular Polygon of Seventeen Sides," *Amer. Math. Month.*, XXVII, 323, with bibliography.

[2] Peet, *Rhind Papyrus*, p. 90; Eisenlohr, *Ahmes Papyrus*, p. 117; G. Vacca, "Sulla quadratura del circolo secondo l' Egiziano Ahmes," *Bollettino di bibliografia e storia delle scienze matematiche*, XI, 65.

[3] *E.g.*, in the *Chóu-peï*. See Mikami, *China*, pp. 8, 46, 135. It should be mentioned again that there are doubts as to the reliability of ancient Chinese texts.

[4] "And he made a molten sea, ten cubits from the one brim to the other: it was round all about . . . and a line of thirty cubits did compass it round about" (1 Kings, vii, 23. See also 2 Chronicles, iv, 2).

[5] In both Mishna and Talmud the value is always 3, the reason being traditional, based upon Solomon's "molten sea."

in the early Hindu works,[1] and in the medieval manuscripts, so that it was generally accepted in all countries and until relatively modern times.

The Greeks were not content with results that were merely empirical, however, and so the rectification of the circumference or the related problem of the squaring of the circle attracted the attention of their philosophers. For example, Anaxagoras (*c.* 440 B.C.) is said by Plutarch[2] to have been put in prison in Athens, and while there to have first attempted the solution. The results of his work are, however, unknown.

Methods of Attacking the Quadrature. There are three methods of attacking the problem: first, by the use of the ruler and compasses only; second, by the use of higher plane curves; third, by such devices as infinite series, leading to close approximations. The leading Greek mathematicians seem to have found the futility of the first method, although they did not prove that it is impossible; with the second method they were successful; with the third method they were less skillful.[3]

Method of Exhaustion. Antiphon (*c.* 430 B.C.) attempted the quadrature by inscribing a polygon (some early writers say a square and others a triangle), and then doubling the number of sides successively until he approximately exhausted the area between the polygon and the circle. By finding the area of each polygon he was thus able to approximate the area of the circle.[4]

Attempts of Hippocrates. Hippocrates of Chios (*c.* 460 B.C.) attempted the solution and was the first to actually square a curvilinear figure. He constructed semicircles on the three


[1] "The diameter and the square of the semidiameter, being severally multiplied by three, are the practical circumference and area. The square-roots extracted from ten times the squares of the same are the neat values." Colebrooke, *Brahmagupta*, p. 308. See also Mahāvīra, p. 189; Colebrooke, *Bhāskara*, p. 87.

[2] *De exilio*, cap. 17, ed. Dübner-Didot of the *Moralia*, I, 734 (Paris, 1885). See also the Leipzig edition of 1891, III, 573.

[3] Heath, *History*, I, 220.


[4] F. Rudio, "Der Bericht des Simplicius über die Quadraturen des Antiphon und des Hippokrates," in *Bibl. Math.*, III (3), 7, and also in book form, with Greek text (Leipzig, 1907); Allman, *Greek Geom.*, pp. 64, 81. With respect to Bryson see *ibid.*, pp. 77, 82; but compare also Volume I, page 84.

sides of an isosceles right-angled triangle and showed that the sum of the two lunes thus formed is equal to the area of the triangle itself. Having a triangle equal in area to a lune, he had only to construct a square equal to the triangle. His proof involves the proposition that the areas of circles are proportional to the squares of their diameters,



—a proposition which Eudemus (*c.* 335 B.C.) tells us that Hippocrates proved.[1] To the quadrature problem as such, however, his contribution was not important. His method of attack was substantially as follows:

In a semicircle *ABCD*, center *O*, he inscribed half of a regular hexagon, *h*. On the three sides and on *OB* he described semicircles as here shown. Then the four small semicircles are together equal to the large semicircle. Subtracting the common shaded parts, the three lunes together with the semicircle on *OB* are equal to *h*, the half of the regular hexagon. Now take from *h* a surface equal to the sum of the lunes, which can be found by the method already given (and here is the fallacy), and there remains a rectilinear figure equal to the semicircle on *OB*. It will be observed that Hippocrates assumed that every lune can be squared, whereas he has shown, as we have seen, that this is possible only in the special case of a right triangle.[2]



[1] *Eudemi fragmenta*, ed. Spengel, p. 128. The proposition is equally true for any right-angled triangle, but Hippocrates proved it only for the isosceles case. See Allman, *Greek Geom.*, p. 66; W. Lietzmann, *Der Pythagoreische Lehrsatz*, p. 32 (Leipzig, 1912); E. W. Hobson, *Squaring the Circle*, p. 16 (Cambridge, 1913); Heath, *History*, I, 183, with a summary of recent literature on the subject.

[2] For details relating to the work of Hippocrates on lunes in general, see Heath, *History*, I, 183–201; Allman, *Greek Geom.*, p. 69.


The Quadratrix. The next noteworthy attempt was made by Deinostratus (*c.* 350 B.C.). Pappus[1] makes this statement:

> For the quadrature of the circle a certain curve was employed by Deinostratus, Nicomedes, and some other more recent geometers, which has received its name from the property that belongs to it; for it is called by them the quadratrix.[2]

This is the curve used by Hippias in the trisection of an angle. In the figure given below it can be shown that

$$\frac{CXA}{CO} = \frac{CO}{OQ};$$

and since these terms are all straight lines except the quadrant CXA, it is possible to construct a straight line equal in length to the quadrant, and hence to rectify the circumference.



To prove the proposition Pappus states that the *reductio ad absurdum* was employed. If

$$\frac{CXA}{CO} \neq \frac{CO}{OQ},$$

then the proportion can be made true by increasing or by decreasing OQ; but it will be shown that this cannot be done without leading to an absurdity.

First, suppose that $\dfrac{CXA}{CO} = \dfrac{CO}{OA'}$. $OA' > OQ$

Then, since $\dfrac{CXA}{C'PA'} = \dfrac{CO}{C'O} = \dfrac{CO}{OA'}$,

it follows that $C'PA' = CO$.

And since, from the property of the curve,

$$\frac{CXA}{XA} = \frac{CO}{PA''},$$

we have $$\frac{CXA}{XA} = \frac{C'PA'}{PA'} = \frac{CO}{PA'};$$


[1] Pappus, *Collectiones*, IV, cap. xxx (ed. Hultsch, I, 253); Hankel, *Geschichte*, p. 151; Cantor, *Geschichte*, I (2), 233; Heath, *History*, I, 226.

[2] Τετραγωνίζουσα (*tetragoni'zousa*).

whence arc $PA' = PA''$, which is impossible, since an arc cannot be equal to its chord, and since their halves cannot be equal.

Next, suppose that $\dfrac{CXA}{CO} = \dfrac{CO}{OA''}$. $OA'' < OQ$

Draw the quadrant $C''MA''$.



Then $\dfrac{C''MA''}{MA''} = \dfrac{CXA}{XA} = \dfrac{CO}{PA''}$.

Therefore $\dfrac{CXA}{C''MA''} = \dfrac{CO}{C''O} = \dfrac{CO}{OA''}$.

But $\dfrac{CXA}{CO} = \dfrac{CO}{OA''}$. By hyp.

Hence CO must be equal to $C''MA''$ if the hypothesis is correct.

But $$\frac{C''MA''}{MA''} = \frac{CXA}{XA},$$

and because CPQ is a quadratrix we have

$$\frac{CXA}{XA} = \frac{CO}{PA''};$$

whence $$\frac{C''MA''}{MA''} = \frac{CO}{PA''}.$$

But in the same way as it was shown in the first part of the proof that

$$C'PA' = CO$$

on the hypothesis there made, so it may be shown here that

$$C''MA'' = CO$$

on this hypothesis. It then follows from this proportion that

$$MA'' = PA'',$$

which leads to the absurdity that the circumference of a circle is equal to the perimeter of a circumscribed polygon.

Hence the second hypothesis is also untenable.

Hence $$\frac{CXA}{CO} = \frac{CO}{OQ},$$

and CXA, a quadrant, can be constructed.

Method of Archimedes. The next noteworthy contribution was that of Archimedes (*c.* 225 B.C.), who asserted that:

1. The area of a circle is equal to the area of a right-angled triangle one of whose sides forming the right angle is equal to the circumference of the circle and the other to the radius.[1]

2. The ratio of the area of a circle to the square on the diameter is approximately 11 : 14.

3. The ratio of the circumference of a circle to the diameter is less than $3\frac{1}{7}$ and greater than $3\frac{10}{71}$.

To prove the third proposition Archimedes inscribed and circumscribed regular polygons, found their areas up to polygons of 96 sides, and showed that the area of the circle lies between these results. These limits, expressed in modern decimal form, are 3.14285714 · · · and 3.14084507 · · ·. If our present notation and our methods of finding a square root had been known, the result would have been closer, since the geometric method permitted of any desired degree of approximation.

The Romans were little concerned with accurate results in such matters as this, and so it is not surprising that Vitruvius (*c.* 20 B.C.) speaks of the circumference of a wheel of diameter 4 feet as being $12\frac{1}{2}$ feet, thus taking π as $3\frac{1}{8}$.[2]

Other Greek Approximations of π. After the time of Archimedes the value $3\frac{1}{7}$ became recognized as a satisfactory approximation and appeared in the works of Heron (*c.* 50?), Dominicus Parisiensis (1378), Albert of Saxony (*c.* 1365), Nicholas Cusa (*c.* 1450), and others. Since one of the common approximations for a square root in the Middle Ages was

$$\sqrt{n} = \sqrt{a^2 + r} = a + \frac{r}{2a + 1},$$

and since this gives $\sqrt{10} = 3 + \frac{1}{2 \times 3 + 1} = 3\frac{1}{7}$, it is natural to expect that $\sqrt{10}$, which is 3.1623 · · ·, would often have been given as the value of π, and this was in fact the case.

[1] Heath, *Archimedes*, p. 231–233; ed. Heiberg, I, 258; Heath, *History*, II, 50–56.

[2] *De Architectura*, X, cap. 14. Rose's edition (1889) gives the diameter as $4\frac{1}{2}$ feet, which would make π less than 3. See *Bibl. Math.*, I (3), 298.

Ptolemy (*c.* 150) seems to have taken the Archimedean limits and to have expressed them in sexagesimals, obtaining substantially $3\frac{1}{7} = 3\ 8'\ 34.28''$ and $3\frac{10}{71} = 3\ 8'\ 27.04''$. He then improved upon the mean between these results by taking $3\ 8'\ 30''$ as the approximate value of π, although a still closer approximation is $3\ 8'\ 29.73355''$. Since $3\ 8'\ 30'' = 3.1416$, his result was very satisfactory.[1]

Hindu Values of π. The Hindu mathematicians took various values of π, and no writer among them seems to have been uniform in his usage.

Āryabhaṭa (*c.* 510), or possibly Āryabhaṭa the Younger, gave the equivalent of 3.1416, his rule being:

> Add 4 to 100, multiply by 8, add 62,000, and you have for a diameter of two *ayutâs* the approximate value of the circumference.[2]

Brahmagupta (*c.* 628) criticized Āryabhaṭa for taking the circumference as 3393 for both diameters 1080 and 1050, which would make π either $3\frac{17}{120}$ or $3\frac{81}{350}$; that is, 3.1416 or 3.2314.

A certain astronomer, Puliśa,[3] to whom Brahmagupta refers, gives $3\frac{177}{1250}$, which is 3.18 +, and Ya'qûb ibn Ṭâriq (*c.* 775) mentions certain Hindu astronomical measurements which give the same value. He also states that Puliśa used a value equivalent to 3.14183, and Brahmagupta a value equivalent to 3.162. For himself Ya'qûb ibn Ṭâriq used in one case a value equivalent to 3.1411.[4]

In case the value 3.1416 is due to either of the Āryabhaṭas, it may have been obtained from the Alexandrian scholars,

[1] Heath, *History*, I, 233.

[2] L. Rodet, "Leçons de Calcul d'Âryabhaṭa," *Journal Asiatique* as cited, reprint, p. 11. There is some doubt as to whether this rule is due to either of the Āryabhaṭas; see G. R. Kaye, "Notes on Indian Mathematics, No. 2, Āryabhaṭa," in *Journ. and Proc. of the Asiatic Soc. of Bengal*, IV, reprint. The word *ayutâs* means myriads, that is, 10,000's. The rule is translated more simply in Volume I, page 156.

[3] The name appears as Paulisa, Pulisa, and Pauliśa. For a discussion, see Sachau's translation of Albêrûnî's *India*, II, 304 (London, 1910); hereafter referred to as Albêrûnî's *India*. On Brahmagupta's criticism of Āryabhaṭa, see I, 168. Nothing is known concerning the life of Puliśa.

[4] Albêrûnî's *India*, II, 67.

by whom it was then known and whose works may well have reached India, or it may have been found independently.

Brahmagupta[1] (*c.* 628) used 3 as the "practical" value and $\sqrt{10}$ as the "exact" value, and these values are also given by Mahāvīra[2] (*c.* 850) and Srīdhara[3] (*c.* 1020).

Bhāskara (*c.* 1150) used $\frac{3927}{1250}$ for the "near" value and $\frac{22}{7}$ in finding the "gross circumference adapted to practice,"[4] the former being the same as the value $3\frac{177}{1250}$ of Pulisa.[5]

Chinese Values of π. The Chinese found various values of π, but the methods employed by the early calculators are unknown. The value 3 was used probably as early as the 12th century B.C.[6] and is given in the *Chóu-peï* and the *Nine Sections*.[7] Ch'ang Höng (*c.* 125) used $\sqrt{10}$, and Wang Fan (*c.* 265) used $\frac{142}{45}$, which is equivalent to 3.1555 ···. Liu Hui (263) gives us the first intimation of the method used by the Chinese in finding the value. He begins with a regular inscribed hexagon, doubles the number of sides repeatedly, and asserts that "if we proceed until we can no more continue the process of doubling, the perimeter ultimately comes to coincide with the circle."[8]

Among other early Chinese values of no high degree of accuracy are those of Men (*c.* 575), who gave 3.14, and Wu (*c.* 450), whose value was 3.1432 +.

Tsu Ch'ung-chih (*c.* 470) was able, by starting with a circle of diameter 10 feet, to obtain 3.1415927 and 3.1415926 for the limits of π, and from these, by interpolation, he obtained the "accurate and inaccurate" values $\frac{355}{113}$ and $\frac{22}{7}$. No closer approximations were made in China until modern times.[9]

[1] Colebrooke's translation, p. 308. [2] Mahāvīra, pp. 189, 200.

[3] Colebrooke's translation, p. 87.

[4] Colebrooke's translation, p. 87.

[5] On the general subject of the Hindu quadratures see C. M. Whish, *On the Hindú Quadrature of the Circle*, a paper read before the Madras Literary Society, December 15, 1832. [6] Mikami, *China*, p. 46.

[7] For discussion of the dates of these works, see Volume I, page 31.

[8] For his computations, see Mikami, *China*, p. 48.

[9] On the work of the later writers, after the introduction of European mathematics, see Mikami, *China*, p. 135. In none of these early approximations was the decimal fraction used.

The Japanese did no noteworthy work in this field until the 17th century. They then developed a kind of native calculus and also made use of European methods which gave them fair approximations to the required ratio.[1]

Later Approximations of π. The following is a brief summary of some of the later European approximations of π, with the names of those who used them:

Franco of Liège[2] (*c.* 1066), $\pi = \frac{22}{7} = 3.142857+$.

Fibonacci[3] (1220), $\pi = \frac{864}{274} = 3.141818$. He also gave the limits as 3.1427 and 3.1410.

Al-Kashî[4] (*c.* 1430), 3.14159265358979 32.

Tycho Brahe[5] (*c.* 1580), $\pi = \frac{88}{\sqrt{785}} = 3.1409$.

Simon Duchesne[6] (*c.* 1583), $\pi = 3\frac{69}{484} = (\frac{39}{22})^2 = 3.14256198$.

Vieta (*c.* 1593), $3.1415926535 < \pi < 3.1415926537$.

Adriaen van Roomen (1561–1615) gave π to 17 decimal places.

Ludolf van Ceulen (1540–1610) gave π to 35 decimal places, and German textbooks still speak of π as the "Ludolphische Zahl."

Adriaen Anthoniszoon (*c.* 1600) and his son Adriaen Metius (1571–1635), $\pi = \frac{355}{113}$, the Chinese value.[7]

J. H. Lambert[8] (*c.* 1770),

$$\pi = (\tfrac{7}{4})^2,\ (\tfrac{16}{9})^2,\ (\tfrac{62}{35})^2,\ (\tfrac{39}{22})^2,\ (\tfrac{248}{123})^2,\ (\tfrac{296}{167})^2,\ \cdots.$$

[1] Smith-Mikami, pp. 60, 63, *et passim.*

[2] *Abhandlungen*, IV, 139. The decimal equivalents are modern in all these cases.

[3] "Practica geometriae," in his *Scritti*, II, 90. See also H. Weissenborn, "Die Berechnung des Kreisumfanges bei Archimedes und Leonardo Pisano," *Berliner Studien für klassische Philologie und Archäologie*, XIV.

[4] See Volume I, pages 289, 290; Volume II, pages 238, 240.

[5] Original name, Tyge Ottesen. It is not known how he came to give this curious value. See F. J. Studnička, *Bericht über die Astrologischen Studien des . . . Tycho Brahe*, p. 49 (Prag, 1901).

[6] Cantor, *Geschichte*, II (2), 592.

[7] They took the approximation $3\frac{15}{106} < \pi < 3\frac{17}{120}$, added the numerators $(15 + 17 = 32)$ and the denominators $(106 + 120 = 226)$, took the means (16 and 113), and gave $\pi = 3\frac{16}{113} = \frac{355}{113} = 3.1415929$, a very close approximation for the time. Priority for this is claimed for Valentinus Otto (*c.* 1550–1605).

[8] *Vorläufige Kenntnisse für die, so die Quadratur und Rektifikation des Circuls suchen*, II, 140 (Berlin, 1772).

The value of π was carried to 140 decimal places (136 correct) by Georg Vega (1756–1802), to 200 by Zacharias Dase (1824–1861), to 500 by Richter (died in 1854), and to 707 by William Shanks (*c.* 1853).

Continued Products and Series. Vieta (*c.* 1593) gave another interesting approximation for π, using continued products for the purpose. His value may be obtained from the following equation:

$$\frac{2}{\pi} = \sqrt{\tfrac{1}{2}} \cdot \sqrt{\tfrac{1}{2} + \tfrac{1}{2}\sqrt{\tfrac{1}{2}}} \cdot \sqrt{\tfrac{1}{2} + \tfrac{1}{2}\sqrt{\tfrac{1}{2} + \tfrac{1}{2}\sqrt{\tfrac{1}{2}}}} \cdots.$$

John Wallis (1655)[1] gave the form

$$\frac{4}{\pi} = \frac{3 \cdot 3 \cdot 5 \cdot 5 \cdot 7 \cdot 7 \cdot 9 \cdot 9 \cdot 11 \cdot 11 \cdots}{2 \cdot 4 \cdot 4 \cdot 6 \cdot 6 \cdot 8 \cdot 8 \cdot 10 \cdot 10 \cdot 12 \cdots}.$$

This is related to Lord Brouncker's value (*c.* 1658) in which use is made of continued fractions, as follows:

$$\frac{4}{\pi} = 1 + \cfrac{1}{2 + \cfrac{9}{2 + \cfrac{25}{2 + \cfrac{49}{2 + \cdots}}}}.$$[2]

Leibniz[3] (1673),

$$\frac{\pi}{4} = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \frac{1}{9} - \cdots.$$

Abraham Sharp (*c.* 1717),

$$\frac{\pi}{6} = \sqrt{\tfrac{1}{3}} \cdot \left(1 - \frac{1}{3 \cdot 3} + \frac{1}{3^2 \cdot 5} - \frac{1}{3^3 \cdot 7} + \frac{1}{3^4 \cdot 9} - \cdots\right),$$

from which he found the value of π to 72 decimal places.

[1] *Arithmetica Infinitorum* (1655), included in his *Opera*, I, 469.

[2] See L. Euler, *Opuscula analytica*, Vol. I (Petrograd, 1783); also (1785), II, 149.

[3] A special case of Gregory's (1671) series. De Lagny (1682) discovered it independently.

John Machin[1] (*c.* 1706),

$$\frac{\pi}{4} = 4 \cdot \left(\frac{1}{5} - \frac{1}{3 \cdot 5^3} + \frac{1}{5 \cdot 5^5} - \frac{1}{7 \cdot 5^7} + \cdots\right) - \left(\frac{1}{239} - \frac{1}{3 \cdot 239^3} + \frac{1}{5 \cdot 239^5} - \frac{1}{7 \cdot 239^7} + \cdots\right).$$

Matsunaga Ryōhitsu[2] (1739), a Japanese writer,

$$\frac{\pi}{3} = 1 + \frac{1^2}{4 \cdot 6} + \frac{1^2 \cdot 3^2}{4 \cdot 6 \cdot 8 \cdot 10} + \frac{1^2 \cdot 3^2 \cdot 5^2}{4 \cdot 6 \cdot 8 \cdot 10 \cdot 12 \cdot 14} + \cdots.$$

The Symbol π. The symbol $\frac{\delta}{\pi}$ was used by Oughtred (1647) to represent the ratio of the diameter to the circumference.[3] Isaac Barrow (from *c.* 1664) used the same symbolism, and David Gregory (1697) used $\frac{\pi}{\rho}$ for the ratio of circumference to radius.

The first to use π definitely to stand for the ratio of c to d was an English writer, William Jones. In his *Synopsis Palmariorum Matheseos* (1706) he speaks (p. 243) of "Periphery (π)"; but on p. 263 he is more definite, giving

"3.14159, &c. $= \pi$,"

"$d = c \div \pi = \overline{a \div \frac{1}{4}\pi}\,|^{\frac{1}{2}}$,"

"$c = d \times \pi = \overline{a \times 4\pi}\,|^{\frac{1}{2}}$,"

and "$a = \frac{1}{4}\pi \times d^2 = c^2 \div 4\pi$."

Euler adopted the symbol in 1737, and since that time it has been in general use.

[1] W. Jones, *Synopsis Palmariorum Matheseos*, p. 243 (London, 1706), gives π "True to above a 100 Places; as Computed by the Accurate and Ready Pen of the Truly Ingenious Mr. John Machin."

[2] For a further list of values of π consult D. E. Smith, "The History and Transcendence of π," on page 396 of J. W. A. Young, *Monographs on . . . Modern Mathematics* (New York, 1911); Tropfke, *Geschichte*, IV (2), 195; F. Rudio, *Archimedes, Huygens, Lambert, Legendre. Vier Abhandlungen über die Kreismessung*, Leipzig, 1892.

[3] "Si in circulo sit 7.22 :: $d \cdot \pi$ :: 113.355: erit $\delta \cdot \pi$ :: 2R.P: periph." This symbolism appears first in the 1647 edition of the *Clavis Mathematicae* (1631). This quotation is from the 1652 edition. See Cajori, *Oughtred*, p. 32.

The proof of the transcendence of π was first given by F. Lindemann (1882), thus showing the impossibility of squaring the circle by the use of ruler and compasses alone.[1]

Duplication of the Cube.[2] Hippocrates of Chios (*c.* 460 B.C.) showed that the problem of duplicating the cube resolves itself into the finding of two mean proportionals between two given lines. If $a:x = x:y = y:b$, then $x^2 = ay$, $y^2 = bx$, and hence $x^4 = a^2y^2 = a^2bx$, or $x^3 = a^2b$. If $b = 2a$, then $x^3 = 2a^3$. That is, the cube of edge x will then have double the volume of a given cube with edge a. Since we have the three equations $x^2 = ay$ (parabola), $y^2 = bx$ (parabola), and $ab = xy$ (hyperbola), we can evidently solve the problem by finding the intersection of two parabolas or of a parabola and a hyperbola. These methods are credited to Menæchmus (*c.* 350 B.C.).[3]

Archytas (*c.* 400 B.C.) had already found the two mean proportionals, solving the problem by means of two cylindric sections, for Eratosthenes (*c.* 230 B.C.) tells us that he "is said to have discovered them by means of his semicylinders."[4] It is possible that Archytas led Menæchmus to discover a solution by means of conics.

Eratosthenes also tells us that Eudoxus (*c.* 370 B.C.) solved the problem "by means of the so-called curved lines," but what these lines were we do not know.[5] The two statements here attributed to Eratosthenes are contained in a letter formerly (but incorrectly) credited to him. In the main, however, this letter sets forth facts with which he was familiar, as is shown from other sources.

[1] D. E. Smith, "The History and Transcendence of π," *loc. cit.*, p. 387; consult this work also with respect to transcendental numbers in general.

[2] J. S. Mackay, *Proc. of the Edinburgh Math. Soc.*, IV, 2; F. G. Teixeira, *Obras sobre Mathematica*, VII, 283–415 (Coimbra, 1915); C. H. Biering, *Historia Problematis Cubi Duplicandi*, Copenhagen, 1844; Heath, *History*, I, 244; F. Enriques, *Fragen der Elementar-Geometrie*, II. Teil (Leipzig, 1907).

[3] On this point see Heath, *History*, I, 251–255; Allman, *Greek Geom.*, p. 160.


[4] In his letter to Ptolemy III. See Archimedes, *Opera*, ed. Heiberg, III, 104, 106. On the solution, as Eudemus relates it, see Allman, *Greek Geom.*, p. 111. See also P. Tannery, "Sur les solutions du problème de Délos par Archytas et par Eudoxe," in his *Mémoires Scientifiques*, I, 53 (Paris, 1912); Heath, *History*, I, 246.

[5] Archimedes, *Opera*, ed. Heiberg, III, 66.

Plato (*c.* 380 B.C.) is said to have solved the problem by means of a mechanical instrument[1] but to have rejected this method as not being geometric.

We are told by Joannes Philop'onus[2] that Apollonius (*c.* 225 B.C.) had a method of finding the two mean proportionals. The construction, however, assumes a postulate which begs the whole question.[3]

Cissoid of Diocles. One of the best-known of the ancient solutions was that of Diocles (*c.* 180 B.C.), who used a curve known as the cissoid (from κισσοειδής (*kissoeides'*), ivylike). In this figure, if $OL = OT$ and AQ is drawn, then P is a point on the curve. Similarly, AS will determine on TQ produced a point P' on the curve. The cissoid evidently passes through A, BR is an asymptote, and the curve is symmetric with respect to AB.



By the aid of the curve two mean proportionals can be found in the following manner:[4]

Let $OM = \frac{1}{2}r$, determine P by producing BM to the curve, and draw AP and produce it to R, letting it cut the circle at Q.

Through P and Q respectively draw SL and QT perpendicular to AB.

Let $BL = a$, $SL = y$, $AL = x$.

Then $\frac{a}{PL} = \frac{BO}{OM} = \frac{r}{\frac{1}{2}r} = 2$; whence $\frac{1}{2}a = PL$.

[1] For figure and description, based on a statement of Eutocius (*c.* 560), see Tropfke, *Geschichte*, II (1), 42; Allman, *Greek Geom.*, p. 173; Archimedes, *Opera*, ed. Heiberg, III, 66; Heath, *History*, I, 255.

[2] Ἰωάννης ὁ Φιλόπονος (*Ioan'nes ho Philop'onos*), also known as ὁ Γραμματικός (*ho Grammatikos'*), an Alexandrian scholar of the 7th century; not a very reliable source.

[3] The proof appears in Heath, *Apollonius*, p. cxxv.

[4] For a somewhat different proof see Heath, *History*, I, 264.

Because $\triangle ALP$ is similar to $\triangle ATQ$, which is congruent to $\triangle BLS$, which is similar to $\triangle ALS$, we have

$$\frac{a}{y} = \frac{y}{x} = \frac{x}{PL} = \frac{x}{\frac{1}{2}a}.$$

We therefore have two mean proportionals between a and $\frac{1}{2}a$.

Hence $$ax = y^2,\ a^2x^2 = y^4,$$

and $$x^2 = \frac{y^4}{a^2}.$$

Also, $$\tfrac{1}{2}ay = x^2 = \frac{y^4}{a^2};$$

whence $$\tfrac{1}{2}a^3 = y^3$$

and $$a^3 = 2y^3.$$

Therefore in the above figure we simply have to make $PL = \frac{1}{2}a$, which we get by having made $OM = \frac{1}{2}r$. Then SL is the side of a cube equal to $\frac{1}{2}a^3$, or $a^3 = 2y^3$. Hence

$$\frac{SL}{\text{side of given cube}} = \frac{a}{\text{side of required cube}}.$$

Later Methods. Several modern writers have suggested methods for duplicating the cube. Among these are Vieta,[1] Descartes,[2] Fermat,[3] de Sluze,[4] and Newton.[5] Descartes considered not only the question of finding two mean proportionals, as required in solving the problem, but also that of finding four; and Fermat[6] went so far as to consider certain cases involving n mean proportionals, a line of work which was later followed by Clairaut.

Viviani[7] solved the problem by the aid of a hyperbola of the second order ($xy^2 = a^3$). Huygens (1654) gave three methods

[1] *Opera Mathematica*, ed. Van Schooten, p. 242. Leyden, 1646.

[2] Suggested in *La Géométrie*, Book III.

[3] In his memoir *Ad locos planos et solidos Isagoge*, written before Descartes published his work, but not made known until after Fermat's death.

[4] In his *Mesolabium*, 1668.

[5] *Arithmetica Universalis*, 1707, p. 309.

[6] *Œuvres*, I, 118.

[7] *Quinto libro di Euclide o Scienze universale delle proposizioni spiegate colla dottrina del Galileo* (Florence, 1647).


of solving. Newton (1707) suggested several methods but preferred one which made use of the limaçon of Pascal. One of the comparatively recent methods is that employed by Montucci,[1] who made use of the curve defined by the equation

$$y = \sqrt{ax} + \sqrt{ax - x^2}.$$

8. Analytic Geometry

Three Principal Steps. There are three principal steps in the development of analytic geometry: (1) the invention of a system of coordinates; (2) the recognition of a one-to-one correspondence between algebra and geometry; and (3) the graphic representation of the expression $y = f(x)$. Of these, the first is ancient, the second is medieval, and the third is modern.

Ancient Idea of Coordinates. The idea of coördinates in the laying out of towns and lands seems unquestionably to have occurred to the Egyptian surveyors. It is to them that Heron was apparently indebted for his fundamental principles, and from them the Roman surveyors acquired their first knowledge of the science. Indeed, the districts (*hesp*) into which Egypt was divided[2] were designated in hieroglyphics by a symbol derived from a grid, as here shown, quite as we designate a survey today.[3]



Latitude and Longitude. The first definite literary references to the subject appear, however, in the works of the early Greek geographers and astronomers. Hipparchus (*c.* 140 B.C.) located points in the heavens and on the earth's surface by means of their longitude[4] and latitude,[5] the former being reck-

[1] *Résolution de l'équation du cinquième degré*, Paris, 1869.

[2] Known to the Greeks as νομοί (*nomoi'*, nomes) and to Pliny as *praefecturae oppidorum*.

[3] E. W. Budge, *The Mummy*, p. 8 (Cambridge, 1893). See also Cantor, *Geschichte*, I (2), 67.

[4] Μῆκος (*me'kos*, length; Latin, *longitudo*), *i.e.*, distance from east to west; so called because the length of the known world was along the Mediterranean Sea.

[5] Πλάτος (*pla'tos*, width; Latin, *latitudo*), *i.e.*, distance from north to south; so called because the width of the known world was north and south.

oned from the meridian of Rhodes, where Hipparchus took his observations. He also located the stars by means of coordinates.

In the second century Marinus of Tyre (*c.* 150) took his prime meridian through the *Fortunatae Insulae*,[1] and perhaps through the most western point, this being the end of the earth as then known, and Ptolemy (*c.* 150) used the same line.[2]

Ancient Surveyors. The ancient surveyors located points in much the same way as the geographers. Heron (*c.* 50?), apparently following the Egyptian surveyors, laid out a field with respect to one axis quite as we do at present, although, strictly speaking, two coordinates are used.[3]

The Romans brought the science of surveying to the highest point attained in ancient times.[4] They laid out their towns with respect to two axes, the *decimanus*, which was usually from east to west, and the *cardo*, an axis perpendicular to the *decimanus*. They then arranged the streets on a rectangular coordinate system, much as in most American cities laid out in the 19th century.[5]

Rectangular Axes in Greek Geometry. In their treatment of geometric figures the Greeks made use of what were substantially two rectangular axes. Menæchmus (*c.* 350 B.C.), for example, may have used that property of the parabola expressed by the equation $y^2 = px$, and also that property of the rectangular hyperbola expressed by the equation $xy = c^2$. Archimedes (*c.* 225 B.C.), who no doubt was indebted to the lost work of Euclid on conics in general, used the same relation for the parabola, his results being expressed as usual in the form of a proportion.[6]

[1] Probably, as stated in Volume I, including the Canary, Madeira, and Azores groups.

[2] Halma's edition of Ptolemy, VI, 17 (Paris, 1828).

[3] See his *Opera quae supersunt omnia*, V, 5 (Leipzig, 1899–1914), on his stereometry and mensuration.

[4] M. Cantor, *Die römischen Agrimensoren*, Leipzig, 1875.

[5] As Frontinus states it in his Liber I, "ager . . . decimanis et cardinibus continetur"; "Ager per strigas [rows] et per scamna [steps] diuisus." He also used oblique coordinates. See the Lachmann and Rudorff edition for diagrams.

[6] On all this discussion see Heath, *Apollonius*, p. cxv seq.; *History*, II, 122.

Apollonius carried the method much farther, as may be seen by the following statement:

If straight lines are drawn from a point so as to meet at given angles two straight lines given in position, and if the former lines are in a given ratio, or if the sum of one of them and of such a line as bears a given ratio to the second is given, then the point will be on a given straight line.[1]

This is only a nonsymbolic method of stating that the equation $x = ay - b$ represents a straight line, a and b being positive.

Sir Thomas Heath calls attention to another essential difference between the Apollonian and Cartesian points of view:

The essential difference between the Greek and the modern method is that the Greeks did not direct their efforts to making the fixed lines of the figure as few as possible, but rather to expressing their equations between areas in as short and simple a form as possible. Accordingly they did not hesitate to use a number of auxiliary fixed lines, provided only that by that means the areas corresponding to the various terms in x^2, xy, . . . forming the Cartesian equation could be brought together and combined into a smaller number of terms. . . . In the case, then, where two auxiliary lines are used in addition to the original axes of coordinates, and it appears that the properties of the conic (in the form of equations between areas) can be equally well expressed relatively to the two auxiliary lines and to the two original axes of reference, we have clearly what amounts to a transformation of coordinates.[2]

Ordinate and Abscissa. As to technical terms, the Greeks used an equivalent of "ordinate."[3] For "abscissa"[4] they used such expressions as "the [portion] cut off by it from the diameter towards the vertex." Apollonius uses the word "asymptote,"[5] but the word had a broader meaning than with us, referring to any lines which do not meet, in whatever direction they are

[1] On the work of Apollonius see Volume I, page 116; Heath, *Apollonius*.

[2] Heath, *Apollonius*, p. cxviii. See also his *History*, II, 126–196.

[3] That is, τεταγμένως (*tetagmen'os*, ordinate-wise). The same term is used for the tangent at the extremity of a diameter. *Ibid.*, p. clxii.

[4] Latin *ab* + *scissa*, from *ab* (off) + *scindere* (to cut).

[5] Ἀσύμπτωτος (*asym'ptotos*, from ἀ privative + σύν, together, + πτωτός, falling).

produced. The names "ellipse," "parabola," and "hyperbola" are probably due to Apollonius,[1] although two of them are found in late manuscripts of the works of Archimedes.

Oresme's Contribution. In the Middle Ages Nicole Oresme (c. 1360) wrote two works[2] in which he took a decided step in advance. He considered a series of points which have uniformly changing *longitudines* and *latitudines*, the first being our abscissas and the second our ordinates.[3] The series of points determined by the ends of the *latitudines* was called a *forma*, and the difference between two successive *latitudines* was called a *gradus*. If the *latitudines* are constant, the series of points was described as *uniformis eiusdem gradus*; but if the *latitudines* varied, the *forma* of the series of points was *difformis per oppositum*. The difference between two successive *latitudines* was the *excessus graduum*, and this might or might not be constant. In the former case the *forma* was *uniformiter difformis*; in the latter case, *difformiter difformis*. The *formae* considered were series of points arranged in rectilinear, circular, or parabolic order. Of course only positive *latitudines* were considered. Here, then, we find the first decided step in the development of a coordinate system, apart from the locating of points on a map of some kind; but we also find a lack of any idea of continuity in the point systems. The method was the subject of university lectures at Cologne as early as 1398,[4] as witness the statutes of that period.[5] Kepler and Galileo recognized its value, and the former was influenced to

[1] Heath, *Apollonius*, p. clxiii; *History*, II, 138.

[2] *Tractatus de latitudinibus formarum* and *Tractatus de uniformitate et difformitate intensionum*. See the *Zeitschrift* (Hl. Abt.), XIII, 92. The first of these tractates was printed at Padua in 1482 and again in 1486, in Venice in 1505, and in Vienna in 1515.

[3] Tropfke, *Geschichte*, II (1), 409; H. Wieleitner, *Bibl. Math.*, XIV (3), 210.

[4] Hankel, *Geschichte*, p. 351.


[5] "Item statuimus quod Bacalarius temptandus debet audivisse libros infrascriptos . . . aliquem tractatum de latitudinibus formarum." See F. J. von Bianco, *Die alte Universität Köln*, I, Anlagen, p. 68 (Cologne, 1885); S. Günther, "Die Anfänge und Entwickelungsstadien des Coordinatenprincipes," *Abhandlungen d. naturf. Gesellsch. zu Nürnberg*, VI (1877); reprint, p. 16; hereafter referred to as Günther, *Die Anfänge*.

make much use of it in his astronomical work.[1] Indeed, the use of a kind of coordinate paper for the graphic representation of the course of the planets is found much earlier even than Oresme, for Günther[2] has called attention to a manuscript of the 10th century[3] in which the graphs closely resemble similar forms of the present day.

Relation of Algebra to Geometry. The second step in the development of analytic geometry has to do with the relation of algebra to geometry. If we consider such a proposition of Euclid (*c.* 300 B.C.) as the one relating to the square on the sum of two lines,[4] we see that it is the analogue of the algebraic identity $(a+b)^2 = a^2 + 2ab + b^2$. Euclid, however, had no algebraic symbolism, and while of course he recognized the analogy to the square of the sum of two numbers, it cannot be said that he related algebra to geometry in the way that we do with our modern symbols.

When Archimedes (*c.* 225 B.C.), Heron (*c.* 50?), and Theon of Alexandria (*c.* 390) found square roots, they used this proposition of Euclid; but, again, they can hardly be said to have grasped the relation that is so familiar to us today.

It is among the Arab and Persian writers that we first find geometric figures used in works devoted solely to algebra. Thus al-Khowârizmî (*c.* 825) considered numerous cases such as the following: "A square and ten Roots are equal to thirty-nine Dirhems."[5] Here he uses the annexed figure, the square AB having for its side one of the roots of the given equation



$$x^2 + 10x = 39,$$

a favorite equation with subsequent writers.

Omar Khayyam (*c.* 1100) made continued use of geometric figures in his work on algebra,[6] thus recognizing the one-to-one

[1] *Opera omnia*, Frisch ed., IV, 610 seq. (Frankfort a. M., 1863).
[2] *Die Anfänge*, p. 19 and Fig. 2 in the plates.
[3] Munich Cod. Lat. 14,436: "Macrobius Boetius in Isagog. Saec. X."
[4] *Elements*, II, 4. [5] Rosen's translation, p. 13.
[6] *L'Algèbre d'Omar Alkhayyâmî*, translated by F. Woepcke, Paris, 1851. So Woepcke remarks: "Il est une particularité de cette algèbre qui mérite d'être

correspondence between algebra and geometry even more than al-Khowârizmî had done before him.[1]

The Hindu algebraists also used geometric figures in their work. For example, Bhāskara (*c.* 1150)[2] has such problems as this: "Tell two numbers, such, that the sum of them, multiplied by four and three, may, added to two, be equal to the product." In such cases he gives two solutions, one algebraic and the other geometric.

Europeans relate Algebra to Geometry. Among the Europeans, Fibonacci was the first mathematician of prominence who recognized the value of relating algebra to geometry. In his *Practica geometriae* (1220) he uses algebra in solving geometric problems relating to the area of a triangle.[3]

In the early printed books there was more or less use of geometric figures in connection with algebraic work. Thus Vieta tells us that Regiomontanus solved algebraically problems which he could not solve by geometry. Pacioli (1494) continually uses geometric figures in his solution of quadratics,[4] and Cardan (1545) does the same.[5] After the publication of Cardan's work the recognition of the relationship became common. Vieta, for example, generalized the idea of the ancients as to representing points on a line, although adhering to the use of proportion in most of his geometric work instead of using the equation form.

The first textbook on algebraic geometry was that of Marino Ghetaldi (1630),[6] who may have been influenced by Vieta.[7] In his solution of geometric problems he freely brought algebra

remarquée et discutée d'abord. C'est que l'auteur se fait une loi, pour toutes les équations dont il s'occupe, de joindre la résolution numérique ou arithmétique à la construction géométrique" (*Préface*, p. vij).

[1] Rosen ed., p. 13 and elsewhere. [2] Colebrooke translation, p. 270.

[3] "Quare quadratum lateris .cg. erit $\frac{1}{3}$ rei; et multiplicabo .cf. in dimidium .cg., hoc est radicem rei in radicem $\frac{1}{2}$ rei, ueniet radix $\frac{1}{2}$ census. . . ." *Scritti*, II, 223.

[4] *E.g.*, *Sūma*, 1494 ed., fol. 146, *v.*, *et passim*.

[5] For the use of geometric figures in his first solution of a cubic, see *Ars Magna*, fol. 29, *v.*

[6] See E. Gelcich, "Eine Studie über die Entdeckung der analytischen Geometrie mit Berücksichtigung eines Werkes des Marino Ghetaldi. . . ." *Abhandlungen*, IV, 191. [7] Tropfke, *Geschichte*, II (1), 414.

to his aid, but it cannot be said that he in any way anticipated the work of the makers of analytic geometry.

Invention of Analytic Geometry.[1] The invention of analytic geometry is commonly attributed to Descartes, he having published (1637) the first treatise on the subject. There seems to be no doubt, however, that the idea occurred to Fermat at about the same time as to Descartes, and to have occurred to Harriot even earlier (*c.* 1600). In the British Museum there are eight volumes of Harriot's manuscripts, and among these may be found "a well-formed analytical geometry, with rectangular coordinates and a recognition of the equivalence of equations and curves."[2]

Fermat on Analytic Geometry. In a letter to Roberval, written September 22, 1636, and hence in the year before Descartes published *La Geometrie*, Fermat shows that he had the idea of analytic geometry some seven years earlier;[3] that is, in 1629. The details of this work appear in his *Isagoge ad locos planos et solidos*, which was published posthumously.[4] He used rectangular axes and followed Vieta in representing the unknowns by vowels (in this case only A and E) and the knowns by consonants. A general point on the curve was represented by I, and the foot of the ordinate from I to the axis of abscissas was represented by Z. The equation of a straight line through the origin was indicated by

$$D \text{ in } A \text{ aequetur } B \text{ in } E,^{5}$$

[1] G. Loria, *Passato-Presente Geom.*; M. Chasles, *Aperçu historique sur l'origine et le développement des méthodes en géométrie*, 3d ed., Paris, 1889 (hereafter referred to as Chasles, *Aperçu*); M. Chasles, *Rapport sur les Progrès de la Géométrie*, Paris, 1870; A. Cayley and E. B. Elliott, "Curve," *Encyc. Britannica*, 11th ed.; Günther, *Die Anfänge*; E. Picard, *Bulletin of the Amer. Math. Soc.*, XI, 404; H. Wieleitner, *Zeitschrift für math. und naturw. Unterr.*, XLVII, 414; W. Dieck, *Mathematisches Lesebuch*, 4. Band, Sterkrade, 1920.

[2] F. V. Morley, "Thomas Hariot," *The Scientific Monthly*, XIV, 63. These manuscripts should be carefully studied. The spelling "Hariot" was used by some of his contemporaries.

[3] ". . . il y a environ sept ans étant à Bourdeaux."

[4] In his *Varia Opera*, p. 2 (Toulouse, 1679); Tropfke, *Geschichte*, II (1), 418; Günther, *Die Anfänge*, p. 43; *Œuvres de Fermat*, ed. P. Tannery and Ch. Henry, Vol. I (Paris, 1891), Vol. II (Paris, 1894).

[5] *I.e.*, $D \cdot A = B \cdot E$, which we should write as $ax = by$.

and that of a general straight line was given by the proportion

ut B ad D, ita $R - A$ ad E.[1]

The equation of a circle appears as

$Bq. - Aq.$ aequetur $Eq.$[2]

If the ratio of $Bq. - Aq.$ to $Eq.$ is constant, Fermat asserted that the resulting figure is an ellipse;[3] and if the ratio of $Bq. + Aq.$ to $Eq.$ is constant, the figure is a hyperbola.[4] He also knew that $xy = a^2$ is the asymptotic equation of a hyperbola[5] and that $x^2 = ay$ is the equation of a parabola.[6]

Descartes publishes La Géométrie. Descartes published his Geometry in 1637, although he had been working upon it for some years,—even as early as 1619.[7] The treatise formed an appendix to his *Discours de la Methode* and was divided into three books. The first book treats of the meaning of the product of lines.[8] The second book defines two classes of curves, the geometric and the mechanic. We might now define the former as curves in which dy/dx is an algebraic function, and the latter as curves in which it is a transcendental function. In this book there is also much attention given to tangents and normals to a curve. The third book is largely algebraic, being entitled, "On the construction of solid or hypersolid problems." It treats particularly of such topics as the number of roots of an equation, "false roots," the increasing or decreasing of the roots, and the transformation of equations.

[1] *I.e.*, $a : b = c - x : y$, or $ay = b\ (c - x)$.

[2] *I.e.*, $B^2 - A^2 = E^2$, or $r^2 - x^2 = y^2$.

[3] "$Bq. - Aq.$ ad $Eq.$ habeat rationem datam, punctum I erit ad ellipsin." *I.e.*, $r^2 - x^2 = ky^2$ is the equation of an ellipse.

[4] "Si $Bq. + Aq.$ est ad $Eq.$ in data ratione, punctum I est ad hyperbolen."

[5] "A in E aeq. Z pl., quo casu punctum I est ad hyperbolen."

[6] "Si $Aq.$ aequatur D in E, punctum I est ad parabolen."

[7] J. Millet, *Descartes. Sa vie, ses travaux, ses découvertes, avant 1637*, p. 100 (Paris, 1867); E. S. Haldane, *Descartes*, p. 59 (London, 1905); C. Rabuel, *Commentaires sur la Geometrie de M. Descartes*, Lyons, 1730; *La Geometrie*, various editions from 1637. See the author's facsimile edition with translation (Chicago, 1925).

[8] "Des Problêmes qu'on peut construire n'y employant que des cercles & des lignes droites."

Instead of using the name "coordinates," Descartes spoke of roots or unknowns. The name "coordinate" is due to Leibniz,[1] as are also the terms "abscissa" and "ordinate," although, as we have seen, the Greeks used terms that were similar to them. Newton, Euler, Cramer, and various other writers used "applicate" to represent an ordinate.

Descartes had an idea of oblique coordinates, but he used only the x-axis and positive perpendicular ordinates in common practice.

Later Writers. In 1658 Jan (Johan) de Witt[2] wrote a work on curve lines[3] in which he set forth a number of typical equations and gave the geometric character of each.

Further work was done by Lahire,[4] after which the elements of plane analytic geometry may be considered as having become established. The most noteworthy single contributor to the elements of the subject thereafter was Newton.[5] In his work on cubic curves he showed that a cubic has at least one real point at infinity, that any cubic belongs to one of four characteristic types, and that there are seventy-two possible forms of a cubic, a number since increased by six.[6] The discussion of the subject was nearly exhaustive, and was the most elaborate one of the kind that had been made up to that time.

The idea of polar coordinates seems due to Gregorio Fontana (1735–1803), and the name was used by various Italian writers of the 18th century.[7]

[1] *Acta Eruditorum* (1692), p. 170.

[2] Born at Dordrecht, September 12/24, 1625; died at The Hague, August 20, 1672. For biography see *The Insurance Cyclopœdia*, Vol. II (London, 1873).

[3] *Elementa Curvarum Linearum*, Leyden, 1659; Amsterdam, 1683.

[4] *Les Lieux Géométriques*, Paris, 1679; *Construction des Equations Analytiques*, Paris, 1679.

[5] In his *Principia* (London, 1687) and his *Arithmetica Universalis* (Cambridge, 1707), but chiefly in his *Enumeratio linearum tertii ordinis*, which probably dates from 1668 or 1669 and which was published as an appendix to his *Optics* in 1704. See also W. W. R. Ball, "On Newton's classification of cubic curves," *Transactions of the London Mathematical Society* (1891), p. 104.

[6] G. Loria, *Ebene Kurven, Theorie und Geschichte*, p. 20 (Leipzig, 1902; 2d ed., 1910–1911), hereafter referred to as Loria, *Kurven*.

[7] For a discussion of the later types of coordinates see the *Encyklopädie*, III, 596, 656; Cantor, *Geschichte*, IV, 513.

Solid Analytic Geometry. Descartes clearly mentioned solid analytic geometry, but he did not elaborate it. Frans van Schooten the younger suggested the use of coordinates in three-dimensional space (1657), and Lahire (1679) also had it in mind. Jean Bernoulli (1698) thought of equations of surfaces in terms of three coordinates, but published nothing upon the theory at that time.

The first work on analytic geometry of three dimensions was written by Antoine Parent and was presented to the French Académie in 1700.[1] A. C. Clairaut (1729) was the first to write on curves of double curvature.[2] The third great contributor to the theory was Euler (1748), with whose work the subject advanced beyond the elementary stage.

Euler also laid the foundations for the analytic theory of curvature of surfaces, attempting to do for the classification of surfaces of the second degree what the ancients had done for curves of the second order. Monge introduced the notion of families of surfaces and discovered the relation between the theory of surfaces and the integration of partial differential equations, enabling each to be advantageously viewed from the standpoint of the other.

Modern Theory. Möbius began his contributions to geometry in 1823, and four years later published his *Barycentrische Calcul.* In this great work he introduced homogeneous coordinates. Of modern contributors to analytic geometry, however, Plücker stands easily foremost. In 1828 he published the first volume of his *Analytisch-geometrische Entwickelungen,* in which there appeared the modern abridged notation. In the second volume (1831) he set forth the present analytic form of the principle of duality. To him is due (1833) the general treatment of foci for curves of higher degree, and the complete classification of plane cubic curves (1835) which had been so frequently attempted before him. He also gave (1839) an

[1] "Des effections des superficies." This appears in his *Essais et Recherches,* Paris, 1705 and 1713.

[2] *Recherches sur les courbes à double courbure,* printed in 1731. It was presented to the Académie when Clairaut was only sixteen years old.

enumeration of plane curves of the fourth order. In 1842 he gave his celebrated "six equations," by which he showed that the characteristics of a curve (order, class, number of double points, number of cusps, number of double tangents, and number of inflections) are known when any three are given. To him is also due the first scientific dual definition of a curve, a system of tangential coordinates, and an investigation of the question of double tangents. The theory of ruled surfaces, begun by Monge, was also extended by him. Possibly the greatest service rendered by Plücker was the introduction of the straight line as a space element, his first contribution (1865) being followed by his well-known treatise on the subject (1868–1869).[1]

Certain Well-Known Curves. There are certain curves that are so frequently met in textbooks on analytic geometry as to deserve mention in an elementary history. Several of these have been considered elsewhere in this work, and a few others, with additional notes on those already given, will now be mentioned,[2] and for convenience will be given alphabetically.

Brachistochrone,[3] the curve of quickest descent, was studied by Galileo, Leibniz, Newton, and the Bernoullis, and was shown to be the cycloid. The name is due to the Bernoullis.[4]

Cardioid, the epicycloid $(x^2 + y^2 - 2ax)^2 = 4a^2(x^2 + y^2)$. The name is due to Giovanni Francesco M. M. Salvemini, called from his birthplace de Castillon (1708–1791), *De curva cardioide* (1741). It had already been studied by Ozanam.[5]

[1] On the various coordinate systems, see the *Encyklopädie*, III, 221, 596.

[2] H. Brocard, *Notes de Bibliographie des Courbes*, lith. autog., Bar-le-Duc, 1897; Partie complémentaire, 1899. See also Chasles, *Aperçu*, and the *Encyklopädie*, III, 185, 457; E. Pascal, *Repertorium der höheren Mathematik*, German translation by A. Schepp, Leipzig, 1902, especially Vol. II, chap. xvii; Loria, *Kurven*; Joaquin de Vargas y Aguirre, "Catálogo General de curvas," *Memorias de la Real Acad. de Ciencias exactas*, XXVI, Madrid, 1908; F. G. Teixeira, *Traité des courbes spéciales remarquables planes et gauches*, 3 vols., Coimbra, 1908, 1909, 1915.

[3] From βράχιστος (*brach'istos*, shortest) and χρόνος (*chron'os*, time). Formerly spelled *brachystochrone* by a confusion of the superlative βράχιστος with its positive βραχύς.

[4] Cantor, *Geschichte*, III, chap. 92.

[5] It is a special case of the limaçon.

Catenary, the French *chaînette*, $y = \frac{1}{2}a\left(e^{\frac{x}{a}} + e^{-\frac{x}{a}}\right)$. The name of the curve (*catenaria*) and the discovery of the equation and its properties are due to Leibniz.[1]

Cissoid of Diocles, the "ivy-shaped" curve, $y^2 = x^3/(2a - x)$, due to Diocles (*c.* 180 B.C.).[2]

Cochlioid,[3] $r = (a \sin\theta)/\theta$, a spiral curve discussed by J. Perk, *Phil. Trans.*, 1700, this form being a late one, due to J. Neuberg, a Belgian geometer. The name originated (1884) with two recent writers, Bentham and Falkenburg.

Cocked hat, the French *bicorne*, $x^2 + \left(\frac{a^2 - x^2}{y} - 2a\right)^2 - a^2 = 0$, a modern name, due to Sylvester.[4]

Conchoid of Nicomedes (*c.* 180 B.C.), the "shell-shaped" curve.[5] The Cartesian equation is $(x-a)^2(x^2+y^2) - b^2x^2 = 0$ and the polar equation is $r = a/\cos\theta + b$.

Conchoid of de Sluze, the cubic curve $a\ (r\cos\theta - a) = k^2\cos^2\theta$ or $a\ (x-a)\ (x^2+y^2) - k^2x^2 = 0$, first constructed by René de Sluze (1662).

Curve of Pursuit, French *courbe du chien* as a special case. The name *ligne de poursuite* seems due to Pierre Bouguer[6] (1732), although the curve had been noticed by Leonardo da Vinci.

Cycloid, the transcendental curve

$$x = a \arccos\frac{a-y}{a} - \sqrt{2ay - y^2}.$$

This curve, sometimes incorrectly attributed to Nicholas Cusa (*c.* 1450), was first studied by Charles de Bouelles (1501). It then attracted the attention of Galileo (1599), Mersenne

[1] On the history of this curve see C. A. Laisant, *Association Française pour l'avancement des Sciences, Congrès de Toulouse*, p. 64 (1887); Loria, *Kurven*, II, 204.

[2] See Volume I, page 118; Volume II, page 314.

[3] From κοχλίας (*cochlí'as*, snail) and εἶδος (*eí'dos*, form).

[4] See also *Educational Times*, quest. 12,978 (Matz).

[5] See Volume I, page 118.

[6] Born at Croisic, Brittany, February 16, 1698; died in Paris, August 15, 1758. He was one of the French geodesists sent to Peru in 1735 to measure an arc of a meridian. See articles by F. V. Morley, R. C. Archibald, H. P. Manning, and W. W. Rouse Ball, *Amer. Math. Month.*, XXVIII.

(1628), and Roberval (1634). Pascal (1659) called it the "roulette," completely solved the problem of its quadrature, and found the center of gravity of a segment cut off by a line parallel to the base.[1] Jean and Jacques Bernoulli showed that it is the brachistochrone curve, and Huygens (1673) showed how its property of tautochronism might be applied to the pendulum.

Devil's Curve, French *courbe du diable*, in general represented by the equation $y^4 - x^4 + ay^2 + bx^2 = 0$, and in particular by $y^4 - x^4 - 96\,a^2y^2 + 100\,a^2x^2 = 0$. The polar equation is

$$r = 2a\sqrt{(25 - 24\tan^2\theta)/(1 - \tan^2\theta)}.$$

It was studied by G. Cramer (1750)[2] and Lacroix (1810)[3] and is given in the *Nouvelles Annales* (1858), p. 317.

Elastic Curve, French *courbe élastique*, the differential equation of which is

$$dy = x^2dx/\sqrt{a^4 - x^4}.$$

It was first studied by Jacques Bernoulli (1703).

Epicycloid, literally "epicycle-shaped," a curve traced by a point on a circle which rolls on the convex side of a given circle. The equation is $(x^2 + y^2 - a^2)^2 = 4a\,[(x - a)^2 + y^2]$. The curve was recognized by Hipparchus (*c.* 140 B.C.) in his astronomical theory of epicycles. Albrecht Dürer (1525) was the first to describe it in a printed work. It was next studied by Desargues (1639), but it first received noteworthy consideration by Lahire (1694) and Euler (1781).

Folium of Descartes, a curve represented by the equation $x^3 + y^3 = 3axy$. The problem was proposed to Roberval to determine the tangent to this curve, and through an error he was led to believe that the curve had the form of a jasmine flower, and hence he gave it the name *fleur de jasmin*, which was afterwards changed. It is also known as the *nœud de ruban*.

[1] H. Bosmans, "Pascal et les premières pages de l'"Histoire de la Roulette,'" *Archives de Philosophie*, I (1923), cah. 3.

[2] *Introduction a l'analyse des lignes courbes algébriques*, p. 19 (Geneva, 1750).

[3] *Traité du calcul différentiel et . . . intégral*, I, 391 (Paris, 1797; 1810 ed.).

Helix, the name given by Archimedes (*c.* 225 B.C.) to a spiral already studied by his friend Conon.[1] It is now known as the spiral of Archimedes. The equation is $r = a\theta$, or $\tan \phi = \theta$. It is one of the class of which the general equation is $r = a\theta^n$. The name is now usually applied to a curve traced upon a cylinder and cutting the generatrices under a constant angle. There are also the conical helix, the spherical helix (or loxodrome), and other types.

Lemniscate,[2] a curve first mentioned by Jacques Bernoulli (1694).[3] Its principal properties were discovered by Fagnano (1750). The analytic theory of the curve is due to Euler (1751, 1752). The general lemniscate has for its equation $(x^2 + y^2)^2 = 2a^2 (x^2 - y^2) + b^4 - a^4$, while that of Bernoulli is represented algebraically by $(x^2 + y^2)^2 = 2a^2(x^2 - y^2)$, and is called the hyperbolic lemniscate. The general lemniscate is also known as Cassini's oval, after Giovanni Domenico (Jean Dominique) Cassini, who described it in 1680.

Limaçon, French *limaçon* (a snail), Italian *lumaca*, from Latin *limax*, called also by the French the *conchoïde du cercle*. The curve is

$$(x^2+y^2-ax)^2 = b^2(x^2+y^2), \quad \text{or} \quad \rho = a + b\cos\phi.$$

Roberval called it the limaçon of Pascal, Étienne Pascal (father of Blaise) having discovered it. German writers speak of it as the *Pascal'sche Schnecke*.

Lituus (the Latin word for an augur's staff), the curve $r^2\theta = a^2$. The name is due to Cotes (*c.* 1710).

Logarithmic or Equiangular Spiral, the curve $r = ae^{k\theta}$, or $k\theta = \log\frac{r}{a}$, studied by Jacques Bernoulli (1692), who spoke of it as a *spira mirabilis*. It is still to be seen, in rude form, upon his tomb in Basel. The logarithmic spiral was the first non-algebraic plane curve to be rectified.

[1] But see Volume I, page 107.

[2] From λημνίσκος (*lemnis'kos*, Latin *lemniscus*), a ribbon on which a pendant is hung.

[3] ". . . formam refert jacentis notae octonarii ∞, seu complicitae in nodum fasciae, sive lemnisci." See F. Cajori, *Hist. of Math.*, 2d ed., p. 221.

Pearls, a name given by Pascal and de Sluze to the curves whose equation is $a^{p+q-r}y^r = x^p(a-x)^q$,

or, in particular, $a^2y = x^2(a-x)$.

De Sluze proposed their consideration to Huygens (1658), and the latter made a careful study of them.

Roseate Curve, *Rosace*, or *Rhodonea*, the curve whose general polar equation is $r = a\cos m\theta$. The name *Rhodonea* is due to Guido Grandi (1713). The *Rosace à quatre feuilles*, or *Quadrifolium*, has for its polar equation $r = a \sin 2\theta$, and for the Cartesian form $(x^2+y^2)^3 = 4a^2x^2y^2$.

Semicubic Parabola, or *Neile's Parabola*, $y^3 = ax^2$. It was the second curve to be rectified. William Neile discovered the curve in 1657. The method of rectification was published by Wallis in 1659, credit being given by him to his pupil, Neile, although there is still some dispute as to whether it was due to him, to Fermat, or to the Dutch writer Van Heuraet.[1]

Serpentine Curve, a name proposed by Newton for the curve $y = abx/(a^2+x^2)$.

Spiral of Archimedes, the curve $r = a\theta$, mentioned under *Helix*.

Spiral of Fermat, the curve $r^2 = \theta$, proposed by Fermat in a letter to Mersenne, June 3, 1636.

Strophoid, French *strophoïde*,[2] a name proposed by a modern writer, Montucci (1846), for the curve $y = x\sqrt{(a-x)/(a+x)}$. Lehmus had already proposed (1842) the name *kukumaeïde* (cucumber seed), and various other names have been used. The curve has been studied by Barrow, Jean Bernoulli, Agnesi (1748), James Booth (1858), and various others.

Tractrix, the *tractoria* of Huygens (1693). The differential equation is

$$\frac{dy}{dx} = \frac{y}{\sqrt{a^2-y^2}}.$$

[1] Hendrik van Heuraet, born at Haarlem in 1633. His brief *Epistolae de curvarum linearum in rectas transmutatione* was published by Van Schooten in 1659.

[2] From στρόφος (*stroph'os*, a twisted band, a cord) + εἶδος (*ei'dos*, form).

Witch of Agnesi, Versiera, Cubique d'Agnesi, or *Agnésienne*, the curve $y^2x + r^2(x - r) = 0$, discussed by Maria Gaetana Agnesi in 1748 in her *Istituzioni Analitiche*.[1]

9. Modern Geometry

Four Periods of Geometry. In order to appreciate the historical setting of modern geometry it is well to remember that the history of geometry in general may be roughly divided into four periods: (1) the synthetic geometry of the Greeks, including not merely the geometry of Euclid but the work on conics by Apollonius and the less formal contributions of numerous other writers; (2) the birth of analytic geometry, in which the synthetic geometry of Desargues, Kepler, Roberval, and other writers of the 17th century merged into the coordinate geometry already set forth by Descartes and Fermat; (3) the application of the calculus to geometry,—a period extending from about 1650 to 1800, and including the names of Cavalieri, Newton, Leibniz, the Bernoullis, l'Hospital, Clairaut, Euler, Lagrange, and d'Alembert, each one, especially after Cavalieri, being primarily an analyst rather than a geometer; (4) the renaissance of pure geometry, beginning with the 19th century and characterized by the descriptive geometry of Monge, the projective geometry of Poncelet, the modern synthetic geometry of Steiner and Von Staudt, the modern analytic geometry of Plücker, the non-Euclidean hypotheses of Lobachevsky, Bolyai, and Riemann, and the foundations of geometry.

Descriptive Geometry. Descriptive geometry as a separate science begins with Monge. He had been in possession of the theory for over thirty years before the publication of the *Géométrie Descriptive* (1794),[2]—a delay due to the jealous

[1] G. Loria, *Bibl. Math.*, XI (2), 7. See the English translation by J. Hellins, I, 222 (London, 1801). See also Volume I, page 519.

[2] G. Loria, *Storia della Geometria Descrittiva* (Milan, 1921), the leading authority on the subject; Chr. Wiener, *Lehrbuch der darstellenden Geometrie*, Leipzig, 1884–1887; *Geschichte der darstellenden Geometrie, ibid.*, 1884. See *Encyklopädie*, III, 517; F. J. Obenrauch, *Geschichte der darstellenden und projectiven Geometrie*, Brünn, 1897.

desire of the military authorities to keep the valuable secret. Certain of its features can be traced back to Frézier, Desargues, Lambert, and other writers of the preceding century, but it was Monge who worked it out in detail as a science, although Lacroix (1795), inspired by Monge's lectures in the École Polytechnique, published the first work on the subject. After Monge's work[1] appeared, Hachette (1812, 1818, 1821) added materially to the theory.

Period of Projective Geometry. It is also in this period that projective geometry has had its development, even if its origin is more remote. The origin of any branch of science can always be traced far back in human history, and this fact is patent in the case of this phase of geometry. The idea of the projection of a line upon a plane is very old. It is involved in the treatment of the intersection of certain surfaces, due to Archytas (*c.* 400 B.C.), and appears in various later works by Greek writers. Similarly, the invariant property of the anharmonic ratio was essentially recognized both by Menelaus (*c.* 100) and by Pappus (*c.* 300). The notion of infinity was also familiar to several Greek geometers and to the Latin writer Lucretius (*c.* 100), so that various concepts that enter into the study of projective geometry were common property long before the science was really founded.

Desargues, Pascal, Newton, and Carnot. One of the first important steps to be taken in modern times, in the development of this form of geometry, was due to Desargues. In a work published in 1639 Desargues set forth the foundation of the theory of four harmonic points, not as done today but based on the fact that the product of the distances of two conjugate points from the center is constant. He also treated of the theory of poles and polars, although not using these terms. In the following year (1640) Pascal, then only a youth of sixteen or seventeen, published a brief essay on conics in which he set forth the well-known theorem that bears his name.

[1] *Essais sur les plans et les surfaces*, Paris, 1795; *Complément des Élémens de Géométrie ou Élémens de Géométrie descriptive*, Paris, 1796; *Essais de Géométrie sur les plans et les surfaces courbes*, Paris, 1812.

In the latter part of the 17th century Newton investigated the subject of curves of the third order and showed that all such curves can be derived by central projection from five fundamental types. In the 18th century relatively little attention was given to the subject, but at the close of this period, as already stated, the descriptive geometry of Monge was brought into prominence,—itself a kind of projective geometry, although not what is technically known by this name.

Inspired by the general activity manifest in the 18th century, and following in the footsteps of Desargues and Pascal, Carnot treated chiefly of the metric relations of figures. In particular he investigated these relations as connected with the theory of transversals,—a theory whose fundamental property of a four-rayed pencil goes back to Menelaus and Pappus, and which, though revived by Desargues, was set forth for the first time in its general form by Carnot in his *Géométrie de Position* (1803), and supplemented in his *Théorie des Transversales* (1806). In these works Carnot introduced negative magnitudes, the general quadrilateral, the general quadrangle, and numerous other similar features of value to elementary geometry.

Poncelet on Projective Geometry. The origin of projective geometry as we know it today is generally ascribed to Poncelet.[1] A prisoner (1813–1814) in the Russian campaign, confined at Saratoff on the Volga, with no books at hand,[2] he was able in spite of all such discouragement to plan the great work[3] which he published in 1822. In this work he made prominent for the first time the power of central projection in demonstration and the power of the principle of continuity in research. His leading idea was the study of projective properties, and as a foundation principle he introduced the anharmonic ratio, a

[1] On the whole question consult the *Encyklopädie*, III, 389.

[2] "Privé de toute espèce de livres et de secours, surtout distrait par les malheurs de ma patrie et les miens propres."

[3] J. V. Poncelet, *Traité des propriétés projectives des figures*, Paris, 1822; *ibid.*, 1865–1866; *Applications d'analyse et de géométrie*, ed. Mannheim and Moutard, 2 vols., Paris, 1862, 1864. On the general subject of the development of modern geometric methods see J. G. Darboux, *Bulletin of the Amer. Math. Soc.*, XI, §17. See also Volume I, page 496.

concept which possibly dates back to the lost porisms of Euclid, and which Desargues (1639) had used. The anharmonic point-and-line properties of conics have since then been further elaborated by Brianchon, Chasles, Steiner, Plücker, Von Staudt, and other investigators. To Poncelet is also due the theory of "figures homologiques," the perspective axis and perspective center (called by Chasles the axis and center of homology), an extension of Carnot's theory of transversals, and the "cordes idéales" of conics which Plücker applied to curves of all orders. Poncelet also considered the circular points at infinity and completed the first great principle of modern geometry, the principle of continuity. Following upon the work of Poncelet, Möbius made much use of the anharmonic ratio in his *Barycentrische Calcul* (1827), but he gave it the name "Doppelschnitt-Verhältniss (ratio bisectionalis)," a term now in common use under Steiner's abbreviated form "Doppelverhältniss." The name "anharmonic ratio" or "anharmonic function" ("rapport anharmonique" or "fonction anharmonique") is due to Chasles, and "cross-ratio" was suggested by Clifford.

Gergonne, Steiner, and Von Staudt. Joseph-Diez Gergonne[1] (1813) introduced the term "polar" in its modern geometric sense, although Servois (1811) had used the expression "pole." Gergonne was the first (1825–1826) to grasp completely the principle to which he gave the name of "Principle of Duality," the most important principle, after that of continuity, in modern geometry. He used the word "class" in describing a curve, explicitly defining class and degree (order) and showing the duality between them. He and Chasles were among the first to study surfaces of higher order by modern methods.

Jacob Steiner, the most noted of the Swiss geometers of the 19th century, gave the first complete discussion[2] of the projective relations between rows, pencils, etc. and laid the foundation for the subsequent development of pure geometry. For the present, at least, he may be said to have closed the theory of conic sections, of the corresponding figures in three-dimensional

[1] See Volume I, page 495.

[2] *Systematische Entwickelungen* . . ., Berlin, 1832. See Volume I, page 524.

space, and of surfaces of the second order, and hence there opens with him the period of the special study of curves and surfaces of higher order.

Between 1847 and 1860 Karl Georg Christian von Staudt set forth a complete system of a pure geometry[1] that is independent of metrical considerations. All projective properties are here established independently of number relations, number being drawn from geometry instead of conversely, and imaginary elements being systematically introduced from the geometric side. A projective geometry, based on the group containing all the real projective and dualistic transformations, is developed, and imaginary transformations are introduced.

Non-Euclidean Geometry. The question of Euclid's fifth postulate, relating to parallel lines, has occupied the attention of geometers ever since the *Elements* was written.[2] The first scientific investigation of this part of the foundation of geometry was made by Girolamo Saccheri[3] (1733), a work which was

[1] *Geometrie der Lage*, Nürnberg, 1847; *Beiträge zur Geometrie der Lage*, 3 parts, Nürnberg, 1856, 1857. See M. Noether, *Zur Erinnerung an K. G. C. von Staudt*, Erlangen, 1901, and Volume I, page 505.

[2] F. Engel and P. Stäckel, *Die Theorie der Parallellinien von Euklid bis auf Gauss*, Leipzig, 1895; G. B. Halsted, various contributions, including "Bibliography of Hyperspace and Non-Euclidean Geometry," *American Journal of Mathematics*, Vols. I, II; *Amer. Math. Month.*, Vol. I; translations of Lobachevsky's Geometry, Vassilief's address on Lobachevsky, Saccheri's Geometry, Bolyai's work and his life; "Non-Euclidean and Hyperspaces," *Mathematical Papers of Chicago Congress*, p. 92; G. Loria, *Die hauptsächlichsten Theorien der Geometrie*, Leipzig, p. 106; A. Karagiannides, *Die Nichteuklidische Geometrie vom Alterthum bis zur Gegenwart*, Berlin, 1893; E. McClintock, "On the Early History of Non-Euclidean Geometry," *Bulletin of New York Mathematical Society*, II, 144; W. B. Frankland, *Theories of Parallelism*, Cambridge, 1910 (particularly valuable); H. Poincaré, "Non-Euclidean Geometry," *Nature*, XLV, 404; P. Stäckel, *Wolfgang und Johann Bolyai, Geometrische Untersuchungen*, 2 vols., Leipzig, 1913. See also Volume I, Chapter X, under the several names mentioned. On the general question of the modern synthetic treatment of elementary geometry, see the *Encyklopädie*, III, 859; for the analytic treatment, *ibid.*, 771. See also C. J. Keyser, *Mathematical Philosophy*, p. 342 (New York, 1922). For an excellent bibliography up to the time it was printed see D. M. Y. Sommerville, *Bibliography of Non-Euclidean Geometry*, London, 1911.

[3] Born at San Remo, September 4 or 5, 1667; died at Milan, October 25, 1733. The work was *Euclides ab omni naevo vindicatus*, Milan, 1733; English translation by G. B. Halsted, Chicago, 1920. Saccheri was a Jesuit and taught mathematics in Turin, Pavia, and Milan.

not looked upon as a precursor of Lobachevsky, however, until Beltrami (1889) called attention to the fact. Johann Heinrich Lambert (1728–1777)[1] was the next to question the validity of Euclid's postulate, in his *Theorie der Parallellinien* (posthumous, Leipzig, 1786), the most important treatise on the subject between the publication of Saccheri's work and the works of Lobachevsky and Bolyai. Legendre (1794) also contributed to the theory, but failed to make any noteworthy advance.

During the closing years of the 18th century Kant's[2] doctrine of absolute space, and his assertion of the necessary postulates of geometry, were the object of much scrutiny and attack. At the same time Gauss was giving attention to the fifth postulate, although at first on the side of proving it. It was at one time surmised that Gauss was the real founder of the non-Euclidean geometry, his influence being exerted on Lobachevsky through his friend Bartels,[3] and on János Bolyai through the father Farkas, who was a fellow student of Gauss, and it will presently be seen that he had some clear ideas of the subject before either Lobachevsky or Bolyai committed their theories to print.

Lobachevsky. Bartels went to Kasan in 1807, and Lobachevsky was his pupil. The latter's lecture notes fail to show that Bartels ever mentioned the subject of the fifth postulate to him, so that his investigations, begun even before 1823, seem to have been made on his own motion, and his results to have been wholly original. Early in 1826 he set forth the principles of his famous doctrine of parallels, based on the assumption that through a given point more than one straight line can be drawn which shall never meet a given straight line coplanar with it. The theory was published in full in 1829–1830, and he contributed to the subject, as well as to other branches of mathematics, until his death.

[1] D. Huber, *Lambert nach seinem Leben und Wirken*, Basel, 1829. See Volume I, page 480. [2] E. Fink, *Kant als Mathematiker*, Leipzig, 1889.

[3] Johann Martin Christian Bartels, born at Braunschweig, August 12, 1769; died at Dorpat, December 19, 1836. He was professor of mathematics at Kasan and later at Dorpat.

The Bolyais and Gauss. János Bolyai received, through his father, Farkas, some of the inspiration to original research which the latter had received from Gauss. When only twenty-one he discovered, at about the same time as Lobachevsky, the principles of non-Euclidean geometry, and he refers to them in a letter of November, 1823. They were committed to writing in 1825 and were published in 1832. Gauss asserts in his correspondence with Schumacher[1] (1831–1832) that he had thought out a theory along the same lines as Lobachevsky and Bolyai, but the publication of their works seems to have put an end to his investigations. His statement on the subject is as follows:

I will add that I have recently received from Hungary a little paper on non-Euclidean geometry in which I rediscover all my own ideas and results worked out with great elegance. . . . The writer is a very young Austrian officer, the son of one of my early friends, with whom I often discussed the subject in 1798, although my ideas were at that time far removed from the development and maturity which they have received from the original reflections of this young man. I consider the young geometer Von Bolyai a genius of the first rank.[2]

This was not, however, the first statement of Gauss upon the subject, for in a letter written on November 8, 1824, he remarked:

The assumption that the sum of the 3 angles is smaller than 180° leads to a new geometry entirely different from ours [the Euclidean] —a geometry which is throughout consistent with itself, and which I have elaborated in a manner entirely satisfactory to myself, so that I can solve every problem in it with the exception of the determining of a constant which is not *a priori* obtainable.[3]

[1] Heinrich Christian Schumacher (1780–1850), the astronomer.

[2] Sedgwick and Tyler, *A Short History of Science*, p. 338 (New York, 1917).

[3] P. Stäckel, *Wolfgang und Johann Bolyai*, I, 95 (Leipzig, 1913). The letter was written to one Taurinus, who, two years later, published a *Geometriae prima elementa* (1826), in which he gives evidence of having thought upon a non-Euclidean trigonometry. See Volume I, page 527.

Riemann's Theory. Of all the contributions which appeared after Bolyai's publication the most noteworthy, from the scientific standpoint, is that of Georg Friedrich Bernhard Riemann. In his Habilitationsschrift (1854) he applied the methods of analytic geometry to the theory and suggested a surface of negative curvature, which Beltrami called "pseudo-spherical," thus leaving Euclid's geometry on a surface of zero curvature midway between his own and Lobachevsky's. He thus set forth three kinds of geometry, Bolyai having noted only two. These Klein (1871) called the elliptic (Riemann's), parabolic (Euclid's), and hyperbolic (Lobachevsky's) geometry.

10. Perspective and Optics

Relation of Perspective to Mathematics. While all painters seek to secure proper perspective in their pictures, the most successful of the painters of the Renaissance made an effort to base their treatment of the subject on mathematical principles. Of late these principles have interested architects more than painters, but in any case the subject is largely a mathematical one.[1]

The Greeks included perspective in their science of optics, and the Arabs in their science of appearances, their title being translated into Medieval Latin as *De aspectibus*.[2] Therefore, while there is a manifest difference between perspective and optics as we consider these terms today, it is necessary to treat of them as closely related.

Ancient Works. While several Greek writers wrote on the subject of perspective, the earliest mathematical work that has come down to us is the *Optics* of Euclid.[3] In this work Euclid

[1] On the history of the subject a beginning can be made with N. G. Poudra, *Histoire de la Perspective,* Paris, 1864, a rather poorly arranged work with no index.

[2] The first translation (1505) of Euclid's *Optics,* however, used the term *perspectiva.*

[3] The latest Latin edition of the *Optics* is that of J. L. Heiberg, in *Euclidis opera omnia,* Vol. VII, Leipzig, 1895. There are various translations of the text from Greek into Latin. The first is that of Zamberto (Venice, 1505), in the collected works of Euclid; the second, that of J. Pena (Péna, de la Pêne), Paris, 1557, or

lays down a series of axioms,[1] quite as he does in his *Elements*, the first being: "Therefore it is assumed that [visual] rays emitted from the eye are carried in a straight line, whatever may be the distance."[2]

On the axioms Euclid bases his propositions, sixty-one in number, proving them geometrically after the plan used by him in the *Elements*.

There is also a work on catoptrics containing thirty-one propositions and attributed to Euclid, but it is doubtful if the text published by Gregory[3] and Heiberg is his.

Some idea of the nature of Euclid's work may be obtained from a single proposition in his *Optics*: "If from the center of a circle a line be drawn at right angles to the plane of the circle, and the eye be placed at any point on this line, the diameters of the circle will all appear equal."[4]

Later Classical Writers. The only Roman writer who paid any attention to the subject is Vitruvius (*c.* 20 B.C.), who, in his work on architecture, has something to say on the plans and elevations of buildings. He seems to have had the idea of two projections, these being on two planes perpendicular to each other and arranged as in descriptive geometry.

Heron of Alexandria (*c.* 50?) is known to have written on dioptrics, but only a fragment of the work exists.[5] His theory of light involved the usual error of most of the Greek scientists, that the rays of light proceed from the eye to the object instead of from the object to the eye.

that of Dasypodius which appeared at Strasburg in the same year. See also G. Ovio, *L'ottica di Euclide*, Milan, 1918; D. Gregory, *Euclidis quae supersunt omnia*, Oxford, 1703, p. [599], with parallel Greek and Latin texts; *La prospettiva di Evclide, . . . tradotta dal R. P. M. Egnatio Danti*, Florence, 1573; *La perspective d'Euclide, traduite en français . . . par R. Fréart de Chanteloup*, Mans, 1663.

[1] In the Gregory edition (1703, p. 604) *θέσεις* (*the'seis*) and *positiones*.

[2] That the eye emitted the visual rays was Plato's idea. Aristotle held a view more in accordance with our own, asking why, if the older idea were correct, we cannot see in the dark.

[3] *Loc. cit.*, p. 643.

[4] Prop. XXXV.

[5] *Opera quae supersunt omnia*, Leipzig, 1899–1914; *Traité de la dioptre*, ed. A. J. H. Vincent, Paris, 1858, in the *Bibliothèque Nationale, Notices et extraits*, XIX, Pt. 2, pp. 157–347.

Ptolemy (*c.* 150) is said to have written upon the subject, but it is not certain that he did so.[1] The work attributed to him contains five books, the first dealing with the properties of light, the second with the nature of vision, the third with reflection, the fourth with concave mirrors and with two or more mirrors, and the fifth with refraction.

The next Greek writer on the subject was Heliodorus of Larissa,[2] whose date is uncertain but who lived after Ptolemy. His work is little more than a commentary on Euclid.[3]

Medieval Writers. One of the greatest of the medieval writers on perspective was the Arab scholar Alhazen (*c.* 1000).[4] His work was the basis of Peckham's *Perspectiva* mentioned below. The following well-known problem relating to optics bears his name: "From two given points within a circle to draw to a point on the circle two lines which shall make equal angles with the tangent at that point."[5]

Of the European writers the first one of importance was Roger Bacon (*c.* 1250). In his *Opus Majus* he devotes Part V (*De scientia perspectiva*) to perspective,[6] dividing it into three parts. Part I explains the general principles of vision, Part II deals with direct vision, and Part III discusses reflection and refraction. In the *Opus Tertium* there is also a brief *tractatus*

[1] There is a MS. in Paris beginning: *Incipit Liber Ptholemaei de Opticis sive Aspectibus translatus ab Ammiraco* [or *Ammirato*] *Eugenio Siculo*, consisting originally of five books. For a discussion, see W. Smith, *Dict. of Greek and Roman Biog.*, III, 573 (London, 1864). See also N. G. Poudra, *Histoire de la Perspective*, p. 28 (Paris, 1864).

[2] Possibly his name was Damianus. At any rate some of the MSS. bear the title Δαμιανοῦ φιλοσόφου τοῦ Ἡλιοδώρου Λαρισσαίου περὶ ὀπτικῶν ὑποθέσεων βιβλία β′.

[3] *La Prospettiva di Eliodoro Larisseo, Tradotto Dal Reverendo Padre M. Egnatio Danti*, Florence, 1573, bound with *La Prospettiva di Evclide*. There are other translations.

[4] Al-Ḥasan . . . ibn al-Ḥaiṭam. See Volume I, page 175. A Latin translation, under the title *Opticae Thesauri Libri VII*, was published at Basel in 1572.

[5] For a discussion of the problem see *American Journal of Mathematics*, IV, 327.

[6] *Rogerii Baconis angli, viri eminentissimi, Perspectiva*, Frankfort, 1614. This is best found, however, in the editions of the *Opus Majus* by S. Jebb (London, 1733; Venice, 1750) and J. H. Bridges (2 vols., Oxford, 1897; suppl. vol., London, 1900). See also E. Wiedemann, "Roger Bacon und seine Verdienste um die Optik," in A. G. Little, *Roger Bacon Essays*, p. 185 (Oxford, 1914).

de perspectiva.[1] Besides this, Bacon wrote two other brief treatises[2] on the subject, and still others are attributed to him without historic sanction.[3]

The work that had the greatest influence upon the subject of perspective in the Middle Ages was the *Perspectiva communis* of John Peckham[4] (*c.* 1280). This work was the recognized standard for three hundred years. It was edited and published by Cardan's father and went through various editions. As already stated, Peckham drew largely upon Alhazen's work. The work is divided into three parts, the second containing fifty-six propositions on reflection, and the third containing twenty-two on refraction.

About the same time as Peckham, the German (or possibly Polish) scholar Witelo (*c.* 1270)[5] was called to Rome and there became conversant with the works of the ancients as well as those of the Arabs in the science of perspective. Georg Tanstetter von Thannau[6] (1480–1530) and Apianus[7] prepared editions of his work which were published at Nürnberg in 1533 and 1551. The treatise is divided into ten books, the first four being a summary of the works of earlier writers; the fifth, a treatment of reflection; the sixth, reflection by convex spheric mirrors; the seventh, cylindric and compound mirrors; the eighth, concave spheric mirrors; the ninth, concave conic mirrors and irregular mirrors; and the tenth, refraction.

Among the other medieval writers on perspective were William of Moerbecke[8] (*c.* 1250) and Campanus[9] (*c.* 1260).

Renaissance Writers. The first writers of the Renaissance to take up the subject were the painters and engravers. Pietro

[1] This in a Paris MS., formerly attributed to Alpetragius, discovered by Duhem and not yet printed. See Little, *loc. cit.*, p. 390.

[2] *De speculis comburentibus* and *Notulae de speculis*, both published at Frankfort (1614) in Combach's *Specula mathematica*, pp. 168–207.

[3] See Little, *loc. cit.*, p. 409 seq.

[4] See Volume I, page 224. It was often printed. For editions, see Kästner, *Geschichte*, II, 264; for Kästner's history of optics in general, *ibid.*, p. 237.

[5] See Volume I, page 228. On his work at Padua see A. Birkenmajer, *Witelo e lo Studio di Padova*, reprint, Padua, 1922.

[6] Professor of astronomy at Vienna.

[7] See Volume I, page 333.

[8] William Fleming. See Volume I, page 229.

[9] See Volume I, page 218.

Franceschi (or Della Francesca), for example, who died in 1492, wrote the work *De corporibus regularibus* and a work *De perspectiva pingendi*,[1] which is still extant in manuscript, and in which he takes up the theory of perspective.[2] There were also such artists as Leonardo da Vinci[3] (*c.* 1500), many of whose ideas on perspective, and particularly on the nature of vision and the camera obscura, were a distinct advance in knowledge; Benvenuto Cellini,[4] whose work on perspective was largely taken from Leonardo;[5] and Albrecht Dürer, whose work on drawing[6] includes some treatment of perspective.

One of the first men in this period to write a work of any note, devoted solely to optics, was Ramus[7] (*c.* 1550). This work was published by his pupil, Friedrich Risner[8] (died 1580), who also published the works of Alhazen (*c.* 1000) and Witelo (*c.* 1270).[9] The work of Ramus is in four books, but it contains little that Witelo did not give.

Optics in the 17th Century. In the 17th century the science of optics took a great step forward, notably through the efforts of Kepler. These efforts first appear in his unpretentious work of 1604, the *Paralipomena ad Vitellionem*, this Vitello (Witelo) being the German or Polish scholar already mentioned. In this little work Kepler explained the mechanism of the eye, comparing the retina to the canvas on which images were depicted. He showed that imperfect vision is caused by the failure of the rays of light to converge properly on the retina. In 1611 he published a work on dioptrics in which he set forth his ideas,

[1] G. Pittarelli, "Intorno al libro 'de perspectiva pingendi' di Pier dei Franceschi," *Atti del Congresso internazionale di scienze storiche*, XII (Rome, 1904), 262.

[2] H. Wieleitner, "Zur Erfindung der verschiedenen Distanzkonstruktionen in der malerischen Perspektive," *Repertorium für Kunstwissenschaft*, XLII (1920), 249.

[3] *Trattato della pittura*, Paris, 1651. See Volume I, page 294.

[4] Born 1500; died *c.* 1571. Various dates of his death are given, ranging from December 13, 1569, to February 25, 1571.

[5] P. Duhem, *Études sur Léonard de Vinci*, sér. I, p. 225 (Paris, 1906); G. P. Carpani, *Memoirs of B. Cellini*, English translation by Roscoe, London, 1878; *Œuvres complètes de Benvenuto Cellini*, 2d ed., 2 vols., Paris, 1847.


[6] *Underweysung der messung*, Nürnberg, 1525; see Volume I, page 326.

[7] See Volume I, page 309.

[8] *Opticae libri quatuor, ex voto Petri Rami novissimo, per Fr. Risnerum . . .*, Cassel, 1606 (posthumous).

[9] Basel, 1572. See page 341.

imperfect though they were, upon the law of refraction. He also gave a scientific explanation of the telescope, then recently invented. In the same year (1611) Antonio de Dominis, archbishop of Spalato, published his *De Radiis Lucis in Vitris Perspectiva et Iride*, in which he explained more fully than his



DESCARTES'S EXPLANATION OF THE RAINBOW

From his *Meteora*, 1656 ed., p. 214

predecessors the phenomena of the rainbow, basing them upon principles of refraction. It was Descartes, however, who in his *Dioptrica* (1637) gave the law that the sine of the angle of incidence has a constant ratio to the sine of the angle of refraction, the ratio being a function of the medium. The law was, in fact, known to Snell twenty years earlier, but he had failed to set it forth in print, although he had taught it. Nevertheless,

Descartes was living in Holland at that time, and there is some suspicion that he had there heard of Snell's discovery. In his *Dioptrica*, Descartes completed the theory of the rainbow by giving an explanation of the outer bow.

Just before Descartes produced his work, François Aguillon (1566–1617), a Belgian Jesuit, published a treatise[1] of some importance. In this he used the term "stereographic projection," although the idea was known to the Greeks.

Frans van Schooten the Younger published in 1656–1657 a book of mathematical exercises[2] in which he treated of perspective,[3] but it contained little that was original.

Newton's Work. Newton began to work seriously on optics about 1666. In his treatise[4] of 1704 he states that part of the treatise was written in 1675; and in his posthumous work the editor states that Newton "first found out his Theory of Light and Colours" as early as 1666, lecturing upon it in 1669. By this time the elementary theory of optics was well established.

11. Instruments in Geometry

Early Instruments. Before the invention of the telescope, microscope, and vernier there can hardly be said to have been any instruments of precision. For practical land measure, however, for leveling, and for the measuring of heights, the world developed several interesting instruments worthy of mention.

In general, the ancient surveyors measured distances by the use of a rope or a wooden rod, the units of measure varying in different localities. They laid off right angles by the use of an

[1] *Francesci Aguilonii e societate Jesu Opticorum libri VI*, Antwerp, 1613.


[2] *Exercitationum Mathematicarum libri V*, Amsterdam, 1656–1657; Dutch edition, *ibid.*, 1659.

[3] "Een korte verhandeling van de Fondementen der Perspective." It was also separately printed, Amsterdam, 1660.

[4] *Opticks: or, a Treatise of the Reflexions, Refractions, Inflexions, and Colours of Light . . .*, London, 1704, with various later editions and translations. His second work in point of publication, but not of composition, was his *Optical Lectures Read in . . . 1669*, published posthumously, London, 1728.

instrument resembling the carpenter's square of the present time, by a kind of cross placed horizontally on a staff, or by the 3–4–5 relation applied to a stretched cord. For finding a level they ordinarily used a right-angled isosceles triangle with a plumb line. Illustrations of such instruments are found on monuments to certain ancient surveyors.[1]

Early Printed Books. The early printed books give us much information as to the nature of the instruments inherited from the Middle Ages. Of these there may be mentioned the mirror for the measuring of heights by the forming of similar triangles, the geometric square (*quadratum geometricum*), the quadrant,



THE QUADRATUM GEOMETRICUM

From Oronce Fine's *De re & praxi geometrica*, Paris, 1556. The two triangles being similar, *AB* is easily found from the distances *AC* and *AF*

the astrolabe, and the cross-staff (*baculum*, also called the *baculus*). The method of using most of these instruments is evident, but a brief description of some of them will be helpful.[2]

The Square. The simplest of all the instruments of this class was the ordinary carpenter's square, known in some of the works on mensuration as the geometric square. Its use in finding short distances by means of the principle of similar triangles will be easily understood from the above illustration.

[1] See page 357.

[2] W. E. Stark, "Measuring Instruments of Long Ago," *School Science and Math.*, X, 48, 126; M. Curtze, "Ueber die im Mittelalter zur Feldmessung benutzten Instrumente," *Bibl. Math.*, X (2), 65; M. Cantor, *Die Römischen Agrimensoren*, Leipzig, 1875; E. N. Legnazzi, *Del Catasto Romano*, Verona, 1887; G. Rossi, *Groma e squadro ovvero Storia dell' Agrimensura Italiana dai tempi antichi al secolo XVII°*, Turin, 1877.

The Baculum. In its simplest form the baculum, arbalète (crossbow), geometric cross, cross-staff, or Jacob's staff[1] was a rod about 4 feet long, of rectangular cross section, and having a crosspiece that could slide upon it and always remain perpendicular to it. The staff was marked off in sections each equal in length to the crosspiece. In actual use the crosspiece was



PRACTICAL MATHEMATICS IN THE 17TH CENTURY


From Simon Jacob's work of 1560 (1565 ed.)

first placed at one of the division marks of the staff, the observer then facing approximately the mid-point of a line that he wished to measure and standing at a distance such that, when he sighted along the staff, the crosspiece should be parallel to the line and just cover it. The crosspiece was then moved to the next division on the staff, the observer taking a position where the first process of covering the line with the crosspiece could be repeated, as shown in the illustration on page 347.

[1] This name had various other uses, however.

The length of the line to be measured was then the same as the distance between the two positions of the observer. There were also various other methods of using the instrument.[1]

Sector Compasses. About the year 1597 Galileo invented the proportional compasses,[2] or sector compasses, an ingenious device for solving a variety of problems often met by architects, engineers, and others who have much to do with applied mathematics. The instrument consists ordinarily of two brass rules



THE BACULUM, OR CROSS-STAFF


From Oronce Fine's *De re & praxi geometrica*, Paris, 1556, showing the methods of measuring distances

hinged at one end. There are usually six pairs of lines, three on each face, radiating from the pivot. One pair might, for example, represent equal parts; another, squares; and the third, lines of polygons; but this varied according to the purpose of the particular instrument.

To give a single illustration of its use, suppose that each line of equal parts is divided into 200 equal segments, numbered by tens, beginning at the pivot. Then, to divide any given line

[1] For a brief résumé see G. Bigourdan, *L'Astronomie, évolution des idées et des méthodes*, p. 116 (Paris, 1911; 1920 ed.), hereafter referred to as Bigourdan, *Astronomie*. [2] *Le operazioni del compasso geometrico e militare*, Padua, 1606.

segment into any number of equal parts, say nine, open a pair of ordinary dividers to the length of the segment, then open the sector compasses so that one point of the dividers rests on




ASTROLABE OF CHAUCER'S TIME

Fine piece of medieval workmanship now in the British Museum. It may well be that Chaucer himself made use of this in preparing his treatise on the astrolabe


90 on one face and the other point rests on 90 on the other face; then the distance from the 10 on one face to the 10 on the other is one ninth of the length of the given line segment.

Astrolabe. Of all the early astronomico-mathematical instruments none was better known than the astrolabe. The name



ITALIAN ASTROLABE OF 1558


It bears the inscription "Patavii Bernardinvs Sabevs faciebat MDLVIII."
From the author's collection



THE ASTROLABE IN SIMPLE MENSURATION

From Bartoli's *Del Modo di Misvrare*, Venice, 1589, showing simple work in a crude kind of trigonometry

is Greek and means the taking of the stars.[1] Hence any instrument for measuring the angles by which a star was "taken" (as a sailor today speaks of "taking" the sun) was, strictly speaking, an astrolabe. One of the early forms was the armillary sphere, so called from the *armillae*,[2] or rings, which were so arranged as to form two, or sometimes three, circles, ordinarily placed at right angles to one another. One ring usually corresponded to the plane of the equator and the other to the plane of the meridian. By these two circles the ancients determined the two coordinates of a star. The astrolabe described by Ptolemy the astronomer is a kind of armillary sphere,[3] and furthermore these spheres are first heard of in connection with the school with which he was associated. It is asserted by early writers that Eratosthenes, through his interest in geodesy and astronomy, induced King Ptolemy III to have such instruments made and placed in the museum at Alexandria.



CHAMPLAIN'S ASTROLABE


Found near the Ottawa River about 1870. It was made in Paris in 1603. This is the type of astrolabe known as the planisphere. From the collection of Samuel V. Hoffman, New York

[1] From ἄστρον (*as'tron*, a heavenly body) + λαμβάνειν, λαβεῖν (*lamba'nein, labein'*, take). Ptolemy spoke of the two circles that he used in locating a star as ἀστρολάβοι κύκλοι (*astrola'boi ky'kloi*) and spoke of the whole instrument as ἀστρολάβον ὄργανον (*astrola'bon or'ganon*) or, commonly, as ὁ ἀστρολάβος (*ho astrola'bos*). See J. Frank, "Zur Geschichte des Astrolabs," *Erlangen Sitzungsberichte*, 50–51. Band, p. 275; R. T. Gunther, *Early Science in Oxford*, II, 181.

[2] *Armilla* means an armlet, bracelet, hoop, or ring. It is probably a diminutive of *armus*, the shoulder or upper arm.

[3] *Almagest*, VII, 2, 4.

Planisphere. Another ancient and common form of the astrolabe consisted simply of a disk upon the rim of which were marked the units of angle measure. Such instruments were probably well known in ancient times among all who made any scientific study of the stars. That they were familiar in ancient Babylon we have definite proof.[1] Fragments of several such instruments have been found and the inscriptions interpreted. They go back to the 2d millennium B.C., which goes to show that the early Greeks undoubtedly knew of their value and made



THE QUADRANT

From the *Protomathesis* of Oronce Fine, Paris, 1530–1532


use of them in angle measure. These astrolabes are in the form of planispheres and are made of clay, baked like the tablets.[2] A planisphere may be defined as a stereographic projection of the celestial sphere either upon the plane of the equator or upon the plane of the meridian.

Such instruments were used in various practical ways in which angle measure was the chief purpose, and this use continued until recent times. Even now they are seen in the Orient in the hands of the astrologers.

[1] E. F. Weidner, *Handbuch der Babylonischen Astronomie*, Lieferung I, 62; with bibliography, Leipzig, 1915.

[2] For a photographic reproduction, see Weidner, *loc. cit.*, p. 107, from A. Jeremias, *Handbuch der altorientalischen Geisteskultur*, Leipzig, 1913. There is a good specimen in the British Museum.

The planisphere in common use in later times represents the stereographic projection of the celestial sphere upon the plane of the equator, the eye being at the pole. Planispheres of




BRASS QUADRANT

Austrian work of the 18th century. The original is 29.5 cm. square. From the author's collection

various types were used by early navigators for the purpose of finding the elevation of the north star, or for other angle measurements, and were often furnished with several plates which could be so adjusted as to allow the instrument to be used in different latitudes.

The Astrolabe in the East. From Babylon[1] the astrolabe may have passed to China and India, or vice versa. At any rate, Mesopotamia seems to have been the source from which the Greeks derived their knowledge of the instrument. It is probable that Thales used it in measuring the distances of ships, since the Babylonian astronomy was already becoming known in the Greek civilization. It may be inferred from Plato's *Timæus* that some such instrument was in use in his day, but in any case an astrolabe of some type was known to Eratosthenes, Hipparchus, and other Greek astronomers even before Ptolemy described the armillary sphere.[2]



SMALL IVORY QUADRANT


Italian work of the 18th century. If we sight through holes in the two projections on the upper right-hand edge, the angle of elevation is indicated by the plumb line and the arc. The original is 5 cm. by 5.9 cm. From the author's collection

Arab Treatises on the Astrolabe. Led by their study of Greek astronomy, the Arabs, beginning in the 9th century, wrote numerous works upon the astrolabe, and these, in turn, influenced the medieval scholars of Europe. Thus we find Messahala (*c.* 800) composing a work upon the subject, which formed the basis of two manuscripts by Rabbi ben Ezra (*c.* 1140). From one of these manuscripts Chaucer (*c.* 1380) seems to have drawn his information for his treatise upon the astrolabe.

[1] A. H. Sayce and R. H. M. Bosanquet, "Babylonian Astronomy," *Monthly Notices of the Royal Astron. Society*, XL, No. 3, with illustrations.

[2] R. Wolf, *Geschichte der Astronomie*, p. 160 (Munich, 1877).


INSTRVCTIO
CAPVT XII.
QVA RATIONE ELICIENDA SIT ALTITVDO ALICVIVS REI, QVAE ACCESSVM NON admittit: Vt sunt Mœnium, arcium, &c.



EXPLANATION OF THE QUADRANT

From *De Quadrante Geometrico*, usually referred to Cornelius de Judeis, Nürnberg, 1594, but in fact written by Levinus Hulsius. Cornelius made the drawings with the help of Martin Geet


Quadrant. Closely related to the astrolabe is the quadrant, an instrument in which only a quarter of a circle is used. It



USE OF THE QUADRANT


From Ottavio Fabri's *L' Uso della Squadra Mobile*, Trent, 1752

appears in various forms, sometimes without an arc, the angles being read on the sides of a square. The earliest description that we have is given in the *Almagest*, and on this account the




DRUMHEAD TRIGONOMETRY

A common method of triangulating in the 16th century. From Belli's *Libro del Misvrar*, Venice, 1569



EARLY APPROACH TO THE PLANE TABLE


The plane table in various forms was probably developed from such an instrument as the one here shown. In this case the table was used merely for taking horizontal angles. From Cosimo Bartoli, *Del Modo di Misvrare*, Venice, 1589



USE OF THE SHADOW AND THE MIRROR IN MEASURING HEIGHTS

From Giovanni Pomodoro's *La Geometria Prattica*, Rome, 1624

honor of its invention is usually awarded to Ptolemy. He used a stone cube, on one of the faces of which the quadrant was cut. On this was mounted a small cylindric pipe, as we should mount a telescope, and by this device he was able to take the height of the sun, evidently by means of the ray of light which shone through the cylinder. There is no indication of the size of Ptolemy's quadrant, but if we judge by the later specimens in use in the East, and by the incomplete records, it was probably a large one. He says that he used his quadrant in taking many astronomical observations; but he gives no results, and it is rather doubtful whether he did more than suggest the instrument, depending upon the results secured by Eratosthenes and others of his predecessors.




ANCIENT LEVELS AND SQUARE

From the tomb of Marcus Æbutius Macedo. Like the tomb of Lucius Æbutius Faustus (page 361), it is of uncertain date

The quadrant is described in many works of the 16th, 17th, and 18th centuries, but with the invention of the telescope all devices of this kind gradually gave way to the transit in astronomical work. The sextant was invented by Thomas Godfrey, of Philadelphia, in 1730.

Drumhead Trigonometry. The continual warfare of the Renaissance period shows itself in many ways in the history of mathematics. Some of these manifestations are mentioned from time to time in this work, and one of them is related to the subject now under consideration. Several writers of the 16th century give illustrations of the use of the drumhead as a simple means of measuring angles of elevation in computing distances to a castle or in finding the height of a tower. Such an illustration is shown on page 355 and is self-explanatory.




EARLY METHODS OF LEVELING

From Pomodoro's *La Geometria Prattica*, Rome, 1624. This was an early Egyptian method and was transmitted through the Greek and Roman surveyors

Somewhat related to this crude instrument is one for taking horizontal angles, as illustrated on page 356. From this device the plane table was probably developed.

The Mirror. In the early printed works on applied geometry there are frequent references to the speculum, a horizontal mirror used in measuring heights by the aid of similar triangles. The method is still in use for certain purposes, but in the 16th and 17th centuries it seems to have been extensively employed. On account of the difficulty of obtaining a satisfactory level, and the fact that one triangle was




JAPANESE LEVELING INSTRUMENT

From Murai Masahiro's *Ryōchi Shinan*, a work on surveying, about 1732

small and not easily measured with accuracy, the method was not of much value. The plan of using the speculum was based upon the principle of similar triangles and is illustrated on page 356.

Leveling Instruments. The common leveling instrument of ancient times was the isosceles triangle with a plumb line from the vertex. This is found in Egyptian remains, is represented on the monuments of Roman surveyors,[1] is referred to by medieval writers, and is still in general use in various parts of the world. Until the invention of the telescope, and the consequent increase in accuracy of observation, it satisfied all ordinary needs. There are many reasons for believing that the early Egyptian surveyors who laid out the pyramids made use of this instrument for establishing their levels. An interesting variant of this instrument is seen in the quadrilateral which the Japanese scholars developed before the free influx of Western mathematics. Such a device is shown on page 358.



EARLY JAPANESE SURVEYING INSTRUMENT

From a drawing in a manuscript of a work (see page 358) by Murai Masahiro, about 1732

The principles underlying the later forms of leveling instruments were not numerous; in fact, the fundamental ones were only two in number,[2] the older one depending upon the plumb line, and the later one upon the state of

[1] On the leveling instruments of the Romans see C. G. de Montauzan, *Essai sur la science et l'art de l'ingénieur aux premiers siècles de l'Empire Romain*, pp. 46, 62, 74 (Paris, 1908). On Greek and Roman engineering instruments see Pauly-Wissowa; R. C. Skyring Walters, *Transactions of the Newcomen Society*, II, 45, and T. East Jones, *ibid.*, p. 61; E. N. Legnazzi, *Del Catasto Romano*, Verona, 1887.

[2] For a discussion of the subject see N. Bion, *Traité de la Construction et des principaux Usages des Instrumens de Mathematique*, p. 285 (Paris, 1713; ed. of The Hague, 1723). This is the best of the early classical treatises upon the subject of mathematical instruments, and is profusely illustrated.

equilibrium of some kind of liquid. The former was used in the ancient triangle illustrated on page 358, and in related types, while the latter is still seen in the ordinary level used by carpenters and in the leveling instrument used by engineers,—the *niveau à l'eau* described by Bion (1713). The triangle level had various special forms, such as an inverted T (that is, ⊥) with a plumb line along the vertical arm. In this form it was called by French writers the *niveau d'air*. The horizontal part was usually a tube through which the observer could sight when running a level line.

After the telescope was invented the tube was fitted with lenses,[1] and the instrument became, either with the plumb line or with the water level attached, not unlike the instrument in common use today. Sometimes the plumb line and the level were attached to the same instrument. Huygens invented a level in the form of a cross on which an inverted T was hung by a ring at the top, the telescope being kept horizontal by means of a weight.[2]

Until the advent of a new type of engineering, made possible by the commercial use of structural steel, the level was used chiefly for two purposes. The first of these was the construction of canals for purposes of irrigation, particularly in Mesopotamia and Egypt, and of aqueducts as a result of the Roman demand for pure water. The second use was seen in the building of fortifications, particularly during and as a result of the wars of the 17th century. The textbooks of that century on applied geometry (mensuration) gave much attention to the subject. The general practice in leveling was not unlike that of the present time, the chief difference being in the degree of precision of the instruments used. It is evident that in construction work of any extent the level was always necessary, but its elaborate use in modern engineering—as in railway gradients, tunnels, and bridges, and as in the erecting of modern office buildings of great height—surpasses anything conceived of in ancient times.

[1] The *niveau d'air à lunette* of Bion's treatise of 1713.
[2] Bion, *loc. cit.*, p. 110.

Other Surveying Instruments. It is not possible, in the limited space that should be allowed the subject in a work of this kind, to mention all the simple surveying instruments and devices that have come down to us from the Egyptian, Greek, and Roman civilizations. The simple staff, with a crude diopter through which to sight in running a line, is found in all parts of the world and is probably very ancient.


From such a humble origin sprang the *groma* used by the Roman surveyors and here illustrated from the tomb of Lucius Æbutius Faustus. He is mentioned in the third line as a *mensor*, but the term more commonly used in the case of a land surveyor was *agrimensor* (field measurer) or *gromaticus* (one who used the *groma*). The *groma* (*cruma*, *gruma*) consisted of the *stella* (the star-shaped part) and the *pondera* (the plumb lines). Surveyors in the time of the empire often spoke of the *machina* or *machinula* which they used and which consisted of two parts, the *groma* or *stella* and the *ferramentum* (the iron standard). Hence Hyginus (*c.* 120) says "ferramento groma superponatur" (let the groma be placed upon the iron standard).



ANCIENT SURVEYING INSTRUMENT

From the tomb of Lucius Æbutius Faustus, a *mensor* (*agrimensor*, surveyor, or perhaps a measurer for architects and builders). The entire inscription was as follows: Tribv Clavdia Lvcivs Aebvtivs Lvcii libertvs Favstvs mensor sevir sibi et Arriae Qvinti libertae avctae vxori et svis et Zepyre libertae vivvs fecit. The instrument shown is the *groma*

In the 17th and 18th centuries, stimulated by the metal work of the Italian artists, the instrument-makers of France and Italy produced many beautiful pieces of workmanship designed with much ingenuity. These pieces are occasionally seen in museums, and one is shown in the following illustration.[1]




ELABORATE MATHEMATICAL INSTRUMENT

Showing artistic metal work of the 18th century. Now in the Metropolitan Museum of Art, New York

Such instruments were often elaborately engraved and some of those apparently made for the noble patrons of the sciences were even gold-plated. One of the elaborate forerunners of the range finder is shown on page 363.

[1] R. T. Gunther, *Early Science in Oxford* (Vols. I and II, Oxford, 1922, 1923), with a catalogue of the early mathematical instruments belonging to the University and colleges of Oxford. See particularly II, 192–233.



EARLY FORM OF RANGE FINDER

From Danfrie's *Declaration de l'Vsage du Graphometre*, Paris, 1597; appendix on trigonometry, p. 11

Other Astronomical Instruments. While this work is only indirectly concerned with astronomy, many astronomical instruments are distinctly mathematical, and some of them are particularly interesting as works of art. Such are the elaborate bronze pieces on the walls of the city of Peking, mostly due to the Jesuit influence which began about 1600, but partly native in their design and general plan. It was through the devising and use of instruments like these that such missionaries as F. Verbiest and J. A. Schall von Bell (*c.* 1660) were able to make observations that demonstrated, even to the hostile critics, the superiority of European astronomy over that of the Chinese. One of these pieces is shown in the illustration.[1]



BRONZE QUADRANT ON THE WALLS OF PEKING


One of several elaborate bronze instruments, most of them made under the influence of Jesuit missionaries

In Persia and India there are still to be found celestial spheres of great beauty, generally dating from the 17th century. These are usually of bronze, sometimes with silver stars.

The Hindus, Persians, and Arabs have also left many astrolabes of beautiful workmanship, some of them with constellations or particular stars represented in silver. Until the invention of the telescope their smaller types of astronomical instruments were unsurpassed both in beauty and in accuracy.

[1] See Volume I, page 272.

The most interesting of the Hindu instruments are found in the five observatories built by the Maharajah Jai Singh between 1728 and 1734.[1] These observatories were located at Delhi, Jaipur, Benares, Ujjain, and Mathurā, and represent the Arab astronomico-astrological science instead of the native Hindu or the European. Jai Singh was a Sikh by birth and was so interested in astronomy that he translated Ulugh Beg's




HINDU CELESTIAL SPHERE

This piece is of bronze, the stars being inlaid in silver. It was made *c.* 1600. From the author's collection

catalogue of the stars (*c.* 1435). He was of the opinion that the small brass instruments used in Samarkand were not accurate enough, and hence he determined to construct pieces so large and substantial as to leave no doubt about the validity of the observations. The results were monumental and are still the object of admiration to those interested in the science of India. An illustration showing one of the most elaborate and carefully preserved of these observatories (the one at Jaipur) will be found on page 366.

[1] G. R. Kaye, *The Astronomical Observatories of Jai Singh*, Calcutta, 1918; see also the review in the *Journal of the Royal Asiatic Soc.*, July, 1919, p. 427.

The Jaipur observatory was constructed by the Maharajah Jai Singh about 1734. The Jesuit missionary Joseph Tieffenthaler, in a work published in 1785, speaks of it as follows:




OBSERVATORY AT JAIPUR, INDIA

Showing the kinds of instruments generally used before the days of the telescope. This observatory, although relatively modern, is based upon ancient models

It is such a work as is never seen in this part of the world and, by the novelty and grandeur of the instruments, strikes one with astonishment. . . . What attracts most attention is a gnomon (*axis mundi*), remarkable for its height of 70 Paris feet. . . . There are three very large astrolabes, cast in copper, suspended by iron rings.

This is all quite as impressive to the visitor now as it was then. The instruments, which had become damaged through age and neglect, were restored in 1902.



JAPANESE CELESTIAL SPHERE

From a wood engraving in Baba Nobutake's *Shogaku Tenmon* (1706)

The Chinese influence shows itself in the Japanese works of the 17th and 18th centuries, as is seen in the illustration from Baba Nobutake's work of 1706. We also find in Japan in this same period the use of the pierced sphere in astronomical observations and in the work of the astrologers. This device was common in Europe in the latter part of the Middle Ages and is found in various printed works of the 16th century.



JAPANESE FIGURE OF AN ASTRONOMER

Caricature in ivory. From the author's collection

12. The Problem of Earth Measure

Need for Instruments of Precision. The need for instruments of a high degree of precision was first felt in connection with astronomy and the measure of the earth. The subject is too extensive to be considered at length in a work of this kind, but its general nature will be understood by a brief reference to the history of the measure of the earth's circumference and density.

Circumference determined from Arc. It should first be understood that the solution of this problem did not involve the ratio of the circumference to the diameter; it required the finding of the circumference when the diameter was unknown. When first undertaken it had nothing to do with navigation, economics, or military conquest; it developed as a purely abstract contribution to human knowledge. The plan adopted by the Greeks was the same, in basic principle, as the one used today, namely, that of measuring the amplitude and the length of an arc of a great circle (generally

a meridian) and from these data computing the circumference. This led to one of the many branches of geodesy, a subject into the history of which we cannot enter at length in this work.

Application of Circle Measure to Geodesy. Pythagoras (*c.* 540 B.C.) was the first, so tradition asserts, to teach that the earth is a sphere and that it is situated in the center of the universe.[1] This idea was accepted by various Greek philosophers, and Aristotle (*c.* 340 B.C.) states that "the mathematicians who have attempted to calculate the circumference of the earth say that it may be forty myriads of stadia,"[2] that is, 400,000 stadia. The stadium varied so much with ancient writers that this does not give us any very satisfactory information.[3] Taking a rough approximation, however, say ten stadia to an Anglo-American mile, this makes the circumference 40,000 miles. Aristotle gives us no information as to the names of the mathematicians who made the calculations, and none as to the method employed, but it has been thought that the approximation is due to Eudoxus (*c.* 370 B.C.). It is evident, however, that the circumference was found by multiplying the length of a known arc, and not by using the ratio of the circumference to the diameter.

Four Greek Computations of the Earth's Circumference. From the time of Aristotle to that of Ptolemy (*c.* 150) there were four noteworthy attempts at measuring the earth's circumference. Of these the first is referred to by Archimedes (*c.* 225 B.C.), who speaks of certain writers as having stated that the circumference is 30 myriads of stadia, say about 30,000 miles. He does not mention the writers, and it is possible that he may have referred to some of the earlier attempts made by his friend Eratosthenes (*c* 230 B.C.). In his computation of the number of grains of sand in the universe, however, he takes the circumference as ten times this distance, so as to be on the safe side.[4]

[1] On this entire subject see Bigourdan, *Astronomie*, p. 144 seq.

[2] *De Cœlo*, II, 16.

[3] One of the stadia was 125 paces (double steps), or, say, 625 Roman feet, equal to 606¾ Anglo-American feet.

[4] *Archimedes*, ed. Heiberg, I, 221.

The third important attempt at the measure of the earth's circumference is definitely known to have been made by Eratosthenes, and the fourth by Poseidonius (*c.* 100 B.C.).

Eratosthenes on the Measure of the Earth. The first attempt of which we have any details is this third one,—the one briefly described as due to Eratosthenes. Supplementing the description given in Volume I, page 110, it may be said that Eratosthenes used the arc of a great circle extending from Syene (the modern Assouan) to Alexandria. He took the length of this arc as 5000 stadia, but how this length was ascertained is not stated in any ancient writings. It is probable that the official pacers,[1] employed by Alexander and other military leaders in planning their campaigns, had made reports of all such standard distances, and Eratosthenes, as librarian at Alexandria, doubtless had access to their records.

It was well known that on the day of the summer solstice the sun's rays lighted up completely the wells of Syene at noontime, and that a body like an obelisk cast no shadow. On the other hand, Eratosthenes found that the zenith distance of the sun on this day, as measured at Alexandria, was $\frac{1}{50}$ of the circumference. It is not known how this angle was found, but it has been thought that Eratosthenes used certain armillary spheres which tradition says were furnished by the king, Ptolemy Euergetes. We are told by Cleomedes (*c.* 40 B.C.), however, who wrote on the *Circular Theory of the Heavenly Bodies,* that he used the *sca'phe,*[2] a concave sundial, and this may easily have been the case. Since the zenith distance of the sun changed $\frac{1}{50}$ of the circumference in 5000 stadia, Eratosthenes concluded that the circumference was 50×5000 stadia, or 250,000 stadia, roughly equivalent to 25,000 miles, or 40,000 kilometers. This would make $694\frac{4}{9}$ stadia to a degree; and since Eratosthenes was naturally aware that his measurements were merely approximate, he felt it allowable to take 700

[1] Βηματισταί (*bematistai'*), singular βηματιστής (*bematistes'*), from βῆμα (*be'ma*), a step.

[2] Σκάφη, originally anything dug out; hence a bowl, and then the bowl of a hemispherical sundial.

stadia as a more convenient measure for 1°. He had crude instruments with which to work, he did not take into consideration the difference of longitude of his two stations, and the stadium was a varying unit at best, so that his assumption of 700 stadia was not an unreasonable one.

Poseidonius on the Measure of the Earth. Poseidonius (*c.* 100 B.C.) was, as we have seen,[1] a Stoic philosopher, well known as an astronomer, a geographer, a historian, and a statesman. After having traveled extensively in all the Mediterranean countries, he opened a school at Rhodes and had among his pupils both Cicero and Pompey. Although his works are lost, Cleomedes (*c.* 40 B.C.) has given us a certain amount of information as to his method of measuring the circumference of the earth. Like Eratosthenes he took a known arc, selecting the one from Rhodes to Alexandria and estimating its length as 5000 stadia. He then observed that when the star Canopus was on the horizon at Rhodes, it was $\frac{1}{4}$ of a sign (that is, $\frac{1}{4}$ of 30°, or $\frac{1}{4}\frac{1}{8}$ of 360°) above the horizon at Alexandria. He concluded that the circumference is 48 × 5000 stadia, or 240,000 stadia. This made the length of the degree $666\frac{2}{3}$ stadia.

It is hardly probable that Poseidonius considered these results as close approximations, since neither the length of his arc nor the elevation of the star could be measured with any approach to accuracy by instruments then available.

Ptolemy on the Measure of the Earth. The last of the noteworthy attempts of the ancient Greeks to find the circumference of the earth was made by Ptolemy (*c.* 150). He took a degree as 500 stadia, thus finding the circumference to be 180,000 stadia. He asserted that it was unnecessary to take the arc of a meridian, an arc of any other great circle being sufficient. We are without information, however, as to his method of measuring the arc selected. It will be noticed that his result is to that of Poseidonius as 3:4; and since this is the ratio between two of the stadia employed by the ancients, it is possible that he simply used the latter's computations.

[1] Vol. I, p. 118. See O. Viedebantt, "Poseidonius," *Klio*, XVI, 94.

The theory has been advanced that all these results set forth by the Greeks were due to Egyptian or other measurements which are no longer extant, but there is no scientific basis for the conjecture.

Arab Measure of the Earth. It was some seven centuries after the last of the Greek geodesists that the Arabs engaged in the work of measuring the circumference of the earth. By order of al-Mâmûn certain mathematicians of Bagdad undertook the necessary surveys on the plain of Sinjar in Mesopotamia. They formed two groups, one party going to the north and one to the south, each proceeding to a point at which the elevation of the pole changed 1° from that of the base station. They then measured the respective distances, one being found to be 57 miles and the other $56\frac{1}{4}$ miles; this mile was given as 4000 "black cubits," but the length of this cubit is now unknown. The difference in the two measurements illustrates the lack of the necessary instruments of precision, even among a people who had brought the construction of such instruments to the highest degree of perfection known at that time.

Invention of the Telescope. With respect to instruments used in astronomy and geodesy the greatest improvement is due to the invention of the telescope, although much is also due to modern technique in manufacture and to the use of such devices as the vernier and the micrometer.

Roger Bacon (*c.* 1250) stated that it was possible to construct tubes by means of which distant objects could be seen as if they were near at hand,[1] but we have no evidence that this was other than a prophetic statement by a man who seemed peculiarly gifted in this respect. Possibly he was led to this prophecy by a knowledge of spectacles, which appeared sometime in the 13th century, for they were certainly known in 1299, and a certain Salvino degli Armati, a Florentine (died in 1317), is mentioned as their inventor.

[1] "Ita ut in incredibili distancia videremus arenas et litteras minias minutas, et ut altissima videantur infima et e contrario." Sloane Ms., fol. 84, a, 2. Cf. *Opus Majus* (ed. Bridges), ii, 164; *Opus Tertium* (ed. Little), 41.

The possibility of the telescope is also mentioned by Fracastorius, who, in his *Homocentricorum seu de stellis Liber Unus* (1538), speaks of using two superposed lenses in looking at a distant object. Somewhat similar statements were made by Giambattista della Porta in his *Magia Naturalis* (Naples, 1558) and by Kepler in his *Paralipomena* (Frankfort, 1604).

The invention seems due, however, not to the work of scientists like those mentioned, but largely to chance. It is uncertain who was the actual inventor, the claims of three artisans being about equal. These men are Zacharias Janszoon (Jansen), Johann Lippersheim (Lippershey, Lipperseim, Laprey, *c.* 1608), and Jacob Metius Adriaenszoon[1] (*c.* 1608).

Janszoon lived at Middelburg, was a grinder of lenses, and, apparently with the aid of his father, improved the microscope in 1590 and is known to have had a telescope in 1610.

Lippersheim was also a lens grinder of Middelburg. He is known to have asked for a patent in 1608 for an instrument intended to see distant objects, the lenses being of rock crystal.

Descartes and others attributed the invention to Jacob Metius Adriaenszoon (*c.* 1608), who happened to make the necessary combination of certain lenses and burning mirrors. He also asked for a patent in 1608, a few days after Lippersheim had made his request.[2]

The invention is known to have been made public in October, 1608, and the knowledge of the instrument spread all through Europe with astonishing rapidity. Even in 1608, and still more in 1609, instruments were made in France, England, Italy, and Germany. Hearing of the new device, Galileo, in 1609, invented an instrument of his own and by its aid at once made remarkable discoveries in astronomy; and from this time on a precision of observation unknown to earlier scientists, although the instrument was not accepted by all astronomers, became possible. It was only after the invention of achromatic lenses, however, that satisfactory results were obtained.

[1] He was a brother of the Adriaen Metius (1571–1635) mentioned in Volume I, page 340. He was born at Alkmaar and was interested in the grinding of lenses.

[2] Bigourdan, *Astronomie*, p. 124.

Modern Measures of the Earth. The first noteworthy modern attempt at measuring the earth was made by Jean Fernel (*c.* 1528), physician to Henri II of France. Fernel took the arc determined by Paris and Amiens, two stations being taken on the same meridian. Knowing the altitude of the sun at Paris, he proceeded northward to a point where the altitude was 30′ less than that at Paris. He then measured the arc by taking the number of revolutions of a wheel of known circumference. No record is available as to his method of allowing for errors, but certain compensations were made and the conclusion was reached that $1° = 57{,}099$ toises, 1000 toises being a little more than a geographic mile. This result is remarkable, the mean afterwards obtained by Lacaille[1] and Delambre in the latter part of the 18th century being 57,068 toises.

In 1617 Snell undertook the measurement of an arc by an elaborate system of triangulation, and although his results were satisfactory as to length of arc, they were not so as to amplitude.

Further French Attempts. In 1669 and 1670 Jean Picard[2] carried on an elaborate system of triangulation, measured an arc from a point near Corbeil to one near Amiens, and found that 1°21′54″ corresponded to 68,347 toises 3 pieds, which gave 57,060 toises to 1°. He estimated that the amplitude was correct to within 2″ or 3″.

In 1686 Newton proved that the earth is an oblate spheroid, a result not generally accepted by French scientists, chiefly owing to the conclusions reached by Jacques Cassini (Cassini II) as mentioned below. It was therefore decided that France should undertake a more elaborate and careful survey, not confined to that country alone, but including arcs nearer to and more remote from the equator.

[1] Nicolas Louis de Lacaille (La Caille); born at Rumigny, May 15, 1713; died in Paris, March 21, 1762. He wrote upon mathematics and physics, but chiefly upon astronomy.

[2] Born at La Flèche, Anjou, July 21, 1620; died in Paris, July 12, 1682. He wrote upon physics and astronomy and was particularly well known for his work on the measure of the earth.

The degree of accuracy reached by Picard was increased through the efforts of Giovanni Domenico (Jean Dominique) Cassini (Cassini I), who extended Picard's meridian in 1701 southward to the Pyrenees. It was afterwards extended northward to Dunkirk (Dunkerque), although the results, published by Jacques Cassini (Cassini II) in 1720, provoked great opposition because of their lack of precision and the incorrect conclusions reached with respect to the elongated form of the earth. In 1735 France sent a mission to Peru, and an elaborate survey was made for the purpose of measuring an arc. This work was carried on under the direction of Bouguer,[1] Condamine,[2] and Godin.[3] By 1745 they had completed the measurement of an arc of 3°. D'Alembert spoke of the work as the greatest scientific enterprise that had thus far been undertaken. In the following year another mission, including such French scientists as Maupertuis and Clairaut, and the Swedish scientist Celsius,[4] began a similar work in Lapland. The result of this survey was the measurement of an arc of 1°. The conclusions reached in Peru and Lapland confirmed Newton's assertion of the flattening of the earth at the poles and led to Voltaire's reference to Maupertuis, against whom he had a personal grudge, as the "great flattener" (*grand aplatisseur*). The form of the earth appears from the fact that degrees of latitude increase in length as we approach the poles.

In 1739 and 1740, owing chiefly to the work of Lacaille, an arc of the meridian was again measured in France, the result being a correction of the errors published in 1720 and a new confirmation of Newton's theory of the shape of the earth.

Toward the close of the 18th century France undertook a third great survey, this time for the purpose of determining the

[1] See page 327.

[2] Charles Marie de la Condamine, born in Paris, January 28, 1701; died in Paris, February 4, 1774. He wrote extensively on geodesy.

[3] Louis Godin, born in Paris, February 28, 1704; died at Cadiz, September 11, 1760. He wrote chiefly on astronomy.

[4] Anders Celsius; born at Upsala, November 27, 1701; died at Upsala, April 25, 1744. He was professor of astronomy at Upsala, but spent some years in France, Germany, and Italy.

length of the standard meter. In this undertaking a number of the greatest French scientists were engaged, but for the geodetic work Delambre and Méchain were chiefly responsible.

In the 19th and 20th centuries extensive triangulations have been made, and with the methods employed there have been connected such prominent names as those of Biot, Arago, Schumacher, Legendre, Laplace, Gauss, and Bessel. The Ordnance Survey of Great Britain, begun in 1783 and completed in 1858, resulted in the measurement of an arc of 10° 13′, extending from the Isle of Wight to one of the Shetland group; the triangulation of India (1790–1884) gave an arc of about 24°; and the Russo-Scandinavian measurements, begun in 1817, resulted in an arc of 25° 20′. The arc recently measured in Africa, extending over a distance of about 65°, will, joined to the Russo-Scandinavian arc, give an arc of about 106°.

Mass of the Earth. The determination of the earth's density depends on the law of gravitation, and so it began in the work of Newton, who estimated it as five or six times that of water.

The first of the later methods depends upon the deflection of a plumb line due to the attraction of a mountain. This was first used by Pierre Bouguer, in Peru (*c.* 1740). By this plan Maskeleyne[1] (1774–1776) placed the density between 4.5 and 5.

The second method is based upon a comparison of the vibrations of a pendulum at sea level with those at the top of a high mountain. Francesco Carlini, the Italian astronomer, used the method in 1821 and obtained a density of 4.84.

The third method is due to Henry Cavendish[2] (1798) and is based upon the mutual attraction of known masses. Francis Baily[3] (1843) obtained the result of 5.67 by this method.

The fourth method uses a finely graduated balance to determine the attraction of known masses. By its use results of 5.69 were obtained by Von Jolly in 1881, and 5.49 by Poynting in 1891. The latest experiments give the result as about 5.53.

[1] Nevil Maskeleyne, born in London, October 5 (O. S.), 1732; died at Greenwich, February 9, 1811. He became astronomer royal in 1765.

[2] Born at Nice, October 10, 1731; died in London, February 24, 1810.

[3] Born at Newbury, Berkshire, April 28, 1774; died in London, August 30, 1844.

TOPICS FOR DISCUSSION

1. Intuitive geometry as it shows itself in the primitive decoration used by various peoples.

2. Intuitive geometry as it shows itself in the early stages of mathematics in various countries.

3. The rise of demonstrative geometry and the six most important contributors to the science in ancient Greece.

4. The various names used for geometry and the special significance of each.

5. The development of the terminology of elementary geometry, especially among the Greeks.

6. The development of the postulates and axioms of elementary geometry before the 19th century.

7. Propositions of elementary geometry of which the origin is known or which have any history of special interest.

8. The various methods of solving each of the Three Famous Problems of antiquity.

9. The historical development of methods for finding the approximate value of π.

10. The principal steps taken by the Greeks in the development of geometric conics.

11. The principal steps in the development of plane analytic geometry, with special reference to the 17th century.

12. A discussion of the history of solid analytic geometry.

13. The history of the most important higher plane curves commonly found in the study of elementary analytic geometry, together with the applications of these curves.

14. The nature, purpose, and history of descriptive geometry.

15. The relation of the fine arts to geometry in the 15th century.

16. The development of projective geometry.

17. The development of the non-Euclidean geometries, with special reference to the work of Bolyai, Lobachevsky, and Riemann.

18. The development of perspective and optics considered as mathematical subjects.

19. A study of the most interesting of the primitive instruments.

20. The general development of geodesy, particularly among the Greeks and in modern times, and with reference to the measure of the circumference of the earth.

CHAPTER VI

ALGEBRA

1. General Progress of Algebra

Nature of Algebra. When we speak of the early history of algebra it is necessary to consider first of all the meaning of the term. If by algebra we mean the science which allows us to solve the equation $ax^2 + bx + c = 0$, expressed in these symbols, then the history begins in the 17th century; if we remove the restriction as to these particular signs, and allow for other and less convenient symbols, we might properly begin the history in the 3d century; if we allow for the solution of the above equation by geometric methods, without algebraic symbols of any kind, we might say that algebra begins with the Alexandrian School or a little earlier; and if we say that we should class as algebra any problem that we should now solve by algebra (even though it was at first solved by mere guessing or by some cumbersome arithmetic process), then the science was known about 1800 B.C., and probably still earlier.[1]

A Brief Survey proposed. It is first proposed to give a brief survey of the development of algebra, recalling the names of those who helped to set the problems that were later solved by the aid of equations, as well as those who assisted in establishing the science itself. These names have been mentioned in Volume I and some of them will be referred to when we consider the development of the special topics of algebra and their application to the solution of elementary problems.

[1] For a brief study of the early history see H. G. Zeuthen, "Sur l'origine de l'Algèbre," in the *Kgl. Danske Videnskabernes Selskab, Math.-fysiske Meddelelser*, II, 4, Copenhagen, 1919; M. Chasles, "Histoire de l'Algèbre," *Comptes rendus*, September 6, 1841; but the subject is treated of in any general history of mathematics and in the leading encyclopedias.

It should also be stated as a preliminary to this discussion that Nesselmann[1] (1842) has divided the history of algebra into three periods: the rhetorical, in which the words were written out in full; the syncopated, in which abbreviations were used; and the symbolic, in which the abbreviations gave place to such symbols as occur in statements like $\sqrt{x} - x^2 = a^{\frac{2}{3}}$. There are no exact lines of demarcation by which to establish these divisions, Diophantus, for example, having made use of certain features of all three; but the classification has some advantages and the student will occasionally find the terms convenient.

It should be borne in mind that most ancient writers outside of Greece included in their mathematical works a wide range of subjects. Ahmes (*c.* 1550 B.C.),[2] for example, combines his algebra with arithmetic and mensuration, and even shows some evidence that trigonometry was making a feeble start. There was no distinct treatise on algebra before the time of Diophantus (*c.* 275).

Algebra in Egypt. The first writer on algebra whose works have come down to us is Ahmes. He has certain problems in linear equations and in series, and these form the essentially new feature in his work. His treatment of the subject is largely rhetorical, although, as we shall see later, he made use of a small number of symbols.

There are several other references to what may be called algebra in the Egyptian papyri, these references consisting merely of problems involving linear or quadratic equations. There is no good symbolism in any of this work and no evidence that algebra existed as a science.

Algebra in India. There are only four Hindu writers on algebra whose names are particularly noteworthy. These are Āryabhaṭa,[3] whose *Āryabhaṭiyam* (*c.* 510) included problems in

[1] G. H. F. Nesselmann, *Alg. Griechen*, p. 302.

[2] As already stated, the period may have been *c.* 1600 B.C. or earlier.

[3] See Volume I, page 153, and remember that there were two Āryabhaṭas and that we are not certain which one of them is entitled to the credit for various contributions.

series, permutations, and linear and quadratic equations; Brahmagupta, whose *Brahmasiddhānta* (*c.* 628) contains a satisfactory rule for solving the quadratic, and whose problems include the subjects treated by Āryabhaṭa; Mahāvīra, whose *Gaṇita-Sāra Sangraha* (*c.* 850) contains a large number of problems involving series, radicals, and equations; and Bhāskara, whose *Bija Gaṇita* (*c.* 1150) contains nine chapters and extends the work through quadratic equations.[1]

Algebra in China. It is difficult to say when algebra as a science began in China. Problems which we should solve by equations appear in works as early as the *Nine Sections*[2] and so may have been known by the year 1000 B.C. In Liu Hui's commentary on this work (*c.* 250) there are problems of pursuit, the Rule of False Position, explained later in this chapter, and an arrangement of terms in a kind of determinant notation.[3] The rules given by Liu Hui form a kind of rhetorical algebra.

The work of Sun-tzï[4] (perhaps of the 1st century, but the date is very uncertain and may be several centuries earlier) contains various problems which would today be considered algebraic. These include questions involving indeterminate equations of which the following is a type:

> There are certain things whose number is unknown. If they are divided by 3 the remainder is 2; by 5, the remainder is 3; and by 7, the remainder is 2. Find the number.

Sun-tzï solved such problems by analysis and was content with a single result, even where several results are admissible.

The Chinese certainly knew how to solve quadratics as early as the 1st century B.C., and rules given even as early as the *K'iu-ch'ang Suan-shu* above mentioned involve the solution of such equations.

[1] H. T. Colebrooke, *Algebra with Arithmetic and Mensuration, from the Sanscrit*, pp. 129–276 (London, 1817). For the various spellings of *Bija Gaṇita* see Volume I, page 278.

[2] *K'iu-ch'ang Suan-shu.*

[3] Mikami, *China*, pp. 19, 23.

[4] *Sun-tzï Suan-king.*

Liu Hui (*c.* 250) gave various rules which would now be stated as algebraic formulas and seems to have deduced these from other rules in much the same way as we should deduce formulas at the present time.[1]

By the 7th century the cubic equation had begun to attract attention, as is evident from the *Ch'i-ku Suan-king* of Wang Hs'iao-t'ung (*c.* 625).

The culmination of Chinese algebra is found in the 13th century. At this time numerical higher equations attracted the special attention of scholars like Ch'in Kiu-shao (*c.* 1250), Li Yeh (*c.* 1250), and Chu Shï-kié (*c.* 1300),[2] the result being the perfecting of an ancient method which resembles the one later developed by W. G. Horner (1819).

With the coming of the Jesuits in the 16th century, and the consequent introduction of Western science, China lost interest in her native algebra and never fully regained it.

Algebra in Greece. Algebra in the modern sense can hardly be said to have existed in the golden age of Greek mathematics.[3] The Greeks of the classical period could solve many algebraic problems of considerable difficulty, but the solutions were all geometric. Hippocrates (*c.* 460 B.C.), for example, assumed a construction which is equivalent to solving the equation $x^2 + \sqrt{\frac{3}{2}} \cdot ax = a^2$, and Euclid (*c.* 300 B.C.), in his *Data*, solved problems equivalent to the following:

1. $xy = k^2$, $x - y = a$ (Prob. 84).
2. $xy = k^2$, $x + y = a$ (Prob. 85).
3. $xy = k^2$, $x^2 - y^2 = a^2$ (Prob. 86).

In his *Elements* (II, 11) Euclid solved the equivalent of $x^2 + ax = a$, and even of $x^2 + ax = b^2$, substantially by completing the geometric square and neglecting negative roots.

After Euclid there came a transition period from the geometric to the analytic method. Heron (*c.* 50?), who certainly

[1] Mikami, *China*, pp. 35, 36. [2] Mikami, *China*, pp. 63, 79, 89.

[3] Nesselmann, *Alg. Griechen*; Heath, *Diophantus*. On the "application of areas" see Heath, *History*, and R. W. Livingstone, *The Legacy of Greece*, p. 111 (Oxford, 1922).

solved the equation $144\,x\,(14 - x) = 6720$, may possibly have used the analytic method for the purpose of finding the roots of


$$\tfrac{1}{14}\tfrac{1}{4}\,x^2 + \tfrac{2}{7}\tfrac{9}{}\,x = 212.$$

With Diophantus (*c.* 275) there first enters an algebraic symbolism worthy of the name, and also a series of purely algebraic problems treated by analytic methods. Many of his equations being indeterminate, equations of this type are often called Diophantine Equations. His was the first work devoted chiefly to algebra, and on this account he is often, and with much justice, called the father of the science.

Algebra among the Arabs and Persians. The algebraists of special prominence among the Arabs and Persians were Mohammed ibn Musa, al-Khowârizmî, whose *al-jabr w'al muqâbalah* (*c.* 825) gave the name to the science and contained the first systematic treatment of the general subject as distinct from the theory of numbers; Almâhânî (*c.* 860), whose name will be mentioned in connection with the cubic; Abû Kâmil (*c.* 900), who drew extensively from al-Khowârizmî, and from whom Fibonacci (1202) drew in turn; al-Karkhî (*c.* 1020), whose *Fakhrî* contains various problems which still form part of the general stock material of algebra; and Omar Khayyam (*c.* 1100), whose algebra was the best that the Persian writers produced.

Medieval Writers. Most of the medieval Western scholars who helped in the progress of algebra were translators from the Arabic. Among these were Johannes Hispalensis (*c.* 1140), who may have translated al-Khowârizmî's algebra; Gherardo of Cremona (*c.* 1150), to whom is also attributed a translation of the same work; Adelard of Bath (*c.* 1120), who probably translated an astronomical work of al-Khowârizmî, and who certainly helped to make this writer known; and Robert of Chester, whose translation of al-Khowârizmî's algebra is now available in English.[1]

[1] L. C. Karpinski, *Robert of Chester's Latin Translation . . . of al-Khowarizmi.* New York, 1915.



FIRST PAGE OF AL-KHOWÂRIZMÎ'S ALGEBRA

From a MS. of 1456. It begins, "Liber mahucmeti de Algebra et almuchabala." In Mr. Plimpton's library

The greatest writer on algebra in the Middle Ages was Fibonacci, whose *Liber Quadratorum* (*c.* 1225) and *Flos* both relate to the subject. The former work includes the treatment of such problems as $x^2 + y^2 = z^2$ and other well-known types, and shows great ingenuity in the solution of equations.

Of the German algebraists in the Middle Ages the leading writer was Jordanus Nemorarius (*c.* 1225). His *De Numeris Datis*, already described (Vol. I, p. 227), contains a number of problems in linear and quadratic equations of the type still familiar in our textbooks. In general, however, the medieval writers were more interested in mathematics as related to astronomy than in mathematics for its own sake.[1]

The Renaissance. Algebra in the Renaissance period received its first serious consideration in Pacioli's *Sūma* (1494), a work which summarized in a careless way the knowledge of the subject thus far accumulated. By the aid of the crude symbolism then in use it gave a considerable amount of work in equations.

The next noteworthy work on algebra, and the first to be devoted entirely to the subject, was Rudolff's *Coss* (1525). This work made no decided advance in the theory, but it improved the symbolism for radicals and made the science better known in Germany. Stifel's edition of this work (1553–1554) gave the subject still more prominence.

The first epoch-making algebra to appear in print was the *Ars Magna* of Cardan (1545). This was devoted primarily to the solution of algebraic equations. It contained the solution of the cubic and biquadratic equations,[2] made use of complex numbers, and in general may be said to have been the first step toward modern algebra.

The next great work on algebra to appear in print was the *General Trattato* of Tartaglia (1556–1560), although his side of the controversy with Cardan over the solution of the cubic equation had already been given in his *Qvesiti ed invenzioni diverse* (1546).

[1] On the general topic see P. Cossali, *Origine, trasporto in Italia, primi progressi in essa dell' Algebra*, 2 vols., Parma, 1797–1799.

[2] In chapters xi and xxxix et seq.

Algebra in the New World. As already stated,[1] the first mathematical work published in the New World was the *Sumario Compendioso* of Juan Diez. This appeared in the City of Mexico in 1556 and contains six pages on algebra. Some idea of its general nature may be obtained from two of the problems relating to the subject. Of these the first, literally translated and requiring the solution of the quadratic equation $x^2 - 15\frac{3}{4} = x$, is as follows:

Find a square from which if $15\frac{3}{4}$ is subtracted the result is its own root.

Let the number be *cosa* [x]. The square of half a *cosa* is equal to $\frac{1}{4}$ of a *zenso* [x^2]. Adding 15 and $\frac{3}{4}$ to $\frac{1}{4}$ makes 16, of which the root is 4, and this plus $\frac{1}{2}$ is the root of the required number.

Proof: Square the square root of 16 plus half a *cosa*, which is four and a half, giving 20 and $\frac{1}{4}$, which is the square number required. From $20\frac{1}{4}$ subtract 15 and $\frac{3}{4}$ and you have 4 and $\frac{1}{2}$, which is the root of the number itself.

The second problem, also literally translated, requires the solution of the quadratic equation $x^2 + x = 1260$:

A man takes passage in a ship and asks the master what he has to pay. The master says that it will not be any more than for the others. The passenger on again asking how much it would be, the master replies: "It will be the number of pesos which, multiplied by itself and added to the number, gives 1260." Required to know how much the master asked.

Let the cost be a *cosa* of pesos. Then half of a *cosa* squared makes $\frac{1}{4}$ of a *zenso*, and this added to 1260 makes 1260 and a quarter, the root of which less $\frac{1}{2}$ of a *cosa* is the number required. Reduce 1260 and $\frac{1}{4}$ to fourths; this is equal to $\frac{5041}{4}$, the root of which is 71 halves; subtract from it half a *cosa* and there remains 70 halves, which is equal to 35 pesos, and this is what was asked for the passage.

Proof: Multiply 35 by itself and you have 1225; adding to it 35, you have 1260, the required number.

[1] See Volume I, page 353. D. E. Smith, *The Sumario Compendioso of Brother Juan Diez*, Boston, 1921.

First Teachable Textbooks in Algebra. The first noteworthy attempt to write an algebra in England was made by Robert Recorde, whose *Whetstone of witte* (1557) was an excellent textbook for its time. The next important contribution was Masterson's incomplete treatise of 1592–1595, but the work was not up to the standard set by Recorde.

The first Italian textbook to bear the title of algebra was Bombelli's work of 1572. In this book the material is arranged with some attention to the teaching of the subject.[1]

By this time elementary algebra was fairly well perfected, and it only remained to develop a good symbolism. As will be shown later, this symbolism was worked out largely by Vieta (*c.* 1590), Harriot (*c.* 1610), Oughtred (*c.* 1628), Descartes (1637), and the British school of Newton's time (*c.* 1675).

So far as the great body of elementary algebra is concerned, therefore, it was completed in the 17th century.

2. Name for Algebra

Early Names. The history of a few of the most familiar terms of algebra not elsewhere discussed will now be considered, and of these the first is naturally the name of the science itself.

Ahmes (*c.* 1550 B.C.) called his treatise "Rules for inquiring into nature, and for knowing all that exists, [every] mystery, . . . every secret,"[2] and this idea is not infrequently expressed by later writers. Thus Seki (*c.* 1680) called a certain part of algebra the *kigen seihō*, meaning a method for revealing the true and buried origin of things, and we find the same idea in the titles of algebras by Follinus (1622)[3] and Gosselin (1577),[4] and in a note on Ramus written by Schoner in 1586.[5]

[1] *L'Algebra parte maggiore dell' arimetica* . . . , Bologna, 1572. There is a second edition, differing only in the title-page, Bologna, 1579.

[2] He adds: "Behold, this roll was written . . . [under . . . the King of Upper] and Lower Egypt, Aauserrē. . . . It was the scribe Aḥmōse who wrote this copy." Peet, *Rhind Papyrus*, p. 33. Professor Peet gives the probable date as between 1788 and 1580 B.C.

[3] *Algebra sive liber de rebus occultis.*

[4] . . . *de occulta parte numerorum.*

[5] " . . . Almucabalam, hoc est, librum de rebus occultis" (p. 322).

Since the Greeks gave the name "arithmetic" to all the theory of numbers, they naturally included their algebra under that title,[1] and this explains why the algebra of Diophantus went by the name of arithmetic.

The Hindu writers had no uniform name for the science. Āryabhaṭa (*c.* 510) included algebra in his general treatise, the *Āryabhaṭiyam*; Brahmagupta (*c.* 628) placed it in his large treatise, giving a special name (*kuṭaka*, the pulverizer)[2] to his chapter on indeterminate equations. Mahāvīra (*c.* 850) included it in his *Gaṇita-Sāra-Sangraha*, a title meaning a brief exposition of the compendium of calculation. Bhāskara (*c.* 1150) had a name for general arithmetic, *Bija Gaṇita*,[3] meaning the calculation of seeds, that is, of original or primary elements,[4] and a special name for algebra, *Avyakta gaṇita*,[5] or *Avyakta-kriyā*, the former referring to the calculation with knowns and the latter to that with unknowns.

The Chinese used various fanciful titles for their books containing algebra and spoke of the method of the *t'ien-yuen* (celestial element),[6] meaning the algebra that made use of calculating rods (the Japanese *sangi*), to indicate coefficients.[7] Similar fanciful names were used in Japan, as also the name *yendan jutsu* (method of analysis), and the name *kigen seihō* already mentioned.

[1] So Euclid's *Elements*, II, devoted to arithmetic, includes a considerable part of algebra, such as the geometric proofs about $(a \pm b)^2$ and $(a + b)(a - b)$. The fact was recognized by Ramus (1569) when he stated that "Algebra est pars arithmeticae" (1586 ed., p. 322).

[2] Colebrooke (pp. 112, 325) transliterates this as *cuṭṭácára*, *cuṭṭa*, *cuṭṭaca*, *cuṭṭaca-vyavahára*, and *cuṭṭacád'hyára*, meaning the determination of a pulverizing multiplier p such that, if n_1, n_2, and n_3 are given numbers, then $pn_1 + n_2$ shall be divisible by n_3.

[3] Or *Vija Ganita*, *Beej Gunnit*. The spelling as given in the first printed edition (Calcutta, 1846) is *Beej Gunita*. The name *Vija-kriyā*, meaning seed analysis, is also used.

[4] Sir M. Monier-Williams, *Indian Wisdom*, 4th ed., p. 174 (London, 1893).

[5] Nesselmann, *loc. cit.*, p. 44; Colebrooke translation, p. 129. Bhāskara also used *sama-sodanam* (transposition) to include the two terms which had been used by al-Khowârizmî.

[6] *T'ien-yuen-shu*, celestial-element method. The Japanese called it *tengen jutsu*.

[7] Mikami, *China*, p. 157.

Algebra at one time stood a fair chance of being called *Fakhrî*, since this was the name given to the work of al-Karkhî (*c.* 1020), one of the greatest of the Arab mathematicians. Had his work been translated into Latin, as al-Khowârizmî's was, the title might easily have caught the fancy of the European world. Al-Karkhî relates that he was long and sorely hindered in his attempts to complete his work, because of the tyranny and violence endured by the people, until "God, may his name be hallowed and exalted, sent to their aid our protector, the vizir, the illustrious lord, the perfect one in government, the vizir of vizirs, clothed with double authority, Abû Gâlib," whose familiar name was Fakhr al-Mulk. In honor of this patron the name Fakhr gave rise to the title of the book, *al-Fakhrî*.

The Name "Algebra." Our real interest in the name centers around the word *algebra*, a word appearing, as we have seen, in the title of one of the works by al-Khowârizmî (*c.* 825),—*al-jabr w'al-muqâbalah*. It also appears in the early Latin translations under such titles as *Ludus algebrae almucgrabalaeque* and *Gleba mutabilia*. In the 16th century it is found in English as *algiebar and almachabel*, and in various other forms, but was finally shortened to *algebra*.[1] The words mean restoration and opposition,[2] and one of the clearest explanations of their use is given by Behâ Eddîn (*c.* 1600) in his *Kholâsat al-Ḥisâb* (*Essence of Arithmetic*):

The member which is affected by a minus sign will be increased and the same added to the other member, this being *algebra*; the homogeneous and equal terms will then be canceled, this being *al-muqâbala*.

That is, given $bx + 2q = x^2 + bx - q$,
al-jabr gives $bx + 2q + q = x^2 + bx$,
and *al-muqâbalah* gives $3q = x^2$.

[1] An English translation of al-Khowârizmî's work by F. Rosen appeared in London in 1831. A Latin version was published by Libri in his *Histoire*, Vol. I (Paris, 1835), and by Karpinski (1915) from a Scheubel (Scheybl) MS. at Columbia University, as already stated.

[2] Or redintegration and equation. *Jabr* is from *jabara* (to reunite or consolidate), possibly allied to the Hebrew *gābar* (make strong).

This statement was put into verse, as was usual in the East, and thus became generally known in the Arab schools. It may be crudely translated thus:

Cancel minus terms and then
Restore to make your *algebra*;
Combine your homogeneous terms
And this is called *muqâbalah*.[1]

In a general way we may say that *al-jabr* or *al-jebr* has as the fundamental idea the transposition of a negative quantity, and *muqâbalah* the transposition of a positive quantity and the simplification of each member.[2] Al-Khowârizmî's title was adopted by European scholars,[3] appearing both in the Arabic, with many curious variants, and in Latin. The Moors took the word *al-jabr* into Spain, an *algebrista* being a restorer, one who resets broken bones.[4] At one time it was not unusual to see over the entrance to a barber shop the words "Algebrista y Sangrador" (bonesetter and bloodletter), and both the striped pole which is used in America as a barber's sign and the metal basin used for the same purpose in Europe today are relics of the latter phase of the haircutter's work. From Spain the word passed over to Italy, where, in the 16th century, *algebra* was used to mean the art of bonesetting.[5] Thence it found its way into France as *algèbre*, and so on to England, where one writer (1541) speaks of "the helpes of Algebra & dislocations,"

[1] From a Persian algebra written probably after the 12th century. Nesselmann (p. 50) put it into German verse, and the above English quatrain, taken from his translation, gives only a general idea of the wording.

[2] Rollandus (*c.* 1424) has *De arte dolandi*, the art of chipping off or cutting with an ax, probably meaning the chipping off or subtracting of equals from both members.

[3] Thus Robert of Chester's translation (*c.* 1140) begins his *Liber Algebrae et Almucabola* thus: "In nomine Dei χρij et misericordis incipit liber Restaurationis et Oppositionis numeri . . . filius Mosi Algoŭrizim dixit Mahometh."

[4] So in *Don Quixote* (II, chap. 15), where mention is made of "*un algebrista* who attended to the luckless Samson."

[5] Libri, *Histoire*, 1838 ed., II, 80. The question of the connection of *al-jabr* with the Hebrew root *sh-b-r* (from which comes *tishboreth*, fracture), and with the Hindu word for pulverizer, is worthy of study.

and another (1561) says: "This Araby worde Algebra sygnifyeth as well fractures of the bones, etc. as sometyme the restauration of the same."[1]

As already said, the name was much distorted by the Latin translators. Thus Guglielmo de Lunis (*c.* 1250?) gives it as *gleba mutabilia*, and Roger Bacon (*c.* 1250) speaks of the science as *algebra . . . et almochabala*.[2] A 15th century manuscript testifies both to the mystery of the subject and to the uncertainty of name when it speaks of the subtleties of *largibra*.[3] In the early printed books it appeared in equally curious forms, such as *Gebra vnd Almuthabola*.[4]

Some of the late Latin writers attributed the name to one Geber,[5] an Arab philosopher, whom they supposed to be the inventor of the science;[6] and certain Arab writers speak of a Hindu named Argebahr or Arjabahr, a name which may have influenced the Latin translators.[7] Even as good a scholar as Schoner went far astray in his interpretation of the title.[8]

[1] See Oxford Dictionary under *algebra*.

[2] "Algebra quae est negotiatio, et almochabala quae est census." *Opus Majus*, ed. Bridges, I, p. lvii.

[3] Anon. MS. in Boncompagni's library: " . . . di subtili . & . di largibra." Narducci *Catalogo* (2d ed., 1892), No. 397 (2).

[4] A. Helmreich, *Rechenbüch*, 1561 (1588 ed.).


[5] There was an Arab scholar, Jabir ibn Aflah, Abû Mohammed, of Seville (*c.* 1145), whose astronomy was translated by Gherardo of Cremona, his contemporary, and was printed in 1534.

[6] The name appears as *Greber* in Heilbronner's *Hist. Math.*, p. 340 (1742). Ghaligai (1521) spoke of it as "composta da uno home Arabo di grãde intelligentia," adding that "alcuni dicono essere stato uno il qual nome era Geber."

[7] Libri, *Histoire* (I, 122), thinks that this writer was Āryabhaṭa.

[8] Thus in a note on Ramus (1586 ed., p. 322) he says: "Nomen Algebrae Syriacum putatur significans artem & doctrinam hominis excellentis. Nam *geber* Syris significat virum . . . ut apud nos Magister aut Doctor . . . & ab Indis harum artium perstudiosis dicitur *Aliabra* item *Alboret*, tametsi proprium autoris nomen ignoretur."

As an example of still more uncertain history, A. Helmreich (*Rechenbüch*, 1561; 1588 ed., fol. b 2, *r.*, of the *Vorrede*) asserts that algebra was due to "Ylem/ der grosse Geometer in Egypten/zur zeit desz Alexandri Magni, der da war ein Praeceptor oder vorfahrer Euclidis, desz Fürsten zu Megarien." We also find such forms as *Agabar, Algebra muchabila, Reghola della raibre mochabilr^a, regola del acabrem^a, ghabile, dellacibra e muchabile, lacibra umachabille*, all in MSS. of the 15th century; and in the 16th century such forms as *arcibra*.



FIRST PAGE OF AN ALGEBRA MANUSCRIPT OF *C.* 1460

Possibly by Raffaele Canacci, a Florentine mathematician. On the second line may be seen the name for algebra,—*Algebra amucabale.* In Mr. Plimpton's library

Other Names. Because the unknown quantity was called *res* by the late Latin writers, which was translated into Italian as *cosa*,[1] the early Italian writers called algebra the *Regola de la Cosa*, whence the German *Die Coss* and the English *cossike arte*.[2]

The Italians of the 15th and 16th centuries often called algebra the greater art, to distinguish it from commercial arithmetic, which was the lesser art, just as we speak of higher arithmetic and elementary arithmetic. This distinction may have been suggested by the seven *arti maggiori* and the fourteen *arti minori* recognized by the merchants of medieval Florence.[3] Thus we have such names as *Ars Magna*, used by Cardan (1545),[4] *l'Arte Maggiore*, used by various other Italian writers, and *l'arte mayor*, used by Juan Diez, whose book has been mentioned as having appeared in Mexico in 1556. The title as given in the Mexican book is as follows:

Quiſtiones del arte mayor tocantes al algebra.

Vieta (*c.* 1590) rejected the name "algebra" as having no significance in the European languages, and proposed to use the term "analysis," and it is probably to his influence that the popularity of this term in connection with higher algebra is due.

Of the other names for algebra the only one that we need consider is "logistic." Since the term had dropped out of use as a name for computation about 1500, it was employed to

[1] From the Latin *causa*; compare the French *chose*.

[2] Thus Pacioli (1494 ed., fol. 67, *r.*): "Per loperare de larte magiore: ditta dal vulgo la regola de la cosa ouer algebra e amucabala." It will be recalled that Rudolff's title for his book (1525) on algebra was *Die Coss*. Helmreich says: " . . . vnd wird bey den Welschen [the Italians] genent das buch *Delacosa*/welchs wir Deutschē die *Regulā Cos* oder *Algebra* nennen." *Rechenbüch*, 1561 (1588 ed.), fol. b 2.

[3] E. G. Gardner, *The Story of Florence*, p. 42. London, 1900.

[4] Also by Gosselin, *De arte magna, seu de occulta parte numerorum quae & Algebra & Almucabala vulgo dicitur, libri IV*, Paris, 1577. The name was used, however, for various purposes, as in Kircher's *Ars Magna Lucis et Umbrae*, Amsterdam, 1671, and numerous other works.

designate a higher branch, just as "calculus" was appropriated a century or so later. Thus we find it used by Buteo and others[1] to cover advanced arithmetic and algebra, although it never became popular.[2]

3. Technical Terms

Coefficient. Of the terms commonly used in algebra, it is possible at this time to mention only a few typical ones.

The coefficient was called by Diophantus (*c.* 275) the *ple'thos*[3] (multitude), and by Brahmagupta (*c.* 628) the *anca*, or *prakriti*,[4] but most early writers used no special name. The term "coefficient" and the use of literal coefficients are late developments, the former being due to Vieta.

Unknown Quantity. The unknown quantity was called by Ahmes (*c.* 1550 B.C.) *ahe*[5] or *hau* ("mass," "quantity," or "heap"). Diophantus called it "an undefined number of units."[6] Brahmagupta called it the *yāvat-tāvat*,[7] and possibly this suggested to the Arabs the use of *shā* (*śei, chaï*, meaning "thing" or "anything"), whence the medieval use of *res* (thing) for this purpose.[8] The Chinese used *yuen* (element),[9] as already stated, but they also used a word meaning "thing."[10]

Powers. Before the invention of a satisfactory symbol like x^2 it became necessary to have a special name for the square of the unknown, and in the Greek geometric algebra it was called

[1] H. Vitalis, *Lexicon Mathematicvm*, p. 25 (Rome, 1690); Nesselmann, *loc. cit.*, p. 57.

[2] There are many other names used for this purpose. *E.g.*, 'Alî ibn Velî ibn Hamza, a western Arab, then (1590/91) living in Mecca, wrote a work entitled *Tuḥfet al-a'dad li-davî al-roshd ve'l sadâd* (*The Gift of Numbers for the Possessors of Reason and Correct Insight*), relating to elementary algebra. See E. L. W. M. Curtze, *Abhandlungen*, XIV, 184. The name reminds one of Recorde's *Whetstone of witte*, London, 1557.

[3] Πλῆθος. [4] Colebrooke's translation, pp. 246, 348.

[5] By Egyptologists, *ḥ'w*. See Peet, *Rhind Papyrus*, p. 61.

[6] Πλῆθος μονάδων ἄλογον (*ple'thos mona'don a'logon*).

[7] Also given as *jabut tabut*, literally the "so much as," "as far, so far," "as much, so much," or "however much." Compare Bombelli's use of *tanto* (so much).

[8] The Arabs also used *jidr* (*dyizr*, root), whence the Latin *radix*.

[9] Mikami, *China*, pp. 81, 91. [10] Compare *res* and *cosa*, page 392.

a "tetragon[1] number" or a "power."[2] Diophantus called the third power a cube,[3] the fourth power a "power-power,"[4] the fifth power a "power-cube,"[5] and the sixth power a "cube-cube,"[6] using the additive instead of the multiplicative principle.

The Arab writers called the square of the unknown a *mal*, a word meaning "wealth," whence the medieval Latin *census* (evaluation of wealth, tax) was used for the same purpose, appearing in the early Italian algebras as *censo*, sometimes incorrectly written as *zenso*. Therefore algebra was not uncommonly called *Ars rei et census* as well as *Ars rei*[7] and *Regola o l'arte della cosa.*

Equation. The word "equation," while generally used as at present ever since the medieval writers set the standard, has not always had this meaning. It is used by Ramus in his arithmetic and by his commentator Schoner to denote a continued proportion, although in speaking of algebra Ramus (1567)[8] and Gosselin (1577) follow the ordinary usage.

Absolute Term. In the equation

$$a_0x^n + a_1x^{n-1} + \cdots + a_{n-1}x + a_n = 0$$

we speak of a_n as the absolute term. There have been various names for it, Diophantus (*c.* 275) calling it monads.[9] The

[1] Τετράγωνος ἀριθμός (*tetra'gonos arithmos'*, four-angled number). So in Euclid VII, def. 18.

[2] Δύναμις (*dy'namis*), from the same root as "dynamo," "dynamic," and "dynamite." So in Plato (*Timæus*, 31); but he also uses the term (*Theætetus*, 147 D) to mean the square root of a non-square number. When Diophantus speaks of any particular square number, he uses τετράγωνος ἀριθμός, but otherwise δύναμις. Heath, *Diophantus*, 2d ed., p. 38.

[3] Κύβος (*ku'bos*).

[4] Δυναμοδύναμις (*dynamody'namis*).

[5] Δυναμόκυβος (*dynamo'kubos*).

[6] Κυβόκυβος (*kubo'kubos*).

[7] Nesselmann, *Alg. Griechen*, pp. 55, 56.

[8] Thus Ramus, in his arithmetic (1567), says: "AEquatio est quando continuatae rationes continuantur iterum," Schoner giving as an example

10.	15.	12.
20.	30.	24.

meaning 10 : 15 : 12 = 20 : 30 : 24 (1586 ed., p. 188).

[9] Μονάδες (*mona'des*), with the abbreviation μ°. Heath, *Diophantus*, 2d ed., p. 39.

Hindus called it *rupa* or *ru*,[1] and the Chinese gave it the name *tai*,[2] an abbreviation of *tai-kieh* (extreme limit).

Commutative and Distributive Laws. The use of the terms "commutative" and "distributive" in the usual algebraic sense is due to the French mathematician Servois (1814). The use of the term "associative" in this sense is due to Sir William Rowan Hamilton.[3]

4. Symbols of Algebra

Symbols of Operation. The symbols of elementary arithmetic are almost wholly algebraic, most of them being transferred to the numerical field only in the 19th century,[4] partly to aid the printer in setting up a page and partly because of the educational fashion then dominant of demanding a written analysis for every problem. When we study the genesis and development of the algebraic symbols of operation, therefore, we include the study of the symbols used in arithmetic. Some idea of the status of the latter subject in this respect may be obtained by looking at almost any of the textbooks of the 17th and 18th centuries. Hodder,[5] for example, gives no symbols before page 201,[6] then remarking: "Note that a + thus, doth signifie Addition, and two lines thus $\equiv$ Equality, or Equation, but a $\times$ thus, Multiplication," no other symbols being used. Even Recorde, who invented the modern sign of equality, did not use it in his arithmetic, the *Ground of Artes* (*c.* 1542), but only in his algebra, the *Whetstone of witte* (1557).

Earliest Symbols. The earliest symbols of operation that have come down to us[7] are Egyptian. In the Ahmes Papyrus (*c.* 1550 B.C.) addition and subtraction are indicated by

[1] Colebrooke's *Vīja Ganita*, p. 186 n.; E. Strachey, *Bija Ganita*, p. 117 (London, n.d.), where it is given as *roop*.

[2] Mikami, *loc. cit.*, p. 81. The word is also transliterated *taë*.

[3] Cajori, *Hist. of Math.*, 2d ed., p. 273. New York, 1919.

[4] There are, of course, exceptions. The Greenwood arithmetic (1729), for example, used the algebraic symbols.

[5] His was the first English arithmetic to be reprinted in the American colonies (Boston, 1719).

[6] 1672 ed.

[7] Excepting those connected with notation, as in the subtractive principle of the Babylonians, already mentioned.

special symbols, but these are simply hieratic forms from the hieroglyphics and are not symbols in the sense in which we use the term. The symbol was used to designate addition. It appears in the Ahmes Papyrus as  .[1] The symbol  was used to designate subtraction. It appears in the Ahmes Papyrus as .[2]

Diophantus (*c.* 275) represented addition by simple juxtaposition, as in $K^{\upsilon}\bar{\alpha}\Delta^{\upsilon}\overline{\iota\varsigma}$ for $x^3 + 13x^2$. For subtraction he seems to have used the symbol λ, although we are not certain as to its precise form. Since we have no manuscript of his *Arithmetica* earlier than the Madrid copy of the 13th century, we are also uncertain as to the authenticity of the following passage:

"Minus multiplied by minus makes plus, and minus by plus makes minus. The sign of negation is ψ turned upside down, λ."[3] It is fully as probable that the symbol is a deformed Λ (*lambda*), the Greek letter L and the initial for a word indicating subtraction.[4]

The Hindus at one time used a cross placed beside a number to indicate a negative quantity, as in the Bakhshāli manuscript of possibly the 10th century. With this exception it was not until the 12th century that they made much use of symbols of operation.[5] In the manuscripts of Bhāskara (*c.* 1150) a small circle or a dot is placed above a subtrahend, as in $\overset{\circ}{6}$ or $\dot{6}$ for -6,[6] or the subtrahend is inclosed in a circle, just as children, in scoring a game, indicate 6 less than zero[7] by the symbol ⊛

[1] Ahmes wrote, as usual, from right to left, but hieroglyphic sentences are generally printed from left to right. See the Brit. Mus. facsimile, Pl. IX, row 5.

[2] Peet, *Rhind Papyrus*, p. 64; Eisenlohr, *Ahmes Papyrus*, p. 47.

[3] Tannery, *Diophantus*, I, 13, and *Bibl. Math.*, V (3), 5; Heath, *Diophantus*, 2d ed., p. 130. On the relation of this symbol to the symbol $\frown$, which is used in the Ayer Papyrus (*c.* 200–400), see *Amer. Journ. of Philology*, XIX, 25.

[4] Λειπόντες (*lipon'tes*, diminished by), or Λείπειν (*lei'pein*, to be missing).

Of course a further exception is also to be made of the representing of sums by juxtaposition and of division by means of fractions. For the Bakhshāli MS. see Volume I, page 164.

[6] See Colebrooke's edition, p. 131; Taylor's edition, Introduction, p. 11.

[7] C. I. Gerhardt, *Études historiques*, p. 8. Berlin, 1856.

European Symbols for Plus and Minus. The early European symbol for plus, used in connection with the Rule of False Position, was p,[1] P,[2] or $\bar{p}$, the last being the most common of the three.[3] The word *plus*, used in connection both with addition and with the Rule of False Position, was also employed; but, strange to say, it is much later than the word *minus* as indicating an operation. The latter is found in the works of Fibonacci (1202), while the use of *plus* to indicate addition is not known before the latter part of the 15th century.[4]

Since $\bar{p}$ or $\tilde{p}$ was used for *plus*, $\bar{m}$ or $\tilde{m}$ was naturally used for *minus*, and this usage is found in many works of the 15th and 16th centuries. As usual, the bar simply indicated an omission, as in *Sūma* for *Summa*, in the title of Pacioli's work.[5] In the 15th century the symbol $\overline{19}$ was often used for *minus*,[6] but most writers preferred the $\bar{m}$.

Racial Preferences. We now come to one of the many cases of racial habit in determining mathematical custom. In the 16th century the Latin races generally followed the Italian School, using $\bar{p}$ and $\bar{m}$ or their equivalents,[7] while the German School

[1] So in the Rollandus MS. (*c.* 1424), where the terms are so arranged as to require no minus sign. The Rule of False Position is explained on pages 437–441.

[2] Clavius (1583) uses P and M for *plus* and *minus* in his Rule of False Position, which he gives in his arithmetic; but in his algebra (1608) he uses the cross, saying: "Plerique auctores pro signo + ponunt literam P, . . . sed placet nobis uti nostris signis." It must be remembered that the symbols in the Rule of False Position are hardly symbols of operation in the ordinary sense of the term.

[3] *E.g.*, Chuquet (1484) and Pacioli (1494), the latter first using it in his Rule of False Position (fol. 106).

[4] *Bibl. Math.*, XIII (2), p. 105. Fibonacci used it, however, in the Rule of False Position.

[5] Compare the French *hôtel* for *hostel* or *hospital*, and the German *über* for *ueber*. See also Cajori, "Varieties of Minus Signs," *Math. Teacher*, XVI, 295.

[6] This is the case in Mr. Plimpton's MS. of al-Khowârizmî, written in 1456. See also the Regiomontanus-Bianchini correspondence (*c.* 1464) in the *Abhandlungen*, XII, 233, 279; and Curtze in the *Bibl. Math.*, I (3), 506.

[7] *E.g.*, Pacioli (1494) writes $\bar{m}$ for *minus* in his algebra, but he follows the general custom of using *de* in cases like "7. de 9"; Cardan (1539) writes R . V . 7 . p R . 4 for $\sqrt{7+\sqrt{4}}$; Feliciano Lazesio (1526) writes 6. piu. R . 16 for $6+16x$, and 10. $\bar{m}$R . 4 for $10-4x$; Tartaglia (1556) and Cataneo (1546) write *piu* and *men* for *plus* and *minus* in the Rule of False Position; Santa-Cruz (1594) uses the equivalent Spanish words *mas* and *menos*; and Peletier

preferred the symbols + and −, neither of which is found for this purpose, however, before the 15th century.[1]

Origin of our Plus and Minus Signs. In a manuscript of 1456, written in Germany,[2] the word *et* is used for addition and is generally written so that it closely resembles the symbol +. The *et* is also found in many other manuscripts, as in "5 et 7" for 5 + 7, written in the same contracted form,[3] as when we write the ligature & rapidly. There seems, therefore, little doubt that this sign is merely a ligature for *et*.

The origin of the minus sign has been more of a subject of dispute. Some have thought that it is a survival of the bar in $\overline{19}$ or in $\bar{m}$, but it is more probable that it comes from the habit of early scribes of using it as the equivalent of *m*, as in Sūma for Summa. Indeed, it is quite probable that the use of $\bar{X}$ for 10 thousand (X *mille*) is an illustration of the same tendency, the bar (—) simply standing for *m* (*mille*). In the uncial writing we commonly find ⨪ for *m*, and in the Visigothic we find ÷ for the same purpose. It is quite reasonable, therefore, to think of the dash (−) as a symbol for *m* (minus), just as the cross (+) is a symbol for *et*. It is also possible that its use in this sense may have come from the habit of merchants in indicating a missing number in a case like 2 yd. − 3 in., where the number of feet is missing. We have the same habit in writing certain words today, using either a dash or a series of dots.


(1549) uses the French *plus* and *moins*, while Gosselin (1577) uses P and M. There are, of course, exceptions, as when Trenchant (1566) uses + and − in his work in the Rule of False Position, and when Ramus (1569) writes: "At si falsa conjectura sit, notatur excessus cum signo plus sic †, vel defectus cum signo minus −" (*Schol. Math.*, 1569, p. 138). When he comes to algebra, the plus takes the form —+— (*ibid.*, p. 269). Vieta (*c.* 1590) wrote $x^{4*}.\ px^2$ for $x^4 + px^2$, frequently using the dot as a sign of addition. He also used = for subtraction in certain cases. The asterisk denoted an absence of some power of *x*.

[1] Libri's surmise that they are due to Leonardo da Vinci is not warranted. See *Bibl. Math.*, XIII (2), 52, and his MSS. as published by Boncompagni. Similarly, Treutlein's idea that they are due to Peurbach (*Abhandlungen*, II, 29) is not substantiated.

[2] The al-Khowârizmî MS. in Mr. Plimpton's library.

[3] *E.g.*, the Regiomontanus-Bianchini correspondence, *Abhandlungen*, XII, pp. 233, 279; *Bibl. Math.*, I (3), 506. See also I. Taylor, *The Alphabet*, I, 8 (London, 1883); J. W. L. Glaisher, *Messenger of Math.*, LI, 1–148.

The signs + and − first appeared in print in an arithmetic, but they were not employed as symbols of operation. In the latter sense they appear in algebra long before they do in arithmetic. Their first appearance in print is in Widman's arithmetic (1489), the author saying: "Was − ist / das ist minus . . . vnd das + das ist mer."[1] He then speaks of "4 centner + 5 pfund" and also of "4 centner − 17 pfund," thus showing the excess or deficiency in the weight of boxes or bales. He does not use the symbols to indicate operations, but writes, for example, "$\frac{2}{5}$ $\frac{4}{5}$ $\frac{3}{5}$ adir fa $\frac{9}{5}$ ist $1\frac{4}{5}$," as we write $1\frac{4}{5}$ instead of $1 + \frac{4}{5}$, juxtaposition signifying addition.


>2
4 + 5 Wile du das wyſſ-
4——17 ſen oder deßgley-
3 + 30 chen/So ſumier
4——19 die zenttner vnd
3 + 44 lb vnnd was auß
3 + 22 —iſt/das iſt mi-
Zentner 3——11 lb nus d3 ſetz beſon-
3 + 50 der vnnd werden
4——16 4539 lb (So
3 + 44 du die zendtner
3 + 29 zů lb gemachett
3 —1 12 haſt vnnd das /
3 + 9 + das iſt meer
darzů Addiereſt)vnd 75 minus. Nun
ſolt du für holtz abſchlahen allweeg für
ain legel 24 lb. Vnd das iſt 13 mal 24.
vnd macht 312 lb darzů addier das —
das iſt 75 lb vnd werden 387. Dye ſuß-
trahier von 4539. Vnd bleyben 4152
lb. Nun ſprich 100 lb das iſt ein zentner
pio 4 ff½ wie kumen 4152 lb vnd kumē
171 ff 5 ß 4heller? Vnd iſt recht gemacht
Pfeffer
R

FIRST USE OF THE SYMBOLS + AND −, 1489

First printed use of these symbols, from Widman's *Behennde vnd hüpsche Rechnung*, Leipzig, 1489. This facsimile is from the Augsburg edition of 1526

Manifestly the minus sign was more important as a warehouse mark than the plus sign, since mere juxtaposition serves to express excess.[2]

The first one to make use of the signs + and − in writing an algebraic expression was the Dutch mathematician Vander Hoecke (1514),[3] who gave R $\frac{3}{4}$ − R $\frac{3}{5}$ for $\sqrt{\frac{3}{4}} - \sqrt{\frac{3}{5}}$, and R 3 + 5 for $\sqrt{3+5}$. The next writer to employ them to any extent was Grammateus (1518). He first used them in the Rule of False Position, where, as already stated, they expressed excess and deficiency

[1] 1508 ed., fol. 59; in the 1526 edition, "was auss − ist / das ist minus . . . vnnd das / + das ist meer."

[2] *E.g.*, Albert (1534) writes: "Item/Wie komen 11 cent. 3 stein 18 pfund Zien," and "Item / 12 centner 4 stein —— 6 pfund Talg." He frequently uses the long bar to indicate deficiency, but never uses the plus sign.

[3] Such a statement is likely to be invalidated at any time, and it simply means that no case is known to the author that can be placed earlier than that in Vander Hoecke's work. See the facsimile on page 401. There is a copy of the 1514 edition in the British Museum. For the 1537 edition see *Rara Arithmetica*, p. 183.

instead of operations to be performed.[1] When he wrote upon algebra, however, he used them in the modern sense.[2]

These symbols seem to have been employed for the first time in arithmetic, to indicate operations, by Georg Wälckl (1536),[3] who used $+\frac{1}{3}$ 230 to indicate the addition of $\frac{1}{3}$ of 230, and $-\frac{1}{6}$ 460 to indicate the subtraction of $\frac{1}{6}$ of 460. The algebraist who did the most to bring them into general use was Stifel, to whom the credit for their invention was formerly given.[4] For $3x + 2$ he wrote "3 sum: + 2.,"[5] and similarly for polynomials involving the minus sign.[6] From this time on the two symbols were commonly used by both German and Dutch writers, the particular forms of the signs themselves not being settled until well into the 18th century. Thus, for example, the 1752 edition of Bartjens has

$$xx \overline{\cdot\cdot\cdot\cdot} \div\!\!-\!\!- 2375\,x \text{ ✠ } 1785000$$

for $$x^2 = -2375\,x + 1{,}785{,}000.$$

[1] "Ist zu vil / setze + Ist aber zu wenig / setze —" (1535 ed., fol. E 3). Riese (*Rechnung auff der Linien vnd Federn*, 1522) used the symbols —+— and —÷—, the latter being also used by various other writers to indicate subtraction. Thierfelder (1587), for example, has "25 fl. ÷ 232 gl." (pp. 110, 229). There are numerous variants, such as $\div$ (Coutereels, 1599, 1690 edition of the *Cyffer-Boeck*) and $\overset{\cdot\cdot}{-}$ (Wilkens, 1669). On the present use of ÷ for −, see R. Just, *Kaufmännisches Rechnen*, Leipzig, 1901.

[2] "Vnd mañ brauchet solche zeichen als + ist mehr/vnd—/minder." He illustrates by adding $6x + 6$ and $12x - 4$, thus:

$$\begin{array}{r} 6 \text{ pri.} + 6 \text{ N} \\ 12 \text{ pri.} - 4 \text{ N} \\ \hline 18 \text{ pri.} + 2 \text{ N} \end{array}$$

In one sense $6x + 6$ means an excess of 6 over $6x$, and we evidently find here the transition from the excess stage to the addition stage.

[3] *Die Wâlsch practica*, Strasburg (Nürnberg?), 1536.

[4] Probably because of this expression: "Darumb so gedenck nur nicht, das dise ding schwer seyen zu lernen, oder zubehalten, und ist doch die gantz sach diser meiner zeichen hiemit gantz auszgericht unnd an tag gebracht." *Deutsche Arithmetica*, 1545. For a facsimile from his work, see page 403.

[5] " . . . das zeichen + / welches ich setzē muss zwischen sie / als 2 zu 3 sum: machen 3 sum: + 2. das machstu denn also lesen / 3 summen vnd 2." *Ibid.*, fol. 21.

[6] "Denn wo du dises zeichen — findest / magstu darfur lesen / Weniger oder Minder."

Item wildi afrecken Rx $\frac{12}{27}$ van Rx $\frac{27}{48}$ so ad deert beyde de quadraten coēt $1\frac{1}{144}$ daer na multiplicert dye een quadrate metten anderen coemt 1 die treet van 1 $\frac{1}{144}$ rest $\frac{1}{144}$ nu dē Rx daer af is $\frac{1}{12}$

Irrationale.

De irrationale sedt met — oft met + naer den eysch vanden woerde / als trecht Rx $\frac{3}{5}$ van Rx $\frac{3}{4}$ rest Rx $\frac{3}{4}$ — Rx $\frac{3}{5}$

Item wildi afrecken Rx $\frac{1}{5}$ van Rx $\frac{4}{5}$ rest Rx $\frac{1}{5}$

Item wildi afrecken Rx $\frac{3}{4}$ van Rx 3 — $\frac{3}{16}$ reste Rx $\frac{3}{16}$

Item + van + oft — van — subtraheert inde + van — oft — van + addeert inde subtractie so verre als ß addeerlic sijn oft subtraheerlic.

Multiplicacie inden Rx van rationalen.

Wildi multipliceren inden Rx soe weet dat ghy moet stellen alle de nommers van eenden natu re als Rx te multipliceren met simpelen nommer soe moet ghi den nommer multipliceren nae de qualicyt des Rx. Als wildi multipliceren Rx 9 met 4 so set 4 in sinen Rx multiplicert 4 in haer selven coemt Rx 16 nu multiplicert 9 met 16 coēt 144 hier wt treet Rx coēt 12 soe veel is Rx 9 ghemultiplicert met 4 / want Rx 9 so 3 die multiplicert met 4 coemt 12 als vorē.

Wildi multiplicerē Rx 8 met 5 so multiplicert 5 en

VANDER HOECKE'S USE OF THE PLUS AND MINUS SIGNS

Early use of these signs in Belgium and Holland in 1514. This facsimile is from the 1537 edition

England adopts the Symbols. England early adopted the Teutonic forms, and Recorde (*c.* 1542) says "thys fygure +, whiche betokeneth to muche, as this lyne, — plaine without a crosse lyne, betokeneth to lyttle."[1] Baker (1568) made a vain attempt to change the plus sign, saying: "This Figure ×, betokeneth more: and this plaine line —, signifieth lesse."[2] All this was in connection with the Rule of False Position, and not in connection with arithmetic operations. As symbols of operation most of the English writers of this period reserved the + and − for algebra.[3]

Variants of the Symbols. The variants of the plus sign (+) were naturally many, partly because the early printers had to make up the sign by combining lines that they had in their fonts. Occasionally, however, the religious question enters, as in certain Hebrew works of the 19th century, in which the Christian symbol of the cross is changed[4] to ⊥.

The expression "plus or minus" is very old, having been in common use by the Romans to indicate simply "more or less." It is often found on Roman tombstones, where the age of the deceased was given in some such form as AN · LXXXXIIII · P · M; that is, "94 years more or less."

Symbols of Multiplication. Symbols of multiplication were more slow in their development than symbols of addition and subtraction, the reason being the need for the latter as warehouse marks and in the popular Rule of False Position. The absence of a sign as in £5 and 3 ft. led naturally in the 16th century to a similar usage in such algebraic forms as 6 Pri. for $6x$ and 7 3 (7 zenzo or 7 censo) for $7x^2$. The late medieval

[1] *Ground of Artes*, ed. 1558, fol. Z 6.

[2] Ed. 1580, fol. 184 (numbered 194). Thierfelder (1587) uses × twice through a mistake of the printer (pp. 194, 246), and Wilkens (1669) uses it purposely in connection with + (pp. 190, 191).

[3] Thus Digges (1572), in his treatment of algebra: "Then shall you ioyne them with this signe + Plus"; and Hylles (1600) says: "The badg or signe of addition is +," stating the sum of 3 and 4 as "3 more 4 are 7," and writing 10 —— 3 for "10 lesse 3."

[4] This is found in several such works. Among the latest writers to use the symbol was G. J. Lichtenfeld, *Yedeeoth ha-Sheurim*, Warsaw, 1865.

Der Ander theyl

Von diſen zweyen zeichen/ + vnd —. VII.



O ich von zeychen reden werde/ſoltu mich verſtehn von diſen zeichen + vnd —/Deñ ſolliche verzeichnis/ Sum: oder Sum: A. oder ſ. ꝛc. Werde ich nicht zeychen nennen/ſondern/namen/oder benennung der zalen. Wa ich nu rede von gleichen zeichē/ ſoltu es verſtehn von + vnd + / oder von — vnd —. Alſo auch/wa ich von vngleichen zeichen rede/ſo verſtehe es/von + vnd —.

So haben nu diſe zwey zeichen + vnd — / ein ſonderlichen Algorithmum/welchē ich hie ſtellen will auff 4 Regeln. Denn er gehöret zum Algorithmo der vngerechneten zalen wie du woll ſehen wirſt/vnd alles was vorhin geſagt iſt von diſen namen ſum: ſum: A. ꝛc. das gehöret alles hie her/ als vnter ein einigen Algorithmum.

Die erſt Regel von dem Addiren vnd Subtrahiren. VIII.



Wey gleiche zeichen/ machen eben das ſelbig zeichen/ im Addiren vñ Subtrahiren/ ohn allein ſo du im ſubtrahiren die zal/ die du ſolteſt ſubtrahirē/ nicht kanſt ſubtrahirē.

Exempla vom Addiren.

8 Sum: + 7.	8 Sum: — 18.
12 Sum: + 11.	3 Sum: — 6.
20 Sum: + 18.	11 Sum: — 24.

Hie ſiheſt nu vor augen/wie + vnd + mache im erſten exemplo

STIFEL'S USE OF THE SIGNS + AND — IN ALGEBRA

From Stifel's *Deutsche Arithmetica. Inhaltend. . . . Die Deutsche Coss*, Nürnberg, 1545

writers usually arranged their multiplication tables for commercial use in columns, as in the two cases which follow:


2 · 43 · 86	17	18	306
2 · 44 · 88	17	19	323

In this arrangement no symbols of operation or equality were used,[1] the dot serving for both purposes, being really nothing but a symbol of separation, like the ruled lines.

In the first printed books no such symbols appear, the Treviso arithmetic, for example, giving the multiplication table in the form

2 via 5 fa 10.

Development of the Symbol ×. The common symbol × was developed in England about 1600. In the second edition of Edward Wright's translation of Napier's *Mirifici logarithmorum canonis descriptio* (London, 1618)[2] is "An Appendix to the



Logarithmes," and this contains the statement (p. 4): "The note of Addition is (+) of subtracting (−) of multiplying (x)," a statement that is very likely due to Samuel Wright. The larger symbol (×) is probably due to Oughtred.[3] It was not a new mathematical sign, having long been used in cross

[1] The first of these examples is from a MS. of Benedetto da Firenze written *c.* 1460, and the second from one of Luca da Firenze (*c.* 1475), both in Mr. Plimpton's library.

[2] But not in the 1616 edition.

[3] See F. Cajori, in *Nature*, (December 3) 1914, p. 364; *William Oughtred*, p. 27 (Chicago, 1916); "A List of Oughtred's Mathematical Symbols," *University of California Publications in Mathematics*, I, 171. This monograph should be consulted on the entire question of symbols. It contains a careful study of various algebraic signs. Samuel Wright was the son of Edward Wright. He entered Caius College, Cambridge, in 1612 and died *c.* 1616.

multiplication, in the check of nines,[1] in connection with the multiplication of terms in the division[2] or addition[3] of fractions, for the purpose of indicating the corresponding products in proportion,[4] and in the "multiplica in croce" of algebra as well as in arithmetic.[5] It was probably because of this last use that the symbol was suggested for multiplication, but we have no positive evidence on the subject. It was not readily adopted by arithmeticians, however, being of no practical value to them. In the 18th century some use was made of it in numerical work, but it was not until the second half of the 19th century that it became popular in elementary arithmetic. On account of its resemblance to x it was not well adapted to use in algebra, and so the dot came to be employed, as in $2 \cdot 3 = 6$ (America) and $2 \,.\, 3 = 6$ (Europe). This device seems to have been suggested by the old Florentine multiplication tables; at any rate Vlacq, the Dutch computer (1628), used it in some of his work, thus:

Factores	Faci
7.17	119

although not as a real symbol of operation.[6] Clavius (1583) had an idea of the dot as a symbol of multiplication, for he writes $\frac{3}{5} \cdot \frac{4}{7}$ for $\frac{3}{5} \times \frac{4}{7}$;[7] and Harriot (posthumous work of 1631) actually used the symbol in a case like $2.aaa$ for $2a^3$. The first

[1] In this connection Hylles (1600) speaks of it as the "byas crosse."

[2] As in $\frac{2}{3} \times \frac{3}{5} \frac{9}{10}$, for $\frac{3}{5} \div \frac{2}{3} = \frac{9}{10}$. See page 226.

[3] As in $\frac{4}{5} \times \frac{5}{6} \frac{24}{30} \frac{25}{}$, for $\frac{4}{5} + \frac{5}{6} = \frac{24 + 25}{30}$.

[4] As in the case of $2:3 = 4:6$, as shown at the right. See Buteo, *De Qvadratvra circuli*, p. 67, *et passim* (1559). 2×4 / 3×6

[5] Thus Ghaligai (1521; 1552 ed., fol. 76) gives

$$\begin{matrix} 7 \text{ piu } \text{R}\!\!\text{x} \ 48 \\ 7 \text{ piu } \text{R}\!\!\text{x} \ 48 \end{matrix}$$

to indicate $(7 + \sqrt{48})(7 + \sqrt{48})$.

[6] In his text he uses a rhetorical form, thus: "3041 per 10002 factus erit 30416082."

[7] " . . . minutia minutiae ita scribēda est $\frac{3}{5} \cdot \frac{4}{7}$ pronūciaturque sic. Tres quintae quatuor septimarū vnius integri" (*Epitome*, 1583).

writer of prominence to employ the dot in a general way for algebraic multiplication seems to have been Leibniz (who also used the symbol ⌢)[1] or possibly his contemporary, Christian Wolf, and subsequent algebraists have commonly used it where the absence of a sign does not suffice.[2]

The Symbol ÷. The Anglo-American symbol for division (÷), as already stated, has long been used on the continent of Europe to indicate subtraction. Like most elementary combinations of lines and points, the symbol is old,[3] and toward the close of the 15th century the Lombard merchants used it to indicate a half, as in $4 \div$, $4 \frac{\div}{}$, and similar expressions.[4] There is even a possibility that it was used by some Italian algebraists to indicate division,[5] but it first appeared in print in the *Teutsche Algebra*, by Johann Heinrich Rahn[6] (1622–1676), which appeared in Zürich in 1659. John Pell had been Cromwell's political agent in Switzerland (1654–1658), and Aubrey[7] tells us that "Rhonius was Dr. Pell's pupil at Zürich." He further asserts that "Rhonius's Algebra, in High Dutch, was indeed Dr. Pell's." At any rate, Rahn used the symbol and Pell made it known in England through his translation (London, 1688) of the work.

Symbol (:) for Ratio. The symbol (:) to indicate ratio seems to have originated in England early in the 17th century. It appears in a text entitled *Johnsons Arithmetick; In two*

[1] Gerhardt's edition of his works, II, 239; VII, 54.

[2] Wolf (1713) makes frequent use of the dot in cases like $1 . 2 . 3 . 4$ and $m - 2 . m - 3$, for $4!$ and $(m - 2)(m - 3)$ respectively. See the second edition of his *Elementa Matheseos*, I, 322 (Halle, 1730); also the facsimile of Leibniz's letter, Volume I, page 420.

[3] It was used for *est* as early as the 10th century, as in $i \div$ for *id est* and $it \div$ for *interest*. If used in a case like *divisa* ÷, for *divisa est*, it might possibly have suggested its independent use as a symbol of division.

[4] A. Cappelli, *Dizionario di abbreviature latine ed italiane*, 2d ed., pp. 415, 425. Milan, 1912.

[5] In a MS. in Mr. Plimpton's library, the *Aritmetica et Prattica*, by Giacomo Filippo Biōdi (Biondi) dal Anciso, copied in 1684, the symbol $\frac{\div}{}$ stands for division, so that various forms of this kind were probably used.

[6] Latin Rhonius; see Volume I, page 412.

[7] *Brief Lives*, Oxford edition of 1898, II, 121.

Bookes,[1] but to indicate a fraction, $\frac{3}{4}$ being written 3 : 4. To indicate a ratio it appears in an astronomical work, the *Harmonicon Coeleste* (London, 1651), by Vincent Wing and an unknown writer, "R. B." In this work the forms $A : B :: C : D$ and $A. B :: C. D$ appear frequently as equivalent in meaning.[2] It is possible that Leibniz, who used it as a general symbol of division in 1684,[3] took it from these writers. The hypothesis that it came from $\div$ by dropping the bar has no historical basis. Since it is more international than $\div$, it is probable that the latter symbol will gradually disappear.

Various other symbols have been used to indicate division, but they have no particular interest at the present time.

The Radical Sign. The ancient writers commonly wrote the word for root or side,[4] as they wrote other words of similar kind when mathematics was still in the rhetorical stage. The symbol most commonly used by late medieval Latin writers to indicate a root was ℞,[5] a contraction of *radix*, and this, with numerous variations, was continued in the printed books for more than a century.[6] The symbol was also used for other pur-

[1] Title as in F. Cajori, "Oughtred's Mathematical Symbols," *Univ. of Calif. Pub. in Math.*, I, 181. De Morgan (*Arith. Books*, p. 104) gives it as *Johnson's Arithmatick In 2 Bookes*, 2d ed., London, 1633.

[2] F. Cajori, "Oughtred's Math. Symbols," *loc. cit.*, p. 181. See also W. W. Beman, in *L'Intermédiaire des math.*, IX, 229; F. Cajori, *William Oughtred*, p. 75. In his *Clavis Mathematicae* (1631) Oughtred used a dot to indicate either division or ratio, but in his *Canones Sinuum* (1657) the colon (:) is used for ratio, possibly by some editor or assistant. It appears in the proportion 62496 : 34295 :: 1 : 0/54.9 —. Oughtred ordinarily used the dot for ratio, as in $A. B :: C. D$.

[3] Gerhardt, edition of his works, 3. Folge, V, 223: "$x : y$ quod idem est ac x divis. per y seu $\frac{x}{y}$."


[4] As in Euclid, X, 96. Schoner used *l* for the square root: "Quadrati latus inexplicabile retextum significatur praenota litera *l*" (*De numeris figuratis liber*, 1569; 1586 ed., p. 263). On the Egyptian symbol see Peet, *Rhind Papyrus*, p. 20.

[5] Thus Chuquet (1484) used both ℞ and ℞^{2} for square root, ℞^{3} for cube root, ℞^{4} for fourth root, and so on. See Boncompagni's *Bullettino*, XIII, 655. Regiomontanus (*c.* 1464) has "$\frac{2}{2}\frac{5}{9}$ ig ℞ de $6\frac{2}{4}\frac{1}{7}$ for $\frac{2}{2}\frac{5}{9} - \sqrt{6\frac{2}{4}\frac{1}{7}}$, as in the *Abhandlungen*, XII, 234.

[6] Thus Pacioli (1494) has "E cosi la .R. de .20$\frac{1}{2}$. e .4$\frac{1}{2}$" (fol. 45, *v.*). He also uses ℞. 2^a for square root and ℞. 3^a for cube root, as on fol. 46, *r*. E. de la Roche (1520) used ℞ and ℞^{2} for square root, ℞□ and ℞^{3} for cube root, and ℞ and ℞^{1}

poses, including *responsе*,[1] *res*,[2] *ratio*,[3] *rex*,[4] and the familiar *recipe* in a physician's prescription.[5]

Meanwhile the Arab writers had used various symbols for expressing a root, among them ؟, as in the case of



for $\sqrt{4\frac{1}{2}} + \sqrt{1\frac{1}{2}}$,[6] but none of these signs seem to have influenced European writers.

European Symbols for Roots. The symbol $\sqrt{\ }$ first appeared in print in Rudolff's *Coss* (1525),[7] but without our modern indices. When Stifel edited this work,[8] in 1553, he varied this symbolism, using $\mathfrak{r}/$ for $\sqrt{\ }$, $\mathfrak{z}/$ for $\sqrt[3]{\ }$, $\mathfrak{Z}/$ for $\sqrt[4]{\ }$, and so on. It is frequently said that Rudolff used $\sqrt{\ }$ because it resembled a small *r*, for *radix*, but there is no direct evidence that this is true. The symbol may quite as well have been an arbitrary invention. It is a fact, however, that in and after the 14th century we find in manuscript such forms as  used for the letter *r*.[9]

It was a long time after these writers that a simple method was developed for indicating any root, and then only as a result of many experiments. For example, Vlacq[10] used $\sqrt{\ }$ for square root, $\sqrt{③}$ for cube root, $\sqrt{\sqrt{\ }}$ for fourth root, and so on; Rahn[11]

for fourth root. See the *Abhandlungen*, I, 63. Cardan (1539) and Tartaglia (1556) used $\mathfrak{R}$ for square root and $\mathfrak{R}$ cu. for cube root, while Ghaligai (1521) used $\mathfrak{R}\square$ and $\mathfrak{R}\square\square$, and Bombelli (1572) used $\mathfrak{R}.q$ and $\mathfrak{R}.c.$ respectively for the same purposes. There were also the usual run of eccentricities, as illustrated by the use of Rad. 300 for $\sqrt{300}$ by an Italian arithmetician, Bonini, in 1517.

[1] Trenchant (1566).

[2] For the unknown quantity, as in the Rollandus MS. (*c.* 1424). As representing *res* in general, it is found as early as the 8th century.

[3] As early as the 8th century.

[4] As early as the 14th century.

[5] Also as early as the 14th century.

[6] F. Woepcke, *Recherches*, p. 15. The Arabic forms are read from right to left.

[7] " . . . vermerkt von kürtz wegen radix quadrata mit solchem character $\sqrt{\ }$ . . . radix cubica würt bedeut durch solchen character $C\ \sqrt{\ }\sqrt{\ }$."

[8] *Die Coss Christoph Rudolffs*, fols. 61, 62 (Königsberg i. Pr., 1553). The title-page bears the date 1553; the colophon, 1554.

[9] For these and other forms consult A. Cappelli, *Dizionario*, 2d ed., p. 318.

[10] *Arithmetica Logarithmica*, p. 4. Gouda, 1628.

[11] *Teutsche Algebra*, 1659.

(1622–1676) used $\surd$, $\surd C$, $\surd\surd$, $\surd CC$, and $\surd\surd\surd$ for the square, cube, fourth, sixth, and eighth roots respectively, and various writers used $\surd.\,\overline{3}.$, $\surd.$ cc., $\surd\overline{3}.\overline{3}.$, $\surd S\overline{3}$, and $\surd.\overline{3}.$ cc. for the square, cube, fourth, fifth, and sixth roots respectively.[1]

French, English, and Italian writers of the 16th century were slow in accepting the German symbol, and indeed the German writers themselves were not wholly favorable to it. The letter *l* (for *latus*, side; that is, the side of a square)[2] was often used. Thus we find the Ramus-Schoner work of 1592 using $l4$ for $\surd 4$, lc5 for $\sqrt[3]{5}$, lsq6 and ll6 for $\sqrt[4]{6}$, l ſ 3 for $\sqrt[5]{3}$, and other similar forms, and using the related forms 1 l., 1 q., 1 c., 1 bq., and 1 qc. for a, a^2, a^3, a^4, and a^5 respectively. For the binomial $12+\surd 32$ the work has $b\ 12+1\ 32$, and for the residual $12-\surd 32$ it has $r\ 12-1\ 32$. In a somewhat similar way Gosselin, in his *De Arte Magna* (1577), uses L 9 for $\surd 9$, LC 8 for $\sqrt[3]{8}$, LL 16 for $\sqrt[4]{16}$, and LV 24 PL 9 for $\surd 24+\surd 9$ (the V standing for *universale* and the P for *plus*).

General Adoption of the Radical Sign. In the 17th century our common square-root sign was generally adopted, of course with many variants. Thus Stevin[3] has substantially the same symbols as those used by Rudolff, but with $\surd ③$ for cube root, $\mathbf{W}③$ for the fourth root of the cube root, and so on, with $\surd 3)(2$ for $\surd 3\cdot x^2$ and $\surd 3(2)$ for $\surd 3x^2$. Antonio Biondini, whose algebra appeared in Venice in 1689, has such symbols as $\mathrm{V}8\overline{x}$ for $\surd 8\,x$ and $\underset{c}{\mathrm{V}}24\,xx$ for $\sqrt[3]{24\,x^2}$. The different variants of the root sign are too numerous to mention in detail in this work, particularly as they have little significance. Such forms as

$$\frac{\surd a\times 100}{\lceil 100} \quad \text{for} \quad \frac{\surd a\times 100}{\surd 100}$$

are not uncommon. Newton used $\overset{3}{\surd}8$, $\overset{4}{\surd}16$, . . . for $\sqrt[3]{8}$, $\sqrt[4]{16}$, . . .,[4] but he also used $\sqrt[3]{a}$.


[1] *E.g.*, Cardinael, *Arithmetica*, Bk. I. Amsterdam, 1659.

[2] See page 407, n. 4. [3] *Arithmetiqve* (1585), Girard edition of 1634, pp. 10, 19.

[4] *Arithmetica Universalis*, p. 37 (Cambridge, 1707). Among other statements he has "quod $\surd^6$ valeat $\surd^2\times 3$." Later, as on page 273, he has $\sqrt[3]{-4\pm\surd 8}$.

By the close of the 17th century the symbolism was, therefore, becoming fairly well standardized. We have, however, in Ozanam's *Dictionnaire Mathematique* (Paris, 1691) such forms as $\sqrt{C.aab}$ for $\sqrt[3]{a^2b}$ and $\sqrt{C.a^3+3\,abb}$ for $\sqrt[3]{a^3+3ab^2}$, so that there still remained some work to be done. The 18th century saw this accomplished, and it also saw the negative and fractional exponent come more generally into use. The early history of these forms is considered later.

Symbols of Relation. One of the earliest known symbols of algebra is a sign of equality. This may be said to have appeared in the Ahmes Papyrus (*c.* 1550 B.C.), although Ahmes simply used a hieratic form for a hieroglyphic. He commonly wrote $-$ for the hieroglyphic  , *temt*, meaning "together," the result of addition. In hieroglyphics, for example, we should have  for $10+1=11$.[1] The Egyptians also used  , *er*, meaning "it makes," as in



meaning $\frac{2}{3}\quad\frac{1}{5}\quad\frac{1}{10}\quad\frac{1}{30}\quad er\quad 1,$

or $\frac{2}{3}+\frac{1}{5}+\frac{1}{10}+\frac{1}{30}=1.$[2]

There is no evidence of the use of a generally recognized symbol for equality until the Greeks employed the initials ι^σ or $\overline{\iota\sigma}$ for $\overline{\iota\sigma o\varsigma}$ (*i'sos*),[3] equal. This symbol is found in the *Arithmetica* of Diophantus (*c.* 275).[4] The Arabs, contrary to the Greek custom, used for this purpose the final letter of their word for equality.

In general the classical and medieval writers used the full word.[5] In the Middle Ages a general shorthand was adopted

[1] Eisenlohr ed., p. 39. But on all this consult Peet, *Rhind Papyrus*.

[2] Eisenlohr, *loc. cit.*, p. 41. They also had other forms, for which consult both Peet and Eisenlohr. The symbols used in equations are given on page 422.

[3] As in *isosceles, isoperimetry, isogonal*, etc.

[4] He also used $\overline{\iota\sigma o\varsigma}$ $\overline{\epsilon\sigma\tau\iota}$. For a discussion of the symbol see Heath, *Diophantus*, 2d ed., p. 47. The small Greek letters here shown are modern.

[5] *E.g.*, Fibonacci, in his *Flos* (*c.* 1225), used such forms as *equantur* and *equabitur*. *Scritti*, II, 235.

by university students in copying their texts. Partly as a result of this movement there slowly developed a set of mathematical symbols, other contributing causes being a commercial shorthand and the advantage of expressing an equation in a form easily held by the eye. Thus we have such symbols as ÷ for *est*, p̲ for per, c̊ for *cento*, and ∝ or ∞ for equality.[1] This symbol for equality, ∝ or ∞, was used by Descartes (1637) and is found in various manuscripts of his period. It has generally been thought to come from æ, for *aequalis*, *aequales*, *aequalia*, or *aequantur*. This may be the case, although it is by no means certain.[2]

Various other symbols were used for the same purpose. Thus Buteo (1559) used [; Xylander (1575), ‖; and Hérigone (1634), 2/2. Leibniz (*c.* 1680) used =, ⌐, and other symbols with nearly the same meaning.

Modern Symbol of Equality. As a printed symbol our sign (=) is due to Recorde,[3] who says: "I will sette as I doe often in woorke vse, a paire of paralleles, or Gemowe[4] lines of one lengthe, thus: =, bicause noe .2. thynges, can be moare equalle." If he had used shorter lines (=), there might be some reason for thinking that the symbol was suggested by the medieval use of = for *esse*,[5] but Recorde's clear statement of its arbitrary invention in the form = is conclusive.

The symbol was not immediately popular. When Rahn (1622–1676) wrote his algebra, a century later, he felt obliged

[1] The bar, ——, indicating equality, as used in the correspondence of Regiomontanus, can hardly be considered a symbol in the ordinary sense. See *Bibl. Math.*, I (3), 506.

[2] *E.g.*, in the algebra of Clavius (Rome, 1608, p. 39 seq.) there are expressions like "æquatio inter $\frac{5}{6}$ ℵ+7, & 1 ℵ," and "sit æquatio inter 4 ℵ, & 72 −8 ℵ," so that the ∞ may possibly have come from &. It is quite as reasonable to think that it was a purely arbitrary invention.

[3] *Whetstone of witte*, London, 1557. See the facsimile on page 412.

[4] From O. F. *gemeus*, twins, from Lat. *gemellus*, twin. Recorde uses *gemowe* in his *Pathewaie to knowledg* (1551) to mean parallel, speaking of "Paralleles, or Gemowe lynes." The various zodiacal signs for the gemini may have suggested all these forms.

[5] But not for *est*, where ÷ was commonly used. We find it also in compounds like =nt for *essent*. See A. Cappelli, *Dizionario*, 2d ed., p. 407.

to explain its meaning as not familiar to mathematicians,[1] and the use of ∞ continued until well along in the 18th century.

The Arte

as their woꝛkes doe extende) to diſtincte it onely ſnto twoo partes. Whereof the firſte is, *when one nomber is equalle vnto one other.* And the ſeconde is, *when one nomber is compared as equalle vnto. 2. other nombers.*

Alwaies willyng you to remẽber, that you reduce your nombers, to their leaſte denominations, and ſmalleſte foꝛmes, beſoꝛe you pꝛocede any farther.

And again, if your *equation* be ſoche, that the greateſte denomination *Coſſike*, be ioined to any parte of a compounde nomber, you ſhall tourne it ſo, that the nomber of the greateſte ſigne alone, maie ſtande as equalle to the reſte.

And this is all that neadeth to be taughte, concernyng this woꝛke.

Howbeit, foꝛ eaſie alteratiõ of *equations*. I will pꝛopounde a feiwe exãples, bicauſe the extraction of their rootes, maie the moꝛe aptly bee wꝛoughte. And to auoide the tediouſe repetition of theſe woꝛdes: is equalle to: I will ſette as I doe often in woꝛke vſe, a paire of paralleles, oꝛ Gemowe lines of one lengthe, thus: =, bicauſe noc. 2. thynges, can be moare equalle. And nowe marke theſe nombers.

1. 14.ze. + .15.9 = 71.9.
2. 20.ze. — .18.9 = .102.9.
3. 26.z + 10ze = 9.z — 10ze + 21.9.
4. 19.ze + 192.9 = 10z + 108.9 — 19ze
5. 18.ze + 24.9. = 8.z. + 2.ze.
6. 345 — 12ze = 40ze + 480.9 — 9.z

RECORDE'S SIGN OF EQUALITY

From Recorde's *Whetstone of witte* (1557)

[1] "Bey disem anlaasz hab ich das namhafte gleichzeichen = zum ersten gebraucht, bedeutet *ist gleich*" (*Teutsche Algebra*, 1659). It was probably suggested to him by Pell, who was familiar with Recorde's works.

Symbol of Proportion. The symbol for the equality of ratios (: :), now giving way to the common sign of equality, was introduced by Oughtred (*c.* 1628),[1] and Dr. Pell gave it still more standing when he issued Rahn's algebra in English (1668). It seems to have been arbitrarily chosen.

The symbol $\div\!\!=$ for continued proportion was used by English writers of the 17th and 18th centuries[2] and is still commonly seen in French textbooks.

Symbols of Inequality. The symbols $>$, $<$, for greater and less, are due to Harriot[3] (1631). They were not immediately accepted, for many writers preferred $\sqsubset$ and $\sqsupset$, symbols which Oughtred (1631) had suggested.[4]

The symbols $\neq$, $\not<$, and $\not>$ are modern and are not international, but in the 1647 edition of Oughtred's *Clavis* the somewhat analogous symbols $\sqsubset\!\!\!-$ and $-\!\!\!\sqsupset$ appear for *non majus* and *non minus* respectively. On the Continent the symbols $\gtrdot$ and $\lessdot$, or some of their variants, apparently invented by Pierre Bouguer[5] (1734), are commonly used.

Symbol for Infinity. The symbol for infinity (∞) is first found in print in the *Arithmetica Infinitorum* published by Wallis in 1655,[6] and may have been suggested by the fact that the Romans commonly used this symbol for a thousand, just as we use "myriad" for any large number, although in the Greek it meant ten thousand.

[1] In his *Elementi decimi Euclidis declaratio*, added to the 1648 edition of his *Clavis*, he gives the symbol for "proportio, sive ratio aequalis : :." F. Cajori, *William Oughtred*, p. 26; "Oughtred's Math. Symbols," *loc. cit.*, p. 181, n. 8. It appears also in the 1631 edition of the *Clavis* itself.

[2] *E.g.*, Barrow (*Lectiones Mathematicae*, Lect. XXVII, London, 1683). J. Ward (*c.* 1706) says: "The character made Use of to signify continued Proportionals is $\div\!\!=\!\!\div$" (*The Young Mathematician's Guide*, London, 12th ed., 1771, p. 77). It also appears in the American Greenwood arithmetic (1729).

[3] *Artis Analyticae Praxis*. London, 1631 (posthumous).

[4] *E.g.*, Barrow: "A$\sqsubset$B. A major est quam B. A$\sqsupset$B A minor est quam B" (*Lectiones Opticae & Geometricae* (London, 1674), preface; and English edition (1735), p. 310).

[5] See biographical note on page 327.

[6] This is seen, for example, in such expressions as "jam numerus incrementorum est ∞" (*Opera*, I, 453 (1695)).

Integral Exponents. Our present integral exponents may be said to have begun with Descartes (1637), although Hérigone (1634) had nearly anticipated him. Since the early methods of indicating powers relate naturally to the writing of equations, these are more appropriately considered in connection with that topic (page 421). It may simply be said at this time that Harriot (who died in 1621), in the transition period from the use of forms like Aq to forms like x^2, used *aa* for a^2 and *aaa* for a^3. This symbolism was commonly employed until well into the 18th century, even in writing a polynomial involving a^4; that is, before *c.* 1750 it was common to find expressions like $a^5 + a^4 + aaa + aa + 1$, or even $a^5 + aaaa + aaa + aa + 1$.

In his *Cursus Mathematicus* (1634–1637) Hérigone used *a*2, *a*3, and *a*4 for a^2, a^3, a^4, no doubt influenced by the fact that Girard (1629) used forms like 5 (2) for $5x^2$; and some of his contemporaries, like Dechales (*c.* 1660) and Jacques de Billy (1602–1679), did the same. Descartes (1637), however, wrote the exponents in the present manner;[1] but even without this symbolism Stevin (1585) had already given a systematic discussion of integral exponents.[2]

General Exponents. The general exponent was known in theory long before it came into practical use. Oresme (*c.* 1360) wrote

$\frac{1}{2}\ 2^{\prime}$ for $2^{\frac{1}{2}}$

and

$1^{\prime}\ \frac{1}{2}\ 4$ for $4^{1\frac{1}{2}}$,

and used other similar forms, as already stated. He also gave rules for fractional exponents.

Chuquet (1484) used[3] 12^0 for 12, 12^1 for 12 times a "nombre linear," 12^2 for $12x^2$, and so on. For $9x^{-3}$ he wrote[4] .9.^{3} $\tilde{\text{m}}$,

[1] "Et *aa*, ou a^2, pour multiplier *a* par soy-même; Et a^3, pour le multiplier encore une fois par *a*, & ansi à l'infini" (1705 ed., p. 4).

[2] *L'Arithmetique*, Girard edition of 1634, p. 53. See also the general discussion by Hérigone, *loc. cit.*

[3] As he says, "cõe nõbre simplemẽt pris sans aulcune denomĩacion ou dont sa denoĩacĩo est .0." (Boncompagni's *Bullettino*, XIII, 737).

[4] *Ibid.*, p. 742. See also Ch. Lambo, "Une Algèbre Française de 1484. Nicolas Chuquet," *Revue des Questions Scientifiques*, October, 1902.

thus showing that he had an idea of negative exponents, but it was more than two centuries before the theory was understood. As to fractional exponents, certain evidences show that the idea was developing during the 16th century. This is seen especially in Stifel's *Arithmetica Integra*[1] (1544), where there is given what amounts to the relation

$$(\tfrac{2}{8}\tfrac{7}{})^{1:\frac{3}{4}} = (\tfrac{2}{8}\tfrac{7}{})^{\frac{4}{3}} = \tfrac{8}{1}\tfrac{1}{6}.$$

Albert Girard[2] (1629) employed the fractional exponent, representing it by such forms as $(\frac{1}{6})$ 2000 for $\sqrt[6]{2000}$, and $(\frac{3}{2})$ 49 for $49^{\frac{3}{2}}$; and the study of logarithms from the standpoint of exponents, undertaken at about the same time, tended to bring these general forms into wider use.

Wallis on General Exponents. The first of the writers of this period to explain with any completeness the significance of negative and fractional exponents, however, was Wallis (1655). He showed that x^0 should signify 1, and established relations of the following nature:[3]

$$x^{-1} = \frac{1}{x} \qquad x^{-n} = \frac{1}{x^n}$$

$$x^{\frac{1}{2}} = \sqrt{x} \qquad x^{\frac{1}{q}} = \sqrt[q]{x}$$

$$x^{\frac{2}{3}} = \sqrt[3]{x^2} \qquad x^{\frac{p}{q}} = \sqrt[q]{x^p}$$

Newton supplemented the work of Wallis and in 1669 made use[4] of such forms as $x^{\frac{5}{3}}$ and x^{-3}, and after this time the symbolism became universally recognized.

[1] H. Wieleitner, "Gebrochene Exponenten bei Michael Stifel," *Unterrichtsblätter für Mathematik und Naturwissenschaften*, 1922, No. 5.

[2] *Invention nouvelle en l'algebre*, pp. 97–101. Amsterdam, 1629.

[3] Thus he speaks of $\frac{1}{8}$ "cujus index -3," that is, $2^{-3} = \frac{1}{8}$; of $\frac{1}{\sqrt{2}}$ "cujus index $-\frac{1}{2}$," that is, $2^{-\frac{1}{2}} = \frac{1}{\sqrt{2}}$ ("Arithmetica Infinitorum," in the *Opera* (1695), I, 410, 459, and in the earlier edition).

[4] In the "De Analysi per aequationes numero terminorum infinitas" sent by Collins to Barrow, July 31, 1669. See the *Commercium Epistolicum*, p. 67 (London, 1725). For interesting comments on Newton's use of exponents see G. A. Lecchi, *Arithmetica Universalis Isaaci Newtoni*, Liber II, Pars III, p. 118 (Milan, 1752).

Symbols of Aggregation. Symbols of aggregation first developed to any considerable extent in the 16th century, and in connection with the study of radicals. Tartaglia (1556) writes "22 men (22 men ℞ 6" for $22-(22-\sqrt{6})$. Bombelli (1572) used L, for *legato*, as a kind of symbol of aggregation, as in the squaring of $2+x+\sqrt{20-6x+x^2}$, which appears as

$$2.\ \text{p.}\ \overset{1}{\text{ı}}.\ \text{p. R. q. L 20. m.}\ \overset{1}{6}.\ \text{p.}\ \overset{2}{\text{ı}}.\ \text{⌋},$$

the result being given as

$$\overset{2}{2}.\ \text{p. 24. m.}\ \overset{1}{2}.\ \text{p. R. q. L}\ \overset{4}{4}.\ \text{m.}\ \overset{8}{8}.\ \text{p.}\ \overset{1}{224}.\ \text{p. 320 ⌋},$$

in which the L and the reversed L are clearly symbols of aggregation and may naturally have suggested our square parentheses, first used by Girard (1629) for this purpose.

Other Italian writers frequently employed the letter V, the initial of *universalis*, to indicate that a root sign applied to all the expression which followed. Thus Cardan, in his first printed solution of the cubic equation, has

$$\text{℞ V : cu. ℞ 108 p : 10}$$
$$\text{m : ℞ V : cu. ℞ 108 m : 10}$$

for $$\sqrt{\sqrt[3]{108}+10}-\sqrt{\sqrt[3]{108}-10}.$$

By the time Clavius published his algebra (1608) the parentheses had apparently become common, for he uses them freely without any explanation.

5. Fundamental Operations

Number of Operations. While there were certain operations in arithmetic that were looked upon as fundamental, the number varying from time to time, this was not the case in the early printed algebras. It was only when textbooks, based upon the early arithmetics, came into use, that such operations as addition and subtraction were given as distinct topics. For example, Pacioli (1494) begins his work on algebra[1] by consid-

[1] *Sūma*, fol. 111, *v*.

ering a few definitions, then the laws of signs, and then the operations with monomials, taking up the operations with polynomials somewhat incidentally as they arise;[1] and the same may be said of the other Italian algebraists[2] of the 16th century. Clavius was one of the first to consider the subject somewhat as we do at present.[3] He early introduces a chapter *De additione et svbtractione numerorum Cossicorum*, this being followed by *De mvltiplicatione & diuisione numerorum Cossicorum*.

The reason for this early neglect was that algebra was looked upon as a study for mathematicians, not for boys and girls in their school years. For any mature mind that is interested in mathematics these operations are too simple to require any special attention.

Amount of Work. For this reason the amount of work assigned to topics of this kind by those algebraists who gave them any attention was very slight. For example, Pacioli gives no examples involving numerical cases like $(-2)\ (-3) = +6$, except a few that are completely worked out,[4] and similarly when he comes to surds.[5] An illustration of his problems is seen in the following:

via. 4. p. R. 6.
4. m. R. 6.
16. m. 6.
Productum 10

meaning that $(4 + \sqrt{6})\ (4 - \sqrt{6}) = 16 - 6 = 10$.[6] Similarly, Tartaglia[7] solves a few typical problems involving signs, but gives no exercises for original work.

Difficulties Due to Poor Symbolism. A good idea of the general difficulties which characterized this period of poor symbolism is seen in the algebra of Pedro Nunes, of which the second

[1] *E.g.*, fol. 127, *v.*, seq.
[2] Tartaglia, *General Trattato*, II, fol. 81 (1556); Bombelli, *Algebra, libro primo* (1572); Cardan, *Ars Magna*, cap. i (1545).
[3] *Algebra*, p. 16 (Rome, 1608).
[4] Fol. 111, *v.*, seq.
[5] Fol. 115, *v.*, seq.
[6] Fol. 123, *r.*
[7] *General Trattato, La seconda parte*, fol. 81 (Venice, 1556).

edition appeared at Antwerp in 1567. His multiplication of $3x^2 + 2\frac{1}{4}x + 5\frac{1}{16}$ by $4x + 3$ appears as follows:

$$\begin{array}{l} 3 \cdot ce \cdot \tilde{p} \cdot \quad 2 \cdot co \cdot \frac{1}{4} \cdot \tilde{p} \cdot 5\frac{1}{16} \\ \qquad\qquad 4 \cdot co \cdot \tilde{p} \cdot 3. \\ \hline 12 \cdot cu \cdot \tilde{p} \cdot \quad 9 \cdot ce \cdot \tilde{p} \cdot 20 \cdot co \cdot \frac{1}{4} \\ \qquad\qquad 9 \cdot ce \cdot \tilde{p} \cdot \quad 6 \cdot co \cdot \frac{3}{4} \cdot \tilde{p} \cdot 15\frac{3}{16} \\ \hline 12 \cdot cu \cdot \tilde{p} \cdot 18 \cdot ce \cdot \tilde{p} \cdot 27 \cdot co \cdot \quad \tilde{p} \cdot 15\frac{3}{16} \end{array}$$

Bombelli (1572) sets forth the work more after the modern plan, but gives no cases to be solved independently. The following is a type:

$$\begin{array}{l} \quad \underline{1}\ \mathrm{I}\ \mathrm{p}\ 2 \\ \quad \underline{1}\ \mathrm{I}\ \mathrm{p}\ 2 \\ \underline{2}\ \mathrm{I}\ \mathrm{p}\ 4\ \underline{1}\ \mathrm{p}\ 4 \\ \underline{2}\ \mathrm{I}\ \mathrm{p}\ 4\ \underline{1}\ \mathrm{p}\ 4 \\ \underline{4}\ \mathrm{I}\ \mathrm{p}\ 8\ \underline{3}\ \mathrm{p}\ 24\ \underline{2}\ \mathrm{p}\ 32\ \underline{1}\ \mathrm{p}\ 16 \\ \underline{1}\ \mathrm{I}\ \mathrm{p}\ 2 \\ \hline \underline{5}\ \mathrm{I}\ \mathrm{p}\ 10\ \underline{4}\ \mathrm{p}\ 40\ \underline{3}\ \mathrm{p}\ 80\ \underline{2}\ \mathrm{p}\ 80\ \underline{1}\ \mathrm{p}\ 32, \end{array}$$

meaning that $(x + 2)^2 = x^2 + 4x + 4$,

$$(x^2 + 4x + 4)^2 = x^4 + 8x^3 + 24x^2 + 32x + 16,$$

and this multiplied by $x + 2$ gives the fifth power of $x + 2$.[1]

6. Continued Fractions

Early Ideas. It is not necessary to speak of the history of simple algebraic fractions, since these forms were transferred from arithmetic. When Euclid found the greatest common measure of two lines,[2] or when the same principle was applied to the finding of the greatest common divisor of two numbers,[3]

[1] 1572 ed., p. 69. The I seems to have been the coefficient of the highest power.
[2] *Elements*, X, 3 and 4, for commensurable magnitudes in general.
[3] *Elements*, VII, 2 and 3.

a process was used that is similar to that of converting a fraction into a continued fraction, as is evident from the following:

$$\begin{array}{r} 12)\,38\,(3 \\ \underline{36} \\ 2)\,12\,(6 \\ \underline{12} \end{array} \qquad \frac{12}{38} = \frac{6}{19} = \frac{1}{3+\frac{1}{6}}.$$

This is the earliest important step in the theory of continued fractions.[1] Further traces of the general idea are found occasionally in the Greek and Arab writings.

Beginning of the Modern Theory. Although the Greek use of continued fractions in the case of greatest common measure was well known in the Middle Ages, the modern theory of the subject may be said to have begun with Bombelli (1572). In his chapter relating to square root[2] he considered the case of $\sqrt{13}$. Substituting our modern symbolism, he showed that this number is equal to

$$3+\cfrac{4}{6+\cfrac{4}{6+\cdots}}.$$

In other words, he knew essentially that

$$\sqrt{a^2+b} = a+\cfrac{b}{2a+\cfrac{b}{2a+\cdots}}.$$

The next writer to consider these fractions, and the first to write them in substantially the modern form, was Cataldi[3] (1613), and to him is commonly assigned the invention of the theory. His method was substantially the same as Bombelli's,

[1] On the history in general, see S. Günther, *Beiträge zur Erfindungsgeschichte der Kettenbrüche*, Prog., Weissenburg, 1872; Italian translation, Boncompagni's *Bullettino*, VII.

[2] "Modo di formare il rotto nella estrattione delle Radici quadrate," *Algebra*, p. 35.

[3] *Trattato del modo brevissimo di trovare la radice quadra delli numeri* Bologna, 1613.

but he wrote the result of the square root of 18 in the following form:

$$4.\ \&\ \frac{2}{8.}\ \&\ \frac{2}{8.}\ \&\ \frac{2}{8}.$$

This he then modified, for convenience in printing,[1] into the form[2]

$$4\ \&\ \frac{2}{8.}\ \&\ \frac{2}{8.}\ \&\ \frac{2}{8.}.$$

The third writer to take up the theory was Daniel Schwenter (1618). In attempting to find approximate values for $\frac{177}{233}$ he found the greatest common divisor of 177 and 233, and from this he determined the convergents[3] as $\frac{79}{104}$, $\frac{19}{25}$, $\frac{3}{4}$, $\frac{1}{1}$, and $\frac{0}{1}$.

The next writer of prominence to use these forms was Lord Brouncker,[4] who transformed the product

$$\frac{4}{\pi} = \frac{3 \cdot 3 \cdot 5 \cdot 5 \cdot 7 \cdot 7 \cdots}{2 \cdot 4 \cdot 4 \cdot 6 \cdot 6 \cdot 8 \cdots},$$

which had been discovered by Wallis, into the fraction

$$\frac{4}{\pi} = 1 + \cfrac{1}{2 + \cfrac{9}{2 + \cfrac{25}{2 + \cfrac{49}{2 + \cdots}}}},$$

as already stated on page 311. He made no further use of these forms, Wallis then taking up the work and using the name "continued fraction."[5]

[1] "Notisi, che nõ si potendo cõmodamẽte nella stampa formare i rotti . . . ." See Tropfke, *Geschichte*, II (1), 362.

[2] ". . . facendo vn punti all' 8. denominatore de ciascun rotto, à significare, che il sequente rotto è rotto d' esso denominatore" (p. 70).

[3] For details see Tropfke, *Geschichte*, II (1), 363.

[4] J. Wallis, *Opera Mathematica*, I, 469 (Oxford, 1695). See also *Commercium Epistolicum*, London, 1725 ed., p. 215.

[5] ". . . quae denominatorem habeat continue fractum" (*Opera*, I, 469). His symbolism is

$$1\tfrac{1}{2}\ \tfrac{8}{2}\ \tfrac{25}{2}\ \tfrac{49}{2}\ \tfrac{81}{2}\ \&c.$$

See also Euler, *Introductio in Analysin Infinitorum*, ed. nova, I, 305 (Lyons, 1797).

The next advance was made by Huygens in his work on the description of a planetarium,[1] the ratio 2,640,858 : 77,708,431, for example, being written as a continued fraction.

In some manner, perhaps through the missionaries in China, the idea of the continued fraction found its way to Japan at about this time.[2] Takebe Hikojirō Kenkō (1722) used such forms for the value of π, stating that the plan was due to his brother, Takebe Kemmei. The first few convergents given by him are $\frac{3}{1}$, $\frac{22}{7}$, $\frac{333}{106}$, $\frac{355}{113}$, and $\frac{103993}{33102}$.

Euler founds the Modern Theory. The first great memoir on the subject was Euler's *De fractionibus continuis* (1737), and in this work the foundation for the modern theory was laid. Among other interesting cases Euler developed e as a continued fraction,[3] thus:

$$e = 2 + \cfrac{1}{1 + \cfrac{1}{2 + \cfrac{1}{1 + \cfrac{1}{4 + \cfrac{1}{1 + \cdots}}}}}.$$

Of the later contributors to the theory, special mention should be made of Lagrange[4] (1767) and Galois.[5]

7. The Writing of Equations

Equations in One Unknown. In speaking of the symbols for unknown quantities we are brought directly in touch with the symbols for integral exponents and with the writing of equations, and so it is convenient to treat of these topics in their relation to one another.


[1] *Descriptio automati planetarii*, The Hague, 1698 (posthumous).

[2] Smith-Mikami, p. 145.


[3] *Comm. Acad. Petrop.* for 1737, IX, 120 (Petrograd, 1744). See Tropfke, *Geschichte*, II (1), 337. See also Euler's *Introductio*, I, 293, "De fractionibus continuis," especially the forms on pages 117, 307; and his "De formatione fractionum continuarum," in the *Acta Petrop.* for 1779, I, 3 (Petrograd, 1782), and other essays.

[4] See Serret's edition of his works, II, 539, and VII, 3 (Paris, 1868).


[5] Gergonne's *Annales de Math. Pures et Appliquées*, XIX, 294; posthumous (1828–1829). *Œuvres mathématiques d'Evariste Galois*, pp. 1–8 (Paris, 1897).

As already stated, the Egyptians called the unknown quantity *ahe* or *hau*, meaning "mass."[1] This word was represented in the hieroglyphic[2] as  .

For example, the equation $x(\frac{2}{3}+\frac{1}{2}+\frac{1}{7}+1)=37$ would appear in hieroglyphics as



and in the hieratic of the Ahmes Papyrus[3] (*c.* 1550 B.C.) as



It will be observed that, although Ahmes knew of symbols for plus, minus, and equality, they are not commonly used in his equations. They are found, however, in No. 28 of both the Peet and the Eisenlohr translation.

Symbolism of Diophantus. The first writer to make much effort toward developing a symbolism for the powers of algebraic expressions was Diophantus (*c.* 275). He used the following abbreviations for the various powers of the unknown:

Modern		Diophantus[4]	Late Editions
x^0	$\overset{o}{\mathrm{M}}$	μονάδες, units	$\mu\hat{o}$
x^1	ς	ἀριθμός, number	ς' or $\varsigma^{o\acute{}}$
x^2	Δ^{Y}	δύναμις, power	$\delta^{\hat{\upsilon}}$
x^3	K^{Y}	κύβος, cube	$\kappa^{\hat{\upsilon}}$
x^4	$\Delta^{\mathrm{Y}}\Delta$	δυναμοδύναμις, power-power	$\delta\delta^{\hat{\upsilon}}$
x^5	$\Delta\mathrm{K}^{\mathrm{Y}}$	δυναμόκυβος, power-cube	$\delta\kappa^{\hat{\upsilon}}$
x^6	$\mathrm{K}^{\mathrm{Y}}\mathrm{K}$	κυβόκυβος, cube-cube	$\kappa\kappa^{\upsilon}$

[1] On this term see page 393, n. 5.

[2] There are several variants. See Eisenlohr, *Ahmes Papyrus*, p. 42.

[3] Eisenlohr, *ibid.*, p. 54, No. 33; in the British Museum facsimile, Plate X, row 5.

[4] See Heath, *Diophantus*, 2d ed., p. 32. Much has been written about the symbol ς for x, and Heath gives a careful discussion of the various theories and a statement of the various forms for the symbol as they occur in different MSS. He concludes that the original symbol was a contraction of the initial letters αρ of ἀριθμός (*arithmos'*, number), instead of being the final sigma. Originally the capitals Δ^{Y} were used for δ^{υ}, and similarly K^{Y} instead of κ^{υ}, and so on.

Diophantus wrote his equations quite as we do, except for the symbols; thus the equation


$$(2x+1)^2 = 4x^2 + 4x + 1$$

appears, in modern Greek letters, as

$$\varsigma\ \overline{\beta}\ \ddot{\mathrm{M}}\ \overline{\alpha}\ \square^{\mathrm{os}}\ \grave{\epsilon}\sigma\tau\acute{\iota}\ \Delta^{\Upsilon}\ \overline{\delta}\ \varsigma\ \overline{\delta}\ \mathring{\mathrm{M}}\ \overline{\alpha},$$

and the equation $8x^3 - 16x^2 = x^3$

appears[1] as $\mathrm{K}^{\Upsilon}\ \overline{\eta}\ \bigwedge\ \Delta^{\Upsilon}\ \overline{\iota\varsigma}\ \ddot{\iota}\sigma\ \mathrm{K}^{\Upsilon}\ \overline{\alpha}.$




DIOPHANTUS ON EQUATIONS

From a manuscript of the 14th century showing the symbolism then in use. It begins: "We call the square *δύναμις*, and it has for its symbol a *delta* (Δ) surmounted by a *upsilon* (Υ)." [This character is seen in the middle of line 2.] The symbol $\Delta\Delta^{\nu}$ (for x^4) appears in line 5, $\Delta\kappa^{\nu}$ (for x^5) in line 8, and $\kappa\kappa^{\nu}$ (for x^6) in the last line. From Rodet, *Sur les Notations Numériques*, Paris, 1881

In speaking of Diophantus, however, it should again be stated that the most ancient manuscript of his *Arithmetica* now extant was written in the 13th century,—about a thousand years after the original one appeared. We are therefore quite

[1] In the first equation *ἐστί* (*estí*) stands for "is equal to," and in the second case *ἴσ* stands for *ἴσος* (*ísos*, equal). See Tannery's *Diophantus*, I, 230–231, 258–259.

uncertain as to the symbols used by Diophantus himself and as to the various interpolations that may have been made by the medieval copyists.



ALGEBRAIC SOLUTION ACCORDING TO DIOPHANTUS

From a manuscript of the 14th century. The problem[1] is to find two numbers such that their sum is equal to 20 and the difference of their squares to 80

Oriental Symbols. It was from this Greek method of expressing the equality of the two members that the Arabs seem to have derived theirs, as in the case of



for $$38 = 19x + x^2,$$

the equation being written from right to left after the Semitic custom, which obtains in writing Arabic.[2] From this form, reversed in order of writing, came the one that we use.

The Chinese and Hindus, however, had methods of writing their equations that were very different from those which the


[1] The solution reads substantially as follows:

Let there be	$x + 10$	$10 - x.$
Squares,	$x^2 + 20x + 100$	$x^2 + 100 - 20x.$
Difference of squares,	$40x$	$= 80.$
Division,	x	$= 2,$
whence	$x + 10 = 12$	$10 - x = 8.$

From Rodet

[2] The above is from al-Qalasâdî. See L. Matthiessen, *Grundzüge der antiken und modernen Algebra*, 2d ed., p. 269 (Leipzig, 1896); hereafter referred to as Matthiessen, *Grundzüge*.

Arabs and Persians adapted from the Greek works. The Far East depended more upon position. The Chinese commonly represented the coefficients by sticks, their so-called "bamboo rods" which they used in calculating, and these they placed in squares on a ruled board. Ch'in Kiu-shao (*c.* 1250), for example, represented the equation $x^3 + 15x^2 + 66x = 360$ as here shown; and if he needed to write the equation, he did so in the same manner.[1] The positive terms were represented by red sticks or marks, and the negative terms either by black ones or (as in the illustration) by a stick placed diagonally across some part of the numeral. The system is simply one of detached coefficients, the place values of the coefficients being indicated sometimes by the squares running horizontally, but ordinarily as shown in the illustration. The native Japanese mathematicians used the same method, having imported it from China.[2]



The Hindu method was better than the Chinese, and in one respect was the best that has ever been suggested. Bhāskara (*c.* 1150) represented the equation $18x^2 = 16x^2 + 9x + 18$ as follows:

न्यासः यावत्तावत्या १८ या ० रू ०
यावत्तावत्या १६ या ९ रू १८

This may be transliterated as

ya v 18 *ya* 0 *ru* 0
ya v 16 *ya* 9 *ru* 18

which means

$18x^2$ $0x$ 0
$16x^2$ $9x$ 18

or $18x^2 + 0x + 0 = 16x^2 + 9x + 18,$

or $2x^2 - 9x = 18.$

[1] The Chinese word *tai* (from *tai-kieh* or *tai-chi*, extreme limit, or great extreme) means the absolute term, and *yuen* (element) means the first power of the unknown. See Mikami, *China*, pp. 81, 82, 91; L. Vanhée, "La notation algébrique en Chine au XIII^e siècle," *Revue des Questions Scientifiques*, October, 1913.

[2] Smith-Mikami, p. 50.

The word for the first power of the unknown, the *yāvat-tāvat*, already explained, is abridged to *ya*; and the word for the second power, *yāvat-varga*, to *ya v.*[1]


Such a plan glances the similar terms one above another, and permits of easy transposition.

When the Arabic algebras were translated into Latin, the rhetorical form was used. Thus Robert of Chester (*c.* 1140), in his translation of al-Khowârizmî, wrote "Substantia et 10 radices 39 coaequantur drachmis" for "A square and 10 roots are equal to 39 units"; that is,

$$x^2 + 10x = 39.^2$$

Al-Khowârizmî himself wrote his equations in rhetorical form, thus: "A square, multiply its root by four of its roots, and the product will be three times the square, with a surplus of fifty dirhems."[3]

Medieval Manuscripts. In the manuscript period of the Middle Ages we find



PAGE FROM BHĀSKARA'S BIJA GANITA

From the first printed edition, showing the method of writing equations. Lines 8 and 9 give the equation shown in the text. The next equation is interesting for the use of the dot above the Sanskrit 9 to indicate subtraction, thus:

ya v 2 *ya* 9̇ *ru* 0
ya v 0 *ya* 0 *ru* 18

[1] Colebrooke, *loc. cit.*, p. 140; E. Strachey, *Bija Ganita* (often bound with Colebrooke), p. 117.

[2] In the Scheubel MS., translated by Karpinski, pp. 70–73. Somewhat the same form is used in the transcription in Libri's *Histoire*, I, 255, although the exact wording is "Census et decem radices equantur triginta novem dragmis."

[3] Rosen translation, p. 56; Arabic text in the Rosen edition, p. 40.

letters coming into use to represent algebraic as well as geometric quantities. This is seen in the work of Jordanus Nemorarius (*c.* 1225), a contemporary of Fibonacci.[1] This was not common, however, most writers preferring to use some such symbols as R for *res* (thing, the unknown), *ce.* for *census* (the second power of the unknown), and *cu.* for *cubus* (the third power of the unknown), with other shorthand abbreviations. Such symbolism is seen in the manuscripts of Regiomontanus (*c.* 1463), one problem from which is reproduced in facsimile on page 429. In an Italian manuscript of about the same period[2] the quadratic equation $x^2+10x=39$ appears in the rhetorical form as follows: "lo censo e·10·sue cose cioe·10·sue rā. sono igualj a·39·dramẽ."

Equations in Printed Form. The following examples will suffice to show the general development of the symbolism of the equation from the first printed work containing algebra to the time when our present symbolism was fairly well settled:

Pacioli (1494)[3]: "Trouame .1. n°. che giōto al suo q̄drat° facia .12." Modern form: $x+x^2=12$.

Vander Hoecke (1514)[4]: 4 Se. − 51 Pri. − 30 N. dit is ghelijc $45\frac{3}{5}$. Modern form: $4x^2-51x-30=45\frac{3}{5}$.

Ghaligai (1521)[5]: 1□e 32 c° − 320 numeri. Modern form: $x^2+32x=320$.

[1] For example, in the second problem in his *De Numeris Datis* he says: "Datus numerus sit .a. qui diuidatur in .b.c.d.e. . . ." *Abhandlungen*, II, 135. See also A. Favaro, Boncompagni's *Bullettino*, XII, 129; M. Curtze, *Abhandlungen*, XII; A. Witting and M. Gebhardt, *Beispiele zur Gesch. der Math.*, II, 26 (Berlin, 1913).

[2] In Mr. Plimpton's library. See *Rara Arithmetica*, p. 459. The equation is from the MS., fol. 279, *v*.

[3] Fol. 145, *r*. In his solutions, but not in his problems, he used .co. (*cosa*, thing, as already explained) for x; .ce. (*census* or *zensus*; in Italian, *censo*, evaluation of wealth, tax) for x^2; .cu. (*cubus*) for x^3; .ce.ce. (*census census*) for x^4; and .p°.r°. (*primo relato*) for x^5. The Latin *census* (a registering of citizens and property) was conducted by the censors, who gave *censura*, censure, to those who incurred their disfavor, *census* coming probably from *centere*, to number by the *centum*, hundred.

[4] He used Pri., Se., 3^a, 4^a, and 5^a for x, x^2, x^3, x^4, and x^5 respectively. The system failed because of the difficulty in writing coefficients. See his 1537 edition, fol. 64, *v*.

[5] Fol. 96, *v*.

Rudolff (1525)[1]: Sit 1 ȝ aequatus 12 ℵ − 36. Modern form: $x^2 = 12x - 36$.

Cardan (1545)[2]: cub^o p: 6 reb^o aeq̄lis 20. Modern form: $x^3 + 6x = 20$.

Scheubel (1551)[3]: 4 sex. aequantur 108 ter. Modern form: $4x^7 = 108x^4$.

Tartaglia (1556)[4]: "Trouame uno numero che azontoli la sua radice cuba uenghi ste, cioe .6." Modern form: $x + \sqrt[3]{x} = 6$.

Buteo (1559)[5]: 1 ◊ P6ρP9L1 ◊ P3ρP24. Modern form: $x^2 + 6x + 9 = x^2 + 3x + 24$.

Bombelli (1572)[6]: $\overset{6}{\breve{1}}$. p. $\overset{3}{8}$. Eguale à 20. Modern form: $x^6 + 8x^3 = 20$. In the text proper he would write this equation 1^{6}p. 8^3 eguale à 20. In the same way he would write 1^3 eguale à R. q. 108. p. 10. for $x^3 = \sqrt{108} + 10$, which may be compared with Cardan's form on page 463.

[1] From Stifel's edition of 1553 (1554), fol. 243, *v*. The symbols for the first five powers of the unknown, beginning with the first, are ℵ (contraction of *radix*?), ȝ (*zensus*), ↁ (contraction of cs, for *cubus*), ȝȝ (*zensus zensus*), ȝ (*sursolidus*). The German writers in general used this system until well into the 17th century. Although the symbol for the unknown is usually taken as a contraction for *radix*, it is quite as probable that it is the common ligature of the Greek γ and ρ. This stood for *gram'ma* (γράμμα), a letter, or for *gramme'* (γραμμή), a line. In the medieval works it was a common thing to represent the unknown by a line. This is seen in an algebra as early as al-Khowârizmî's (*c.* 825) and in one as late as Cardan's (1545). For evidence of the frequent use of the symbol for *gramma*, see Michael Neander, ΣΥΝΟΨΙΣ *mensvrarvm et Pondervm*, Basel, 1555. It may, indeed, have suggested to Descartes the use of x, this being the letter most nearly resembling it.


[2] *Ars Magna*, 1545 ed., fol. 30, *r*. His names or abbreviations for x, x^2, x^3, x^4, and x^5 are *res*, $\tilde{q}d$ (*quadratum*), *cu'* or *cub^o* (*cubus*), $\tilde{q}d'$ $\tilde{q}d'$, $\tilde{q}d$ $\tilde{q}d$, or $\tilde{q}d^{ri}$ qd^m (*quadrati quadratum*), and *relatum primum*, with necessary variants of these forms. For a facsimile, see pages 462, 463.

[3] Also 1 pri. + 12 N aequales 8 ra., for $x^2 + 12 = 8x$. See *Abhandlungen*, IX, 455.

[4] *La Noua Scientia*, 1554 ed., fol. 114, *r*. His problems are in rhetorical form, and his symbolism is substantially that of Cardan and other Italian contemporaries. For Recorde's equations (1557) see the facsimile on page 412.

[5] The next step is 3ρ[15], sometimes with both brackets, sometimes with only the first. Buteo, *Logistica, quae & Arithmetica vulgò dicitur*, Lyons, 1559.

[6] *Algebra*, p. 273. He indicated x, x^2, x^3, . . . by $\breve{1}$, $\breve{2}$, $\breve{3}$, ... It is in this work that there appears for the first time in Italian any important approach to the modern symbolism of the equation.



SYMBOLISM OF THE EQUATION AS USED BY REGIOMONTANUS, *c.* 1463

From a letter written by Regiomontanus. The problem is to find a number x such that

$$\frac{100}{x} + \frac{100}{x+8} = 40 \quad \text{or} \quad x^2 + 3x = 20,$$

from which $x = \sqrt{\frac{89}{4}} - \frac{3}{2}$, as stated in the third line from the bottom. The conclusion is that the first divisor is $\sqrt{22\frac{1}{4}}$ minus $1\frac{1}{2}$. See page 427

Gosselin (1577)[1]: 12LM1QP48 aequalia 144M24LP2Q. Modern form: $12x - x^2 + 48 = 144 - 24x + 2x^2$.

Stevin (1585)[2]: 3 ② + 4 egales à 2 ① + 4. Modern form: $3x^2 + 4 = 2x + 4$.

Ramus and Schoner (1586)[3]: 1q—+—81 aequatus sit 65. Modern form: $x^2 + 8x = 65$.

Vieta (*c.* 1590)[4]: 1 Q C − 15 Q Q + 85 C − 225 Q + 274 N, aequatur 120. Modern form:

$$x^6 - 15x^4 + 85x^3 - 225x^2 + 274x = 120.$$

Clavius (1608)[5]: "Sit aequatio inter 1.3ce & 800ce−156751." Modern form: $x^6 = 800x^3 - 156{,}751$.

Girard (1629)[6]: 1 (4) + 35 (2) + 24 = 10 (3) + 50 (1), or with the several exponents inclosed in circles. Modern form: $x^4 + 35x^2 + 24 = 10x^3 + 50x$.

Oughtred (1631)[7]: $\frac{1}{2}$ Z ± √q: $\frac{1}{4}$ Zq − AE = A. Modern form: $\frac{1}{2}Z \pm \sqrt{\frac{1}{4}Z^2 - AE} = A$.

Harriot (1631)[8]: aaa − 3 · bba = + 2 · ccc. Modern form: $x^3 - 3b^2x = 2c^3$.

[1] Guillaume Gosselin was a native of Caen, but we know almost nothing of his life. He published an algebra, *De arte magna, seu de occulta parte numerorum, quae & Algebra, & Almucabala vulgo dicitur; Libri QVATVOR*, Paris, 1577; and a French translation of Tartaglia's arithmetic, Paris, 1578. See H. Bosmans, *Bibl. Math.*, VII (3), 44. In the above equation he uses L for *latus* (the side of the square), Q for *quadratus* (square), and P and M for *plus* and *minus*.

[2] *L'Arithmetiqve*, p. 272. See also his *Œuvres*, Girard ed., p. 69 (Leyden, 1634). He used ①, ②, ③, ··· for x, x^2, x^3, ···.

[3] *Algebrae Liber Primus*, 1586 ed., p. 349.

[4] For a discussion of the dates of his monographs, see Cantor, *Geschichte*, II (2), 582. He also used capital vowels for the unknown quantities and capital consonants for the known, thus being able to express several unknowns and several knowns. The successive powers of A were then indicated by A, Aq, Acu, Aqq, Aqcu, and so on, the additive principle of exponents being followed. The above example is from his *Opera Mathematica*, ed. Van Schooten, p. 158 (Leyden, 1646).

[5] *Algebra*, p. 62. For his symbols, see *ibid.*, p. 11.

[6] *Invention nouvelle en l'algèbre*, p. 131 (Amsterdam, 1629), with rules for the symmetric functions of the roots.

[7] *Clavis*, p. 50; Cajori, *William Oughtred*, p. 29.

[8] See his *Artis Analyticae Praxis*, London, 1631. He represented the successive powers of the unknown by *a*, *aa*, *aaa*, . . . ; Tropfke, *Geschichte*, III (2), 143, a work which should be consulted (pp. 119–148) on this entire topic.

Hérigone (1634)[1]: 154a⌣71a2 +− 14a3⌣a4 2/2 120. Modern form: $154a - 71a^2 + 14a^3 - a^4 = 120$.

Descartes (1637)[2]: yy ∞ cy $-\frac{cx}{b}$ y + ay − ac. Modern form: $y^2 = cy - \frac{cx}{b}\, y + ay - ac$.

Wallis (1693): $x^4 + bx^3 + cxx + dx + e = 0$, which is the modern form, with the exception of xx; this, as already stated, was commonly written for x^2 until the close of the 18th century.[3]

Equating to Zero. It is difficult to say who it was who first recognized the advantage of always equating to zero in the study of the general equation. It may very likely have been Napier, for he wrote his *De Arte Logistica* before 1594 (although it was first printed in Edinburgh in 1839), and in this there is evidence that he understood the advantage of this procedure.[4] Bürgi (*c.* 1619) also recognized the value of making the second member zero, Harriot (*c.* 1621) may have done the same, and the influence of Descartes (1637) was such that the usage became fairly general.[5]

Several Unknowns. The ancients made little use of equations with several unknown quantities. The first trace that we find of problems involving such equations is in Egypt. There are

[1] *Cursus Mathematicus*, 5 vols., Paris, 1634–1637; 2d ed., 6 vols. in 4 (Paris, 1644), Vol. II, chap. xiv. He represented the successive powers of the unknown by *a*, *a*2, *a*3, *a*4, ...

[2] *La Geometrie*, 1637; 1705 ed., p. 36. It will be seen that this form does not differ much from our own. Descartes used the last letters of the alphabet for the unknown quantities and the first letters for the known, and this usage has persisted except in the case of those formulas in which the initial letter serves a better purpose. That it was not immediately accepted, however, is seen by the fact that Rahn (Rhonius) used final letters for unknowns and large letters for knowns, as in his algebra of 1659 (English translation, 1688).

[3] *E.g.*, in Euler's *Algebra*, French ed., Petrograd, 1798, where such forms as $xx + yy = n$ are found.

[4] For example, on page 156 he takes the equation $4\,\mathfrak{R} - 6 = 5\,\mathfrak{R} - 20$ and reduces it to $-\mathfrak{R} + 14 = 0$, "quae aequatio ad nihil est." In general all his higher equations have zero for the second member. See also Eneström, *Bibl. Math.*, III (3), 145.

[5] For a discussion see Tropfke, *Geschichte*, III (2), 26; Kepler's *Opera*, ed. Frisch, V, 104. The credit is often claimed for Stifel (*c.* 1525), but he, like Harriot, made no general practice of equating to zero.

three papyri[1] of the Middle Kingdom (*c.* 2160–1700 B.C.) which contain problems of this nature. One of these problems is to divide 100 square measures into two squares such that the side of one of the squares shall be three fourths the side of the other; that is,

$$x^2 + y^2 = 100,$$
$$y = \tfrac{3}{4}x.$$

The other problems also involve quadratics, one found in 1903 being substantially[2]

$$x^2 + y^2 = 400,$$
$$x : y = 2 : 1\tfrac{1}{2}.$$

Simultaneous Linear Equations. The earliest of the Greek contributions to the subject of simultaneous linear equations are, according to the testimony of Iamblichus (*c.* 325) in his work on Nicomachus, due to Thymaridas of Paros (*c.* 380 B.C.?). He is said to have given a rule called ἐπάνθημα (*epan'thema*, flower), which he seems to have used in solving n special types of equations, namely, $x_0 + x_1 + x_2 + \cdots + x_{n-1} = s$, $x_0 + x_1 = a_1$, $x_0 + x_2 = a_2$, $\cdots x_0 + x_{n-1} = a_{n-1}$, the method being the ordinary one of adding. Iamblichus applied the rule to other cases.[3]

Some use of simultaneous linear equations is also found in the work of Diophantus (*c.* 275), who spoke of the unknowns as the first number, the second number, and so on,[4] a method that was too cumbersome to admit of any good results.

Chinese and Japanese Methods. The subject was greatly extended by the Chinese. Using the "bamboo rods" as calculating sticks, they placed these in different squares on the table so as to represent coefficients of different unknowns, and hence

[1] The Petrie Papyrus, published by F. Ll. Griffith in 1897; the Berlin Papyrus No. 6619, published by H. Schack-Schackenburg, *Zeitschrift für ägyptische Sprache*, XXXVIII (1900), 135; and the Kahun Papyrus, also studied by Schack-Schackenburg in 1903.

[2] M. Simon, *Geschichte der Mathematik im Altertum*, pp. 41, 42 (Berlin, 1909); hereafter referred to as Simon, *Geschichte*.

[3] Heath, in R. W. Livingstone's *The Legacy of Greece*, p. 110 (Oxford, 1922).

[4] That is, ὁ πρῶτος ἀριθμός, ὁ δεύτερος ἀριθμός, and so on. *E.g.*, Book II, Prop. 17; Book IV, Prop. 37.

they needed no special symbols.[1] Indeed, we are quite justified in saying that the first definite trace that we have of simultaneous linear equations is found in China. In the *Arithmetic in Nine Sections*[2] there are various problems that require the solution of equations of the type $y = ax - b$, $y = a'x + b'$. A rule is given for the solution which amounts substantially to the following:

Arrange coefficients, $\begin{matrix} a & a' \\ b & b' \end{matrix}$

Multiply crosswise, $\begin{matrix} ab' & a'b \\ b & b' \end{matrix}$

Add, $\begin{matrix} ab' + a'b \\ b + b' \end{matrix}$

Result, $\dfrac{ab' + a'b}{a - a'}$,

and $\dfrac{b + b'}{a - a'}$.

The method of reasoning is not stated, but the work was probably done by the aid of bamboo rods to represent the coefficients.[3]

The next step of which we have evidence was taken much later by Sun-tzï, the date being uncertain but probably in the 1st century. He solved what is equivalent to the system $2x + y = 96$, $2x + 3y = 144$, and his method of elimination was substantially first to multiply the members of each equation, when necessary, by the coefficient of x in the other equation.[4]

From this time on, the solution of simultaneous linear equations was well known in China. The only improvement made upon the early methods consisted in the arrangement of the bamboo rods in such a way as to allow for a treatment of the coefficients similar to that found in the simplification of determinants. This was finally carried over to Japan and was amplified by Seki Kōwa (1683) into what may justly be called the first noteworthy advance in the theory of these forms.[5]

[1] Mikami, *China*, p. 73.

[2] *K'iu-ch'ang Suan-shu*, of uncertain date, possibly as early as 1100 B.C., and certainly pre-Christian. See Volume I, page 31.

[3] Mikami, *China*, p. 16. [4] *Ibid.*, p. 32. [5] *Ibid.*, p. 191.

Hindu Symbolism. The Hindus represented the various unknowns by the names of colors, calling them "black,"[1] "blue,"[2] "yellow,"[3] "red," and so on. They wrote the coefficients at the right of the abridged words and represented a negative term by a dot placed above the coefficient. For example,[4]

yā 5 kā 8 nī 7 rū 90
yā 7 kā 9 nī 6 rū 62

means $5x + 8y + 7z + 90 = 7x + 9y + 6z + 62$

and gives rise to $\dfrac{kā\ 1\ nī\ 1\ rū\ 28}{yā\ 2}$,

as it appears in the Colebrooke version, which means

$$-y + z + 28 = 2x.$$

Problems involving several unknowns did not possess much interest for the Arab and Persian writers, as may be seen from the algebras of al-Khowârizmî[5] and Omar Khayyam.

Early European Symbolism. The algebraists of the 16th century gave relatively little attention to simultaneous linear equations. The use of x, y, and z for unknown quantities was not suggested until the 17th century, and so it was the custom of some writers to use ordinary capital letters. For example, we find Buteo (1559) using

$$1\ A, \tfrac{1}{2}\ B, \tfrac{1}{2}\ C \lceil\ 17,$$
$$1\ B, \tfrac{1}{3}\ A, \tfrac{1}{3}\ C \lceil\ 17,$$
$$1\ C, \tfrac{1}{4}\ A, \tfrac{1}{4}\ B \lceil\ 17,$$

for what we should now write as $x + \frac{1}{2}y + \frac{1}{2}z = 17$, etc. By multiplication he reduces these to three equivalent equations in

[1] *Kālaca*, abridged to *ka* (क). See Colebrooke's translation of Bhāskara, pp. 184 n., 227. The known number was *rūpa*, abridged to *rū*.

[2] *Nīlaca*, abridged to *nī* (नी).

[3] *Pītaca*, abridged to *pī* (पी). [4] Colebrooke's translation, p. 231.

[5] But see E. Wiedemann, *Sitzungsberichte der Physikalisch-medizinischen Sozietät zu Erlangen*, 50.–51. Bd. (1920), p. 264.

which, however, the original symbolism changes slightly, the period replacing the comma to indicate addition, thus:

$$2A \cdot 1B \cdot 1C \lceil 34,$$
$$1A \cdot 3B \cdot 1C \lceil 51,$$
$$1A \cdot 1B \cdot 4C \lceil 68.$$

He then eliminates in the usual manner.[1] Gosselin, in his *De arte magna* (Paris, 1577), uses a similar arrangement.[2]

Literal Equations. The equations considered by the ancient and medieval writers were numerical. Even the early Renaissance algebraists followed the same plan, their crude symbolism allowing no other. It was not until the close of the 16th century that the literal equation made its appearance, owing largely to the influence of the new symbolism invented by Vieta and his contemporaries. For example, Adriaen van Roomen published in 1598 a commentary on the algebra of al-Khowârizmî[3] in which he distinguished between two types of equation, the *numerosa* and the *figurata*. The former was applied to problems with numerical data, while the latter resulted in general formulas.[4] Van Roomen asserts that writers on algebra up to his time used the *numerosa* method only, whereas he was the first to use the *figurata* one, although as a matter of fact Vieta seems to have preceded him. The actual dates of invention, but not of publication, are, however, obscure.

8. The Solution of Equations

Linear Equations. The earliest solutions of problems involving equations were doubtless by trial. In the time of Ahmes (*c.* 1550 B.C.), however, the methods of making the trials

[1] G. Wertheim, "Die Logistik des Johannes Buteo," *Bibl. Math.*, II (3), 213.

[2] H. Bosmans, "Le 'De arte magna' de Guillaume Gosselin," *Bibl. Math.*, VII (3), 44.

[3] H. Bosmans, "Le fragment du Commentaire d'Adrien Romain sur l'algèbre de Mahumed ben Musa El-chowârezmî," *Annales de la Société Scientifique de Bruxelles*, XXX (1906), second part, p. 266.

[4] "Differentia igitur inter has duas talis statui potest, quod figurata inveniat regulam solvendi problema propositum; numerosa vero duntaxat regulae illius exemplum."

were fairly well simplified. Thus, his equation[1] $x+\frac{1}{7}x=19$ is solved substantially as follows: Assume 7 as the number. Then, to use the form of the text,

"Once	gives	7
$\frac{1}{7}$	gives	1
$1\frac{1}{7}$	gives	8

"As many times as 8 must be multiplied to make 19, so many times must 7 be multiplied to give the required result.

"Once	gives	8
Twice	gives	16
$\frac{1}{2}$	gives	4
$\frac{1}{4}$	gives	2
$\frac{1}{8}$	gives	1
"Together, $2, \frac{1}{4}, \frac{1}{8}$	gives	19 [in which he selects the addends making 19].

"Multiply $2, \frac{1}{4}, \frac{1}{8}$ by 7 and obtain the required result.

"Once	gives	$2, \frac{1}{4}, \frac{1}{8}$
Twice	gives	$4, \frac{1}{2}, \frac{1}{4}$
4 times	gives	$9, \frac{1}{2}$
"Together, 7	gives	$16, \frac{1}{4}, \frac{3}{8}$, the result."[2]

The Greek methods are discussed later in connection with the quadratic.

The chief contribution to the solution of linear equations made by the Arab writers was the definite recognition of the application of the axioms to the transposition of terms and the reduction of an implicit function of x to an explicit one, all of which is suggested by the name given to the science by al-Khowârizmî (*c.* 825).

[1] In this discussion all equations will be given in the modern form. On the general history of solutions two of the best works for the student to consult are A. Favaro, "Notizie storico-critiche sulla Costruzione delle Equazioni," *Atti della R. Accad. di Scienze, Lettere ed Arti in Modena*, Vol. XVIII, 206 pages with extensive bibliography; Matthiessen, *Grundzüge*.

[2] For the translation I am indebted to Dr. A. B. Chace, of Providence, Rhode Island. For a slightly different version see Peet, *Rhind Papyrus*, p. 61.

False Position. To the student of today, having a good symbolism at his disposal, it seems impossible that the world should ever have been troubled by an equation like $ax + b = 0$. Such, however, was the case, and in the solution of the problem the early writers, beginning with the Egyptians, resorted to a method known until recently as the Rule of False Position. The ordinary rule as used in the Middle Ages seems to have come from India,[1] but it was the Arabs who made it known to European scholars. It is found in the works of al-Khowârizmî (*c.* 825), the Christian Arab Qosṭâ ibn Lûqâ al-Ba'albekî (died *c.* 912/13), Abû Kâmil (*c.* 900), Sinân ibn al-Fatḥ (10th century), Albanna (*c.* 1300),[2] al-Ḥaṣṣâr (*c.* 12th century),[3] and various others. The Arabs called the rule the *ḥisab al-Khataayn*[4] and so the medieval writers used such names as *elchataym*.[5] When Pacioli wrote his *Sūma* (1494) he used the term *el cataym*,[6] probably taking it from Fibonacci. Following Pacioli, the European writers of the 16th century used the same term, often with a translation into the Latin or the vernacular.[7]

[1] There is a medieval MS., published by Libri in his *Histoire*, I, 304, and possibly due to Rabbi ben Ezra. It refers to this rule, "quem Abraham compilavit et secundum librum qui Indorum dictus est composuit." See M. Steinschneider, *Abhandlungen*, III, 120; F. Woepcke, "Mémoire sur la propagation des chiffres indiens," *Journal Asiatique* (Paris, 1863), I (6), 34, 180; Matthiessen, *Grundzüge*, p. 275; C. Košt'ál, *Regula falsae positionis*, Prog., Braunau, 1886.

[2] See Volume I, page 211. For the original and a translation of his process see F. Woepcke, *Journal Asiatique*, I (6), 511.

[3] See Volume I, page 210. For a translation of his arithmetic see H. Suter, *Bibl. Math.*, II (3), 12; on this rule see page 30.

[4] Rule of Two Falses. There are various transliterations of the Arabic name.

[5] Leonardo Fibonacci, in the *Liber Abaci*, cap. XIII, under the title *De regulis elchatayn* says: "Elchataieym quidem arabice, latine duarum falsarum posicionum regula interpretatur. . . . Est enim alius modus elchataym; qui regula augmenti et diminucionis appelatur." See the Boncompagni edition, p. 318; M. Steinschneider, *Abhandlungen*, III, 122; G. Enestrom, *Bibl. Math.*, IV (3), 205.

[6] He speaks of it as a "certa regola ditta El cataym. Quale (secondo alcuni) e vocabulo arabo." Fol. 98, *v*.

[7] Thus we have " . . . per il Cataino detto alcuni modo Arabo" (Cataneo, *Le Pratiche*, Venice, 1567 ed., fol. 58); "Delle Regole del Cattaino ouero false positioni" (Pagani, 1591, p. 164); "Regola Helcataym (vocabulo Arabo) che in nostra lingua vuol dire delle false Positioni" (Tartaglia, *General Trattato*, I, fol. 238, *v*. (Venice, 1556)); "La Reigle de Faux, que les Arabes appellent la Reigle Catain" (Peletier, 1549; 1607 ed., p. 253); "La regola del Cataino" (G. Ciacchi, *Regole generali d' abbaco*, p. 278 (Florence, 1675)).

This name was not, however, the common one in the European books, and in the course of the 16th century it nearly disappeared. In general the method went by such names as Rule of False,[1] Rule of Position,[2] and Rule of False Position.[3]

Rule of Double False explained. The explanation of this rule, as related to the equation $ax + b = 0$, is as follows:

Let g_1 and g_2 be two guesses as to the value of x, and let f_1 and f_2 be the failures, that is, the values of $ag_1 + b$ and $ag_2 + b$, which would be equal to 0 if the guesses were right. Then

$$ag_1 + b = f_1 \tag{1}$$

and
$$ag_2 + b = f_2; \tag{2}$$

whence
$$a(g_1 - g_2) = f_1 - f_2. \tag{3}$$

From (1), $ag_1g_2 + bg_2 = f_1g_2$,

and from (2), $ag_1g_2 + bg_1 = f_2g_1$;

whence
$$b(g_2 - g_1) = f_1g_2 - f_2g_1. \tag{4}$$

Dividing (4) by (3), $$-\frac{b}{a} = \frac{f_1g_2 - f_2g_1}{f_1 - f_2}.$$

But, since $$-\frac{b}{a} = x,$$

we have here a rule for finding the value of x.[4]

[1] "La Reigle de Faux" (Trenchant, 1566; 1578 ed., p. 213); "Falsy" (Van der Schuere, 1600, p. 185); "Regula Falsi" (Coutereels, 1690 edition of the *Cyffer-Boeck*, p. 541).

[2] "Auch Regula Positionum genant" (Suevus, 1593, p. 377); "Reigle de Faux, mesmes d'une Position" (Peletier, 1549; 1607 ed., p. 269).

[3] "Rule of falshoode, or false positions" (Baker, 1568; 1580 ed., fol. 181); "False Positie" and "Fausse Position" (Coutereels, Dutch-French ed., 1631, p. 329); "Valsche Positie" (Eversdyck's Coutereels, 1658 ed., p. 360); "Reghel der Valsches Positien" (Wilkens, 1669 ed., p. 353).

[4] The formula is more elegantly derived by taking the eliminant of

$$\begin{aligned} ax + b + 0 &= 0 \\ ag_1 + b - f_1 &= 0 \\ ag_2 + b - f_2 &= 0, \end{aligned}$$

which is
$$\begin{vmatrix} x & 1 & 0 \\ g_1 & 1 & -f_1 \\ g_2 & 1 & -f_2 \end{vmatrix} = 0,$$

by the expansion of which the result at once appears.

Suppose, for example, that

$$5x - 10 = 0.$$

Make two guesses as to the value of x, say $g_1 = 3$ and $g_2 = 1$.

Then $$5 \cdot 3 - 10 = 5 = f_1,$$

and $$5 \cdot 1 - 10 = -5 = f_2.$$

Then $$x = \frac{f_1 g_2 - f_2 g_1}{f_1 - f_2} = \frac{5 \cdot 1 - (-5) \cdot 3}{5 - (-5)} = \frac{20}{10} = 2.$$

Awkward as this seems, the rule was used for many centuries, a witness to the need for and value of a good symbolism. We have here placed two false quantities in the problem, and from these we have been able to find the true result.

Recorde's Rule in Verse. From the above formula for x it will be possible to interpret the doggerel rule given by Robert Recorde in his *Ground of Artes* (*c.* 1542):


Gesse at this woorke as happe doth leade.
By chaunce to truthe you may procede.
And firste woorke by the question,
Although no truthe therein be don.
Suche falsehode is so good a grounde,
That truth by it will soone be founde.
From many bate to many mo,
From to fewe take to fewe also.
With to much ioyne to fewe againe,
To to fewe adde to manye plaine.
In crossewaies multiplye contrary kinde,
All truthe by falsehode for to fynde.[1]

Recorde thought highly of the rule, and it was appreciated by writers generally until the 19th century.[2]


[1] *Ground of Artes*, 1558 ed., fol. Z 4.

[2] Thus Thierfelder (1587, p. 226) says: "Für allen Regeln der gantzen Arithmetic (ohn allein die Regel Cosz auzgenommen) ist sie die Kunstreichste / weytgegreifflichste vñ schönste"; and Peletier (1549; 1607 ed., p. 269) remarks: "Gemme Phrissien a inuenté l'artifice de soudre par la Reigle de Faux, mesmes d'une Position, grand' partie des exemples subjects à l'Algebre." Even as late as 1884, in the *Instruction für den Unterricht an den Gymnasien in Österreich* (Vienna, 1884, p. 315), the rule is recommended.

Method of the Scales. The Arabs modified the rule by what they called the Method of the Scales,[1] a name derived from the following figure, used in the solution:



Suppose, for example, that we wish to solve the equation $x + \frac{2}{3}x + 1 = 10$, a problem set by Behâ Eddîn (*c.* 1600). We may make as our guesses $g_1 = 9$, whence $f_1 = 6$; $g_2 = 6$, whence $f_2 = 1$. Then place the figures thus:



The lines now aid the eye to write the result according to the rule already set forth, as follows:[2]

$$x = \frac{6 \cdot 6 - 1 \cdot 9}{6 - 1} = \frac{27}{5} = 5\tfrac{2}{5}.$$

Rule of Single False. Thus far we have considered the Rule of Double False, where a double guess was made, but there was also a modification of the method known as the Rule of Single False.[3] Albanna (*c.* 1300) gives the latter in the form of a rule which, worked out in modern symbols, is as follows: Given that

$$ax + b = 0.$$

Make a guess, g, for the value of x, the failure being f; that is,

$$ag + b = f.$$

[1] *'Alm bi'l kaffatain.* This name was translated into Latin as the *Regula lancium* or *Regula bilancis.*

[2] For variations of the method see Matthiessen, *Grundzüge*, p. 278; for the Arab proof by geometry see *ibid.*, p. 281.

[3] Tartaglia called it "Position Sempia," as distinct from "Position Doppia" (*General Trattato*, I, fol. 239, *v.*, and 266, *r.*); the Spanish had rules "De vna falsa posicion" and "De dos falsas posiciones" (Santa-Cruz, 1594; 1643 ed., fols. 210, 212); Clavius (1586 Italian ed., pp. 195, 203) has "Regola del falso di semplice positione" and "di doppia positione"; and Chuquet (1484, MS., fols. 32, 42) has "De la Rigle de vne posicion" and "de deux posicions."

Then to obtain the required rule we may proceed as follows:

$$a = \frac{f}{g - x},$$

whence
$$\frac{fx}{g - x} + b = 0$$

and
$$x = \frac{gb}{b - f} = \frac{g(ag) - gf}{ag} = \frac{g(f - b) - gf}{f - b},$$

this last indicating the rule used.

For example, in the equation $\frac{1}{5}x + \frac{1}{6}x = 20$, if we take $g = 30$, we have

$$\tfrac{1}{5} \cdot 30 + \tfrac{1}{6} \cdot 30 = 11,$$

which is 9 too small, whence $f = -9$. Then

$$x = \frac{30(-9 + 20) - 30(-9)}{-9 + 20} = 54\tfrac{6}{11}.$$

Apologies for the Name of Rule of False. The name "Rule of False" was thought to demand an apology in a science whose function it is to find the truth, and various writers made an effort to give it. Thus Humphrey Baker (1568) says:

> The Rule of falsehoode is so named not for that it teacheth anye deceyte or falsehoode, but that by fayned numbers taken at all aduentures, it teacheth to finde out the true number that is demaunded, and this of all the vulgar Rules which are in practise) is ỹ most excellence.[1]

Besides the "Rule of False" the method was also called the "Rule of Increase and Diminution,"[2] from the fact that the error is sometimes positive and sometimes negative. Indeed, as already stated, in the 16th century the symbols + and − were much more frequently used in this connection than as symbols of operation.

[1] 1580 ed., fol. 181. Similar excuse is offered by Thierfelder (1587, p. 225): "Darum̃ nicht dasz sie falsch oder vnrecht sey"; by Apianus (1527): "Vnd heisst nit darum falsi dass sie falsch vnd unrecht wehr, sunder, dass sie auss zweyen falschen vnd vnwahrhaftigen zalen, vnd zweyen lügen die wahrhaftige vnd begehrte zal finden lernt." Like explanations are given by many other writers.

[2] "Regula augmenti et decrementi" or "diminutionis."

Regula Infusa. Rabbi ben Ezra (*c.* 1140) tells us of a substitution method due to another Hebrew writer, Job ben Salomon, of unknown date, which was called in Latin translation the *Regula infusa.*[1] This may be illustrated as follows:

Given $m(ax+b)+c=0,$

let $ax+b=y,$

and then $my+c=0,$

whence $y=-c/m,$

and so $ax+b=-c/m,$

which can now be solved. Rabbi ben Ezra illustrates this by taking

$$x-\tfrac{1}{3}x-4-\tfrac{1}{4}(x-\tfrac{1}{3}x-4)=20$$

and letting $x-\frac{1}{3}x-4=y;$

whence $y-\frac{1}{4}y=20$

and $y=26\frac{2}{3},$

and so $x-\frac{1}{3}x-4=26\frac{2}{3},$

which can now be solved.[2] Although the method is very artificial, it is occasionally found in the algebras of today, especially in connection with radical equations.

Classification of Equations. Our present method of classifying equations according to their degree is a modern one. The first noteworthy attempt at a systematic classification is found in the algebra of Omar Khayyam (*c.* 1100), but the classification there given is not our present one. Omar considers equations of the first three degrees as either simple or compound. The simple equations are of the type $r=x$, $r=x^2$, $r=x^3$, $ax=x^2$, $ax=x^3$, $ax^2=x^3$. Compound equations are first classified as trinomials, and these include the following twelve forms: (1) $x^2+bx=c$, $x^2+c=bx$, $bx+c=x^2$; (2) $x^3+bx^2=cx$,

[1] ". . . secundum regulam que vocatur infusa. Et ipsa est regula Job, filii Salomonis." Libri, *Histoire*, 1838 ed., I, 312. There is some doubt, however, as to whether Libri was right in referring this work to Rabbi ben Ezra.

[2] Matthiessen, *Grundzüge*, p. 272.

$x^3 + cx = bx^2$, $cx + bx^2 = x^3$, $x^3 + cx = d$, $x^3 + d = cx$, $cx + d = x^3$; $x^3 + bx^2 = d$, $x^3 + d = bx^2$, $bx^2 + d = x^3$. They are then classified as quadrinomials, as follows: (3) $x^3 + bx^2 + cx = d$, $x^3 + bx^2 + d = cx$; (4) $x^3 + bx^2 = cx + d$, $x^3 + cx = bx^2 + d$, $x^3 + d = bx^2 + cx$. Of this early plan of classifying equations by the number of terms we still have a trace in our chapter on binomial equations.

Classification according to Degree. Such was the general method of classifying equations, naturally with variations in details, until after books began to be printed. Pacioli (1494), for example, has a similar system.[1] It was not until about the beginning of the 17th century that the classification according to degree, with a recognition that a literal coefficient might be either positive or negative, was generally employed, and this was due in a large measure to the influence of such writers as Stevin (1585), Vieta (*c.* 1590), Girard (1629), Harriot (1631, posthumous), Oughtred (1631), and Descartes (1637). In particular, Descartes set forth in his *Géométrie* the idea of the degree of an equation, or, as he says, of the dimensions of an equation,[2] reserving the word "degree" for use with respect to lines.[3]

Quadratic Equations. The first known solution of a quadratic equation is the one given in the Berlin Papyrus mentioned on page 432. The problem reduces to solving the equations

$$x^2 + y^2 = 100, \qquad y = \tfrac{3}{4}x,$$

and the solution is substantially as follows:

Make a square whose side is 1 and another whose side is $\frac{3}{4}$. Square $\frac{3}{4}$, giving $\frac{9}{16}$. Add the squares, giving $2\frac{5}{16}$, the square root of which is $\frac{5}{4}$. The square root of 100 is 10. Divide 10 by $\frac{5}{4}$, giving 8, and $\frac{3}{4}$ of 8 is 6. Then

$$8^2 + 6^2 = 100 \quad \text{and} \quad 6 = \tfrac{3}{4} \text{ of } 8,$$

[1] *Sūma*, 1494 ed., fol. 145 seq.

[2] "Sçachez donc qu'en chaque Equation, autant que la quantité inconnuë a de dimensions, autant peut-il y avoir de diverses racines, c'est à dire de valeurs de cette quantité" (1705 ed., p. 106). He then speaks of "$x^3 - 9xx + 26x - 24 \propto 0$, . . . en laquelle x ayant trois dimensions a aussi trois valeurs qui sont 2, 3, & 4."

[3] ". . . distingué divers degrez entre ces lignes" (*ibid.*, p. 27).

so that the roots of the two implied equations are 6 and 8. The solution is therefore a simple case of false position.[1]

The Greeks were able to solve the quadratic equation by geometric methods. As already stated, Euclid (*c.* 300 B.C.) has in his *Data* three problems involving quadratics. Of these the first (Prob. 84) is as follows:

> If two straight lines include a given area in a given angle and the excess of the greater over the less is given, then each of them is given.

Expressed in algebraic form with reference to the rectangle, if $xy = k^2$ and $x - y = a$, then x and y can be found. Euclid solves the problem geometrically.[2] He also gives in the *Elements* such geometric problems as the following:

> To cut a given straight line so that the rectangle contained by the whole and one of the segments shall be equal to the square on the remaining segment.[3]

This may be represented algebraically by the equation $a(a - x) = x^2$ or by $x^2 + ax = a^2$.

Quadratics among the Hindus. It is possible that the altar constructon of the Hindus involved the solution of the equation $ax^2 + bx = c$, and this may date from the *Śulvasūtra* period (roughly speaking, say 500 B.C.); but whether or not this is the case, we have no record of the method of solution.[4]

When we come to the time of Āryabhaṭa (*c.* 510), we find a rule, relating to the sum of a geometric series, which shows that the solution of the equation $ax^2 + bx + c = 0$ was known, but we have no rule for the solution of the equation itself.[5]

It should be repeated, however, that up to the 17th century an equation of the type $x^2 + px = q$, for example, was looked

[1] Schack-Schackenburg, *Zeitschrift für ägyptische Sprache*, XXXVIII, 135; XL, 65. See also Cantor, *Geschichte*, I (3), 95, and Simon, *Geschichte*, p. 41.

[2] The other two have already been given on page 381.

[3] *Elements*, II, 11. See also VI, 28, 29.

[4] G. Milhaud, "La Géométrie d'Apastamba," in the *Revue générale des sciences*, XXI, 512–520.

[5] The rule for the summation is No. XX in Rodet's *Leçons de Calcul d'Âryabhaṭa*, pp. 13, 33 (Paris, 1879). In all such cases the possibility of the younger Āryabhaṭa must be considered.

upon as distinct from one of the type $x^2 - px = q$; the idea that p might be either positive or negative did not occur to algebraists until some time after the invention of a fairly good symbolism. This accounts for the special rules for different types that are found in the Middle Ages and the early Renaissance.

Brahmagupta's Rule. Brahmagupta (*c.* 628) gave a definite rule for the quadratic. For example, he gave the equation

ya v 1 *ya* 1ȯ
ru 9̇,

that is, $x^2 - 10x = -9$,[1] with the solution substantially as follows:

Here absolute number (9̇) multiplied by (1) the [coefficient of the] square (9̇), and added to the square of half the [coefficient of the] middle term, namely, 25, makes 16; of which the square root 4, less half the [coefficient of the] unknown (5̇), is 9; and divided by the [coefficient of the] square (1) yields the value of the unknown 9.

Expressed in modern symbols,

$$x = \frac{\sqrt{-9 \cdot 1 + (-5)^2} - (-5)}{1} = 9.$$

Mahāvīra's Rule. Mahāvīra (*c.* 850) gave no rule for the quadratic, but he proposed a problem involving the equation

$$c + \frac{c}{3} + 3 \cdot \frac{c^2}{36} = m,$$

adding the following statement:

In relation to the combined sum [of the three quantities] as multiplied by 12, the quantity thrown in so as to be added is 64. Of this [second] sum the square root diminished by the square root of the quantity thrown in gives rise to the measure . . .

Expressed in modern symbols, this means that

$$c = \sqrt{12m + 64} - \sqrt{64},$$

which shows that Mahāvīra had substantially the modern rule for finding the positive root of a quadratic.[2]

[1] Colebrooke, p. 347, his transliteration being followed.
[2] See his work, p. 192.

The Hindu Rule. Śrīdhara (*c.* 1025) was the first, so far as known, to give the so-called Hindu Rule for quadratics. He is quoted by Bhāskara (*c.* 1150) as saying:


Multiply both sides of the equation by a number equal to four times the [coefficient of the] square, and add to them a number equal to the square of the original [coefficient of the] unknown quantity. [Then extract the root.][1]

This rule, although stated by Bhāskara, is not the first one given by him. He begins by saying:

[Its re-solution consists in] the elimination of the middle term, as the teachers of the science denominate it. . . . On this subject the following rule is delivered. . . . When a square and other [term] of the unknown is involved in the remainder; then after multiplying both sides of the equation by an assumed quantity, something is to be added to them, so as the side may give a square-root. Let the root of the absolute number again be made equal to the root of the unknown; the value of the unknown is found from that equation.

It will be observed that this is simply a more general form of Śrīdhara's rule. The method has been the subject of much discussion by the various commentators on Bhāskara.[2]

Al-Khowârizmî's Rules. Al-Khowârizmî (*c.* 825) used two general methods in solving the quadratic of the form $x^2 + px = q$, both based upon Greek models. Given $x^2 + 10x = 39$, he constructed a square as here shown. Then the unshaded part is $x^2 + px$, and is therefore equal to q. In order to make it a square we must add the four shaded squares, each of which is $(\frac{1}{4}p)^2$ and the sum of



[1] *Vîja-Gaṇita*, p. 209. That is, given $ax^2 + bx = c$, we have first $4a^2x^2 + 4abx = 4ac$. Then $4a^2x^2 + 4abx + b^2 = b^2 + 4ac$, whence

$$2ax + b = \sqrt{b^2 + 4ac},$$

the negative root being neglected. The purpose of the multiplication by $4a$ was to avoid fractions.

[2] See the *Vîja-Gaṇita*, pp. 207–209.

which is $\frac{1}{4}p^2$, which in this case is 25. Since $25+39=64$, we have

$$x+\tfrac{1}{2}p=8;$$

whence

$$x+5=8$$

and

$$x=3.$$

His statement is as follows:

You halve the number of the roots, which in the present instance yields five. This you multiply by itself; the product is twenty-five. Add this to thirty-nine; the sum is sixty-four. Now take the root of this, which is eight, and subtract from it half the number of the roots, which is five; the remainder is three. This is the root of the square which you sought for; the square itself is nine.[1]

The negative root was neglected, as was regularly the case until modern times.

His second method was similar to our common one. In the figure the unshaded part is x^2+px, and he adds the square of $\frac{1}{2}p$. He then has

$$x^2+px+\tfrac{1}{4}p^2=\tfrac{1}{4}p^2+q,$$

whence

$$x=\sqrt{\tfrac{1}{4}p^2+q}-\tfrac{1}{2}p,$$


of which he takes only the positive root.[2]

Al-Khowârizmî also considers other forms, his solution[3] of the type $x^2+q=px$ being based upon the identity

$$(\tfrac{1}{2}p)^2-(\tfrac{1}{2}p-x)^2=x(p-x)=px-x^2=q,$$

from which it follows that

$$x=\tfrac{1}{2}p+\sqrt{\tfrac{1}{4}p^2-q}.$$



Omar Khayyam's Rule. Omar Khayyam's rule (*c.* 1100) for solving the quadratic $x^2+px=q$ is as follows:

Multiply half of the root by itself; add the product to the number and from the square root of this sum subtract half the root. The remainder is the root of the square.[4]

[1] Rosen ed., p. 8.

[2] For a discussion of his methods see Matthiessen, *Grundzüge*, p. 299.

[3] For discussion and for the geometric proof see Rosen's edition, p. 16; Matthiessen, *Grundzüge*, p. 304; Libri, *Histoire*, I, 236.

[4] That is, $x=\sqrt{\frac{1}{4}p^2+q}-\frac{1}{2}p$. By "half the root" is meant $\frac{1}{2}p$, and by "the number" is meant q. He used the equation $x^2+10x=39$, which was the one

He also gave rules for other types, that for $x^2 + q = px$ being based upon the identity

$$x(p - x) + (x - \tfrac{1}{2}p)^2 = (\tfrac{1}{2}p)^2,$$

and that for $px + q = x^2$ upon the identity[1]

$$x(x - p) + (\tfrac{1}{2}p)^2 = (x - \tfrac{1}{2}p)^2.$$

Chinese Work in Quadratics. The Chinese gave some attention to quadratic equations in the Middle Ages, including those of the form

$$x^2 - sx - \left(\frac{s^2 - H^2}{2}\right) = 0,$$

but how far they were original in their work has not yet been scientifically determined.[2]

With respect to the quadratic equation the medieval algebraists added nothing of importance to the work of the Arabic writers from whom they derived their inspiration, and the Renaissance algebraists did little except in their improvement of the symbolism. It was not until the close of the 16th century that the next noteworthy contribution was made.

Harriot treats of Equations by Factoring. The first important treatment of the solution of quadratic and other equations by factoring is found in Harriot's *Artis Analyticae Praxis* (1631). He takes as his first case the equation

$$aa - ba + ca = + bc$$

and writes it in the form

$$\begin{array}{l|l} a - b & = aa - ba \\ \underline{a + c} & \quad + ca - bc, \end{array}$$

also used by al-Khowârizmî and was apparently a favorite problem of the schools. He also considered the arithmetically impossible solutions. For a discussion of his methods and proofs see Woepcke's translation, p. 17; Matthiessen, *Grundzüge*, p. 301.

[1] See Woepcke's translation, pp. 20, 23; Matthiessen, *Grundzüge*, pp. 305, 309.

[2] L. Vanhée, in *T'oung-pao*, XIII, 291; XII, 559; XV, 111.

where the first member stands for $(a - b)\ (a + c)$ and the equation becomes $(a - b)\ (a + c) = 0$. From this fact he finds that $b = a$.[1]

In a similar way the equation

$$aaa + baa - cda = + bcd$$

is factored into $(a + b)\ (aa - cd) = 0$, and the solution is given that $aa = cd$.[2]

Vieta advances the Theory. In the work of Vieta the analytic methods replaced the geometric, and his solutions of the quadratic equation were therefore a distinct advance upon those of his predecessors. For example, to solve the equation[3] $x^2 + ax + b = 0$ he placed $u + z$ for x. He then had

$$u^2 + (2z + a)u + (z^2 + az + b) = 0.$$

He now let $2z + a = 0$, whence $z = -\frac{1}{2}a$, and this gave

$$u^2 - \tfrac{1}{4}(a^2 - 4b) = 0,$$

$$u = \pm \tfrac{1}{2}\sqrt{a^2 - 4b},$$

and

$$x = u + z = -\tfrac{1}{2}a \pm \tfrac{1}{2}\sqrt{a^2 - 4b}.$$

[1] "Nam si ponatur $a = b$ erit $a - b = 0$," p. 16. The relation $a + c = 0$ is neglected.

[2] P. 19. The relation $a = -b$ is neglected.

[3] The symbolism used here is, of course, modern. Vieta's own solution is as follows:

"Si A quad. + B 2 in A, aequatur Z plano. A + B esto E. Igitur E quad. aequabitur Z plano + B quad.

"Consectarium. Itaque $\sqrt{Z \text{ plani} + B \text{ quad.}} - B$ fit A, de qua primum quaerebatur.

"Sit B 1. Z planum 20. A 1 N. 1 Q + 2 N, aequatur 20. et fit 1 N $\sqrt{21} - 1$."

That is, if $A^2 + 2BA = Z$, we may represent this in modern form as $x^2 + 2ax = b$, where $A = x$, $B = a$, $Z = b$.

Let $A + B = E$, that is, let $x + a = u$.

It follows that $u^2 = x^2 + 2ax + a^2 = b + a^2$, and so $x = \sqrt{b + a^2} - a$.

In particular, he says, let $B = 1$ and $Z = 20$. The equation is then $x^2 + 2x = 20$, whence $x = \sqrt{21} - 1$.

He has similar solutions for the following:

"Si A quad. — B in A 2, aequatur Z plano,"

and "Si D 2 in A — A quad., aequatur Z plano,"

showing that at this time in his work he had not grasped the idea of such a general quadratic equation as $x^2 + a_1x + a_2 = 0$. In his *De numerosa potestatum*

Modern Methods. Of the modern methods[1] for obtaining the formula for the solution of the quadratic, interesting chiefly from the standpoint of theory, a single one may be mentioned. This method uses determinants and is due to Euler and Bézout, but was improved by Sylvester (1840) and Hesse (1844).

Given $$x^2+px+q=0,$$

let $$x=u+z;$$

whence $$x^2=(u+z)x$$

Then $$x^3+px^2+qx=0,$$

$$x^2-(u+z)x=0,$$

and $$x^3-(u+z)x^2=0;$$

whence $$\begin{vmatrix} 1 & p & q \\ 0 & 1 & -(u+z) \\ 1 & -(u+z) & 0 \end{vmatrix}=0.$$

Expanding, $$-p(u+z)-(u+z)^2-q=0,$$

and hence $$u^2+(2z+p)u+(z^2+pz+q)=0.$$

Letting $$2z+p=0,$$

we find that $$u=\pm\tfrac{1}{2}\sqrt{p^2-4q},$$

and $$x=-\frac{p}{2}\pm\tfrac{1}{2}\sqrt{p^2-4q}.$$

Simultaneous Quadratic Equations. Problems involving the combination of a linear and a quadratic equation were, as we have seen, familiar to the Egyptians, and the Greeks were fully able to apply their geometry to such cases. The algebraic treatment of two quadratics was not seriously considered, however, until it was taken up by Diophantus (*c.* 275) for indeterminate forms. He speaks of equations like

$$y^2=ax^2+bx+c$$

and $$z^2=a'x^2+b'x+c'$$

. . . *resolutione tractatus* (Paris, 1600), however, he uses the terms "affected" and "pure" with respect to quadratic equations. See Volume I, page 311. See also his *De aequationum recognitione et emendatione libri duo*, Tract. II, cap. i (Paris, 1615); Matthiessen, *Grundzüge*, p. 311.

[1] For a list of modern methods consult Matthiessen, *Grundzüge*, p. 315 seq.

as "double equations."[1] Among his more difficult equations of this type is the pair

$$x^2 + x - 1 = u^2,$$
$$x^2 - 1 = w^2,$$

which Diophantus expresses as follows:

To find three numbers such that their solid content minus any one gives a square.[2]

The subject never interested the medieval writers particularly, and not until the 17th century do we find much attention paid to it. By that time the symbolism was such that the only question involved was that of stating the cases in which a solution is possible.

Indeterminate Quadratic Equations. The study of indeterminate quadratic equations begins with such cases as $x^2 + y^2 = z^2$. The finding of formulas for these sides of a Pythagorean Triangle occupied the attention of various Greek writers. Proclus (*c.* 460) tells us that Pythagoras (*c.* 540 B.C.) himself gave a rule, and tradition says that it was, as expressed in modern symbols,

$$n^2 + \left(\frac{n^2-1}{2}\right)^2 = \left(\frac{n^2+1}{2}\right)^2$$

where n is an odd number. Plato (*c.* 380 B.C.) gave the rule

$$(2n)^2 + (n^2-1)^2 = (n^2+1)^2,$$

which, like the one attributed to Pythagoras, is connected with Euclid's proposition[3] to the effect that

$$\left(\frac{a+b}{2}\right)^2 - \left(\frac{a-b}{2}\right)^2 = ab,$$

a relation that forms the basis of the theory of quarter squares.

[1] Διπλοϊσότης, διπλῆ ἰσότης, διπλῆ ἴσωσις. See Heath, *Diophantus*, 2d ed., p. 73.

[2] Book IV, 23. That is, the first number is x, the second is 1, and the third is $x + 1$, the "solid content" being $x \cdot 1 \cdot (x + 1)$. The results are $\frac{1}{8}\frac{7}{8}$, 1, and $\frac{2}{8}\frac{5}{8}$. For further explanation see Heath, *loc. cit.*, p. 184.

[3] Heath, *Diophantus*, 2d ed., pp. 116, 242 n.

Diophantus on Indeterminate Equations. It was Diophantus, however, who may properly be called the father of the study of indeterminate equations, which were generally limited in his *Arithmetica* to quadratic types. With these equations the object was to obtain rational results, while with indeterminate equations of the first degree the object was usually to obtain integral results. The problem proposed by Diophantus is that of solving either one or two equations of the form

$$Ax^2 + Bx + C = y^2.$$

His simpler types may be represented by the following:

To add the same [required] number to two given numbers so as to make each of them a square.[1]

One of the more difficult problems is as follows:

To find three numbers such that their sum is a square and the sum of any pair is a square.[2]

Pell Equation. One of the most famous indeterminate quadratic equations is of the form

$$Ay^2 + 1 = x^2.$$

This form is commonly attributed to John Pell (1668) but is really due to Fermat (*c.* 1640) and Lord Brouncker[3] (1657). The problem itself is apparently much older than this, however, for it seems involved in various ancient approximations to the square roots of numbers. Thus the Greek approximation $\frac{7}{5}$ for the ratio of the diagonal to the side of a square goes back to Plato's time at least, and 7 and 5 are the roots of the equation

$$x^2 - 2y^2 = -1.$$

[1] III, 11. That is, if the given numbers are 2 and 3, then $x + 2$ and $x + 3$ must both be squares. He finds that $x = 8\frac{7}{4}$.

[2] III, 6. His results are 80, 320, 41. For solution see Heath, *Diophantus*, 2d ed., pp. 68, 158.

[3] E. E. Whitford, *The Pell Equation*, New York, 1912; H. Konen, *Geschichte der Gleichung* $t^2 - Du^2 = 1$, Leipzig, 1901; G. Wertheim, "Ueber den Ursprung des Ausdruckes 'Pellsche Gleichung,'" *Bibl. Math.*, II (3), 360; Heath, *Diophantus*, 2d ed., p. 286.

Theon of Smyrna (*c.* 125) considered a relation that would now be written as the equation

$$x^2 - 2y^2 = \pm 1,$$

carrying his computations as far as the case of

$$17^2 - 2 \cdot 12^2 = 1$$

and stating a rule for finding the solutions.

The special case of the Cattle Problem, doubtfully attributed to Archimedes, requires the number of bulls of each of four colors, white (W), blue (B), yellow (Y), and piebald (P), and the number of cows of the same colors (w, b, y, p) such that $B = (\frac{1}{4} + \frac{1}{5})(Y + P)$, $P = (\frac{1}{6} + \frac{1}{7})(W + Y)$, $w = (\frac{1}{3} + \frac{1}{4})(B + b)$, $b = (\frac{1}{4} + \frac{1}{5})(P + p)$, $p = (\frac{1}{5} + \frac{1}{6})(Y + y)$, and $y = (\frac{1}{6} + \frac{1}{7})(W + w)$. Reduced to a single equation, the problem involves the solution of the indeterminate quadratic equation

$$x^2 - 4{,}729{,}494y^2 = 1,$$

and the number of yellow bulls, for example, has 68,848 periods of three figures each.[1]

The general problem may have been discussed in the lost books of Diophantus,[2] perhaps in the form

$$x^2 - Ay^2 = 1,$$

and its equivalent is clearly stated in the works of Brahmagupta[3] (*c.* 628).

Fermat (*c.* 1640) was the first to state that the equation $x^2 - Ay^2 = 1$, where A is a non-square integer, has an unlimited number of integral solutions,[4] and from that time on the problem attracted the attention of various scholars, among the most prominent being Euler (1730), who stated that the solu-

[1] Heath, *History*, II, 97; Whitford, *loc. cit.*, p. 20, with bibliography.

[2] P. Tannery, "L'arithmétique des Grecs dans Pappus," in the *Mémoires de la Soc. des sci. de Bordeaux*, III (2), 370.

[3] Colebrooke ed., p. 363.

[4] *Œuvres*, ed. Tannery and Henry, II, 334 (Paris, 1894); Whitford, *loc. cit.*, p. 46.

tion of the equation $ax^2 + bx + c = y^2$ requires the solution of the equation $x^2 - Ay^2 = 1$.[1] It was he who, through an error, gave to the general type the name of the Pell Equation.

Cubic Equation. The oldest known cubic equation of the form $x^3 = k$ is possibly due to Menæchmus (*c.* 350 B.C.), although tables of cubes had been worked out by the Babylonians two thousand years earlier. It had been recognized since the time of Hippocrates (*c.* 460 B.C.) that the solution of the problem of the duplication of the cube depended on the finding of two mean proportionals between two given lines. Algebraically this means the finding of x and y in the equations

$$\frac{a}{x} = \frac{x}{y} = \frac{y}{b}.$$

From these relations it is evident that

$$y^2 = bx \text{ (a parabola)}$$

and $$xy = ab \text{ (an equilateral hyperbola)};$$

whence $$y^3 = ab^2 \text{ (a cubic equation)}.$$

Menæchmus is said to have solved the cubic by finding the intersection of the two conics. If $b = 2a$, then $y^3 = 2a^3$, and the problem becomes the well-known one of the duplication of the cube, which interested so many Greek writers.[2]

The next reference to the cubic among the Greeks is in a certain problem of Archimedes, to cut a sphere by a plane so that the two segments shall have a given ratio.[3] This reduces to the proportion

$$\frac{c - x}{b} = \frac{c^2}{x^2}$$

and to the equation $x^3 + c^2b = cx^2$.

[1] P. H. von Fuss, *Correspondance mathématique et physique de quelques célèbres géomètres du XVIIIième siècle*, I, 37 (Petrograd, 1843).

[2] For a partial list of these writers see Woepcke, translation of Omar Khayyam, p. xiij. The reference to Menæchmus is not certain.

[3] *De sphaera et cylindro*, Lib. II. See also Heath's *Archimedes*, chap. vi.

Eutocius (*c.* 560) tells us that Archimedes solved the problem by finding the intersection of two conics, namely,

$$x^2 = \frac{a^2}{c} y \text{ (a parabola)}$$

and $$y(c - x) = bc \text{ (a hyperbola)}.$$

Diophantus solved a single cubic equation, $x^3 + x = 4x^2 + 4$. This equation arises in connection with the following problem:

To find a right-angled triangle such that the area added to the hypotenuse gives a square, while the perimeter is a cube.[1]

His method is not given, the statement, expressed in modern language, being that x "is found to be" 4. Possibly Diophantus saw that $x(x^2 + 1) = 4(x^2 + 1)$; whence $x = 4$.

The Cubic among the Arabs and Persians. Nothing more is known of the cubic equation among the Greeks, but the problem of Archimedes was taken up by the Arabs and Persians in the 9th century. In a commentary on Archimedes, Almâhânî (*c.* 860) considered the question, but so far as known he contributed nothing new. He brought the problem into such prominence, however, that the equation $x^3 + a^2b = cx^2$ was known among the Arab and Persian writers as Almâhânî's equation.[2]

One of his contemporaries, Ṯâbit ibn Qorra (*c.* 870), considered special cases of cubic equations, as in the duplication of the cube. These equations he solved by geometric methods, but he was unable to contribute to the general algebraic theory.

A little later Abû Jaʿfar al-Khâzin (*c.* 960), a native of Khorâsân, considered the problem and, as Omar Khayyam tells us, "solved the equation by the aid of conic sections."[3]

The last of the Arabs to give any particular attention to the solution was Alhazen[4] (*c.* 1000). Omar Khayyam[5] refers to

[1] Bk. VI, prob. 17. See also Heath's *Diophantus*, 2d ed., p. 66.

[2] Matthiessen, *Grundzüge*, p. 367; Cantor, *Geschichte*, I, chap. xxxv.

[3] Woepcke translation, p. 3.

[4] Al-Ḥasan ibn al-Ḥasan ibn al-Ḥaiṯam.

[5] Woepcke's translation, p. 73, with discussion; Matthiessen, *Grundzüge*, p. 367.

his method. Alhazen solved the equation by finding the intersection of

$$x^2 = ay \text{ (a parabola)}$$

and

$$y(c - x) = ab \text{ (a hyperbola)},$$

a method not unlike the one attributed to Archimedes.

The last of the Persian writers to consider the cubic equation with any noteworthy success was Omar Khayyam[1] (*c.* 1100). In his list of equations he specified thirteen forms of the cubic that had positive roots, this being a decided advance in the general theory. He solved equations of the type $x^3 + b^2x = b^2c$ by finding the intersection of the conics $x^2 = by$ and $y^2 = x(c - x)$; of the type $x^3 + ax^2 = c^3$ by finding the intersection of $xy = c^2$ and $y^2 = c(x + a)$; and of the type $x^3 \pm ax^2 + b^2x = b^2c$ by finding the intersection of $y^2 = (x \pm a)(c - x)$ and $x(b \pm y) = bc$.

It is said, but without proof from the sources, that Omar Khayyam stated that it was impossible to solve in positive integers the equation $x^3 + y^3 = z^3$, the simplest of the family of equations of the type $x^n + y^n = z^n$ with which Fermat's name is connected.

In general it may be said that the Arab writers believed that the cubic equation was impossible of solution.[2]

Chinese and Hindu Interest in the Cubic. The Chinese algebraists did nothing worthy of note with the general cubic equation. Their interests lay in applied problems, and these all led to numerical equations. The numerical cubic first appears in a work by Wang Hs'iao-t'ung, about 625.[3]

He gave the following problem:

> There is a right-angled triangle the product of the sides of which is $706\frac{1}{8}$ and the hypotenuse of which is greater than one side by $36\frac{9}{10}$. Find the lengths of the three sides.

Wang used a numerical equation of the form $x^3 + ax^2 - b = 0$ and stated the answer incorrectly as $14\frac{7}{20}$, $49\frac{1}{5}$, and $51\frac{1}{4}$, although there is doubt as to the validity of the copy.

[1] Ball, *Hist. of Math.*, 6th ed., p. 159; Woepcke's translation, p. 25 seq.
[2] Cantor, *Geschichte*, I (2), 736.
[3] In the *Ch'i-ku Suan-king*. See Mikami, *China*, p. 54.

Various later Chinese algebraists treated of numerical equations, but it was not until the 18th century, when European influences were powerful, that any attempt was made by them to classify equations of the third degree. In a work prepared under the direction of Emperor Kanghy, who ruled China from 1662 to 1722, nine types are given:

$$x^3 \pm bx = c \qquad x^3 \pm ax^2 = c$$
$$x^3 \pm ax^2 \pm bx = c \qquad -x^3 + ax^2 = c,$$

but in every case the solution is numerical and only a single positive root is given.[1]

The Hindus paid little attention to cubic equations except as they entered into relatively simple numerical problems relating to mensuration. Bhāskara (*c.* 1150) gave one example,

$$x^3 + 12x = 6x^2 + 35,$$

the root being 5,[2] but such a result is easily found by trial, the equation being made for this purpose.

Medieval Interest in the Cubic. In the Middle Ages various sporadic attempts were made by European scholars to solve the cubic equation. Fibonacci, for example, attacked the problem in his *Flos* of *c.* 1225. He states that one Magister Johannes, a scholar from Palermo, proposed to him the problem of finding a cube which, with two squares and ten roots, should be equal to 20.[3] That is, the problem is to solve the equation

$$x^3 + 2x^2 + 10x = 20,$$

a numerical equation discussed later (p. 472). Another attempt was made by an anonymous writer of the 13th century whose work has been described by Libri.[4] He took two cubics, one of the type $ax^3 = cx + k$ and the other of the type $ax^3 = bx^2 + k$.

[1] The work was the *Lü-li Yüan-yüan*. See Mikami, *China*, pp. 117–119.

[2] Colebrooke, *loc. cit.*, p. 214.

[3] "Altera uero questio á predicto magistro Iohanne proposita fuit, vt inueniretur quidam cubus numerus, qui cum suis duobus quadratis et decem radicibus in unum collectis essent uiginti" (*Flos*, Boncompagni ed., p. 228).

[4] *Histoire*, 1838 ed., II, 213, 214. The MS. is probably Florentine.

In each case he displayed great ignorance, possibly because he was unable, on account of his unfamiliarity with radicals to check his results. It is also possible that he sought only approximate results, although this is not stated. His method in the first case was as follows:

Given $$ax^3 = cx + k,$$

we have $$x^3 = \frac{c}{a}x + \frac{k}{a};$$

whence he assumed that $$x = \frac{c}{2a} + \sqrt{\left(\frac{c}{2a}\right)^2 + \frac{k}{a}},$$

which is the root of $ax^2 = cx + k$ but not of the given equation.

His method in the second case was equally fallacious.

Slight attempts at numerical cubics were also made by Regiomontanus,[1] who gave, for example, the equation

$$\tfrac{1}{3}\tfrac{1}{2}\tfrac{1}{4}\,x^3 + \tfrac{5}{2}\tfrac{4}{9}\tfrac{0}{} = 3x;$$

but he contributed nothing of value to the theory.

The Cubic in Printed Books. Pacioli (1494) asserted substantially that the general solution is impossible.[2]

Of the early German writers only one made any noteworthy attempt at the solution, and this was a failure. Rudolff (1525) suggested three numerical equations, each with one integral root and each being easily solved by factoring.[3] His method in connection with one of these equations is interesting. In modern symbols it is substantially as follows:

Given $$x^3 = 10x^2 + 20x + 48,$$

we have $$x^3 + 8 = 10x^2 + 20x + 56;$$

whence $$x^2 - 2x + 4 = 10x + \frac{56}{x+2},$$

[1] Cantor, *Geschichte*, II, chap. lv.

[2] Fol. 149, *r.*, has the following:

> Impossibile. Censo de censo: e cēso equale. a cosa.
> Impossibile. Censo de censo e cosa. equale. a censo.

That is, the solution of equations like $ax^4 + cx^2 = dx$ and $ax^4 + dx = cx^2$ is impossible.

[3] *Die Coss*, 1553 ed., fol. 477, *r.*

all of which is correct. He now assumes that he can, in general, split the two members and say that

$$x^2 - 2x = 10x$$

and

$$4 = \frac{56}{x+2}.$$

Both of these equations are satisfied if $x = 12$, but the method is not otherwise general.

Similar solutions of special cases are found in various works of the 16th century, notably in a work by Nicolas Petri of Deventer,[1] published at Amsterdam in 1567. This writer was highly esteemed by his contemporaries.[2]

A few special cases, such as

$$x^3 = 38\tfrac{1}{2}\,x - 90,$$

he solves by factoring, and he then proceeds to a more elaborate discussion of certain cases that are mentioned later.

The Italian Algebraists and the Cubic. The real interest in the cubic lies, however, in the work of the Italian algebraists of the 16th century, and notably in the testimony of Cardan and Tartaglia. Cardan (1545) says that Scipio del Ferro discovered the solution of the type $x^3 + bx = c$ thirty years earlier (*c.* 1515), revealing the secret to his pupil Antonio Maria Fior (Florido).[3] The source of the solution is unknown. Ferro may have re-

[1] *Arithmetica. Practicque omne cortelycken te lerē chijpherē . . . Door my Nicolaum petri F. Dauentriensem*, Amsterdam, 1567. The name also appears as Nicolas Peetersen or Pietersz (Pieterszoon), Petri F. meaning Petri Filius (son of Peter).

[2] H. Bosmans, "La 'Practiqve om te leeren cypheren' de Nicolas Petri de Deventer," *Annales de la Société scientifique de Bruxelles*, XXXII, 2e Partie, Reprint, 1908.

[3] "Verùm temporibus nostris, Scipio Ferreus Bononiensis, capitulum cubi & rerum numero aequalium inuenit, rem sanè pulchram & admirabilem. . . . Huius emulatiõe Nicolaus Tartalea Brixellensis, amicus noster, cũ in certamẽ cũ illius discipulo Antonio Maria Florido uenisset, capitulum idem, ne uinceretur, inuenit, qui mihi ipsum multis precibus exoratus tradidit" (*Ars Magna*, fol. 3, *r.*). On the general work of the Italians with respect to the development of algebra see E. Bortolotti, "Italiani scopritori e promotori di teorie algebriche," in the *Annuario della R. Università di Modena*, Anno 1918–1919.

ceived it from some Arab writer, or he may have discovered it himself in spite of his apparent lack of mathematical ability. Tartaglia agrees with Cardan's statement except as to time, placing it somewhat earlier (in 1506),[1] a matter of little consequence. Cardan further says that Florido had a contest with Tartaglia which resulted in the latter's discovery of the method for solving this particular type, and that Tartaglia, at Cardan's request, revealed it to him.

Tartaglia states his side of the case rather differently and more explicitly. He says that Zuanne de Tonini da Coi[2] (see Volume I, page 295) sent him, in 1530, two problems, namely,

$$x^3 + 3x^2 = 5$$

and

$$x^3 + 6x^2 + 8x = 1000,$$

neither of which he could solve; but that in 1535[3] he found the method of solving any equation of the type $x^3 + ax^2 = c$. Tartaglia further states that he had a contest with Florido in 1535 and knew that he had only to set problems of this type to defeat his opponent, provided he could first find the latter's method of solving problems of the type $x^3 + bx = c$. He therefore exerted himself and succeeded in discovering it just before the contest,[4] thus being able to solve anything that Florido could set, and being able to propose problems that the latter could not master.

Tartaglia and Cardan. Da Coi now importuned Tartaglia to publish his method, but the latter declined to do so. In 1539 Cardan wrote to Tartaglia, and a meeting was arranged at which,

[1] ". . . se auantaua che gia trenta anni tal secreto gli era stato mostrato da un gran mathematico." From Qvesito XXV, dated December 10, 1536. See the 1554 edition of the *Qvesiti*, fol. 106, *v*.

[2] Also known as Giovanni Colle and Joannes Colla.

[3] In his statement of December 10, 1536 (Qvesito XXV), he says: " . . . & questo fu l'anno passato, cioe del .1535. adi .12. di. Febraro (uero è in Venetia ueneua à esser del .1534.) . . . " See also A. Oliva, *Sulla soluzione dell' equazione cubica di Tartaglia*, Milan, 1909.

[4] "Per mia bona sorte, solamente .8. giorni auanti al termine . . . Io haueua ritrouata la regola generale." *Qvesiti*, libro nono, Qvesito XXV; 1554 ed., fol. 106, *v*.

Tartaglia says, having pledged Cardan to secrecy, he revealed the method in cryptic verse[1] and later with a full explanation.[2]

Cardan admits that he received the solution from Tartaglia, but says that it was given to him without any explanation.[3] At any rate, the two cubics $x^3 + ax^2 = c$ and $x^3 + bx = c$ could now be solved. The reduction of the general cubic $x^3 + ax^2 + bx = c$ to the second of these forms does not seem to have been considered by Tartaglia at the time of the controversy. When Cardan published his *Ars Magna* (1545), however, he transformed the types $x^3 = ax^2 + c$ and $x^3 + ax^2 = c$ by the substitutions $x = y + \frac{1}{3}a$ and $x = y - \frac{1}{3}a$ respectively, and transformed the type $x^3 + c = ax^2$ by the substitution $x = \sqrt[3]{c^2}/y$, thus freeing the equations of the term in x^2. This completed the general solution, and he applied the method to the complete cubic in his later problems.

Cardan's Originality. Cardan's originality in the matter seems to have been shown chiefly in four respects. First, he reduced the general equation to the type $x^3 + bx = c$; second, in a letter written August 4, 1539, he discussed the question of the irreducible case; third, he had the idea of the number of roots to be expected in the cubic; and, fourth, he made a beginning in the theory of symmetric functions.[4]

[1] Quando chel cubo con le cose appresso
Se aggualia à qualche numero discreto — $x^3 + bx = c$
Trouan dui altri differenti in esso. — $u - v = c$
Dapoi terrai questo per consueto
Che 'l lor produtto sempre sia eguale — $uv = \left(\frac{b}{3}\right)^3$
Al terzo cubo delle cose neto,
El residuo poi suo generale
Delli lor lati cubi ben sostratti
Varra la tua cosa principale. — $x = \sqrt[3]{u} - \sqrt[3]{v}$

Qvesiti, 1554 ed., fol. 120, *v*. There are sixteen lines more. See also Gherardi in Grunert's *Archiv*, LII, 143 seq. and 188.

[2] Substantially this: If $x^3 + bx = c$, let $u - v = c$ and $uv = \left(\frac{b}{3}\right)^3$. Then $x = \sqrt[3]{u} - \sqrt[3]{v}$, for $(\sqrt[3]{u} - \sqrt[3]{v})^3 + b(\sqrt[3]{u} - \sqrt[3]{v}) = u - v$. See the second part of his Qvesito XXXV, *Qvesiti*, 1554 ed., fol. 121, *v*.

[3] ". . . ut Nicolaus inuenerit & ipse, qui cum nobis rogantibus tradidisset, suppressa demonstratione . . . " *Ars Magna*, 1545 ed., fol. 29, *v*., shown in facsimile on pages 462, 463.

[4] See also Eneström's summary in *Bibl. Math.*, VII (3), 293.


relinquitur prima 6 m: ℞ $30\frac{6}{7}$, hæ autem quantitates proportionales ſunt, & quadratum ſecundæ eſt æquale duplo producti ſecundæ in primam, cum quadruplo primæ, ut proponebatur.

De cubo & rebus æqualibus numero. Cap. XI.

CIpio Ferreus Bononienſis iam annis ab hinc triginta fermè capitulum hoc inuenit, tradidit uero Anthonio Mariæ Florido Veneto, qui cũ in certamen cũ Nicolao Tartalea Brixellenſe aliquando uenisſet, occaſionem dedit, ut Nicolaus inuenerit & ipſe, qui cum nobis rogantibus tradidiſſet, ſuppreſſa demonſtratione, freti hoc auxilio, demonſtrationem quæſiuimus, eamq; in modos, quod difficillimum fuit, redactam ſic ſubiecimus.

DEMONSTRATIO.

Sit igitur exempli cauſa cubus G H & ſexcuplum lateris G H æquale 20, & ponam duos cubos A E & C L, quorum differentia ſit 20, ita quod productum A C lateris, in C K latus, ſit 2, tertia ſcilicet numeri rerum pars, & abſcindam C B, æqualem C K, dico, quod ſi ita fuerit, lineam A B reſiduum, eſſe æqualem G H, & ideo rei æſtimationem, nam de G H iam ſupponebatur, quod ita eſſet, perficiam igitur per modum primi ſuppoſiti 6 capituli huius libri, corpora D A, D C, D E, D F, ut per D C intelligamus cubum B C, per D F cubum A B, per D A triplum C B in quadratum A B, per D E triplum A B in quadratũ B C. quia igitur ex A C in C K fit 2, ex A C in C K ter fiet 6 numerus rerum, igitur ex A B in triplum A C in C K fiunt 6 res A B, ſeu ſexcuplum A B, quare triplum producti ex A B, B C, A C, eſt ſexcuplum A B, at uero differentia cubi A C, à cubo C K, & exiſtenti à cubo B C ei æq̃le ex ſuppoſito, eſt 20, & ex ſuppoſito primo 6 capituli, eſt aggregatum corporum D A, D E, D F, tria igitur hæc corpora ſunt 20, poſita uero B C m: cubus A B, æqualis eſt cubo A C, & triplo A C in quadratum C B, & cubo B C m: & triplo B C in quadratum A C m: per demonſtrata illic, differentia autem tripli B C in quadratum A C, à triplo A C in quadratum B C eſt productum A B, B C, A C, quare cum hoc, ut demonſtratum eſt, æquale ſit ſexcuplo A B, igitur addito ſexcuplo A B, ad id quod fit ex A C in quadratum B C ter, fiet triplum B C in quadratum A C, cum igitur B C ſit m: iam oſtenſum eſt, quod productum C B



in

CARDAN'S SOLUTION OF THE CUBIC

First page of the solution as given in the first edition of Cardan's *Ars Magna*, Nürnberg, 1545. The solution was slightly expanded in the second edition, Basel, 1570

in quadratum A C ter, eſt m: & reliquum quod ei æquatur eſt p: igitur triplum C B in q̃dratum A B, & triplum A C in q̃dratũ C B, & ſexcuplũ A B nihil faciunt. Tanta igitur eſt differentia, ex cõmuni animi ſententia, ipſius cubi A C, à cubo B C, quantum eſt quod cõflatur ex cubo A C, & triplo A C in quadratum C B, & triplo C B in quadratum A C m: & cubo B C m: & ſexcuplo A B, hoc igitur eſt 20, quia differentia cubi A C, à cubo C B, fuit 20, quare per ſecundum ſuppoſitum 6ti capituli, poſita B C m: cubus A B æquabitur cubo A C, & triplo A C in quadratum B C, & cubo B C m: & triplo B C in quadratum A ç m: cubus igitur A B, cum ſexcuplo A B, per communem animi ſententiam, cum æquetur cubo A C & triplo A C in quadratum C B, & triplo C B in quadratum A B m: & cubo C B m: & ſexcuplo A B, quæ iam æquatur 20, ut probatum eſt, æquabuntur etiam 20, cum igitur cubus A B & ſexcuplum A B æquentur 20, & cubus G H, cum ſexcuplo G H æquentur 20, erit ex communi animi ſententia, & ex dictis, in $3\,5^{2}$ p: & $3\,1^{2}$ undecimi elementorum, G H æqualis A B, igitur G H eſt differentia A C & C B, ſunt autem A C & C B, uel A C & C K, numeri ſeu liniæ continentes ſuperficiem, æqualem tertiæ parti numeri rerum, quarum cubi differunt in numero æquationis, quare habebimus regulam.

Regvla.

Deducito tertiam partem numeri rerum ad cubum, cui addes quadratum dimidij numeri æquationis, & totius accipe radicem, ſcili cet quadratam, quam ſeminabis, uniq; dimidium numeri quod iam in ſe duxeras, adijcies, ab altera dimidium idem minues, habebisq; Binomium cum ſua Apotome, inde detracta ℞ cubica Apotomæ ex ℞ cubica ſui Binomij, reſiduũ quod ex hoc relinquitur, eſt rei eſtimatio.

Exemplum. cubus & 6 poſitiones, æquantur 20, ducito 2, tertiam partem 6, ad cubum, fit 8, duc 10 dimidium numeri in ſe, fit 100, iunge 100 & 8, fit 108, accipe radicem quæ eſt ℞ 108, & eam geminabis, alteri addes 10, dimidium numeri, ab altero minues tantundem, habebis Binomiũ ℞ 108 p: 10, & Apotomen ℞ 108 m: 10, horum accipe ℞ cub. & minue illam quę eſt Apotomæ, ab ea quæ eſt Binomij, habebis rei æſtimationem, ℞ v: cub: ℞ 108 p: 10 m: ℞ v: cubica ℞ 108 m: 10.

cub⁹ p: 6 reb⁹ æq̃lis 20
2 20
8 ——— 10
108
℞ 108 p: 10
℞ 108 m: 10
℞ v: cu. ℞ 108 p: 10
m: ℞ v: cu. ℞ 108 m: 10

Aliud, cubus p: 3 rebus æquetur 10, duc 1, tertiam partem 3, ad cubum, fit 1, duc 5, dimidium 10, ad quadratum, fit 25, iunge 25 & 1,

CARDAN'S SOLUTION OF THE CUBIC

Continuation of the solution as given on page 462. For the meaning of the symbols see page 428

With respect to the irreducible case, his solution of the type $x^3 + bx = c$ is

$$x = \sqrt[3]{\sqrt{\tfrac{1}{4}c^2 + \tfrac{1}{27}b^3} + \tfrac{1}{2}c} - \sqrt[3]{\sqrt{\tfrac{1}{4}c^2 + \tfrac{1}{27}b^3} - \tfrac{1}{2}c},$$

and if b is negative and is such that $\frac{1}{4}c^2 + \frac{1}{27}b^3$ is also negative, then we have the cube root of a complex number, thus reaching an expression that is irreducible even though all three values of x turn out to be real.[1]

With respect to the number of roots to be expected in the cubic, he gave[2] the equations $x^3 + 10x = 6x^2 + 4$ with roots $2, 2 \pm \sqrt{2}$; $x^3 + 21x = 9x^2 + 5$ with roots $5, 2 \pm \sqrt{3}$; and $x^3 + 26x = 12x^2 + 12$ with roots $2, 5 \pm \sqrt{19}$; but before this time only two roots were ever found,[3] negative roots being generally rejected.

As to the question of symmetric functions, he stated that the sum of the roots is minus the coefficient of x^2.[4]

Cardan's solution, with part of his explanation, is shown in facsimile on pages 462 and 463. In the solution he states that the root of the equation $x^3 + 6x = 20$ is

$$x = \sqrt[3]{\sqrt{108} + 10} - \sqrt[3]{\sqrt{108} - 10}.$$

He also gave thirteen forms of the cubic which have positive roots, these having already been given by Omar Khayyam.

[1] The reality of the roots for this case was shown by Kästner (1745) and A. C. Clairaut (1746). As an example of the irreducible case, in the equation $x^3 - 63x - 162 = 0$ the rule gives

$$x = \sqrt[3]{81 + 30\sqrt{-3}} + \sqrt[3]{81 - 30\sqrt{-3}},$$

which we cannot reduce, although as a matter of fact the solution is

$$x = (-3 + 2\sqrt{-3}) + (-3 - 2\sqrt{-3}) = -6.$$

[2] Cap. XVIII, Exemplum quintum: "Cubus & 10 res, aequatur 6 quadratis p: 4 (1545 ed., fol. 39, *r.*). The roots are "2 p: R̃ 2, uel 2 m: R̃ 2, potest etiam esse 2." The folio is incorrectly numbered 36.

[3] It was Euler (1732) who gave the first noteworthy modern discussion of the cubic, insisting on the recognition of all three roots and stating how these roots were found. "De formis radicum aequationum cuiusque ordinis conjectatio," in *Comment. Petropol. ad annos 1732–1733*, printed in 1738; VI, 217.

[4] ". . . uelut in quinto exemplo, 2 p: R̃ 2, & 2, & 2 m: R̃ 2, componunt 6, numerum quadratorum," and so for other cases. That is, in the case of $x^3 + 10x = 6x^2 + 4$ the sum of $2 + \sqrt{2}$, 2, and $2 - \sqrt{2}$ is 6 (fol. 39, *v.*).

Nicolas Petri and the Cubic. In his work of 1567 Nicolas Petri of Deventer, as already mentioned, gave some attention to the cubic equation. This is found in a subdivision on *Cubicq Coss*,[1] in which he gives eight cubic equations such as

$$x^3 = 9x + 28,$$

$$23x^3 + 32x = 905\tfrac{5}{9}, \quad \text{and} \quad x^3 = 3x^2 + 5x + 16,$$

all of which he solves by Cardan's method.

In the same year that Petri's work appeared, Pedro Nuñez (to take the form of his name used in the treatise here mentioned) published his *Libro de algebra en arithmetica y geometria* at Antwerp.[2] In this work he considers such equations as

$$x^3 + 3x = 36 \quad \text{and} \quad x^3 + 9x = 54,$$

and seeks to show that Tartaglia's rule is not practical where one root is easily found by factoring. He shows a familiarity with the works of both Tartaglia and Cardan.

Vieta generalizes the Work. Although Cardan reduced his particular equations to forms lacking a term in x^2, it was Vieta[3] who began with the general form

$$x^3 + px^2 + qx + r = 0$$

and made the substitution $x = y - \frac{1}{3}p$, thus reducing the equation to the form

$$y^3 + 3by = 2c.$$

He then made the substitution

$$z^3 + yz = b, \quad \text{or} \quad y = \frac{b - z^3}{z},$$

which led to the form

$$z^6 + 2cz^3 = b^2,$$

a sextic which he solved as a quadratic.

[1] "Volgen sommighe exempelen ghesolueert deur die Cubicq Coss."

[2] H. Bosmans, "Sur le 'Libro de algebra' de Pedro Nuñez," *Bibl. Math.*, VIII (3), 154. The original name is Nunes. See Volume I, page 348.

[3] *Opera mathematica. IV. De aequationum recognitione et emendatione libri duo*, Tract. II, cap. vii (Paris, 1615). His equation is stated thus: "Proponatur A cubus + B plano 3 in A, aequari Z solido 2"; that is, $A^3 + 3\,BA = 2\,Z$, or, in our symbols, $y^3 + 3by = 2c$. The problem as worked out by Vieta is given in Matthiessen, *Grundzüge*, p. 371.

He also gave two or three other solutions, but the one here shown is particularly clear and simple. In his work in equations he was greatly aided by his new symbolism (p. 430).

Hudde's Contribution. Although Descartes contributed to the solution of the cubic equation by his convenient symbolism and by his work on equations in general, he made no specific contribution of importance. The next writer to materially simplify the work of Vieta was Hudde (*c.* 1658). Taking advantage of Descartes's symbolism, he brought the theory of the cubic equation to substantially its present status. He is also the first algebraist who unquestionably recognized that a letter might stand for either a positive or a negative number.[1]

His method of solving the cubic equation is to begin with

$$x^3 = qx + r$$

and let $x = y + z$,

so that $y^3 + 3y^2z + 3yz^2 + z^3 = qx + r$.

He then lets $y^3 + z^3 = r$

and $3zy^2 + 3z^2y = qx$,

which gives $y = \frac{1}{3}q/z$.

Hence $y^3 = r - z^3 = \frac{1}{27}q^3/z^3$,

and so $z^3 = \frac{1}{2}r \pm \sqrt{\frac{1}{4}r^2 - \frac{1}{27}q^3} = A$

and $y^3 = \frac{1}{2}r \mp \sqrt{\frac{1}{4}r^2 - \frac{1}{27}q^3} = B$.

Hence $x = \sqrt[3]{A} + \sqrt[3]{B}$,

which satisfies both his assumptions.[2]

Equation of the Fourth Degree. After the cubic equation had occupied the attention of Arab scholars, with not very significant results, the biquadratic equation was taken up. Abû'l-Faradsh[3] completed the *Fihrist* *c.* 987, and in this he refers to

[1] Enestrom, in *Bibl. Math.*, IV (3), pp. 208, 216.

[2] The problem, as worked out by Hudde, is given in Matthiessen, *Grundzüge*, p. 374.

[3] Abû'l-Faradsh (Faraj) Mohammed ibn Isḥâq, known as Ibn Abî Ya'qûb al-Nadîm. The title is *Kitâb al-Fihrist* (*Book of Lists*). See the *Abhandlungen*, VI, 1.

the following problem by Abû'l-Wefâ (*c.* 980): "On the method of finding the root of a cube and of a fourth power and of expressions composed of these two powers."[1] The last means that we are to solve the equation $x^4 + px^3 = q$. The equation could have been solved by the intersection of the hyperbola $y^2 + axy + b = 0$ and the parabola $x^2 - y = 0$, but the work in which Abû'l-Wefâ's problem appeared is lost and we do not know what he did in the way of a solution.

Woepcke, a French orientalist (*c.* 1855), has called attention to an anonymous MS. of an Arab or Persian algebraist in which there is given the biquadratic equation

$$(100 - x^2)\ (10 - x)^2 = 8100.$$

This is solved by taking the intersection of $(10 - x)y = 90$ and $x^2 + y^2 = 100$, but there is no evidence that the author was concerned with the algebraic theory.[2]

It may therefore be said that the Arabs were interested in the biquadratic equation only as they were in the cubic, that is, from the standpoint of the intersection of two conics.

The Italian Algebraists and the Biquadratic. The problem of the biquadratic equation was laid prominently before Italian mathematicians by Zuanne de Tonini da Coi, who in 1540 proposed the problem, "Divide 10 into three parts such that they shall be in continued proportion and that the product of the first two shall be 6." He gave this to Cardan[3] with the statement that it could not be solved, but Cardan denied the assertion, although himself unable to solve it. He gave it to Ferrari, his

[1] *Abhandlungen*, VI, 73, note 253. See also Matthiessen, *Grundzüge*, p. 543; F. Woepcke, *Recherches . . . Constructions géom. par Aboûl Wafâ*, p. 36, 8, n. 2 (Paris, 1855).

[2] Woepcke's translation of Omar Khayyam, Addition D, p. 115. The problem was to construct an isosceles trapezium (trapezoid) *ABCD* such that $AB = AD = BC = 10$, and the area is 90.

[3] Cardan states it thus: "Exemplum. Fac ex 10 tres partes proportionales ex quarum ductu primæ in secundam, producantur 6. Hanc proponebat Ioannes Colla, & dicebat solui non posse, ego uero dicebam, eam posse solui, modum tamẽ ignorabam, donec Ferrarius eum inuenit." *Ars Magna*, cap. xxxix, qvæstio v; 1545 ed., fol. 73, *v*.

pupil (Vol. I, p. 300), and the latter, although then a mere youth, succeeded[1] where the master had failed.

Ferrari's method[2] may be summarized in its modern form as follows: Reduce the complete equation

$$x^4 + a_1x^3 + a_2x^2 + a_3x + a_4 = 0$$

to the form $x^4 + px^2 + qx + r = 0$

and thence to $x^4 + 2px^2 + p^2 = px^2 - qx - r + p^2$,

or $(x^2 + p)^2 = px^2 - qx + p^2 - r$.

Write this as

$$(x^2 + p + y)^2 = (p + 2y)x^2 - qx + (p^2 - r + 2py + y^2).$$

Now determine y so that the second member shall be a square. This is the case when

$$4(p + 2y)(p^2 - r + 2py + y^2) = q^2,$$

which requires the solution of a cubic in y, which is possible. The solution then reduces to the mere finding of square roots.

This method soon became known to algebraists through Cardan's *Ars Magna*, and in 1567 we find it used by Nicolas Petri in the work already mentioned. Petri solves four equations, the first being

$$x^4 + 6x^3 = 6x^2 + 30x + 11.$$

Of this he gives only the root $1+\sqrt{2}$, neglecting the roots $1-\sqrt{2}$, $-4 \pm \sqrt{5}$ because they are negative.

[1] The proportion is $\frac{6}{x} : x = x : \frac{1}{6}x^3$, and the other condition is that

$$\frac{6}{x} + x + \frac{1}{6}x^3 = 10,$$

the two conditions reducing to $x^4 + 6x^2 + 36 = 60x$. Ferrari's method makes this depend upon the solution of the equation $y^3 + 15y^2 + 36y = 450$, or, as Cardan (*Ars Magna*, fol. 74, *r.*) states the problem, "1 cubum p: 15 quadratis p: 36 positionibus æquantur 450."

[2] Cardan, *Ars Magna*, 1545, cap. xxxix, qvæstio v, fol. 73, *v.*; Bombelli, *Algebra*, 1572, p. 353; Matthiessen, *Grundzüge*, p. 540. Bombelli's first special case is "1 ♁ p. 20 ♁ eguale à 21"; that is, $x^4 + 20x = 21$.

Vieta and Descartes. Vieta (*c.* 1590) was the first algebraist after Ferrari to make any noteworthy advance in the solution of the biquadratic.[1] He began with the type $x^4 + 2gx^2 + bx = c$, wrote it as $x^4 + 2gx^2 = c - bx$, added $g^2 + \frac{1}{4}y^2 + yx^2 + gy$ to both sides, and then made the right side a square after the manner of Ferrari. This method also requires the solution of a cubic resolvent.

Descartes[2] (1637) next took up the question and succeeded in effecting a simple solution of problems of the type

$$x^4 + px^2 + qx + r = 0,$$[3]

a method considerably improved (1649) by his commentator Van Schooten.[4] The method was brought into its modern form by Simpson (1745).[5]

Equation of the Fifth Degree. Having found a method differing from that of Ferrari for reducing the solution of the general biquadratic equation to that of a cubic equation, Euler had the idea that he could reduce the problem of the quintic equation to that of solving a biquadratic, and Lagrange made the same attempt. The failures of such able mathematicians led to the belief that such a reduction might be impossible. The first noteworthy attempt to prove that an equation of the fifth degree could not be solved by algebraic methods is due to Ruffini (1803, 1805),[6] although it had already been considered by Gauss.

The modern theory of equations in general is commonly said to date from Abel and Galois. The latter's posthumous (1846) memoir on the subject established the theory in a satisfactory manner. To him is due the discovery that to each equation there

[1] *De aequationum recognitione et emendatione libri duo*, Tract. II, cap. vi, prob. iii (Paris, 1615). For solution see Matthiessen, *Grundzüge*, p. 547.

[2] *La Geometrie*, Lib. III; 1649 ed., p. 79; 1683 ed., p. 71; 1705 ed., p. 109.

[3] For examples see Matthiessen, *Grundzüge*, p. 549.

[4] 1649 ed., p. 244.

[5] For the various improvements see Matthiessen, *Grundzüge*, p. 545 seq.

[6] "Della insolubilità delle equazioni algebraiche generali di grado superiore al quarto," *Mem. Soc. Ital.*, X (1803), XII (1805).

corresponds a group of substitutions (the "group of the equation") in which are reflected its essential characteristics. Galois's early death left without sufficient demonstration several important propositions, a gap which has since been filled.

Abel[1] showed that the roots of a general quintic equation cannot be expressed in terms of its coefficients by means of radicals.

Lagrange had already shown that the solution of such an equation depends upon the solution of a sextic, "Lagrange's resolvent sextic," and Malfatti and Vandermonde had investigated the construction of resolvents.

The transformation of the general quintic into the trinomial form $x^5 + ax + b = 0$ by the extraction of square and cube roots only was first shown to be possible by Bring (1786) and independently by Jerrard[2] (1834). Hermite (1858) actually effected this reduction by means of a theorem due to Tschirnhausen, the work being done in connection with the solution by elliptic functions.[3]

Symmetric Functions. The first formulas for the computation of the symmetric functions of the roots of an equation seem to have been worked out by Newton, although Girard (1629) had given, without proof, a formula for a power of the sum, and Cardan (1545) had made a slight beginning in the theory. In the 18th century Lagrange (1768) and Waring (1770, 1782) made several valuable contributions to the subject, but the first tables, reaching to the tenth degree, appeared in 1809 in the Meyer-Hirsch *Aufgabensammlung*. In Cauchy's celebrated memoir on determinants (1812) the subject began to assume new prominence, and both he and Gauss (1816) made numerous and important additions to the theory. It is, however, since the discoveries by Galois that the subject has become one of great

[1] *Mémoire sur les équations algébriques*, Christiania, 1824, and Crelle's *Journal*, 1826.

[2] R. Harley, "A contribution to the history . . . of the general equation of the fifth degree . . .," *Quarterly Journal of Mathematics*, VI, 38.

[3] For a bibliography of much value in the study of the history of equations see G. Loria, in *Bibl. Math.*, V (2), 107.

significance. Cayley (1857) gave a number of simple rules for the degree and weight of symmetric functions, and he and Brioschi simplified the computation of tables.

Harriot's Law of Signs. The law which asserts that the equation $X = 0$, complete or incomplete, can have no more real positive roots than it has changes of sign, and no more real negative roots than it has permanences of sign, was apparently known to Cardan[1]; but the first satisfactory statements relating to the matter are due to Harriot (died 1621)[2] and Descartes.[3]

Numerical Higher Equations. The solution of the numerical higher equation for approximate values of the roots begins, so far as we know, in China. Indeed, this is China's particular contribution to mathematics, and in this respect her scholars were preëminent in the 13th and 14th centuries.[4] In the *Nine Sections*, written apparently long before the Christian era, there is found the "celestial element method."[5] This was a method of solving numerical higher equations; it is found in various early Chinese works, reaching its highest degree of perfection in the works of Ch'in Kiu-shao (1247). Here it appears, as already stated, in a form substantially equivalent to Horner's Method (1819).[6]

Fibonacci on Numerical Equations. The first noteworthy work upon numerical higher equations done in Europe is due to Fibonacci (1225), and relates to the case of the cubic equation

[1] Cantor, *Geschichte*, II (2), 539; Eneström, *Bibl. Math.*, VII (3), 293.

[2] *Artis analyticae praxis. Ad aequationes Algebraicas . . . resolvendas*, London, 1631 (posthumous); Matthiessen, *Grundzüge*, 2d ed., pp. 18, 268.

[3] *La Geometrie*, 1637; 1649 ed., p. 78; 1705 ed., p. 108, with the statement: "On connoît aussi de ceci combien il peut y avoir de vrayes racines, & combien de fausses en chaque Equation; à sçavoir, il y en peut avoir autant de vrayes que les signes + & − s'y trouvent de fois être changez, & autant de fausses qu'il s'y trouve de fois deux signes +, ou deux signes − qui s'entresuivent." The law usually bears the name of Descartes.

[4] Y. Mikami, *China*, 25, 53, 76, *et passim*; L. Matthiessen, "Zur Algebra der Chinesen," in *Zeitschrift für Math. und Phys.*, XIX, Hl. Abt., 270. For doubts as to the originality of this work and as to the authenticity of the text of the *Nine Sections* see G. Loria, "Che cosa debbono le matematiche ai Cinesi," *Bollettino della Mathesis*, XII (1920), 63.

[5] *T'ien-yuen-shu*, the Japanese *tengen jutsu*.

[6] See Volume I, page 270. For a detailed solution see Mikami, *China*, p. 76 seq.

$x^3 + 2x^2 + 10x = 20$, already mentioned. His method of attack was substantially as follows:

Since $x^3 + 2x^2 + 10x = 20,$

we have $10(x + \frac{1}{10}x^3 + \frac{1}{5}x^2) = 20,$

or $x + \frac{1}{10}x^3 + \frac{1}{5}x^2 = 2,$

so that $x < 2.$

But $1 + 2 + 10 = 13 < 20,$

and so $x > 1.$

But x is not fractional; for if $x = a/b$, then

$$\frac{a}{b} + \frac{a^3}{10b^3} + \frac{a^5}{5b^5}$$

cannot be integral, and so x must be irrational.

Further, x cannot be the square root of an integer; for, from the given equation,

$$x = \frac{20 - 2x^2}{10 + x^2},$$

and if x were equal to $\sqrt{a}$ we should have

$$\sqrt{a} = \frac{20 - 2a}{10 + a},$$

which is impossible.

Fibonacci here closes his analysis and simply makes a statement which we may express in modern symbols as

$$x = 1^\circ\ 22'\ 7''\ 42'''\ 33^{iv}\ 4^{v}\ 40^{vi},$$

a result correct to $1\frac{1}{2}^{vi}$; that is, the value is only $\frac{1}{31104000000}$ too large. How this result was obtained no one knows, but the fact that numerical equations of this kind were being solved in China at this time, and that intercourse with the East was possible, leads to the belief that Fibonacci had learned of the solution in his travels, had contributed what he could to the theory, and had then given the result as it had come to him.

Vieta and Newton contribute to the Theory. About the year 1600 Vieta suggested that a particular root of a numerical equation could be found by a process similar to that of obtaining

a root of a number. By substituting in $f(x)$ a known approximate root of $f(x) = n$ he was able to find the next figure by division.[1]

Newton (1669) simplified this method of Vieta's, and the plan of procedure may be seen in his solution of the equation $y^3 - 2y - 5 = 0$. He first found by inspection that $2 < y < 3$. He then let $2 + p = y$;

whence $$y^3 - 2y - 5 = -1 + 10p + 6p^2 + p^3 = 0,$$

and $p = 0.1$, approximately.

Letting $0.1 + q = p$, we have

$$0.061 + 11.23\,q + 6.3\,q^2 + q^3 = 0;$$

whence $q = -0.0054$, approximately.

Letting $-0.0054 + r = q$, we have

$$0.000541708 + 11.16196\,r + 6.3\,r^2 = 0;$$

whence $r = -0.00004854$, approximately.

Similarly, we could let $-0.00004854 + s = r$, and proceed as before. We could then reverse the process and find p.

In this way he finds[2] the approximate value

$$y = 2.09455147.$$

As already stated (page 471), in 1819 William George Horner carried this simplification still farther, the root being developed figure by figure. The process terminates if the root is commensurable, and it may be carried to any required number of decimal places if it is incommensurable.[3]

Fundamental Theorem. The Italian algebraists of the 16th century tacitly assumed that every rational integral equation has a root. The later ones of that century were also aware that a quadratic equation has two roots, a cubic equation three roots,

[1] Burnside and Panton, *Theory of Equations*, 4th ed., I, 275. Dublin, 1899.

[2] "De analysi per aequationes numero terminorum infinitas," extract of 1669 in the *Commercium Epistolicum*, p. 76 (London, 1725). Wallis also gave an approximation method in 1685.

[3] For a simple presentation see Burnside and Panton, *loc. cit.*, I, 227, and consult that work (I, 275) for further information on the subject.

and a biquadratic equation four roots. The first writer to assert positively that every such equation of the nth degree has n roots and no more seems to have been Peter Roth, a Nürnberg Rechenmeister, in his *Arithmetica philosophica* (Nürnberg, 1608).[1] The law was next set forth by a more prominent algebraist, Albert Girard, in 1629.[2] It was, however, more clearly expressed by Descartes (1637), who not only stated the law but distinguished between real and imaginary roots and between positive and negative real roots in making the total number.[3] Rahn (Rhonius), also, gave a clear statement of the law in his *Teutschen Algebra* (1659).[4]

After these early steps the statement was repeated in one form or another by various later writers, including Newton (*c.* 1685) and Maclaurin (posthumous publication, 1748). D'Alembert attempted a proof of the theorem in 1746, and on this account the proposition is often called d'Alembert's Theorem. Other attempts were made to prove the statement, notably by Euler (1749) and Lagrange, but the first rigorous demonstration is due to Gauss (1799, with a simple treatment in 1849).

Trigonometric Solutions. In the 16th century Vieta[5] suggested (1591) the treatment of the numerical cubic equation by trigonometry, and Van Schooten later elaborated the plan. Girard[6] (1629) was one of the first, however, to attack the problem scientifically. He solved the equation 1 ③ ∞ 13 ① + 12, that is, $x^3 = 13x + 12$, by the help of the identity

$$\cos 3\phi = 4\cos^3\phi - 3\cos\phi.$$

[1] In modern works the name also appears as Rothe. See Tropfke, *Geschichte*, III (2), 95, with a quotation from the original work. Roth died at Nürnberg in 1617. See Volume I, page 421.

[2] "Toutes les équations d'algèbre reçoivent autant de solutions, que la denomination de la plus haute quantité le demonstre." *Invention nouvelle en l'algèbre*, Amsterdam, 1629; quoted in Tropfke, *Geschichte*, III (2), 95, to which refer for further details.

[3] "Au reste tant les vrayes racines que les fausses ne sont pas toujours réelles, mais quelquefois seulement imaginaires." *La Geometrie* (1705 ed.), p. 117.

[4] English translation, London, 1668. See Volume I, page 412.

[5] See Van Schooten's edition of his *Opera*, p. 362 (Leyden, 1646).

[6] *Invention nouvelle en l'algèbre*, Amsterdam, 1629. On the primitive Arab method see Matthiessen, *Grundzüge*, p. 894; on Girard, see *ibid.*, p. 896.

A single solution of a quadratic equation by trigonometric methods will show the later development of the subject.[1]

Fischer's Solution. Let

$$x^2 - px + q = 0. \qquad p^2 \geqq 4q$$

Then let $x_1 = p\cos^2\phi$

and $x_2 = p\sin^2\phi.$

Then $x_1 + x_2 = p(\cos^2\phi + \sin^2\phi) = p$

and $x_1 x_2 = p^2(\cos\phi \sin\phi)^2$

$= \frac{1}{4}p^2\sin^2 2\phi.$

The angle ϕ can now be found from the relation

$$\sin 2\phi = 2\sqrt{q}/p.$$

For example, given the equation

$$x^2 - 93.7062\,x + 1984.74 = 0,$$

we find $2\phi = 71^\circ\ 57'\ 44.6''$,

whence $\phi = 35^\circ\ 58'\ 52.3''$;

and hence $x_1 = 61.3607$

and $x_2 = 32.3454.$

Such methods have been extensively used with the cubic and biquadratic equations.[2]

9. Determinants

Among the Chinese. The Chinese method of representing the coefficients of the unknowns of several linear equations by means of rods on a calculating board naturally led to the discovery of simple methods of elimination. The arrangement of the rods was precisely that of the numbers in a determinant. The Chinese, therefore, early developed the idea of subtracting columns and rows as in the simplification of a determinant.[3]

[1] This is due to Fischer, *Die Auflösung der quadratischen und kubischen Gleichungen durch Anwendung der goniometrischen Functionen*, Elberfeld, 1856. See Matthiessen, *Grundzüge*, p. 885, and consult this work for a detailed history of the subject. [2] For a list of writers see Matthiessen, *Grundzüge*, p. 888 seq.

[3] Mikami, *China*, pp. 30, 93.

Among the Japanese. It was not until Chinese science had secured a firm footing in Japan, and Japanese scholars had begun to show their powers, that the idea of determinants began to assume definite form. Seki Kōwa, the greatest of the Japanese mathematicians of the 17th century, is known to have written a work called the *Kai Fukudai no Hō* in 1683. In this he showed that he had the idea of determinants and of their expansion. It is strange, however, that he used the device only in eliminating a quantity from two equations and not directly in the solution of a set of simultaneous linear equations.[1]

Determinants in Europe. So far as Western civilization is concerned, the theory of determinants may be said to have begun with Leibniz[2] (1693), who considered these forms solely with reference to simultaneous equations, as the Chinese had already done.

It was Vandermonde (1771) who first recognized determinants as independent functions. To him is due the first connected exposition of the theory, and he may be called its formal founder. Laplace (1772) gave the general method of expanding a determinant in terms of its complementary minors, although Vandermonde had already considered a special case. Immediately following the publication by Laplace, Lagrange (1773) treated of determinants of the second and third orders and used them for other purposes than the solution of equations.

The next considerable step in advance was made by Gauss (1801). He used determinants in his theory of numbers, introduced the word "determinant"[3] (though not in the present signification,[4] but rather as applied to the discriminant of a quantic), suggested the notion of reciprocal determinants, and came very near the multiplication theorem.

[1] T. Hayashi, "The Fukudai and Determinants in Japanese Mathematics," in the *Proc. of the Tokyō Math. Soc.*, V (2), 257; Mikami, *Isis*, II, 9.

[2] Sir Thomas Muir, *Theory of Determinants in the Historical Order of Development* (4 vols., London, 1890, 1911, 1919; 2d ed., 1906, 1911, 1920, 1923), which consult on the whole question; M. Lecat, *Histoire de la théorie des Déterminants à plusieurs dimensions*, Ghent, 1911.

[3] Laplace had used "resultant."

[4] "Numerum bb — ac, cuius indole proprietates formae (a, b, c) imprimis pendere in sequentibus docebimus, determinantem huius uocabimus."

The next great contributor was Jacques-Philippe-Marie Binet,[1] who formally stated (1812) the theorem relating to the product of two matrices of m columns and n rows, which for the special case of $m = n$ reduces to the multiplication theorem.

On the same day (November 30, 1812) that he presented his paper to the Académie, Cauchy presented one on the same subject. In this paper he used the word "determinant" in its present sense, summarized and simplified what was then known on the subject, improved the notation, and gave the multiplication theorem with a proof more satisfactory than Binet's. He may be said to have begun the theory of determinants as a distinct branch of mathematics.

Aside from Cauchy, the greatest contributor to the theory was Carl Gustav Jacob Jacobi.[2] With him the word "determinant" received its final acceptance. He early used the functional determinant which Sylvester has called the Jacobian, and in his famous memoirs in Crelle's *Journal* for 1841 he considered these forms as well as that class of alternating functions which Sylvester has called alternants.

About the time of Jacobi's closing memoirs Sylvester (1839) and Cayley began their great work in this field. It is impossible to summarize this work briefly, but it introduced the most important phase of the recent development of the theory.

10. Ratio, Proportion, and the Rule of Three

Nature of the Topics. It is rather profitless to speculate as to the domain in which the concept of ratio first appeared. The idea that one tribe is twice as large as another and the idea that one leather strap is only half as long as another both involve the notion of ratio; both are such as would develop early in the history of the race, and yet one has to do with ratio of numbers and the other with ratio of geometric magnitudes. Indeed, when we come to the Greek writers we find Nicomachus including ratio in his arithmetic, Eudoxus in his geometry, and

[1] Born at Rennes, February 2, 1786; died in Paris, May 12, 1856.

[2] See Volume I, page 506.

Theon of Smyrna in his chapter on music.[1] Still later, Oriental merchants found that they could easily secure results to certain numerical problems by a device which, in the course of time, became known as the Rule of Three, and so this topic found place in commercial arithmetics, although fundamentally it is an application of proportion. Since ratio, proportion, and variation are now considered as topics of algebra, however, it is appropriate to treat of these subjects, as well as the Rule of Three, in the present chapter.[2]

Technical Terms. The word "ratio" as commonly used in school, while sanctioned by ancient usage,[3] has never been a favorite outside the mathematical classroom. It is a Latin word[4] and was commonly used in the arithmetic of the Middle Ages to mean computation. To represent the idea which we express by the symbols $a:b$ the medieval Latin writers generally used the word *proportio*, not the word *ratio*; while for the idea of an equality of ratio, which we express by the symbols $a:b=c:d$, they used the word *proportionalitas*.[5] That these terms were thoroughly grounded in the vernacular is seen today in the common use of such expressions as "divide this in the proportion of 2 to 3," and "your proportion of the expense," and in

[1] P. Tannery, "Du rôle de la musique grecque dans le développement de la mathématique pure," *Bibl. Math.*, III (3), 161.

[2] It may be said that medieval writers looked upon ratio and proportion as a branch of mathematics quite distinct from geometry and arithmetic. See also S. Günther, *Geschichte der Mathematik*, p. 180 (Leipzig, 1908); hereafter referred to as Günther, *Geschichte*.

[3] For a discussion of the terms λόγος (*lo'gos*), *ratio*, and *proportio*, see Heath, *Euclid*, Vol. II, pp. 116–129. See also Boethius, ed. Friedlein, p. 3 *et passim*.

[4] From the verb *reri*, to think or estimate; past participle, *ratus*. Hence *ratio* meant reckoning, calculation, relation, reason.

[5] Thus Boethius (*c.* 510): "Proportionalitas est duarum vel plurium proportionum similis habitudo," ed. Friedlein, p. 137; Jordanus Nemorarius (*c.* 1225): "Proportionalitas est si'litudo ꝓportionū" (1496 ed., Lib. 2).

In the Biblioteca Laurenziana at Florence is a MS. (Codex S. Marco Florent. 184) of Campanus (*c.*1260) *De proportione et proportionalitate* with the inscription *Tractatus Campani de proporcione et proporcionabilitate.* Included with it is a MS. of al-Miṣrî (*c.*900), *Epistola Ameti filii Joseph de proporcione et proportionalitate.* See *Bibl. Math.*, IV (3), 241; II (2), 7. The title of Pacioli's work affords another example: *Süma de Arithmetica Geometria Proportioni & Proportionalita*, Venice, 1494.

the occasional use of an expression like "the proportionality of the cost is the same as that of the amount."

That the word "proportion" was commonly used in medieval and Renaissance times to mean ratio is seen in most mathematical works of those periods.[1] It was so used by the American Greenwood (1729)[2] and has by no means died out in our language.[3]

The use of *proportio* for *ratio* was not universal in the early days of printing, however, for various writers used both terms as we use them today.[4]

General Types of Ratio. From the time of the Greeks to the 17th century the writers on theoretical arithmetic employed a set of terms and ideas in connection with ratio that seem to mathematicians of the present time unnecessarily complicated. A few of these have survived in our algebra, most of them have disappeared, and all of them had, under ancient conditions, good reasons for being. Of those which are still found in some of our textbooks there may be mentioned three general types of ratio of integers: namely, a ratio of equality,[5] like $a:a$; a ratio

[1] Thus Campanus (*c.* 1260): "Proportio est duarum quantitatum eiusdem generis ad inuicem habitudo" (Codex S. Marco Florent. 184); Jordanus Nemorarius (*c.* 1225): "Proportio est duaꝝ quātitatum eiusdem generis vnius ad alteram certa in quātitate relatio" (1496 ed., Lib. 2); Leonardo of Cremona: "la proporcion del diametro a la circonferentia" (original MS. in Mr. Plimpton's library; see *Rara Arithmetica*, p. 474); Chuquet (1484): "Proporcion cest labitude qui est entre deux nōbres quant est compare (lung) a laultre" (the Marre MS. in the author's possession, used by Boncompagni, *Bullettino*, XIII, 621); Rudolff (1526): "Die proporcion oder schickligkeit der ersten gegē der andern" (1534 ed., fol. E viij). Barrow (1670) used the expression in his lectures on geometry, and most other writers of the period did the same.

[2] " . . . the Proportion that each Figure bears to its neighbouring Figure" (p. 50).

[3] *E.g.*, Alison and Clark, *Arithmetic*, chap. xxi (Edinburgh, 1903).

[4] Thus Fine (Finæus): "Ratio igitur . . . est duarum quantitatum eiusdem speciei adinuicem comparatarū habitudo. . . . Proportio est, contingens inter comparatas adinuicem quantitates rationum similitudo" (*Protomathesis*, 1530–1532; 1555 ed., fols. 38 and 57). See also L. L. Jackson, *The Educational Significance of Sixteenth Century Arithmetic*, p. 119 (New York, 1906).

[5] The "aequalitatis proportio" of the Latin writers; *e.g.*, Scheubel (1545). In a numerical ratio like $a:b$, both a and b were generally considered integral unless the contrary was stated, but the incommensurable ratio of lines was recognized by the Pythagoreans and by all subsequent geometers.

of greater inequality, like $a:b$ when $a>b$; and a ratio of lesser inequality,[1] like $a:b$ when $a<b$. Of the last two there were recognized various subspecies, such as multiple ratio,[2] like $ma:a$, where m is integral; superparticular ratio, like $(m+1):m$ (which had several types, such as sesquialteran, as in the case of $3:2$, sesquitertian, as in the case of $4:3$, and so on); superpartient, like $(m+n):n$, where $m>n>1$, as in the cases of $5:3$ and $7:3$; multiple superparticular, like $(mn+1):m$, as in the cases of $7:3$ and $15:7$; and multiple superpartient, like $(mn+k):m$, where $m>k>1$, as in the cases of $14:3$ and $19:5$. These terms were capable of a large number of combinations and were essentially, from our present point of view, the result of an effort to develop a science of general fractions at a time when the world had no good symbolism for the purpose. With the introduction of our common notation and the invention of a good algebraic symbolism such terms disappeared.[3] This disappearance was hastened by such writers as Stifel (1546), who spoke out plainly against their further use, although his own acts were not always consistent with this statement.[4] When $\frac{9}{16}$ had to be called "suboctupla subsuperquadripartiens nonas" by a writer as late as 1600,[5] it was evident that the ancient usage must give way, and that ratios must be considered with respect to the modern fractional notation instead of depending upon the ancient Roman method.

[1] Boethius, ed. Friedlein, p. 238.

[2] Boethius (ed. Friedlein, p. 46) speaks of such relations: "Maioris vero inaequalitatis .V. sunt partes. Est enim una, quae vocatur multiplex, alia superparticularis, . . . " It should be observed that the Greeks did not consider ratios as numbers in the way that we do; that is, they did not consider $6:3$ as identical with 2 but as a relation of 6 to 3, this relation being a multiple ratio.

[3] For a full treatment of the subject see Pacioli, *Sūma*, 1494 ed., fol. 72.

[4] In his *Rechenbuch* (p. 35) he says: "Von den Proportzen. Zvm ersten | des Boetius | Stapulensis | Apianus | Christoff Rudolff | vnd andere gelerte Leuth | die proportiones leren | mit solche worten Multiplex | Duplex | Tripla | Superparticularis | Sesquialtera . . . vnnd der gleichen wort ohn zal | ist wol recht vnd nutzlich gelert Aber das man ein Teutschen Leser | dem die Lateinisch sprach ist vnbekant | will man solchen worten beladen | das ist ohn not | vnnd ohn nutz." In his *Coss Christoffs Rudolffs* (chap. 12), however, he gives "die fünferley proportionirte zalen," the multiple, superparticular, superpartient, multiplexsuperparticular, and the multiplexsuperpartient.

[5] Van der Schuere, 1624 ed., fol. 193.

Other Greek Ideas of Ratio. Certain other Greek ideas have come down to us and still find a place in our algebras. For example, we speak of $a^2 : b^2$ as the duplicate ratio[1] of a to b, although to double $a : b$ would give $2a : b$. To the Greek, however, the ratio $a_1 : a_n$ was considered as compounded or composed[2] of the ratios $a_1 : a_2, a_2 : a_3 \cdots, a_{n-1} : a_n$; and since $a^2 : b^2$ is similarly compounded of $a^2 : ab$ and $ab : b^2$ or of $a : b$ and $a : b$, it was called[3] the duplicate of $a : b$.

In like manner we have from the Greeks the idea of ratios compounded by addition when as a matter of fact they have been, according to our conception, multiplied.[4]

In the Middle Ages the distinction between ratios and fractions, or ratio and division, became less marked, and in the Renaissance period it almost disappeared except in cases of incommensurability.[5] An illustration of this fact is seen in the way in which Leibniz speaks of "ratios or fractions."[6]

Proportion as Series. The early writers often used *proportio* to designate a series,[7] and this usage is found as late as the 18th century.[8] The most common use of the word, however, limited it to four terms. Thus the early writers spoke of an arithmetic proportion, meaning $b - a = d - c$, as in 2, 3, 4, 5; and of a geometric proportion, meaning $a : b = c : d$, as in 2, 4, 5, 10. To

[1] Euclid's διπλασιων (*diplasi'on*), but commonly given by other Greek writers as διπλάσιος λόγος (*dipla'sios lo'gos*). See Heath, *Euclid*, Vol. II, p. 133.

[2] Heath, *Euclid*, Vol. II, p. 133.

[3] Euclid, *Elements*, VI, def. 5, apparently an interpolation. See Heath, *Euclid*, Vol. II, p. 189.

[4] See Heath, *Euclid*, Vol. II, p. 168. Similarly, Scheubel (1545, Tract. II) speaks "de proportionum Additione . . . siue ut alij Compositione," saying that the ratios 9 : 4 and 5 : 3 "componunt" 45 : 12.

[5] A modern Arab arithmetic, published at Beirut in 1859, remarks: "This division is called by the Magrebiner [West Arabs] 'the denomination,' but the Persians call it *al-nisbe* [the ratio]." H. Suter, *Bibl. Math.*, II (3), 17.

[6] ". . . aut in rationibus vel Fractionibus." Letter to Oldenburg, 1673.

[7] Thus Pacioli (1494): "che tu prendi li numeri . . . 1. 2. 3. 4. 5. 6. 7. 8. 9. 10 . . . hauerai la prima specie de la proportion" (*Sūma*, fol. 72, *r.*).

[8] When Vitalis (Geronimo Vitale) published his *Lexicon Mathematicvm* (Rome, 1690), the usage was apparently unsettled. He says: "Igitur Proportio Arithmetica est cum tres, vel plures numeri per eandem differentiam progrediuntur; vt 4. 7. 10. 13. 16. 19. 22. & sic procedendo in infinitum" (p. 681); but he also uses *proportio* in the modern sense (p. 732).

these proportions the Greeks[1] added the harmonic proportion

$$\frac{1}{b}-\frac{1}{a}=\frac{1}{d}-\frac{1}{c},$$

as where $a=\frac{1}{2}$, $b=\frac{1}{3}$, $c=\frac{1}{4}$, and $d=\frac{1}{5}$. These three names are now applied to series. To them the Greeks added seven others, all of which go back at least to Eudoxus (*c.* 370 B.C.).[2] The Renaissance writers began to exclude several of these,[3] and at the present time we have only the geometric proportion left, and so the adjective has been dropped and we speak of proportion alone.

Types of Proportion. The fact that geometric proportion has survived, in algebra at least, is largely due to Euclid's influence, since algebraically a proportion is nothing more than a fractional equation and might be treated as such. Especially is this true of such expressions as "by alternation," "by inversion," "by composition," "by division," and "by composition and division," three of which are now misnomers in the modern use of the words. They come to us directly from the Arabs,[4] who received them from Greek sources.[5] There were also various other types of geometric proportion besides the one commonly seen in textbooks,[6] but most of these types are now forgotten,

[1] On the theory in Euclid, see Heath, *Euclid*, Vol. I, p. 137; Vol. II, pp. 113, 119, 292.

[2] Boethius and certain of his predecessors gave all ten forms, those besides the three above mentioned being as follows:

$$a:c=b-c:a-c, \qquad a:c=a-c:b-c,$$
$$b:c=b-c:a-b, \qquad a:c=a-c:a-b,$$
$$a:b=b-c:a-b, \qquad b:c=a-c:b-c,$$
$$b:c=a-c:a-b.$$

See his *Arithmetica*, ed. Friedlein, p. 137 seq.; Günther, *Math. Unterrichts*, p. 85; Cantor, *Geschichte*, I, chap. xi, for the earlier knowledge of these forms.

[3] Thus Ramus: "Genera autē proportionis duo tantum instituimus, quia haec sola simplicia & mathematica sunt. Nicomachus fecit decem. Jordanus addidit undecimam." *Scholarvm Mathematicarvm, Libri vnvs et triginta*, p. 134 (Basel, 1569).

[4] *E.g.*, see al-Karkhî (*c.* 1020), the *Kâfî fîl Hisâb*, ed. Hochheim, II, 15.

[5] Heath, *Euclid*, Vol. I, p. 137; Vol. II, pp. 113, 119, 133, 168, 189, 292.

[6] *E.g.*, Scheubel (1545): "Sex sunt species proportionalitatis, permuta, conuersa, coniuncta, disiuncta, euersa, & aequa" (Tractatus II).

although "continued proportion," but with a change in the older meaning,[1] has survived both in algebra and in geometry.

Terms Used in Proportion. The terms "means," "antecedent," and "consequent" are due to the Latin translators of Euclid.[2] There have been attempts at changing them, as when the antecedent was called a leader and the consequent a comrade,[3] but without success. It would be quite as simple to speak of them as the first, second, third, and fourth terms.[4]

Rule of Three. The mercantile Rule of Three seems to have originated among the Hindus. It was called by this name by Brahmagupta (*c.* 628) and Bhāskara[5] (*c.* 1150), and the name is also found among the Arab and medieval Latin writers.

Brahmagupta and Mahāvīra state the Rule. Brahmagupta stated the rule as follows: "In the Rule of Three, Argument, Fruit, and Requisition are the names of the terms. The first and last terms must be similar. Requisition multiplied by Fruit, and divided by Argument, is the Produce."[6] Mahāvīra (*c.* 850) gave it in substantially the same form; thus: "*Phala* multiplied by *Icchā* and divided by *Pramāṇa* becomes the answer, when the *Icchā* and *Pramāṇa* are similar."[7]

For example: "A lame man walks over $\frac{1}{8}$ of a *krōśa* [32,000 feet] together with $\frac{1}{5}$ [thereof] in $7\frac{1}{2}$ days. Say what [distance] he [goes over] in $3\frac{1}{5}$ years [at this rate],"—a very good illustration of the absurdity of the Oriental problem.

[1] *E.g.*, Fine (Finæus, 1530) defines a *proportio continua* as one like "8/4/2/1: ut enim 8 ad 4, sic 4 ad 2, atq3 2 ad 1" (*De Arithmetica Practica*, 1555 ed., fol. 59), and a *proportio disjuncta* as (to use his symbolism) one like 8/4, 6/3.

[2] Euclid used μεσότητες (*mesot'etes*, means), ἡγούμενα (*hegou'mena*, leading [terms], antecedents), and ἑπόμενα (*hepom'ena*, following [terms], consequents), but he had no need for "extremes." See *Elements*, VII, 19.

[3] Thus Scheubel (1545): " . . . alter antecedens uel dux, alter consequens uel comes appellatur." The use of *dux* (duke, leader) comes from Euclid's term.

[4] *E.g.*, see Clavius (1583), *Epitome Arithmeticae Practicae*, chap. xvii.

[5] One of the scholiasts of Bhāskara called it the *Trairāśica*, the "three rule." See Colebrooke's translation, pp. 33, 283. On the general subject see Taylor's translation, p. 41.

[6] Colebrooke's translation, p. 283.

[7] *Gaṇita-Sāra-Sangraha*, p. 86. The *phala* is the given quantity corresponding to what is to be found; the *pramāna* was a measure of length, but in proportion it is the term corresponding to the *icchā*; the *icchā* is the third term in the rule.

Bhāskara (*c.* 1150) gave the rule in much the same form as that used by Brahmagupta, thus: "The first and last terms, which are the argument and requisition, must be of like denomination; the fruit, which is of a different species, stands between them; and that, being multiplied by the demand [that is, the requisition] and divided by the first term, gives the fruit of the demand [that is, the Produce]."[1]

As an example, Bhāskara gave the following: "Two palas and a half of saffron are purchased for three sevenths of a niska: How many will be purchased for nine niskas?"

His work appears[2] as

3 5 9
7 2 1

In our symbolism it might be represented as $\frac{3}{7}\ N.\ 2\frac{1}{2}\ P.\ 9\ N.$ It is thus seen that the idea of equal ratios is not present, as would be the case if we should write $x : 2\frac{1}{2} = 9 : \frac{3}{7}$. Proportion was thus concealed in the form of an arbitrary rule, and the fundamental connection between the two did not attract much notice until, in the Renaissance period, mathematicians began to give some attention to commercial arithmetic. One of the first to appreciate this connection was Widman[3] (1489), and in this he was followed by such writers as Tonstall[4] (1522), Gemma Frisius[5] (1540), and Trenchant[6] (1566).

Names for the Rule of Three. Recorde (*c.* 1542) calls the Rule of Three "the rule of Proportions, whiche for his excellency is called the Golden rule,"[7] although his later editors called it by the more common name.[8] Its relation to algebra was first strongly emphasized by Stifel[9] (1553–1554).


[1] Colebrooke's translation, p. 33. [2] See Taylor's translation, p. 41.

[3] "Sy ist auch recht genāt regula proportionū/wā in d' regel werdē erkāt vñ erfundē alle ꝑportiones" (1508 ed., fol. 50). [4] *De Arte Supputandi*, Lib. III.

[5] He calls his chapter "De Regvla Proportionvm, siue Trium Numerorum" (1575 ed., fol. C 6).

[6] "La regle de troys, qui est la regle des proportions ou proportionaux" (1578 ed., p. 120). [7] 1558 ed., fol. M 4. [8] *E.g.*, John Mellis, 1594 ed., p. 449.

[9] "Gar wunderbarlich wickeln vnd verknüpffen sich zusammen die Detri vnd die Coss also dass die Coss im grund auch wol möchte genennt werden die Detri. . . . Vñ steckt also die gantz Coss in der Regel Detri/widervmb steckt die Gantz Detri in der Coss." See the *Abhandlungen*, I, 86.



RULE OF THREE IN THE SIXTEENTH CENTURY

From an Italian MS. of 1545. Notice the arrangement of terms; also the early per cent sign as given at the end of the fifth line. From a manuscript in Mr. Plimpton's library

When the rule appeared in the West, it bore the common Oriental name,[1] although the Hindu names for the special terms were discarded. So highly prized was it among merchants, however, that it was often called the Golden Rule,[2] a name apparently in special favor with the better mathematical writers.[3] Hodder, the popular English arithmetician of the 17th century, justifies this by saying: "The Rule of Three is commonly called, *The Golden Rule*; and indeed it might be so termed; for as Gold transcends all other Mettals, so doth this Rule all others in Arithmetick."[4] The term continued in use in England until the end of the 18th century at least,[5] perhaps being abandoned because of its use in the Church.

[1] Thus Pacioli (1494) calls it the "regula trium rerum la regola ditta dl̄ .3. ouer de le .3. cose," and "la regola del .3."; the Treviso arithmetic (1478), "La regula de le tre cose"; Pellos (1492), the "Regula de tres causas." Chuquet (1484) remarks, "La rigle de troys est de grant recōmandacion. . . . La rigle de troys est ansi appellee pource quelle Requiert tousiours troys nombres"; Grammateus (1518) speaks of the "Regula de tre in gantzē" and "in prüchen," and Rudolff (1526) of "Die Regel de Tri," a term often abridged by German writers into "Regeldetri," as in the work of Licht (1500). Klos, the Polish writer of 1538, also calls it the "Regula detri."

[2] Thus Petzensteiner (1483): "Vns habn̄ die meyster der freyn kunst vō d' zal ein regel gefundē die heist gulden regel Dauo das sie so kospar vnd nucz ist. . . . Sie wirdet auch geneñet regula d' tre nach welsischer [*i.e.*, Italian] zungen. . . . Sie hat auch vile āder namē"; and Köbel (Augsburg edition of 1518) speaks of "Die Gulden Regel (die von den Walen de Try genant wirt)." In Latin it often appears as *regula aurea*. The Swedish savant, Peder Månsson, writing in Rome *c.* 1515, speaks of the rule "quam nonnulli regulam auream dixere: Itali vero regulam de tri" (see *Bibl. Math.*, II (2), 17).

The French writers used the same expression. Thus Peletier (1549): "La Reigle de Trois . . . vulgairement ansi dite. . . . Les ancients l'ont appellee la Reigle d'or: parce que l'invention en est tres ingenieuse, & l'usage d'icelle infini" (1607 ed., p. 68).

[3] Thus Ciacchi (1675): "La Regola del Tre così chiamata da' Practici vulgari, e da' Mattematici regola d' oro, o pure delle quattro proporzioni è principalissima, ed apporta vn' inestimabile benefizio, ed vna gran comodità a' Mercanti." *Regole generali d' abbaco*, p. 121 (Florence, 1675).

[4] See the tenth edition, 1672, p. 87. This simile was a common one with writers; thus Petzensteiner (1483): "als golt vbertrifft alle ander metall."

Vitalis (Geronimo Vitale), in his *Lexicon Mathematicvm*, p. 748 (Rome, 1690), says: "Quare meritò *Aurea* appellata est; namque plus auro valet: & non Arithmeticis modo, Geometris . . . necessaria est; sed & vniuerso hominum generi, in commercijs ineundis, . . ."

[5] *E.g.*, the 1771 edition of Ward's *Young Mathematician's Guide* (p. 85) speaks "of Proportion Disjunct; commonly called the Golden Rule."

Regula de Tri
oder gulden Regul.

REgula de Tri MERCATORum/
Genannt aurea proportionum
Darumb das ſie gar bequemlich
Im Kauff/begert drey ding/namlich
Den Kauff/das Werth/die Frag zum dritten/
Das Werth ſoll ſtet ſtehen in der mitten/
Der Kauff vornen/die Frag dahinden/
Wilt du die Frag vnd Facit finden/
Hinden/vornen gleich Namen richt/
Die klein die groß allzeit zerbricht/
Mult plicier die hinder Zahl
Mit der mittleren allemahl/
Theil s Produckt mit dem vordern ab/
So kombt dir dein Frag vnd Auffgab.
Verkehrt Regul wilt dein Prob finden/
Was erſtlich gſtanden iſt dahinden
Muß vornen/s forder hinden ſtehen/
S Facit muß in die mitten gehen/
Die mitler Zahl rauß kommen ſoll/
Kombt ſie/ſo haſts getroffen wol.

16.

RULE OF THREE, OR THE GOLDEN RULE, IN VERSE

From Lautenschlager's arithmetic (1598)

The Merchants' Rule. Its commercial uses also gave to the Rule of Three the name of the Merchants' Key or Merchants' Rule,[1] and no rule in arithmetic received such elaborate praise as this one which is now practically discarded as a business aid.[2]

A Rule without Reason. The rule was usually stated with no explanation; thus Digges (1572) merely remarked, "Worke by the Rule ensueing. . . . Multiplie the last number by the seconde, and diuide the Product by the first number," and similar statements were made by most other early arithmeticians, occasionally in verse.[3]

The arrangement of the terms was the same as in the early Hindu works, the first and third being alike. As Digges expressed it, "In the placing of the three numbers this must be observed, that the first and third be of one Denomination."[4] This custom shows how completely these writers failed to recognize the relation between the Rule of Three and proportion.

[1] Thus Licht (1500, fol. 9) says: "Regula Mercatorum. [Q]uam detri. quia de trib⁹. per apocopam appellamus. Regula Aurea docte ac perite ab omib⁹ appellari videt deberi"; Clavius (1583) calls it the "clavis mercatorum"; Peletier (1549) says, "Mesmes, aucuns l'ont nommee la Clef des Marchands" (1607 ed., p. 68); Wentsel (1599) speaks of the "Regvla avrea mercatoria/ regvla de tri, Regvla van dryen/ regle a trois, &c."; and Lautenschlager, in his arithmetic in rime (1598), has, as shown in the facsimile on page 487,

> Regula de Tri | oder gulden Regul.
> REgnla de Tri MERCATORum |
> Genannt aurea proportionum.


Of the various other names, *Schlussrechnung* has continued among the Germans. See R. Just, *Kaufmännisches Rechnen*, I. Teil, p. 75 (Leipzig, 1901).

[2] A few of the hundreds of eulogies given to the rule are as follows: Gemma Frisius (1540), "Res breuis est & facilis, vsus immensus, cum in vsu communi, tum in Geometria ac reliquis artibus Mathematicis" (1563 ed., fol. 18); Adam Riese (1522), "Ist die furnameste vnder allē Regeln" (1550 ed. of *Rechenung*, fols. 13, 59); Clavius (1583; *Opera*, 1611, II, 35), "Primo autem loco sese offert regula illa nunquam satis laudata, quae ob immensam vtilitatem, Aurea dici solet, vel regula Proportionum, propterea quod in quatuor numeris proportionalibus, quorum priores tres noti sunt . . . vnde & regula trium apud vulgus appellata est," showing that he, like Stifel, recognized the relation to proportion; Van der Schuere (1600), "Den Regel van Drien, die van vele ten rechte den Gulden Regel genaemd word/ overmits zyne weerdige behulpzaemheyd in alle andere Regelen" (1634 ed., fol. 12).

[3] *E.g.*, Lautenschlager (1598) and Sfortunati (1534; 1545 ed., fol. 33).

[4] 1579 ed., p. 29.

Once set down, it was the custom to connect the terms by curved lines, as in the following cases:[1]



The Arabs, however, used such forms as

1	5
3	15

and

2	4
3	6

to indicate a proportion, paying no attention to the labels on the numbers.[2]

Arrangement of Terms. The rule being purely arbitrary, it became necessary to have this arrangement in the proper order, and the early printed books gave much attention to it. Borghi (1484) gave a whole chapter to this point,[3] and Glareanus (1538) arranged[4] an elaborate scheme to help the student.[5] Later writers, however, recognized that if the rule were to be considered as a case of proportion, it would be necessary to rearrange the terms so that the first two should be alike. Thus in place of a form like

12 yards———20s.———6 yards,

[1] The first of these is from a 17th century MS. in the author's library; the second is from Werner's *Rechenbuch*, 1561, fol. B 2.

[2] E. Wiedemann, "Über die Wage des Wechselns von Châzinî und über die Lehre von den Proportionen nach al-Bîrûnî," *Sitzungsberichte der Physik.-med. Societät zu Erlangen*, 48. u. 49. Bd., p. 4.

[3] "¶Como le tre cose contenute in detta regola sono ordinate, e quale debbi esser prima, e qual seconda, e qual terza" (1540 ed., fol. 36).

[4] 1543 ed., fol. 20.

[5] He arranged his rule thus:

Sinistra	Medius locus	Dextera
Res empta	Numer⁹ pretij	Numer⁹ q̄stionis
Diuisor	Multiplicādus	Multiplicator
2	36	7

as given in Hodder's 17th century work,[1] we find Blassière (1769)[2] and others of the 18th century using such forms as

Ellen		Ellen		Guld.		Guld.
3	:	36	=	4	:	x

In the old Rule of Three the result was naturally written at the right, and for this reason the unknown quantity came to be placed at the right in the commercial problems in proportion.[3]

Inverse Proportion. Of the various special forms of the Rule of Three the one known as inverse proportion is the simplest. It results when the ratio of two quantities is equal to the reciprocal of the ratio of two quantities which seem to correspond to them. Bhāskara (*c.* 1150) gives an illustration: "Bullocks which have plowed four seasons cost four niskas: what will bullocks which have plowed twelve seasons cost?"[4]

This rule went by such names as the inverse, converse, or everse Rule of Three.[5] Recorde (*c.* 1542) used a name that became quite common in England, remarking: "But there is a contrarye ordre as thys: That the greater the thyrde summe is aboue the fyrste, the lesser the fourthe summe is beneth the second. and this rule you maye call the Backer rule."[6]

[1] *Arithmetick*, 10th ed. (1672), p. 89. [2] 1790 ed., p. 149.

[3] Thus Rabbi ben Ezra (*c.* 1140) wrote

47	63
7	o

for $47:7 = 63:x$, the o standing for the unknown. In the translation of certain Arab works the unknown is placed first, as in $x:84 = 12:7$, because the Arabs wrote from right to left.

[4] Taylor's *Lilawati*, p. 42, with spelling as given by Taylor. The result is written "niska 1, and fraction $\frac{1}{7}$."

[5] Thus Köbel (1514), "Die Regel de Tri verkert/ im Latein Regula conuersa genant" (1549 ed., fol. 68); Gemma Frisius (1540), "Regvla Trivm Euersa"; Albert (1534), "Regula Detri Conuersa"; Thierfelder (1587), "Regula Conversa/ oder vmbekehrte Regel de Try"; Van der Schuere (1600), "Verkeerden Reghel van Drien"; Digges (1572), "The Rule of Proportion Inuersed."

[6] 1558 ed., fol. M 6. In some later editions it is called "the Backer or Reverse Rule" (1646 ed., p. 180). See also Baker (1568; 1580 ed., fol. 46). In French it appeared as "La regle de troys rebourse" (Trenchant, 1566; 1578 ed., p. 155), "La Reigle de Trois Reuerse ou Rebourse" (Peletier, 1549; 1607 ed., p. 74), and "La Regle Arebourse" (Coutereels, Dutch and French work, 1631 ed., p. 204). It was occasionally called the bastard Rule of Three. Thus Santa-Cruz (1594): "Exemplo de la regla de tres bastarda."

In this rule it was the custom to leave the terms as in simple proportion but to change the directions for solving. Hylles (1592) gives the rule as follows:

> The Golden rule backward or conuerst,
> Placeth the termes as dooth the rule direct:
> But then it foldes[1] the first two termes rehearst,
> Diuiding the product got by that effect.
> Not by the first, but onely by the third,
> So is the product the fourth at a word.[2]

Compound Proportion. What has been called, for a century or two, by the name of compound proportion originally went by such names as the Rule of Five[3] when five quantities were involved, the Rule of Seven if seven quantities were used, and so on. Bhāskara, for example, gives rules of five, seven, nine, and eleven.[4] Peletier (1549) speaks of Ptolemy as the inventor of the Rule of Six, referring, however, to the proposition in geometry relating to a transversal of three sides of a triangle.[5] The names beyond that for five were rarely used;[6] indeed, all beyond that for three were more commonly called by the general name of Double Rule of Three,[7] Compound Rule of

[1] An interesting translation of *plicare* as found in *multiplicare* (to manifold).

[2] *Arithmeticke*, 1600 ed., fol. 135.

[3] It appears in Bhāskara (*c.* 1150) as *pancha-rāśica* (five-rule). See Taylor's translation, p. 43; but the spellings in Colebrooke's translation, p. 37, are here followed. In Europe we find such names as "Regvle de cinqve parte" (Ortega, 1512; 1515 ed.); "Regula Quinque/oder zwyfache Regel de Try" (Thierfelder, 1587); "Regel von fünffen" (Rudolff, 1526); "Die Regel von fünff zalen" (Köbel, 1514; 1531 ed.); "*Regvla dvplex* Auch *Regula Quinque* genant . . . die zwyfache Regel . . . von fünff Zalen" (Suevus, 1593); "den Zaamengestelden Reegel van Drien. Anders Genaamd Den Reegel van Vyven" (Blassière, 1769); and "Den Regel van Vyven, of anders genaamt den Dobbelen Regel van Drien" (Bartjens, 1792 ed.).

[4] *Pancha-rāśica*, *sapta-rāśica*, *nava-rāśica*, and *écádaśa-rāśica*, as spelled by Colebrooke.

[5] "La Reigle de six Quantités a esté inuentee par Ptolemee" (1607 ed., p. 220). On this see also Cardan's *Practica* (1539) with its "Caput 46. de regula 6. quātitatum."

[6] Thus Ciacchi (1675) says: "Non è molto vsitata da' practici Arrimetici la regola del sette."

[7] As by Recorde (*c.* 1542). The name also appears under such forms as "Regula duplex" (Gemma Frisius, 1540) and "la regle double" (Trenchant, 1566).

Three,[1] Conjoint Rule,[2] Plural Proportion,[3] and, finally, Compound Proportion, a term which became quite general in the 18th century.

Artificial Nature of the Problems. The artificial nature of the problems in compound proportion has been evident from the beginning. Thus Mahāvīra (*c.* 850) gives this case: "He who obtains 20 gems in return for 100 gold pieces of 16 *varṇas*—what [will he obtain] in return for 288 gold pieces of 10 *varṇas*?"[4] And Bhāskara (*c.* 1150) has this type: "If eight best, variegated silk scarfs measuring three cubits in breadth and eight in length cost a hundred [nishcas]; say quickly, merchant, if thou understand trade, what a like scarf three and a half cubits long and half a cubit wide will cost."[5]

Practice. There appears in certain English arithmetics of the present day a chapter on Practice, a kind of modification of the Rule of Three. In the manuscripts of the later Middle Ages and in the early printed books of Italy the word is used to mean simply commercial arithmetic in general, whence possibly the origin of our phrase "commercial practice" today.[6] When northern writers of the 16th century spoke of Italian practice they usually referred merely to Italian commercial arithmetic in general.[7] In the 17th century the Dutch writers generally used the term *Practica* (*Practijcke*) to mean that part of arith-

[1] "Regula Trivm Composita" (Clavius, 1583); "The Golden Rule compound" (Recorde, *c.* 1542; 1646 ed., p. 195); "Den menichvuldigen Regel" (Coutereels, 1631 ed., p. 213); "Gecomposeerden dubbelden Regel" (Houck, 1676, with a distinction between this and the mere "Dubbelden Regel" and "Regel van Conjuncte").

[2] "Den Versamelden Reghel" or "La Regle Conjoincte" (Coutereels, 1631 ed., p. 219).

[3] "The Double Rule of Three . . . Under this Rule is comprehended divers Rules of Plural Proportion" (Hodder, *Arithmetick*, 10th ed., 1672, p. 131).


[4] English translation, p. 91.

[5] Colebrooke's translation, p. 37.

[6] So Tartaglia (1556) has a chapter "Della Prattica Fiorentina" for Florentine commercial arithmetic, and Riese (1522) has "Rechenung . . . die practica genandt."

[7] So Stifel (1544): "Praxis Italica Praxis illa quam ab Italis ad nos devolutam esse arbitramur." Even as late as 1714 there was a chapter in Starcken's arithmetic on "Die Italianische Practica/ oder Kurtze Handels-Rechnung."

metic relating to financial problems,[1] and they also used the expression "Italian Practice," as in the work by Wentsel (1599) of which part of the title is here shown in facsimile. The term "Welsh practice" had a similar meaning, the word "Welsh" ("Welsch") signifying foreign.[2] This expression is often found in the German arithmetics of the 16th century.


T' FONDAMENT
Van Arithmetica: mette Ita-
liaensche Practijck / midtsgaders d'aller
noodtwendichste stucken van den Reghel van
Interest.
Beydes in Nederduytsende in Franchois/
met redelicke ouereenstemminghe ofte
Concordantien.
Alles
Door MARTINVM VVENCESLAVM,
AQVISGRANENSEM.

WELSH OR ITALIAN PRACTICE

Part of the title-page of the arithmetic of Wentsel (Wenceslaus) (159

In England practice came to mean that part of commercial arithmetic in which short processes were used. Baker (1568) mentions it in these words:

Some there be, whiche doe call these rules of practise, breefe rules: for that by them many questions may bee done with quicker expedition, then by the Rule of three. There be others which call them the small multiplication, for because that the product is alwayes lesse in quantity, than the number whiche is to be multiplyed.

[1] Thus Eversdyck's edition of Coutereels (1658 ed., p. 91), Stockmans (1676 ed., p. 173), Van der Schuere (1600, fol. 49), and Mots (1640, fol. G 5).

[2] Modern German *wälsch*, foreign; particularly Gallic, Roman, Italian. So we find Rudolff (1526) speaking of "Practica oder Wellisch Rechnung" and Helmreich (1561) having a chapter on "De Welsche Practica oder Rechnung." Dutch writers commonly used "foreign" instead of "Welsch," and so we often find chapters on "Buyten-lantsche Rekeninghe."

This rather indefinite statement gave place to clearer definitions as time went on, and Greenwood (1729) speaks of practice as follows:

THIS *Rule* is a contraction or rather an Improvement of the *Rule of Three*; and performs all those Cases, where *Unity* is the *First Term*; with such Expedition, and Ease, that it is, in an extraordinary manner, fitted to the *Practice* of Trade, and Merchandise; and from thence receives its *Name.*

A single example from Tartaglia (1556) will show the resemblance of the Italian solution to that which was called by American arithmeticians the "unitary method," especially in the 19th century:

If 1 pound of silk costs 9 *lire* 18 *soldi*, how much will 8 ounces cost?[1]

The solution is substantially as follows:

1 lb. costs 9 *lire* 18 *soldi*,
4 oz. cost $\frac{1}{3}$ of this, or 3 *lire* 6 *soldi*,
8 oz. cost twice this, or 6 *lire* 12 *soldi.*

11. SERIES

Kinds of Series. Since the number of ways in which we may have a sequence of terms developing according to some kind of law is limitless, like the number of laws which may be chosen, there may be as many kinds of series or progressions[2] as we wish. The number to which any serious attention has been paid in the development of mathematics, however, is small. The arithmetic and geometric series first attracted attention, after which the Greeks brought into prominence the harmonic series. These three were the ones chiefly studied by the ancients. Boethius (*c.* 510) tells us that the early Greek writers knew these three, but that later arithmeticians had suggested three others which had no specific names.[3]

[1] 1 lb. = 12 oz., 1 *lira* = 20 *soldi.*

[2] For our purposes we shall not distinguish at present between these terms.

[3] "Vocantur autē quarta: quinta: vel sexta" (*Arithmetica*, 1st ed., 1488, II, cap. 41; ed. Friedlein, p. 139).

Occasionally some special kind of series is mentioned, as when Stifel speaks of the "astronomical progression" 1, $\frac{1}{60}$, $\frac{1}{3600}$, $\cdots$,[1] one of the few instances of a decreasing series in the early European books.

Most of the Hindu writers used only two elementary series, but Brahmagupta (*c.* 628), Mahāvīra (*c.* 850),[2] and Bhāskara (*c.* 1150) all considered the cases of the sums of squares and cubes.[3] The Arab[4] and Jewish[5] writers also gave some attention to these several types.

Medieval Treatment of Series. In the medieval works a series was generally considered as ascending, although descending series had been used by Ahmes, Archimedes, and certain Chinese writers long before the time in which these works were written.[6] The same custom was followed by the early Renaissance writers.[7]

Somewhat better known than the classification of series as arithmetic, geometric, and harmonic, at least before the 17th century, was the classification into natural, nonnatural, continuous, and discontinuous, these terms being used rather loosely

[1] *Arithmetica Integra*, 1544, fol. 64, the name being *astronomica progressio*. It is simply the natural series of "astronomical fractions."

[2] English translation, p. 170. His rule for the sum of the squares is substantially

$$s = n\left\{\left[\frac{(2n-1)d^2}{6} + ad\right](n-1) + a^2\right\}.$$

[3] Colebrooke's translation, p. 52. Bhāskara remarks: "Former authors have stated that the sum of the cubes of the terms one, &c. is equal to the square of the summation"; that is, $\Sigma n^3 = (\Sigma n)^2$. (Taylor's translation, p. 60.)

[4] *E.g.*, al-Ḥaṣṣâr (*c.* 1175?); see *Bibl. Math.*, II (3), 32. They are also given by al-Qalaṣadî (*c.* 1475); see Boncompagni's *Bullettino*, XIII, 277.

[5] *E.g.*, Rabbi ben Ezra (*c.* 1140), in the *Sefer ha-Mispar*, Silberberg translation, p. 120.

[6] So Fibonacci (1202) says: ". . . colligere numeros quotcumque ascendentes ab ipso dato numero equaliter, ut per ascensionem unitatis, uel binarii, uel ternarii . . ." (*Liber Abaci*, p. 166 (fol. 70, *r.*)).

[7] Thus Stifel, in his edition of Rudolff's *Coss* (1553): "Es ist aber Progressic (eygentlich zu reden nach der Arithmetica) ein ordnung vieler zalen so nach einander auffsteygen oder absteygen nach eyner rechten richtigen Regel" (fol 7, *v.*). So Trenchant (1566) states definitely that the terms must increase, and Chuquet (1484) says: "Progression est certaine ordonnance de nombre par laquelle le premier est surmonte du second dautant que le second est surmonte du tiers et ꝑ sequēment les ault's se plus en ya" (fol. 20, *r.*).

by early writers. For example, the series 1, 2, 3, . . . was called a natural series,[1] from which we have the expression "natural series of numbers."[2] A discontinuous, or intercised,[3] progression was one in which the difference was not unity.[4]

Name for Series. The Greek name for a series, as used first by the early Pythagoreans, was ἔκθεσις (*ek'thesis*),[5] literally a selling out, and the name for a term of the series was ὅρος (*hor'os*),[6] literally a boundary. Boethius (*c.* 510), like the other Latin writers, used the word *progressio*,[7] and this was generally the custom until modern times.

The Teutonic writers followed their usual plan of avoiding international names based upon the Latin, and so we find various terms used by the Dutch[8] and German[9] mathematicians.

[1]Thus Chuquet (1484): "Et doit on sauoir que progression se fait en plusieurs et diuerses manieres. Car aulcunesfoiz elle cõmance a .1. et progredyst par .1. cõme .1. 2. 3. 4. ℈c. telle est appellee par les anciens progression naturelle ou continue ꝓgression" (fol. 20, *r.*). Similarly, Pellos (1492) speaks of "ꝓgression natural" (1, 2, 3, . . .), "ꝓgression nõ natural" (1, 3, 5, . . .), and "ꝓgression ni part natural ℈ ni part nõ natural" (8, 9, 10, . . .).

Van der Schuere (1600), however, calls any series like 1, 2, 3, . . . or 1, 3, 5, . . . a "natuerlikke overtredinghe /oft Aritmetische Progressio," speaking of a geometric series as "onnatuerlikke overtredinghe."

[2]This is found in Stifel (1544): "naturalis numerorum Progressio, est Progressio Arithmetica prograediens ab unitate per binarium ad reliquos numeros secundum differentiam unitatis. ut

$$1.^{1.}\ 2.^{1.}\ 3.^{1.}\ 4.^{1.}\ 5. \ldots"$$

He also speaks "de Progressione naturali numerorum imparium," viz., 1, 3, 5, 7, . . . (*Arithmetica Integra*, fols. 20, 21).

[3]*Intercissa* (Huswirt, 1501), *int'cise* (Chuquet, 1484), *vnderschnitten* (Köbel, 1514).

[4]"Alcune cõmãce a .1. mais el progredist par aultre nombre que .1. cõe 1. 3. 5. ℈c. ou .1. 4. 7. ℈c. et est ceste appellee Int'cise progression ou prog'ssion discontinuee" (Chuquet (1484), fol. 20, *r.*; Boncompagni's *Bullettino*, XIII, 617).

Santa-Cruz (1594) says that any progression "começando de la vnidad dicha continua."

[5]A word also meaning exhibition or exposition.

[6]A word also meaning a limit, marking stone, rule, standard, or boundary between two objects.

[7]Ed. Friedlein, I, pp. 9, 10, *et passim.*

[8]Dutch writers of the 16th century used *progressio* together with such terms as *overtredinghe* (stepping over) and *opklimminge* (ascending; literally, upclimbing).

[9]Although modern writers use *Reihe*. Köbel (1514), for example, says: "Die acht species ist Progressio zu Latein/vnnd ist Fürzelen geteutscht," in later German *Fürzählung*. The terms *Aufsteigung*, *Fortgehung*, *Reihe*, and *Progression* are also used.

The change to the name "series" seems to have been due to writers of the 17th century. James Gregory, for example, writing in 1671, speaks of "infinite serieses," and it was in connection with infinite sequences that it was at first used by the British algebraists. Even as late as the 1693 edition of his algebra, however, Wallis used the expression "infinite progressions" for infinite series.

Extent of Treatment. Although series was commonly looked upon as one of the fundamental operations,[1] it was rarely accorded much attention in the early printed books. Tzwivel (1505), for example, gives only 32 lines to both arithmetic and geometric progressions, including all definitions and rules; while Huswirt (1501) allows only one page and Digges (1572) only two pages to the subject.

Nearly all the early writers limited the work to finding the sum of the series,[2] although a few gave a rule for finding the last term of an arithmetic or a geometric series. With these writers there was no attempt to justify the rule, the mere statement sufficing. It was only through the influence of a better algebraic symbolism in the 17th century that the various cases could easily be discussed and the development of rules for all these cases made simple.

Relation to Proportion. The ancient writers commonly connected progression with proportion, or rather with proportionality, to use a name which, as already stated, was at one time popular; and they applied the names "arithmetic," "geometric," and "harmonic" to each. Some of the early printed books call attention to this relation, saying that a proportion is merely a progression of four terms.[3]

[1] Thus Pacioli (1494): "la sexta e penultīa specie dilla pratica d' arithcª. laq̈le e chiamata ꝑgressiõe" (*Sūma*, fol. 37).

[2] Johann Albert (1534) distinctly states that this is the sole purpose of the work: "Progredirn leret/wie man viel zaln (welche nach natürlicher ordnung oder durch gleiche mittel/nach einander folgen) in eine Summa/auffs kürtzest vnd behendest bringen sol" (1561 ed., fol. E 4). See also Treutlein, *Abhandlungen*, I, 60.

[3] Thus Trenchant (1566): ". . . car Progression n'est qu'vne continuation des termes d'vne proportion" (1578 ed., p. 274).

Arithmetic Series. The first definite trace that we have of an arithmetic series as such is in the Ahmes Papyrus (*c.* 1550 B.C.), where two problems are given involving such a sequence. The first[1] of the problems is as follows: "Divide 100 loaves among five persons in such a way that the number of loaves which the first two receive shall be equal to one seventh of the number that the last three receive."

The solution shows that an arithmetic progression is understood, in which $n = 5$, $s_5 = 100$, and

$$\frac{(a+4d)+(a+3d)+(a+2d)}{7} = (a+d)+a.$$

Then, by modern methods, $2d = 11a$.

Therefore $$100 = s = \frac{2a+4d}{2} \cdot 5 = 60a,$$

whence $a = 1\frac{2}{3}$ and $d = 9\frac{1}{6}$.

Therefore the series is $1\frac{2}{3}$, $10\frac{5}{6}$, 20, $29\frac{1}{6}$, $38\frac{1}{3}$, although the method here given is not the one followed by Ahmes.

The second problem,[2] with its solution as given by Ahmes, reads as follows:

Rule of distributing the difference. If it is said to thee, corn measure 10, among 10 persons, the difference of each person in corn measure is $\frac{1}{8}$. Take the mean of the measures, namely 1. Take 1 from 10, remains 9. Make one half of the difference, namely, $\frac{1}{16}$. Take this 9 times. This gives to thee $\frac{1}{2}\ \frac{1}{16}$. Add to it the portion of the mean. Then subtract the difference $\frac{1}{2}$ from each portion, [this is in order] to reach the conclusion. Make as shown:

$1\ \frac{1}{2}\ \frac{1}{16}$, $1\ \frac{1}{4}\ \frac{1}{8}\ \frac{1}{16}$, $1\ \frac{1}{4}\ \frac{1}{16}$, $1\ \frac{1}{8}\ \frac{1}{16}$, $1\ \frac{1}{16}$,

$\frac{1}{2}\ \frac{1}{4}\ \frac{1}{8}\ \frac{1}{16}$, $\frac{1}{2}\ \frac{1}{4}\ \frac{1}{16}$, $\frac{1}{2}\ \frac{1}{8}\ \frac{1}{16}$, $\frac{1}{2}\ \frac{1}{16}$, $\frac{1}{4}\ \frac{1}{8}\ \frac{1}{16}$.

This may be stated in modern form as follows: Required to divide 10 measures among 10 persons so that each person shall have $\frac{1}{8}$ less than the preceding one.

[1] Problem 40 in the Eisenlohr translation, p. 72; Peet, *Rhind Papyrus*, p. 78.

[2] No. 64 of the Eisenlohr translation. The version here given is furnished by Dr. A. B. Chace. For another translation see Peet, *loc. cit.*, p. 107.


That is, $n = 10$, $s_n = 10$, $d = -\frac{1}{8}$, so that

$$s_n = 10 = \frac{2a + (n-1)d}{2} \cdot n = (2a - \tfrac{9}{8}) \cdot 5,$$

whence $a = 1\frac{9}{16}$, and the series is the descending progression

$$1\tfrac{9}{16}, \quad 1\tfrac{7}{16}, \quad 1\tfrac{5}{16} \quad \cdots, \quad \tfrac{9}{16}, \quad \tfrac{7}{16}.$$

Connection with Polygonal Numbers. The Greeks knew the theory of arithmetic series, but they usually treated it in connection with polygonal numbers. For example, the following are the first four triangular numbers:



It is evident that each triangular number is the sum of the series $\sum_1^n n$, and the Greeks were well aware of the rule for this summation.[1]

Chinese Work in Series. Nowhere in the very early Chinese works do we find any attempt to sum either an arithmetic or a geometric series.[2] In the *Wu-ts'ao Suan-king*, written about the beginning of the Christian era, or possibly earlier, we find the following problem:

There is a woman who weaves 5 feet the first day, her weaving diminishing day after day until, on the last day, she weaves 1 foot. If she has worked 30 days, how much has she woven in all?

The unknown author then gives this rule:

Add the amounts woven on the first and last days, take half the sum, then multiply by the number of days.

It is interesting to see that this earliest Chinese problem that we have yet found on the subject is, like the second case in the Ahmes Papyrus, one involving a descending series.[3]

[1] Heath, *Diophantus*, 2d ed., 247; Gow, *loc. cit.*, p. 103; Nesselmann, *Alg. Griechen*, chap. xi. [2] Mikami, *China*, p. 18. [3] Mikami, *China*, p. 41.

In Europe the rule for the sum was naturally the same as in the East, allowing for the difference in language,[1] and was occasionally put in verse for easy memorizing.[2]

The rule for finding any specified term is given by Cardan in his *Practica* (1539) and by Clavius in his *Epitome* (1583).

Geometric Series. The first examples of a geometric series yet found are due to the Babylonians, *c.* 2000 B.C., and tablets containing such examples are still extant.[3] In Egyptian mathematics the first problem on this subject thus far found is in the Ahmes Papyrus (*c.* 1550 B.C.)[4] and reads as follows:

The *one* scale.		Household	7
Once gives	2801	Cats	49
Twice gives	5602	Mice	343
Four times gives	11204	Barley [spelt]	2301 [*sic*]
Together	19607	Hekt measures	16807
		Together	19607

The left-hand column seems to be intended as a deduction of a rule for summing a geometric progression. Probably Ahmes saw that if the ratio is equal to the first term, $s_n = (s_{n-1} + 1)r$. Thus he found the sum of four terms to be 2800, and to this he added 1 and multiplied the result by 7 in order to obtain the sum of five terms. Possibly this is the significance of the expression "The *one* scale." Similarly, in the right-hand column

[1] Thus in an old MS. at Munich: "Addir albeg zesam daz erst vnd das leczt, vnd daz selb multiplicir mit dem halben der zal des posicionum" (Curtze, *Bibl. Math.*, IX (2), 113).

[2] Thus Huswirt (1501):

> Si primus numerus cum postremo faciat par
> Eius per mediũ loca singula multiplicabis
> Ast impar medium vult multiplicari locorum.

That is, $s = \frac{1}{2}n\ (a + l)$ if $(a + l)$ is odd, but $s = n \cdot \frac{1}{2}(a + l)$ if $(a + l)$ is even. The rule for the two cases goes back at least to Fibonacci (1202). See the Boncompagni edition, I, 166. By the time of Stifel (1544) a single rule answered for both cases.

[3] Hilprecht, *Tablets*, p. 17.

[4] Eisenlohr translation, p. 184, No. 79. The author has used a MS. translation from the hieratic, by Dr. A. B. Chace. On this section consult Tropfke, *Geschichte*, II (1), 315; Peet, *Rhind Papyrus*, p. 121. As in all such cases, reference to Ahmes means to the original from which he copied.

it is quite possible that Ahmes added four terms, then added 1, making the 2801 of the left-hand column, and finally multiplied by 7; but all this is merely conjectural.

The problem suggests the familiar one of the seven cats, although here stated quite differently. There is some doubt as to the word "household," the original word *pir* (*pr*) possibly having a different meaning. The *hekt* (*hekat*) was a measure of capacity. Essentially, therefore, Ahmes uses a rule based upon the formula $s = a\,(r^n - 1)/(r - 1)$. It is interesting to observe that a similar problem is given by Fibonacci (1202) and is solved in much the same way.[1]

The Greeks had rules for summing such a series,[2] and Euclid gave one that may be expressed as follows:

$$\frac{a_{n+1} - a_1}{a_n + a_{n-1} + \cdots + a_1} = \frac{a_2 - a_1}{a_1},$$

which amounts to saying that

$$\frac{ar^n - a}{s_n} = \frac{ar - a}{a},$$

whence would come our common formula

$$s_n = \frac{ar^n - a}{r - 1}.$$

The Hindus showed their interest in geometric series chiefly in the summation problems. The following typical problem is taken from Bhāskara (*c.* 1150):

A person gave a mendicant a couple of cowry shells first; and promised a twofold increase of the alms daily. How many *nishcas* does he give in a month?[3]

[1] *Scritti*, I, 311; Tropfke, *Geschichte*, VI (2), 15.

[2] Nesselmann, *Alg. Griechen*, p. 160; Euclid, *Elements*, IX, 35, 36; Heath, *Euclid*, Vol. II, p. 420.

[3] The wording and spelling is that of Colebrooke, *Bhāskara*, § 128, p. 55. A *niśka* (to take the better spelling) is 16 × 16 × 4 × 20 cowry shells. The cowry shell was then used as a small unit of value. The answer given in the translation is 2,147,483,646 cowry shells = 104,857 *nishcas*, 9 *drammas*, 9 *paṅas*, 2 *cácinis*, and 6 shells. See also *ibid.*, p. 291.

The Arabs apparently obtained the rule for summation from the Greeks, and it appears in an interesting form in the chessboard problem in the works of Albêrûnî (*c.* 1000).

Medieval European Rule. The medieval writers apparently obtained the rule from the Arabs, for it appears in the *Liber Abaci* of Fibonacci (1202).[1] The first modern treatment of the case is found in the *Algorithmus de Integris* (1410) of Prosdocimo de' Beldamandi.[2] Prosdocimo's treatment is as follows:

$$a + ar + ar^2 + \cdots + ar^{n-1} = ar^{n-1} + \frac{ar^{n-1} - a}{r - 1},$$

which is but little more complicated than our ordinary formula.[3] The same rule is given by Peurbach[4] (*c.* 1460). It is given by Chuquet (1484) in the form

$$s = \frac{rar^{n-1} - a}{r - 1},$$

and this is the plan used by Simon Jacob (1560), Clavius (1583), and others. Stifel (1544) gave the rule in the awkward form of

$$s = \frac{(rar^{n-1} - a)a}{ar - a},$$

a method used by Tartaglia[5] (1556), although he ordinarily preferred the one given by Prosdocimo de' Beldamandi.[6]

The ordinary type of puzzle problem in series, running through all the literature of the subject from the time of the Hindus to the 19th century, may be illustrated by the following from Baker (1568): "A Marchante hath solde 15 yeardes of Satten, the firste yarde for 1 $\bar{s}$, the second 2 $\bar{s}$, the thyrd 4 $\bar{s}$, the

[1] Boncompagni ed., I, 309, under "de duplicatione scacherii."

[2] First printed at Padua in 1483.

[3] As in all such cases it is to be understood that the rule is stated rhetorically in the original work, the modern algebraic notation being then unknown.

[4] It appears in his *Elementa Arithmetices Algorithmvs de numeris integris*, Vienna, 1492.

[5] *General Trattato*, II, fol. 6, *r*.

[6] *Ibid.*; see the last problem on the same page.

fourth $8\bar{s}$, and so increasing by double progression Geometricall . . . ," the total cost being then required.[1]

Other problems relate to the buying of orchards in which the value of the trees increases in geometric series, or to buying a number of castles on the same plan. Problems of this kind are mentioned later.

The rule for the sum of n terms is given by Clavius[2] (1583) and was undoubtedly known to various earlier writers. If we designate the elements by a, r, n, l, and s, and if any three of these elements are known, then the others can be found. This general problem was first stated by Wallis[3] (1657) and was solved for all cases not requiring logarithms. His formula[4] for S, one of the earliest stated in a form analogous to the one used at present, is

$$\frac{VR - A}{R - 1} = S.$$

The first infinite geometric series known to have been summed is the one given by Archimedes (*c.* 225 B.C.) in his quadrature of the parabola.[5] The series summed is

$$1 + \tfrac{1}{4} + (\tfrac{1}{4})^2 + \cdots + (\tfrac{1}{4})^n + \cdots.$$

The general formula for summing the infinite series a, ar, $ar^2, \cdots, ar^n, \cdots$, where $r < 1$, was given by Vieta (*c.* 1590).

Harmonic Series. Pythagoras and his school gave much attention to the cultivation of music, not only as a means of exciting or subduing the passions but as an abstract science. This led to, or at any rate was connected with, the important

[1] 1580 ed., fol. 40. Substantially the same problem is given in Trenchant (1566; 1578 ed., p. 292).

[2] "Detrahatur primus terminus ab vltimo, & reliquus numerus per numerum, qui vna unitate minor sit, quam denominator, diuidatur. Si enim Quotienti vltimus terminus, siue maius extremum adiiciatur, componetur summa omnium terminorum" (*Opera*, 1611, II, 68, of the *Epitome Arithmeticae Practicae*); that is, $s = (l - a)/(r - 1) + l$.

[3] *Opera*, I, cap. xxxi, p. 158 seq.

[4] " . . . si terminus primus seu minimus dicatur A, maximus V, communis rationis Exponens R, & progressionis summa S" (p. 158).

[5] Heath, *Archimedes*, chap. vii; Kliem translation, p. 137.

discovery of the relation of the tone to the length of the vibrating string, and hence to the introduction of harmonic proportion,[1] which later writers developed into harmonic series.

Higher Series. The first instances of the use of arithmetic series of higher order were confined to special cases. The series of squares was the earliest to attract attention. Archimedes[2] used geometry to show that

$$3\,[a^2 + (2\,a)^2 + (3\,a)^2 + \cdots + (na)^2] = (n+1)\,(na)^2 + a\,(a + 2\,a + 3\,a + \cdots + na).$$

For $a = 1$ this reduces to

$$1^2 + 2^2 + 3^2 + \cdots + n^2 = \tfrac{1}{6}\,n\,(n+1)\,(2\,n+1),$$

which appears substantially in the *Codex Arcerianus* (6th century). It is also found in the Hindu literature as shown by the works of Mahāvīra (*c.* 850).[3]

The sum of the cubes appears in the *Codex Arcerianus* in the form

$$1^3 + 2^3 + 3^3 + \cdots + 10^3 = (\tfrac{1}{2} \cdot 10 \cdot 11)^2.$$

The Hindus had rules for finding this sum, and they appear in the works of Brahmagupta (*c.* 628),[4] Mahāvīra (*c.* 850),[5] and Bhāskara (*c.* 1150).[6]

Among the Arabs similar rules are found, as in the works of al-Karkhî (*c.* 1020),[7] where

$$\sum_1^{10} n^2 = (1 + 10) \cdot 10 \cdot (\tfrac{10}{3} + \tfrac{1}{6}) = 385$$

and

$$\sum_1^{10} n^3 = \left(\sum_1^{10} n\right)^2.$$

[1] T. Gomperz, *Les penseurs de la Grèce*, p. 112 (Lausanne, 1904); H. Hankel, *Geschichte*, p. 105; Gow, *Greek Math.*, p. 68.

[2] See Tropfke, *Geschichte*, II (1), 318, on the entire topic. On this point see the Heiberg edition of *Archimedes*, II (1), 34.

[3] English translation, p. 170.

[4] Colebrooke's translation, p. 293.

[5] P. 171.

[6] Colebrooke's translation, p. 53.

[7] See Woepcke's translation of the *Fakhrî*, pp. 60, 61.

Fibonacci[1] (1220) and various other medieval scholars gave the same treatment of the subject. In the *Liber Quadratorum*[2] (1225) Fibonacci also gave the related forms

$$12(1^2+3^2+5^2+\cdots+n^2)=n(n+2)(2n+2) \text{ when } n \text{ is odd}$$

and

$$12(2^2+4^2+6^2+\cdots+n^2)=n(n+2)(2n+2) \text{ when } n \text{ is even.}$$

That the sum of the cubes may be found by adding the odd numbers is apparent from the following relations:

$$1^3=1,$$
$$2^3=3+5,$$
$$3^3=7+9+11,$$

and so on. This method of finding $\sum_1^n n^3$ was known to Nicomachus (*c.* 100). The general formula

$$1^3+2^3+3^3+\cdots+n^3=[\tfrac{1}{2}n(n+1)]^2$$

appears in substance in Pacioli's *Sūma*[3] (1494), but was already known.

A rule for summing the fourth powers, which may be expressed by

$$\sum_1^n n^4=\left(\sum_5^{n-1}+\sum n\right)\sum n^2,$$

appears in the *Key of Computation* of al-Kashî (*c.* 1430).[4]

Bernoulli Numbers. The case of $\sum n^m$ attracted attention in the 17th century, but the rule is first found in the *Ars Conjectandi*[5] (1713) of Jacques Bernoulli and involves what Euler[6]

[1] *Scritti*, I, 167 (fol. 70, *v.*). [2] *Scritti*, II, 263, 264. [3] Fol. 44, *r.*, l. 29.

[4] The *Miftâḥ al-ḥisâb* of Jemshîd ibn Mes'ûd ibn Maḥmud Giyâṯ ed-dîn al-Kashî (died *c.* 1436).

[5] II, cap. 3, p. 97; Tropfke, *Geschichte*, VI (2), 24.

[6] *Institutiones calculi differentialis*, II, § 122 (Petrograd, 1755). Euler's words are "ab inventore Jacobo Bernoulli vocari solent Bernoulliani."

designated as the "Bernoulli Numbers." These numbers (A, B, C, D) appear in the following summation of powers as given by Bernoulli:[1]

$$\int n^c \propto \frac{1}{c+1} n^{c+1} + \frac{1}{2} n^c + \frac{c}{2} A n^{c-1} + \frac{c \cdot c-1 \cdot c-2}{2 \cdot 3 \cdot 4} B n^{c-3} + \frac{c \cdot c-1 \cdot c-2 \cdot c-3 \cdot c-4}{2 \cdot 3 \cdot 4 \cdot 5 \cdot 6} C n^{c-5} + \frac{c \cdot c-1 \cdot c-2 \cdot c-3 \cdot c-4 \cdot c-5 \cdot c-6}{2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot 7 \cdot 8} D n^{c-7},$$

where $A \propto \frac{1}{6}$, $B \propto -\frac{1}{30}$, $C \propto \frac{1}{42}$, $D \propto -\frac{1}{30}$,

and where $\propto$ expresses equality; and the method of deriving these values is also given.

Revival of Infinite Series. The interest in the infinitesimal as an element in analysis, which manifested itself about the beginning of the 17th century, carried with it the notion of an infinite number of elements. Partly, no doubt, on this account the study of series with an infinite number of terms, already known to the Greeks, was revived, and the idea of products with an infinite number of factors was suggested.

The first of these products of any special interest has already[2] been mentioned as due to Vieta (1593). It may be expressed in modern form[3] as

$$\frac{2}{\pi} = \sqrt{\frac{1}{2}} \cdot \sqrt{\frac{1}{2} + \frac{1}{2}\sqrt{\frac{1}{2}}} \cdot \sqrt{\frac{1}{2} + \frac{1}{2}\sqrt{\frac{1}{2} + \frac{1}{2}\sqrt{\frac{1}{2}}}} \cdots,$$

and this, with others of the same nature, has already been considered in this work.[4]

There are three general periods in the later development of infinite series:[5] (1) the period of Newton and Leibniz,—that

[1] *Ars Conjectandi*, p. 97. [2] Vol. I, p. 312.

[3] See the Van Schooten edition of Vieta's works, p. 300.

[4] For logarithmic series, see page 513 and Volume I, page 434.

[5] R. Reiff, *Geschichte der unendlichen Reihen*, Tübingen, 1889. See also, for comparison, H. Wieleitner, "Zur Geschichte der unendlichen Reihen im christlichen Mittelalter," *Bibl. Math.*, XIV (3), 150; Tropfke, *Geschichte*, VI (2), 54.

of its introduction; (2) the period of Euler,—the formal stage; (3) the modern period,—that of the scientific investigation of the validity of infinite series. This third period, which may be designated as the critical one, began in 1812 with the publication of Gauss's celebrated memoir on the series

$$1 + \frac{\alpha \cdot \beta}{1 \cdot \gamma} x + \frac{\alpha \cdot (\alpha+1) \cdot \beta \cdot (\beta+1)}{1 \cdot 2 \cdot \gamma \cdot (\gamma+1)} x^2 + \cdots.$$

Euler had already considered this series, but Gauss was the first to master it, and under the name of "hypergeometric series," due to Pfaff (1765–1825), it has since occupied the attention of a large number of mathematicians. The particular series is not so important as the standard of criticism which Gauss set up, embodying the simpler criteria of convergence and the questions of remainders and the range of convergence.

Cauchy (1821) took up the study of infinite series and elaborated the theory of convergence which James Gregory (1668) had already begun and to which Maclaurin, Euler, and Gauss had made noteworthy contributions.[1] The term "convergent series" is due to Gregory (1668) and the term "divergent series" to Nicolas (I) Bernoulli (1713).[2]

Abel (1826) gave careful study to the series

$$1 + \frac{m}{1} x + \frac{m(m-1)}{2!} x^2 + \cdots,$$

correcting certain of Cauchy's conclusions and giving a scientific summation of the series for complex values of m and x.

Binomial Theorem. The development of $(a+b)^n$ for any integral value of n, or at least a device for finding the coefficients, was known in the East long before it appeared in Europe. The case of $n = 2$ was also known to Euclid (*c.* 300 B.C.),[3] but any evidence of the generalization of the law for other values

[1] On the history of criteria of convergence see F. Cajori, in the *Bulletin of the New York Math. Soc.*, II, 1; see also III, 186.

[2] F. Cajori, *Bulletin of the Amer. Math. Soc.*, XXIX, 55.

[3] *Elements*, II. For a summary of his work on algebraic identities see Nesselmann, *Alg. Griechen*, p. 154.

of n first appears, so far as we know, in the algebra of Omar Khayyam (*c.* 1100). This writer did not give the law, but he asserted that he could find the fourth, fifth, sixth, and higher roots of numbers by a law that he had discovered and which did not depend upon geometric figures.[1] He states that this law was set forth by him in another work, but of this work there seems to be no copy extant.

Pascal Triangle. In one of the works of Chu Shï-kié (1303), the greatest of the Chinese algebraists of his time, the triangular arrangement of the coefficients is given in the following form,

1
1 1
1 2 1
1 3 3 1
1 4 6 4 1
1 5 10 10 5 1
. . . .

a form now commonly known as the Pascal Triangle.[2]

This triangular array first appeared in print on the title-page of the arithmetic of Apianus (1527), as shown in the illustration on page 509.[3] In the form

1 2 1
1 3 3 1
1 4 6 4 1
1 5 10 10 5 1

[1] "J'ai composé un ouvrage sur la démonstration de l'exactitude de ces méthodes. . . . J'en ai, en outre, augmenté les espèces, c'est-à-dire que j'ai enseigné à trouver les côtés du carré-carré, du quadrato-cube, du cubo-cube, etc., à une étendue quelconque, ce qu'on n'avait pas fait précédemment. Les démonstrations que j'ai données à cette occasion ne sont que des démonstrations arithmétiques." Translated by F. Woepcke, *L'Algèbre d'Omar Alkhâyyamî*, p. 13 (Paris, 1851).

[2] Mikami, *China*, p. 106.

[3] On the general subject see H. Bosmans, "Note historique sur le Triangle arithmétique, dit de Pascal," *Annales de la Société scientifique de Bruxelles*, XXXI, October, 1906; Tropfke, *Geschichte*, VI (2), 37.



PASCAL TRIANGLE AS FIRST PRINTED, 1527

Title-page of the arithmetic of Petrus Apianus, Ingolstadt, 1527, more than a century before Pascal investigated the properties of the triangle

it is first found in Stifel's *Arithmetica Integra* (1544), appearing a year later in the *De Nvmeris et Diversis Rationibvs* of Scheubel (1545). It also appears in the various editions of Peletier's arithmetic (Poitiers, 1549 and later). Tartaglia (1556) gave it as his own invention,[1] and soon after his time it became common property. Bombelli (1572), for example, gave the coefficients for all powers of $a + b$ up to the seventh, using them in finding corresponding roots,[2] and Oughtred (1631) gave them up to the tenth power.[3] The triangular array was investigated by Pascal (1654) under a new form, substantially as follows:[4]

	1	2	3	4	5	6	7	8	9	10
1	1	1	1	1	1	1	1	1	1	1
2	1	2	3	4	5	6	7	8	9	
3	1	3	6	10	15	21	28	36		
4	1	4	10	20	35	56	84			
5	1	5	15	35	70	126				
6	1	6	21	56	126					
7	1	7	28	84						
8	1	8	36							
9	1	9								
10	1									

Rangs Paralleles

Rangs Perpendiculaires

He made numerous discoveries relating to this array and set them forth in his *Traité du triangle arithmétique*,[5] published


[1] *General Trattato*, II, fols. 69, *v.*; 71, *v.* (Venice, 1556). [2] *Algebra*, p. 64.

[3] F. Cajori, *William Oughtred*, p. 29 (Chicago, 1916).

[4] This is from the plate in Pascal's *Œuvres*, Vol. V (Paris, 1819). The description is given on pages 1–56. In the original there are diagonals in the above figure. See also Tropfke, *Geschichte*, VI (2), 37.

[5] "Le nombre de chaque cellule est égal à celui de la cellule qui la précède dans son rang perpendiculaire, plus à celui de la cellule qui la précède dans son rang parallèle" (*Œuvres*, V, 3) (Paris, 1819).

posthumously in 1665, and among these was essentially our present Binomial Theorem for positive integral exponents. After this time the triangular array was common in the East as well as in the West.



PASCAL TRIANGLE IN JAPAN

From Murai Chūzen's *Sampō Dōshi-mon* (1781), showing also the sangi forms of the numerals

Generalization of the Binomial Theorem. The generalization of the binomial theorem for negative and fractional values of n is due to Newton, who set it forth in letters which he wrote to Oldenburg on June 13, 1676, and October 24, 1676.[1]

The proof of the Binomial Theorem was slowly developed by later writers. Among those who contributed to a satisfactory demonstration were Maclaurin[2] for rational values of n, Giovanni Francesco M. M. Salvemini (de Castillon)[3] and Kästner (1745)[4] for integral values, Euler[5] (1774) for fractional exponents, and Abel[6] (*c.* 1825) for general values of n, taking n as a complex number.

[1] See *Commercium Epistolicum*, London, 1712; 1725 ed., pp. 131, 142. In his letter of October 24 he proceeds from $(1-x^2)^{\frac{0}{2}}$, $(1-x^2)^{\frac{2}{2}}$, $(1-x^2)^{\frac{4}{2}}$, $\cdots$ to $(1-x^2)^{\frac{1}{2}}$ and $(1-x^2)^{\frac{3}{2}}$, "vel generaliter $\overline{1-xx}]^{m}$," and finds, for example, that "$\overline{1-xx}]^{\frac{1}{2}}$ valeret $1-\frac{1}{2}x^2-\frac{1}{8}x^4-\frac{1}{16}x^6$ &c." On the doubtful assertion that Pascal may have anticipated this discovery, see G. Eneström, *Bibl. Math.*, V (3), 72.

[2] *Treatise of Fluxions*, p. 607 (1742).

[3] Born at Castiglione, 1708; died 1791. See *Phil. Trans.*, XLII (1742), 91. He used the theory of combinations. See page 326.

[4] Cantor, *Geschichte*, III, 660.

[5] *Novi comment. Petrop.*, XIX, 103; see Tropfke, *Geschichte*, II (1), 331. See also the English translation of Euler's *Algebra*, I, 172, 177 (London, 2d ed., 1810).

[6] The article appeared posthumously in Crelle's *Journal*, I (1826), 311. See also Abel's *Œuvres*, I, 219 (Christiania, 1881).

The generalization of the Binomial Theorem into the Polynomial Theorem was due chiefly to Leibniz (1695), Jacques Bernoulli, and De Moivre.[1]

Finite Differences. The treatment of series by the method of finite differences appeared in the 17th century. In 1673 Leibniz wrote to Oldenburg concerning the following scheme of treating the series of cubes:

		0	0	0		
	6	6	6	6		
6	12	18	24	30		
1	7	19	37	61	91	
0	1	8	27	64	125	216

He said that John Pell attributed the discovery to Gabriel Mouton, of Lyons.[2]

Taylor's Formula and Maclaurin's Formula. In 1715 Brook Taylor published the formula which bears his name, and which we now express as follows:

$$f(x+h) = f(x) + hf'(x) + \frac{h^2}{2!}f''(x) + \cdots.$$[3]

It was not until 1742 that Colin Maclaurin published the corresponding formula

$$f(x) = f(0) + xf'(0) + \frac{x^2}{2!}f''(0) + \cdots,$$

a relation that is easily derived from the preceding one.[4]

Trigonometric Series. The development of trigonometric functions in series first attracted the attention of mathematicians

[1] De Moivre's articles appeared in the *Phil. Trans.*, XIX (1697), 619; XX, 190.

[2] *Commercium Epistolicum*, p. 109 (London, 1712; 1725 ed.). Gabriel Mouton, born at Lyons, 1618; died at Lyons, September 28, 1694. He suggested (1670) a system of measures not unlike the metric system.

[3] *Methodus Incrementorum directa et inversa*, prop. 7 (London, 1715). The series had already been announced by him in 1712.

[4] *A Complete System of Fluxions*, Edinburgh, 1742.

in the 17th century. To James Gregory (1671) are due the following:[1]

$$x = \tan x - \tfrac{1}{3}\tan^3 x + \tfrac{1}{5}\tan^5 x - \tfrac{1}{7}\tan^7 x + \cdots,$$

$$\tan x = x + \tfrac{1}{3}x^3 + \tfrac{2}{15}x^5 + \tfrac{17}{315}x^7 + \cdots,$$

$$\sec x = 1 + \tfrac{1}{2}x^2 + \tfrac{5}{24}x^4 + \tfrac{61}{720}x^6 + \cdots.$$

He also gave the important series

$$\text{arc tan } x = x - \tfrac{1}{3}x^3 + \tfrac{1}{5}x^5 - \cdots,$$

but this is easily deduced from the one given above for tan x. Newton[2] gave (*c.* 1669) the antitrigonometric series for arc sin x, essentially as follows:

$$\text{arc sin } x = \sin^{-1} x = x + \tfrac{1}{6}x^3 + \tfrac{3}{40}x^5 + \tfrac{5}{112}x^7 + \cdots.$$

Logarithmic Series. The idea of expressing a logarithm by means of a series seems to have originated with Gregory and to have been elaborated by Nicolaus Mercator[3] (1667), who discovered, for a special case at least, the relation

$$\log(1+a) = a - \tfrac{1}{2}a^2 + \tfrac{1}{3}a^3 - \tfrac{1}{4}a^4 + \cdots,$$

where $$1 \geqq a > -1.$$

The value of Mercator's and Gregory's contributions was recognized by Wallis in reviews which he wrote of their works.[4]

12. Logarithms

Technical Terms. The word "logarithm"[5] means "ratio number" and was an afterthought with Napier. He first used the expression "artificial number," but before he announced his discovery he adopted the name by which it is now known.[6]

[1] These were communicated to Collins in a letter from Gregory. See the *Commercium Epistolicum*, London, 1712; 1725 ed., pp. 98, 210 n.

[2] *Commercium Epistolicum*, pp. 97, 126; Tropfke, *Geschichte*, VI (2), 46.

[3] *Logarithmotechnia sive methodus construendi logarithmos nova*, London, 1668. The theory was worked out the year before.

[4] *Phil. Trans.*, 1668, pp. 640, 753.

[5] From the Greek λόγος (*log'os*), ratio, + ἀριθμός (*arithmos'*), number.

[6] This fact is evident from his *Descriptio*, 1619 ed.

Briggs introduced (1624) the word "mantissa." It is a late Latin term of Etruscan origin, originally meaning an addition, a makeweight, or something of minor value, and was written *mantisa*. In the 16th century it came to be written *mantissa* and to mean "appendix,"[1] and in this sense it was probably considered by Briggs. The name also appears in connection with decimals in Wallis's *Algebra* (1685), but it was not commonly used until Euler adopted it in his *Introductio in analysin infinitorum* (1748). Gauss suggested using it for the fractional part of all decimals.[2]

The term "characteristic" was suggested by Briggs (1624) and is used in the 1628 edition of Vlacq.[3]

The characteristic was printed in the early tables, and it was not until well into the 18th century that the custom of printing only the mantissas became generally established.

Napier's Invention. So far as Napier's invention is concerned, Lord Moulton expressed the fact very clearly when he said:[4]

> The invention of logarithms came on the world as a bolt from the blue. No previous work had led up to it, foreshadowed it or heralded its arrival. It stands isolated, breaking in upon human thought abruptly without borrowing from the work of other intellects or following known lines of mathematical thought.

Napier worked at least twenty years upon the theory. His idea was to simplify multiplications involving sines, and it was a later thought that included other operations, applying logarithms to numbers in general. He may have been led to his discovery by the relation

$$\sin A \sin B = \tfrac{1}{2}(\cos \overline{A - B} - \cos \overline{A + B}),$$

[1] With this meaning it appeared as late as 1701 in J. C. Sturm, *Mathesis juvenalis*.

[2] "Si fractio communis in decimalem convertitur, seriem figurarum decimalium . . . fractionis *mantissam* vocamus . . ." See E. Hoppe, "Notiz zur Geschichte der Logarithmentafeln," *Mittheilungen der math. Gesellsch. in Hamburg*, IV, 52.

[3] ". . . prima nota versus sinistram, quam Characteristicam appellare poterimus . . ." It again appeared in Mercator's *Logarithmotechnia* (1668).

[4] "Inaugural Address: The Invention of Logarithms," *Napier Tercentenary Memorial Volume*, p. 1 (London, 1915).

for, as Lord Moulton says, in no other way can we "conceive that the man to whom so bold an idea occurred should have so needlessly and so aimlessly restricted himself to sines in his work, instead of regarding it as applicable to numbers generally."

Napier published his *Descriptio*[1] of the table of logarithms in 1614. This was at once translated into English by Edward Wright,[2] but with the logarithms contracted by one figure.

In Napier's time sin ϕ was a line, not a ratio. The radius was called the *sinus totus*, and when this was equal to unity the length of the sine was simply stated as sin ϕ. If r was not unity, the length was $r \sin\phi$. With this statement we may consider Napier's definition of a logarithm:

> The Logarithme therefore of any sine is a number very neerely expressing the line, which increased equally in the meane time, whiles the line of the whole sine decreased proportionally into that sine, both motions being equal-timed, and the beginning equally swift.[3]

From this it follows that the logarithm of the *sinus totus* is zero. Napier saw later that it was better to take $\log 1 = 0$.[4]

Napier then lays down certain laws relating to proportions, which may be stated symbolically as follows:

1. If $a:b = c:d$, then $\log b - \log a = \log d - \log c$.
2. If $a:b = b:c$, then $\log c = 2 \log b - \log a$.
3. If $a:b = b:c$, then $2 \log b = \log a + \log c$.
4. If $a:b = c:d$, then $\log d = \log b + \log c - \log a$.
5. If $a:b = c:d$, then $\log b + \log c = \log a + \log d$.
6. If $a:b = b:c = c:d$, then $3 \log b = 2 \log a + \log d$ and $3 \log c = \log a + 2 \log d$.

[1] *Mirifici Logarithmorum Canonis Descriptio*, Edinburgh, 1614.

[2] *A Description of the Admirable Table of Logarithmes*, London, 1616, published after Wright's death.

[3] Wright's translation of the *Descriptio*, pp. 4, 5.

[4] As to the priority of this idea, see G. A. Gibson, "Napier's logarithms and the change to Briggs's logarithms," in the *Napier Tercentenary Memorial Volume*, p. 114 (London, 1915); this volume should be consulted on all details of this kind. See also Dr. Glaisher's article on logarithms in the eleventh edition of the *Encyclopædia Britannica*.

The system was, therefore, designed primarily for trigonometry, but would also have been valuable for purposes of ordinary computation had not a better plan been suggested.

Napier also wrote a work on the construction of a table,[1] which was published posthumously as part of the 1619 edition of the *Descriptio.*

Napier's logarithms are not those of the so-called Napierian, or hyperbolic, system, but are connected with this system by the relation $\log_n a = 10^7 \cdot \log_e 10^7 - 10^7 \cdot \log_e a$. The relation between the sine and its logarithm in Napier's system is

$$\sin \phi = 10^7 \cdot e^{-\log_n \sin \phi / 10^7}$$

so that the sine increases as its logarithm decreases.

Briggs's System. Henry Briggs, professor of geometry at Gresham College, London, and afterwards Savilian professor of geometry at Oxford, was one of the first to appreciate the work of Napier. Upon reading the *Descriptio* (1614) he wrote:

> Naper, lord of Markinston, hath set my head and hands at work with his new and admirable logarithms. I hope to see him this summer, if it please God; for I never saw a book which pleased me better, and made me more wonder.

He visited Merchiston in 1615 and suggested another base, of which, however, Napier had already been thinking. In Briggs's *Arithmetica Logarithmica* the preface, written by Vlacq, contains the following statement[2] by the author of the work itself:

> That these logarithms differ from those which that illustrious man, the Baron of Merchiston published in his *Canon Mirificus* must not surprise you. For I myself, when expounding their doctrine publicly in London to my auditors in Gresham College, remarked that it would be much more convenient that o should be kept for the logarithm of

[1] *Mirifici ipsius canonis constructio.*

[2] *Arithmetica Logarithmica sive Logarithmorum Chiliades Triginta* (London, 1624), preface. The original is in Latin; the translation of the statement is from the *Napier Tercentenary Memorial Volume.*

the whole sine (as in the *Canon Mirificus*). . . . And concerning that matter I wrote immediately to the author himself; and as soon as the season of the year and the vacation of my public duties of instruction permitted I journeyed to Edinburgh, where, being most hospitably received by him, I lingered for a whole month. But as we talked over the change in logarithms he said that he had for some time been of the same opinion and had wished to accomplish it. . . . He was of the opinion that . . . 0 should be the logarithm of unity.

The real value of the proposition made by Briggs at this time was that he considered the values of $\log 10^n a$ for all values of n. The relation between the two systems as they first stood may be indicated as follows:

Napier, $\log y = r(\log_e r - \log_e y)$, where $r = 10^7$;

Briggs, $\log y = 10^{10}(10 - \log_{10} y)$;

Napier (later suggestion), $\log y = 10^9 \log_{10} y$.

The first table of logarithms of trigonometric functions to the base 10 was made by Gunter, a colleague of Briggs at Gresham College, and was published in London in 1620.[1]

The Base *e*. In the 1618 edition of Edward Wright's translation of the *Descriptio* there is printed an appendix, probably written by Oughtred, in which there is the equivalent of the statement that $\log_e 10 = 2.302584$, thus recognizing the base e.

Two years later (1620) John Speidell[2] published his *New Logarithmes*, also using this base. He stated substantially that

$$\log n = 10^{-1}(\text{nap log } 1 - \text{nap log } n),$$

or

$$\log n = 10^5(10 + \log_e 10^{-5} x).$$

Continental Recognition. The same year (1624) that Briggs published his *Arithmetica Logarithmica* Kepler's first table appeared. A year later Wingate's *Arithmetiqve Logarithmiqve* (Paris, 1625) gave the logarithms of numbers from 1 to 1000, together with Gunter's logarithmic sines and tangents.

[1] *Canon Triangulorum, sive Tabulae Sinuum et Tangentium.*

[2] See *Napier Tercentenary Memorial Volume*, pp. 132, 221; F. Cajori, *History of Elem. Math.*, p. 164, rev. ed. (N.Y., 1917), and *History of Math.*, p. 153, rev. ed. (N.Y., 1919).

Holland was the third Continental country to recognize the work of Napier and Briggs. In 1626 there was published a work[1] by Adriaen Vlacq,[2] assisted by Ezechiel de Decker. In 1628 Vlacq republished Briggs's tables,[3] filling the gap from 20,000 to 90,000. The tables in this work were reprinted in London by George Miller in 1631.[4] It is interesting to note that the next complete edition of Vlacq's tables appeared in China.[5]

In Germany the theory was first made known by Johann Faulhaber[6] (1630).

Logarithms in Arithmetic. By the middle of the seventeenth century, logarithms found their way into elementary arithmetics, as is seen in Hartwell's (1646) edition of Recorde's *Ground of Artes*,[7] where it is said that "for the extraction of all sorts of roots, the table of Logarithmes set forth by M. Briggs are most excellent, and ready." Thereafter they were occasionally found in textbooks of this kind, both in Great Britain and on the Continent.

Forerunners of Bürgi. Napier approached logarithms from the standpoint of geometry, whereas at the present time we approach the subject from the relation $a^m a^n = a^{m+n}$. This relation was known to Archimedes[8] and to various later writers.

More generally, if we take the two series

	0	1	2	3	4	5	6	7
and	1	2	4	8	16	32	64	128,

[1] *Eerste Deel van de Nieuwe Telkonst*, Gouda, 1626. See D. Bierens de Haan in Boncompagni's *Bullettino*, VI, 203, 222; J. W. L. Glaisher, "Notice respecting some new facts in the early history of logarithmic tables," *Philosoph. Mag.*, October, 1872.

[2] Born at Gouda, *c.* 1600; died after 1655. The common Dutch spelling is now Vlack.

[3] *Arithmetica Logarithmica*, Gouda, 1628. It was also published with a French title-page.

[4] *Logarithmicall Arithmetike*, London, 1631.

[5] *Magnus Canon Logarithmorum . . . Typis Sinensibus in Aula Pekinensi . . .*, 1721.

[6] *Inginieurs-Schul, erster Theil, darinen durch den Canonem Logarithmicum . . .*, Frankfort, 1630. Faulhaber was born at Ulm, May 5, 1580; died at Ulm, 1635.

[7] Also the editions of 1662 and 1668.

[8] *Opera omnia*, ed. Heiberg, 2d ed., II, 243; Heath, *Archimedes*, p. 230.

the first one being arithmetic and the second one being geometric, we see that the latter may be written as follows:

$$2^0 \quad 2^1 \quad 2^2 \quad 2^3 \quad 2^4 \quad 2^5 \quad 2^6 \quad 2^7.$$

From this it is evident that

$$2^3 \cdot 2^4 = 2^7, \qquad (2^2)^3 = 2^6,$$
$$2^7 : 2^3 = 2^4, \qquad (2^4)^{\frac{1}{2}} = 2^2,$$

which are the fundamental laws of logarithms.

Most writers[1] refer to Stifel as the first to set forth these basal laws, and we shall see that he did set them forth very clearly; but he was by no means the first to do so, nor did they first appear even in his century. Probably the best of the statements concerning them which appeared in the 15th century were those of Chuquet in *Le Triparty en la Science des Nombres*, written in 1484, from which Estienne de la Roche copied so freely in his *Larismethique* of 1520. Chuquet expressed very clearly the relations

$$a^m a^n = a^{m+n}$$

and

$$(a^m)^n = a^{mn}$$

in connection with the double series to which reference[2] has been made, calling special attention to the latter law as "a secret" of proportional numbers.[3]

[1] Among them is Kästner, *Geschichte der Mathematik*, I, 119, who has been generally followed in this matter. See also Th. Müller, *Der Esslinger Mathematiker Michael Stifel*, Prog., p. 16 (Esslingen, 1897), where the author states: "'Dies ist das älteste Buch,' sagt Strobel, 'in welchem die Vergleichung des arithmetischen Reihe mit der geometrischen als der Grund der Logarithmen vorkommt.'"

Much of the work on this topic appeared in the author's paper published in the *Napier Tercentenary Memorial Volume*, p. 81 (London, 1915).

[2] "Il convient poser plus3s nōbres ꝑporcional3 cōmancans a 1. constituez en ordonnance continuee cōme 1.2.4.8.16.32. &c. ou .1.3.9.27. &c. ℭ Maintenant conuient scauoir que .1. represente et est ou lieu des nombres dōt le, denōīa^{on} est .0./2. represente et est ou lieu des premiers dont leur denomīacion est .1./4. tient le lieu des second3 dont leur denomīacion est .2. Et .8. est ou lieu des tiers .16. tient la place des quartz" (fol. 86, *v.*, of the *Triparty*). This is taken from the copy made by A. Marre from the original manuscript. Boncompagni published it in the *Bullettino*, XIII, 593 seq., fol. 86, *v.*, being on page 740.

[3] "ℭ Semblement qui multiplie .4. qui est nombre second par .8. qui est nombre tiers montent .32. qui est nombre quint . . . ℭ En ceste consideration est maīfeste

It is difficult to say when a plan of this kind first appears in print, because it is usually hinted at before it is stated definitely. Perhaps it is safe, however, to assign it to Rudolff's *Kunstliche rechnung* of 1526, where the double series is given and the multiplication principle is clearly set forth;[1] and inasmuch as this work had great influence on Stifel, who in turn influenced Jacob, Clavius, and Bürgi, it was somewhat epoch-making.

The next writer to refer to the matter was probably Apianus (1527), who followed Rudolff so closely as to be entitled to little credit for what he did.

Following Apianus, the first arithmetician of any standing who seems to have had a vision of the importance of this relation was Gemma Frisius (1540), who gave the law with relation to the double array

3	9	27	81	243	729
0	1	2	3	4	5,

saying that the product of two numbers occupies a place indicated by the sum of their places (3×9 occupying the place indicated by $1 + 2$, or 3), and that the square of a number in the fifth place occupies the $2 \times$ 5th place.[2]

The first arithmetician to take a long step in advance of Rudolff was Stifel (1544), the commentator (1553) on *Die*

vng secret qui est es nombres ꝓporcionalz. Cest que qui multiplie vng nombre ꝓporcional en soy Il en viēt le nombre du double de sa denomīacion come qui mltiplie .8. qui est tiers en soy Il en vient .64. qui est sixe. Et .16. qui est quart multiplie en soy. Il en doit venir 256. qui est huyte. Et qui multiplie .128. qui est le .7e. ꝓporcional par .512. qui est le 9e. Il en doit venir 65536. qui est le 16e" (*ibid.*, p. 741).

[1] "Nun merck wenn du zwo zalen mit einander multiplicirst/ wiltu wissen die stat des quocients/ addir die zalen der natürlichen ordnung so ob den zweyen mit einander gemultiplicirten zalen gefunden/ dz collect bericht dich. Als wen ich 8 multiplicir mit 16. muss komen 128. darumb das 3 vnd 4 so vber dem 8 vnnd 16 geschriben zusamen geaddirt 7 machen." He gives several examples, but goes no farther with the law.

[2] "Si enim duos quoscunque ex his numeris inuicem multiplicaueris, productúmque per primum diviseris, producetur numerus eo loco ponendus, quē duo facta indicabunt . . ." (ed. 1553, fol. 17, *r.*, and note by Peletier (Peletarius), fol. 78, *v.*). The relation is not so clear as in some of the other texts, on account of the arrangement of the series.

Coss. It is not, however, in this work that the theory is set forth, but in the *Arithmetica Integra* of 1544. Stifel here refers several times to the laws of exponents. At first he uses the series

0	1	2	3	4	5	6	7	8
1	2	4	8	16	32	64	128	256,

distinctly calling the upper numbers *exponents*, and saying that the exponents of the factors are added to produce the exponent of the product and subtracted to produce the exponent of the quotient.[1] Moreover, he expressly lays down four laws, namely, that addition in arithmetic progression corresponds to multiplication in geometric progression, that subtraction corresponds to division, multiplication to the finding of powers, and division to the extracting of roots. Furthermore, Stifel not only set forth the laws for positive exponents but also saw the great importance of considering the negative exponents of the base which he selected, using the series

−3	−2	−1	0	1	2	3	4	5	6
$\frac{1}{8}$	$\frac{1}{4}$	$\frac{1}{2}$	1	2	4	8	16	32	64

and making the significant remark: "I might write a whole book concerning the marvellous things relating to numbers, but I must refrain and leave these things with eyes closed."[2] What these mysteries were we can only conjecture.

[1] "Qualicunq; facit Arithmetica progressio additione, & subtractione, talis facit progressio Geometrica multiplicatione, & diuisione. ut plene ostendi lib. 1. capite de geomet. progres. Vide ergo,

0.	1.	2.	3.	4.	5.	6.	7.	8.
1.	2.	4.	8.	16.	32.	64.	128.	256.

Sicut ex additione (in superiore ordine) 3 ad 5 fiunt 8, sic (in inferiore ordine) ex multiplicatione 8 in 32 fiunt 256. Est autem 3 exponens ipsius octonarij, & 5 est exponens 32 & 8 est exponens numeri 256. Item sicut in ordine superiori, ex subtractione 3 de 7, remanent 4, ita in inferiori ordine ex diuisione 128 per 8, fiunt 16" (fols. 236, 237).

It will be noticed that he speaks of 8 as "exponens numeri 256," and not as the exponent of 2, but this has no significance with respect to the theory.

[2] "Posset hic fere nouus liber integer scribi de mirabilibus numerorum, sed oportet ut me hic subducã, & clausis oculis abeã."

A number of French writers of this period were also aware of the law, and Peletier[1] (1549) stated it clearly for the case of multiplication. Five years later Claude de Boissière elaborated this treatment and spoke of the "marvellous operations" which can be performed by means of the related series. Two years after Boissière's work was published the theory was again given by Forcadel (1565), with a statement that the idea was due to Archimedes, that it was to be found in Euclid, and that Gemma Frisius had written upon it. Ramus recognized its value but added nothing to it or to its possible applications. When, however, Schoner came to write his commentary on the work of Ramus, in 1586, a decided advance was made, for not only did he give the usual series for positive exponents, but, like Stifel, he used the geometric progressions with fractions as well, although, as stated above, not with negative exponents. Further, he used the word "index" where Stifel had used "exponent," and, like this noteworthy writer, gave evidence of an appreciation of the importance of the law. In general the French writers already named (and in the list should also be included the name of Chauvet) paid no attention to any of the laws except that of multiplication, while the German writers, following the lead of Stifel, took the broader view of the theory. This was not always the case, for Sigismund Suevus, a German arithmetician who wrote as late as 1593, did not go beyond the limits set by most of the French arithmeticians; but in general the German writers were in the lead. This is particularly true of Simon Jacob (1565), who followed Stifel closely, recognizing all four laws, and, as is well known, influencing Jobst Bürgi. These writers did not use the general exponents essential to logarithms, but the recognition of the four laws is significant.

[1] The extract here given is from the 1607 edition of *L'Arithmetiqve*, p. 67. In speaking of the series

3	6	12	24	48	96
0	1	2	3	4	5

he says: "Ie sçauoir qui est le nõbre qui eschet au neufieme lieu en ceste Progression Double Ie diuise 48, qui est sur 4, par le premier nombre de la Progression, 3: prouiennent 16: lesquels je multiplie par 96, qui est sur 5: (car 4 & 5 font 9) prouiendront 1536, qui sera le nombre à mettre au neufieme lieu."

Bürgi and the Progress Tabulen. In 1620 Jobst Bürgi published his Progress Tabulen, a work conceived some years earlier. As stated above, it is well known that he was influenced by Simon Jacob's work. The tables were printed at Prag and are simply lists of antilogarithms with base 1.0001. The logarithm is printed in red in the top line and the left-hand column, and the antilogarithms are in black, and hence Bürgi calls the logarithm *Die Rothe Zahl.* The first part of his table is as follows:

	0	500	1000	1500	2000
0	100000000	100501227	101004966	101511230	102020032
10	10000	11277	15067	21381	30234
20	20001	21328	25168	31534	40437
30	30003	31380	35271	41687	50641

The manuscripts of Bürgi are at the Observatory at Pulkowa, but none seem to be of a date later than 1610, so that he probably developed his theory independently of Napier. It is evident that he approached the subject algebraically, as Napier approached it geometrically.[1]

The only extensive table of antilogarithms is due to James Dodson (London, 1742).

Logarithms in the Orient. Logarithms found their way into China through the influence of the Jesuits. The first treatise upon the subject published in that country was a work by one Sié Fong-tsu, a pupil of the Polish Jesuit John Nicolas Smogolenski (1611–1656). This treatise was published about 1650, although Smogolenski had already mentioned the theory in one of his works.[2] Vlacq's tables (1628) were reprinted in Peking, as already stated, in 1713.

[1] Thus Kepler says: " . . . qui etiam apices logistici Justo Byrgio multis annis ante editionem Neperianam, viam praeiverunt, ad hos ipsissimos Logarithmos. Etsi homo cunctator et secretorum suorum custos, foetum in partu destituit, non ad usus publicos educavit" (*Opera Omnia*, VII, 298) (Frankfort a. M., 1868).

[2] The *T'ien-pu Chên-yüan*, as stated in Volume I, page 436.

13. Permutations, Combinations, Probability

Permutations and Combinations. The subject of permutations may be said to have had a feeble beginning in China in the *I-king* (*Book of Changes*), the arrangements of the mystic trigrams, as in ䷀, furnishing the earliest known example.[1] It is not improbable that it was the *I-king* that suggested to a certain Japanese daimyō of the 12th century that he write a book, now lost, upon permutations.[2]

Greek Interest in the Subject. The subject received some slight attention at the hands of certain Greek writers. Plutarch[3] (1st century) tells us[4] that Xenocrates (*c.* 350 B.C.), the philosopher, computed the number of possible syllables as 1,002,000,000,000, but it does not seem probable that this represents an actual case in combinations.[5] Plutarch also states that Chrysippus (*c.* 280–*c.* 207 B.C.), a Stoic philosopher, found the number of combinations of ten axioms to be more than 1,000,000, and that Hipparchus (*c.* 140 B.C.) gave the number as 101,049[6] if admitted and 310,925 if denied; but we have no evidence of any theory of combinations among the Greeks.[7]

Interest of Latin Writers in the Subject. The Latin writers, having little interest in any phase of mathematics except the practical, paid almost no attention to the theory of combinations. The leading exception was Boethius (*c.* 510). He gives a rule for finding the combinations of n things taken two at a time which we should express as $\frac{1}{2}n(n-1)$.[8]

[1] See Volume I, page 25.

[2] The theory is referred to as *Keishizan* in Volume I, page 274.

[3] *Quaestiones Conviv.*, Lib. VIII, 9, iii, 12; ed. Dübner, II, 893 (Paris, 1877).

[4] Tropfke, *Geschichte*, II (1), 351.

[5] Gow, *Greek Math.*, pp. 71 n., 86; Tropfke, *Geschichte*, II (1), 351.

[6] "Centena millia atque insuper mille et quadraginta novem." The number in Tropfke is incorrect.

[7] With respect to the single possible case in Pappus, see ed. Hultsch, II, 646–649: "Nam ex tribus dissimilibus generis triades diversae inordinatae existunt numero decem."

There is a slight trace of interest in the subject in the works of Plato and Aristotle, but not enough to be worthy of discussion in this chapter. See J. L. Heiberg, *Philologus*, XLIII, 475, with references.

[8] J. L. Heiberg, *ibid.*

Hindu Interest in the Subject. The Hindus seem to have given the matter no attention until Bhāskara (*c.* 1150) took it up in his *Lilāvati.* In this work he considered the subject twice. He asserted that an idea of permutations "serves in prosody . . . to find the variations of metre; in the arts [as in architecture] to compute the changes upon apertures [of a building]; and [in music] the scheme of musical permutations; in medicine, the combinations of different savours."[1] He gave the rules for the permutations of n things taken r at a time, with and without repetition, and the number of combinations of n things taken r at a time without repetition.[2]

Early European Interest in the Subject. Early in the Christian Era there developed a close relation between mathematics and the mystic science of the Hebrews known as the cabala. This led to the belief in the mysticism of arrangements and hence to a study of permutations and combinations. The movement seems to have begun in the anonymous *Sefer Jezira* (*Book of Creation*), and shows itself now and then in later works.

It seems to have attracted the attention of the Arabic and Hebrew writers of the Middle Ages in connection with astronomy. Rabbi ben Ezra (*c.* 1140), for example, considered it with respect to the conjunctions of planets, seeking to find the number of ways in which Saturn could be combined with each of the other planets in particular, and, in general, the number of combinations of the known planets taken two at a time, three at a time, and so on. He knew that the number of combinations of seven things taken two at a time was equal to the number taken five at a time, and similarly for three and four and for six and one. He states no general law, but he seems to have been aware of the rule for finding the combinations of n things taken r at a time.[3]

[1] Colebrooke translation, p. 49. [2] *Ibid.*, p. 123.

[3] D. Herzog, *Zophnath Paneach* (in Hebrew), Cracow, 1911. It is an edition of Josef ben Eliezer's supercommentary (that is, a commentary on a commentary by Rabbi ben Ezra) on the Bible. The passage occurs in an extract from Rabbi ben Ezra's astrological manuscript *ha-Olam*, now in Berlin. The title of the book means "the revealer of secrets." See also J. Ginsburg, "Rabbi ben Ezra on Permutations and Combinations," *Mathematics Teacher*, XV, 347.

Levi ben Gerson, in his *Maassei Choscheb* (*Work of the Computer*), written in 1321, carried the subject considerably farther. He gave rules for the permutation of n things taken all together and also taken r at a time, and for the combination of n things taken r at a time.[1]

A few years later Nicole Oresme (*c.* 1360) wrote a work[2] in which he gave the sum of the numbers representing the combinations of six things taken 1, 2, 3, 4, and 5 at a time. He also gave these combinations in detail, as that ${}_2C_6 = 15$, ${}_3C_6 = 20$, and so on, of course in the rhetorical form, and seems to have known the general law involved, although he did not state it.

First Evidence of Permutations in Print. The first evidence of an interest in the subject to be found in the printed books is given in Pacioli's *Sūma* (1494), where he showed how to find the number of permutations of any number of persons sitting at a table.[3] In England the subject was touched upon by W. Buckley (*c.* 1540), who gave special cases of the combinations of n things taken r at a time. Tartaglia (1523) seems first to have applied the theory to the throwing of dice.[4]

In the 16th century the learned Rabbi Moses Cordovero[5] wrote the *Pardes Rimmonim* (*Orchard of Pomegranates*),[6] in which he gave an interesting treatment of permutations and combinations and showed some knowledge of the general laws.[7]

[1] Eneström, *Bibl. Math.*, XIV (3), 261; G. Lange, German translation of the treatise, published at Frankfort a. M., 1909; Tropfke, *Geschichte*, VI (2), 64.

[2] *Tractatus de figuratione potentiarum et mensurarum difformitatum.* See H. Wieleitner, "Ueber den Funktionsbegriff und die graphische Darstellung bei Oresme," *Bibl. Math.*, XIV (3), 193.

[3] Fol. 43, *v*. He gives the results for $n = 1, 2, \ldots, 11$, and adds "Et sic in infinitum."


[4] In the *General Trattato*, II, fol. 17, *r*., he states that he discovered the rule: "Regola generale del presente auttore ritrouata il primo giorno di quarasima l'anno 1523. in Verona, di sapere trouare in quanti modi puo variar il getto di che quantita di dati si voglia nel tirar quelli." See also *L'Enseignement Mathématique*, XVI (1914), 92.

[5] Born at Safed, Palestine, in 1522; died at Safed, June 25, 1570.

[6] Salonika, 1552, with later editions.

[7] M. Turetsky, "Permutations in the 16th century Cabala," *Mathematics Teacher*, XVI, 29.

At about the same time Buteo not only discussed the question of the number of possible throws with four dice[1] but took up the problem of a combination lock with several movable cylinders like those shown in the illustration of the lock below.



EARLY COMBINATION LOCK

From Buteo's *Logistica*, 1560 ed., p. 313

As would naturally be expected, special cases of combinations of various kinds occur in the works of the 17th century. An illustration is found in the *Artis Analyticae Praxis* (p. 13) of Harriot, where the following symbolism is used for the product of binomials:

$$\begin{array}{l|l} a-b & aaaa-baaa+bcaa \\ a-c & \quad -caaa+bdaa \\ a-d & \quad -daaa+cdaa-bcda \\ \underline{a-f} & \quad -faaa+bfaa-bcfa \\ & \qquad\quad +cfaa-bdfa \\ & \qquad\quad +dfaa-cdfa+bcdf \end{array}$$

The first writer to give the general rule that

$$_nC_r = \frac{n(n-1)(n-2)\cdots(n-r+1)}{r!}$$

was Hérigone[2] (1634).

[1] "Ludens aleator tesseris quatuor, quaero quibus & quot modis inter se diuersis iacere possit?" *Logistica*, Lyons, 1559; 1560 ed., p. 305.

[2] *Cursus mathematicus*, II, 102. Paris, 1634.

In his work on the arithmetic triangle[1] Pascal showed the relation between the formation of the binomial coefficients and the theory of combinations, a subject also treated of by Fermat and others. Among the early writers upon the theory were Huygens, Leibniz,[2] Frénicle,[3] and Wallis,[4] and there is a brief tract on the subject which is thought to be due to Spinoza (1632–1677).[5]

The first work of any extent that is devoted to the subject was Jacques Bernoulli's *Ars Conjectandi*.[6] This work contains the essential part of the theory of combinations as known today. In it appears in print for the first time, with the present meaning, the word "permutation."[7] For this concept Leibniz had used *variationes* and Wallis had adopted *alternationes*. The word "combination" was used in the present sense by both Pascal and Wallis.[8] Leibniz used *complexiones* for the general term, reserving *combinationes* for groups of two elements and *conternationes* for groups of three,—words which he generalized by writing *con2natio*, *con3natio*, and so on.

Probability.[9] The theory of probability was mentioned in connection with the throwing of dice by Benvenuto d' Imola, a commentator on Dante's *Divina Commedia*, printed in the

[1] Written *c.* 1654 but printed posthumously in 1665. Beginning at this point, the reader may profitably consult the *Encyklopädie der math. Wissensch.*, I, 29.

[2] *Ars combinatoria*, 1666.

[3] "Abrégé des combinaisons" (1676), published in the *Mém. de l'acad. royale des sciences*, Paris, V (1729), 167.

[4] *De combinationibus, alternationibus, et partibus aliquotis, tractatus* (1685), in his *Opera*, II, 483 (Oxford, 1693).

[5] D. Bierens de Haan, "Twe zeldzame Werken van Benedictus Spinoza," *Nieuw Archief voor Wiskunde*, Amsterdam, XI (1884), 49. The title of the tract is *Reeckening van Kanssen*, and the work appeared in 1687, ten years after Spinoza's death.

[6] Posthumously printed at Basel in 1713. There is an English edition of 1795 under the title: *Permutations and Combinations: Being an Essential and Fundamental Part of the Doctrine of Chances.*

[7] "*De Permutationibus. Permutationes* rerum voco variationes. . . ."

[8] In the latter's *De Combinationibus*, English ed., 1685; *Opera* (1693), II, 483. His definition of combinations is on page 489 of that work.

[9] I. Todhunter, *History of the Mathematical Theory of Probability*, Cambridge, 1865; C. Gouraud, *Histoire du Calcul des Probabilités*, Paris, 1848.

Venice edition of 1477.[1] The gambling question first appears in a mathematical work, however, in Pacioli's *Sūma*[2] (1494). Here two gamblers are playing for a stake which is to go to the one who first wins n points, but the play is interrupted when the first has made p points and the second q points. It is required to know how to divide the stakes. The general problem also appears in the works of Cardan[3] (1539) and, as already stated, of Tartaglia[4] (1556). It first attracted wide attention in connection with the question proposed to Pascal (*c.* 1654) and by him sent to Fermat. The statement was substantially the one given in Pacioli to the effect that two players of equal skill left the table before completing the game. The stakes, the necessary score, and the score of each person being known, required to divide the stakes. Pascal and Fermat agreed upon the result, but used different methods in solving. As a result of the discussion so much interest was aroused in the theory that the doctrine of probability is generally stated to have been founded by Pascal and Fermat.

The first printed work on the subject was probably a tract of Huygens that appeared in 1657.[5] There also appeared an essay upon the subject by Pierre Rémond de Montmort in 1708.[6] The first book devoted entirely to the theory of probability was the *Ars Conjectandi* (1713) of Jacques Bernoulli, already mentioned. The second book upon the subject was De Moivre's *Doctrine of Chances: or, A Method of Calculating the Probability of Events in Play* (1718); and the third,

[1] Cantor, *Geschichte*, II (2), 327; Tropfke, *Geschichte*, II (1), 356. This is the fifth or sixth printed edition, Hain 5942; Copinger, I, 185, No. 5942.

[2] "Una brigata gioca apalla a .6o. el gioco e .1o ꝑ caccia. e fāno posta duc .1o. acade ꝑ certi accidēti che nō possano fornire e luna ꝑte a .5o. e laltra .2o. se dimanda che tocca ꝑ ꝑte de la posta." Fol. 197, *r.*

[3] *Practica*, Milan, 1539, "Caput 61. De extraordinariis & ludis," No. 17 of the chapter.

[4] *General Trattato* (1556), I, fol. 265, *r.*, where he quotes Pacioli under the title "Error di fra Luca dal Borgo." On a trace of the theory in a writing by Giovanni Francesco Peverone (*c.* 1550), see L. Carlini, *Il Pitagora*, VII, 65.

[5] "De ratiociniis in ludo aleae," in Van Schooten's *Exercitationum mathematicarum libri quinque*, Leyden, 1657. See also the *pars prima* of Bernoulli's *Ars Conjectandi*.

[6] *Essai d'analyse sur les jeux d'hasard*, Paris, 1708; 2d ed., *ibid.*, 1714.

Thomas Simpson's *Laws of Chance* (1740). One of the best-known works on the theory is Laplace's *Théorie analytique des probabilités*, which appeared in 1812. In this is given his proof of the method of least squares.

The application of the theory to mortality tables in any large way may be said to have started with John Graunt, whose *Natural and Political Observations* (London, 1662) gave a set of results based upon records of deaths in London from 1592. The first tables of great importance, however, were those of Edmund Halley, contained in his memoir on *Degrees of Mortality of Mankind*,[1] in which he made a careful study of annuities. It should be said, however, that Cardan seems to have been the first to consider the problem in a printed work, although his treatment is very fanciful. He gives a brief table in his proposition "Spatium vitae naturalis per spatium vitae fortuitum declarare," this appearing in the *De Proportionibvs Libri V*,[2] p. 204.

Although a life-insurance policy is known to have been underwritten by a small group of men in London in 1583, it was not until 1699 that a well-organized company was established for this purpose.

Besides the early work of Graunt and Halley there should be mentioned the *Essai sur les probabilités de la vie humaine* (Paris, 1746; supplementary part, 1760) by Antoine Deparcieux the elder (1703–1768). The early tables were superseded in the 18th century by the Northampton Table. Somewhat later the Carlisle Table was constructed by Joshua Milne (1776–1853). In 1825 the Equitable Life Assurance Society of London began the construction of a more improved table, since which time other contributions in the same field have been made by the Institute of Actuaries of Great Britain in coöperation with similar organizations, by Sheppard Homans (*c.* 1860) of New York,—the so-called American Experience Table,—and by Emory McClintock (1840–1916), also of New York.

[1] *Phil. Trans.*, London, 1693.

[2] Basel, 1570. For a sketch of the later tables see the articles on Life Insurance in the encyclopedias.

TOPICS FOR DISCUSSION

1. Leading steps in the development of algebra.
2. General racial characteristics shown in the early development of algebra.
3. The early printed classics on algebra.
4. Various names for algebra, with their origin and significance.
5. Development of algebraic symbolism relating to the four fundamental operations and to aggregations.
6. Development of symbolism relating to powers and roots.
7. Development of symbolism relating to the equality and to the inequality of algebraic expressions.
8. Methods of expressing equations, with a discussion of their relative merits.
9. Methods of solving linear equations.
10. Methods of solving quadratic equations.
11. History of the discovery of the method of solving cubic and biquadratic equations.
12. History of continued fractions and of their uses.
13. General steps in the development of the numerical higher equation.
14. History of the Rule of False Position, with the reasons for the great popularity of the rule.
15. Development of the idea of classifying equations according to degree instead, for example, according to the number of terms.
16. Development of the indeterminate equation.
17. General steps in the application of trigonometry to the solution of the quadratic and cubic equations.
18. General steps in the early development of determinants.
19. History of the Rule of Three and of its relation to proportion.
20. General nature of series in the early works on mathematics.
21. History of infinite products in the 17th century.
22. The historical development of the Binomial Theorem.
23. History and applications of Taylor's and Maclaurin's formulas.
24. History of the Pascal Triangle and of its applications.
25. The invention of logarithms and the history of their various applications.
26. History of permutations, combinations, and the theory of probability.

CHAPTER VII

ELEMENTARY PROBLEMS

I. Mathematical Recreations

Purpose of the Study. In this chapter we shall consider a few of the most familiar types of problems that have come down to us. Some of these types relate to arithmetic, while others have of late taken advantage of algebraic symbolism, although at one time they were solved without the modern aids that algebra supplies.

Mathematical Recreations. Ever since problems began to be set, the mathematical puzzle has been in evidence. Without defining the limits that mark the recreation problem it may be said that the Egyptians and Orientals proposed various questions that had no applications to daily life, the chief purpose being to provide intellectual pleasure. The Greeks were even more given to this type of problem, and their geometry was developed partly for this very reason. In the later period of their intellectual activity they made much of indeterminate problems, and thereafter this type ranked among the favorite ones.

In the Middle Ages there developed a new form of puzzle problem, one suggested by the later Greek writers and modified by Oriental influences. This form has lasted until the present time and will probably continue to have a place in the schools.

Problems of Metrodorus. So far as the Greeks were concerned, the source book for this material is the *Greek Anthology*.[1] This contains the arithmetical puzzles supposed to be due to

[1] The first noteworthy edition was that of Friedrich Jacobs, Leipzig, 1813–1817. There is an English translation by W. R. Paton, London, 1918, being Volume V of the Loeb Classical Library. In this translation the arithmetic problems begin on page 25, and from these the selections given here have been made.

Metrodorus about the year 500(?). A few of these problems will serve to show the general nature of the collection.

Polycrates Speaks: "Blessed Pythagoras, Heliconian scion of the Muses, answer my question: How many in thy house are engaged in the contest for wisdom performing excellently?"

Pythagoras Answers: "I will tell thee, then, Polycrates. Half of them are occupied with belles lettres; a quarter apply themselves to studying immortal nature; a seventh are all intent on silence and the eternal discourse of their hearts. There are also three women, and above the rest is Theano. That is the number of interpreters of the Muses I gather round me."

The following problem relates to a statue of Pallas:

"I, Pallas, am of beaten gold, but the gold is the gift of lusty poets. Christians gave half the gold,[1] Thespis one eighth, Solon one tenth, and Themison one twentieth, but the remaining nine talents and the workmanship are the gift of Aristodicus."

The following relates to the finding of the hour indicated on a sundial and still appears in many algebras, modified to refer to modern clocks:

"Best of clocks,[2] how much of the day is past?"
"There remain twice two thirds of what is gone."

The next problem involves arithmetic series, as follows:

Crœsus the king dedicated six bowls weighing six minæ,[3] each [being] one drachma heavier than the other.[4]

A type that has long been familiar in its general nature is seen in the following:

A. "Where are thy apples gone, my child?"
B. "Ino has two sixths, and Semele one eighth, and Autonoe went off with one fourth, while Agave snatched from my bosom and carried

[1] It should be recalled that this was written probably about the time of such Christian scholars as Capella and Cassiodorus.
[2] Literally, hour indicator.
[3] A mina contained 100 drachmas.
[4] That is, than the one next smaller. Find the weight of each.

away a fifth. For thee ten apples are left, but I, yes I swear it by dear Cypris, have only this one."[1]

The following problem has more of an Oriental atmosphere:

"After staining the holy chaplet of fair-eyed Justice that I might see thee, all-subduing gold, grow so much, I have nothing; for I gave forty talents under evil auspices to my friends in vain, while, O ye varied mischances of men, I see my enemy in possession of the half, the third, and the eighth of my fortune."[2]

One of the remote ancestors of a type frequently found in our algebras appears in the following form:

"Brick-maker, I am in a great hurry to erect this house. Today is cloudless, and I do not require many more bricks, but I have all I want but three hundred. Thou alone in one day couldst make as many, but thy son left off working when he had finished two hundred, and thy son-in-law when he had made two hundred and fifty. Working all together, in how many days can you make these?"

This collection of puzzles, now attributed entirely to Metrodorus, contains numerous enigmas, one of which is numerical enough to deserve mention:

If you put one hundred in the middle of a burning fire, you will find the son and slayer of a virgin.[3]

Comparison with Oriental Problems. Such problems seem more Oriental than Greek in their general form, but if we could ascertain the facts we should probably find that every people cultivated the somewhat poetic style in the recreations of mathematics. It happens, however, that we have more evidence of it in India and China than we have in the Mediterranean countries, and hence we are led to believe it was more frequently found among the higher class of mathematicians in the East than among those of the West.

[1] There were 120, for $120 = 40 + 15 + 30 + 24 + 10 + 1$.

[2] $480 + 320 + 120 + 40 = 960$.

[3] The answer is Pyrrhus, son of Deidameia and slayer of Polyxena; for if ρ, the Greek symbol for 100, is inserted in the middle of the genitive form πυρός (fire), it becomes πυρρός (Pyrros, Pyrrhus). This is the mythological Pyrrhus (Neoptolemus), son of Achilles and Deidameia.

Medieval Collections. The first noteworthy collection of recreations, after the one in the *Greek Anthology*, is the *Propositiones ad acuendos iuvenes*, of which there is extant no manuscript written before the year 1000. This collection is attributed to Alcuin of York (*c.* 775), who is known to have sent a list of such recreations to Charlemagne.[1] It contains many stock problems such as those of the hare and hound, and the cistern pipes. Rabbi ben Ezra (*c.* 1140), Fibonacci (1202), Jordanus Nemorarius (*c.* 1225), and many other medieval writers made use of these standard types.

Printed Books. The first noteworthy collection of recreative problems to appear in print was that of Claude-Gaspar Bachet (1612).[2] While not so popular as various later works, and containing much that is trivial, it was a pioneer and is much better than some of those that went through many more editions.

From the bibliographical standpoint the most interesting of the printed collections is that of a Jesuit scholar, Jean Leurechon (1624).[3] He published his work under the name of H[endrik] van Etten at Pont-à-Mousson in 1624. It was a poor collection of trivialities,[4] but it struck the popular fancy and went through at least thirty-four editions before 1700, some of these being published under other names.

The next writer of note was Jacques Ozanam (1640–1717), a man who was self-taught and who had a gift for teaching others. He had faith in the educational value of recreations, and this fact, together with his familiarity with the subject and his success as a teacher, enabled him to write one of the most popular works on the subject that has ever appeared.

[1] "Misi aliquas figuras Arithmeticae subtilitatis laetitiae causa" (Cantor *Geschichte*, I (2), 784).

[2] *Problemes plaisans et delectables, qui se font par les nombres. Partie recueillis de diuers autheurs, & inuentez de nouueau auec leur demonstration*, Lyons 1612. There is a copy of this edition in the Harvard Library. Later editions: Lyons, 1624; Paris, 1874, 1879, 1884.


[3] Born at Bar-le-Duc, *c.* 1591; died at Pont-à-Mousson, January 17, 1670. He wrote on astronomy.

[4] Montucla, in his revision of Ozanam, speaks of it as "une pitoyable rapsodie."

The work was first published in 1692 or 1694[1] and since then there have been at least twenty different editions.

There have been many other works on the subject,[2] but none of them has had the popular success of those of Leurechon and Ozanam.

Japanese Geometric Problems. The Japanese inherited from the Chinese a large number of curious geometric problems, and by their own ingenuity and perseverance elaborated these tests of skill until they far surpassed their original teachers. Some of these problems were mentioned in Volume I, and the circle problem will be referred to in Chapter X of this volume; but in this connection it is proper to refer to one type of interesting problems frequently found in the early Japanese works. These problems refer to the inscribing and measuring of circles inscribed in various figures such as semicircles, fans, and ellipses.



A FAN PROBLEM FROM JAPAN
From Takeda Shingen's *Sampō Benran*, 1824


2. Typical Problems

Pipes filling the Cistern. Few problems have had so extended a history as the familiar one relating to the pipes filling a cistern,[3] and the traveler who is familiar with the Mediterranean

[1] The date 1692 is on the testimony of Montucla, in his 1790 edition of Ozanam. It is probable that he was in error on this point. See *L'Intermédiare des Mathématiciens*, VI, 112, and various histories of mathematics.

[2] Bibliographies that are fairly complete may be found in E. Lucas, *Récréations Mathématiques*, 4 vols., I, 237 (Paris, 1882–1894); W. Ahrens, *Mathematische Unterhaltungen und Spiele*, p. 403 (Leipzig, 1901; 2d ed., 1918). These are the leading modern contributors to the subject, the works of Lucas being probably the best that have as yet appeared.

[3] See the author's article in the *Amer. Math. Month.*, XXIV, 64, from which extracts are here made.



PROBLEM OF TANGENT CIRCLES

From a manuscript by Iwasaki Toshihisa (*c.* 1775)

lands cannot fail to recognize that here is its probable origin. Not a town of any size that bears the stamp of the Roman power is without its public fountain into which or from which several conduits lead. In the domain of physics, therefore, this would naturally be the most real of all the problems that came within the purview of every man, woman, or child of that civilization. Furthermore, the elementary clepsydra[1] may also have suggested this line of problems, the principle involved being the same.

The problem in definite form first appears in the Μετρήσεις (*metre'seis*) of Heron (*c.* 50?), and although there is some question as to the authorship and date of the work, there is none as to the fact that this style of problem would appeal to such a writer as he. It next appears in the writings of Diophantus (*c.* 275)[2] and among the Greek epigrams of Metrodorus (*c.* 500?), and soon after this it became common property in the East as well as the West. It is found in the list attributed to Alcuin (*c.* 775); in the *Lilāvati* of Bhāskara[3] (*c.* 1150); in the best-known of all the Arab works on arithmetic, the *Kholâsat al-Ḥisâb* of Behâ Eddîn (*c.* 1600); and in numerous medieval manuscripts. When books began to be printed it was looked upon as one of the standard problems of the schools, and many of the early writers gave it a prominent position, among them being men like Petzensteiner (1483), Tonstall (1522), Gemma Frisius (1540), and Robert Recorde (*c.* 1542).[4]

[1] Attributed to Plato (*c.* 380 B.C.) but improved by Ctesibius of Alexandria in the second century B.C. On the subject of clepsydræ see Chapter IX of this volume.

[2] In Bachet's edition (the Fermat edition of 1670, p. 271) appears this metrical translation:

Totum implere lacum tubulis è quatuor, uno
Est potis iste die, binis hic & tribus ille,
Quatuor at quartus.
Dic quo spatio simul omnes.

[3] See Taylor's translation, p. 50; Colebrooke translation, p. 42.

[4] In Recorde it appears for the first time in English: "Ther is a cestern with iiij. cocks, conteinyng 72 barrels of water, And if the greatest cocke be opened, the water will auoyde cleane in vj howers," etc. (*Ground of Artes*, 1558 ed., fol. A 7, *v.*).

Variants of the Problem. Such, then, was the origin of what was once a cleverly stated problem of daily life. This problem, like dozens of others, went through many metamorphoses, of which only a few will here be mentioned.

In the 15th century, and probably much earlier, there appeared the variant of a lion, a dog, and a wolf, or other animals, eating a sheep,[1] and this form was even more common in the 16th century.[2]

In the 16th century we also find in several books the variant of the case of men building a wall or a house, and this form has survived to the present time. It appeared in Tonstall's *De Arte Supputandi* (1522)[3] and in Cataneo's work (1546),[4] and in due time became modified to the form beginning, "If *A* can do a piece of work in 4 days, *B* in 3 days," and so on.

The influence of the wine-drinking countries shows itself in the variant given by Gemma Frisius (1540),[5] who states that a man can drink a cask of wine in 20 days, but if his wife drinks with him it will take only 14 days, from which it is required to find the time it would take his wife alone.

The influence of a rapidly growing commerce led one of the German writers of 1540 to consider the case of a ship with

[1] Johann Widman (1489) under the chapter title "Eyn fasz mit 3 zapffen." His form is: "Lew Wolff Hunt Itm̄ des gleichen 1 lew vnd 1 hunt vñ 1 wolff diese essen mit einander 1 schaff. Vnd der lew esz das schaff allein in einer stund. Vnd d' wolf in 4 stunden. Vnd der hunt in 6 stunden. Nun ist die frag wan sy dass schaff all 3 mit eināder essen/ in wie lāger zeit sy das essen" (1509 ed., fol. 92; 1519 ed., fol. 112).

[2] Thus Cataneo, *Le Pratiche*, 1546; Venice edition of 1567, fol. 59, *v.*: "Se un Leone mangia in 2. hore una pecora, & l' Orso la mangia in 3. hore, & il Leopardo la mangia in 4. hore, dimandasi cominciando a mangiare una pecora tutti e 3. a un tratto in quanto tempo la finirebbono."

This form is also found in J. Albert's work of 1534 (1561 ed., fol. N viii), in Coutereels (1631 ed., p. 352), and in the works of numerous other writers.


In this chapter a few authors of textbooks will be mentioned whose names are not of sufficient importance to entitle them to further attention. The dates will serve to show their relative chronological position. For names of major importance consult Volume I.

[3] With the statement that it is similar to the one about the cistern pipes: "Questio hæc similis est illi de cisterna tres habēte fistulas: et simili modo soluenda" (fol. f 1).

[4] See fol. 60, *v.*, of the Venice edition of 1567.

[5] 1563 edition of his arithmetic, fol. 38.

3 sails, by the aid of the largest of which a voyage could be made in 2 weeks, with the next in size in 3 weeks, and with the smallest in 4 weeks, it being required to find the time if all




FROM CALANDRI'S WORK OF 1491

The problems of the pipe filling the cistern and of the serpent crawling out of the well. Calandri's was the first arithmetic printed with illustrations

three were used. Unfortunately several factors were ignored, such as that of one sail blanketing the others and the fact that the speed is not proportional to the power.[1]

[1] "Item/ 1 ein Schiff mit 3 Siegeln gehet vom Sund gen Riga/ Mit dem grösten allein/ in 2 wochen/ Mit dem andern/ in 3 wochen/ Vnnd mit dem kleinsten/in 4 wochen," etc. (J. Albert (1540; 1561 ed., fol. N vii)).

The agricultural interests changed the problem to that of a mill with four "Gewercken,"[1] and other interests continued to modify it further until, as is usually the case, the style of problem has tended to fall from its own absurdity. Its varied history may be closed by referring to a writer of the early 19th century,[2] moved by a bigotry which would hardly be countenanced today, who proposed to substitute a problem relating to priests praying for souls in purgatory.



THE TURKS AND CHRISTIANS

From Buteo's *Logistica*, Lyons, 1559 (1560 ed., p. 304). The problem begins: "In naui vectores quindecim Christiani totidēq; Iudei, suborta tēpestate magna"

Turks and Christians. There is a well-known problem which relates that fifteen Turks and fifteen Christians were on a ship which was in danger, and that half had to be sacrificed. It being necessary to choose the victims by lot, the question arose as to how they could be arranged in a circle so that, in counting round, every fifteenth one should be a Turk.


It is probable that the problem goes back to the custom of *decimatio* in the old Roman armies,[3] the selection by lot of every tenth man when a company had been guilty of cowardice, mutiny, or loss of standards in action. Both Livy (II, 59) and Dionysius (IX, 50) speak of it in the case of the mutinous army of the consul Appius Claudius (471 B.C.), and Dionysius further speaks of it as a general custom. Polybius (VI, 38) says that it was a usual punishment when troops had given way to

[1] "Ein Mülmeister hat ein Müle mit vier Gewercken/ Mit dem ersten mehlt er in 23 stūden 35 Scheffel/Mit dem andern 39 Scheffel/ Mit dem dritten 46 Scheffel/Vnnd mit dem vierten 52 Scheffel," etc. The question then is, How long it will take them together to grind 19 Wispel (1 Wispel = 24 Scheffel) (*ibid.*).

[2] R. Hay, *The Beauties of Arithmetic*, p. 218 (1816).

[3] E. Lucas, *Arithmétique Amusante*, p. 17 (Paris, 1895).

panic. The custom seems to have died out for a time, for when Crassus resorted to decimation in the war of Spartacus he is described by Plutarch (*Crassus*, 10) as having revived an ancient punishment. It was extensively used in the civil wars and was retained under the Empire, sometimes as *vicesimatio* (every twentieth man being taken), and sometimes as *centesimatio* (every hundredth man).



THE JOSEPHUS PROBLEM IN JAPAN

From Muramatsu Kudayū Mosei's *Manloku Jinkō-ri* (1665)

Now it is very improbable that those in charge of the selection would fail to have certain favorites, and hence it is natural that there may have grown up a scheme of selection that would save the latter from death. Such customs may depart, but their influence remains.


In its semimathematical form the problem is first referred to in the work of an unknown author, possibly Ambrose of Milan (*c.* 370), who wrote, under the nom de plume of Hegesippus, a work *De bello iudaico*.[1] In this work he refers to the fact that Josephus was saved on the occasion of a choice of this kind.[2] Indeed, Josephus himself refers to the matter of his being saved by lucky chance or by the act of God.[3]

[1] Edited by C. F. Weber and J. Caesar, Marburg, 1864. See W. Ahrens, *Math. Unterhaltungen und Spiele*, p. 286 (Leipzig, 1901; 2d ed., 1918).

[2] "Itaque accidit ut interemtis reliquis Iosephus cum altero superesset neci" (quoted from Ahrens, *loc. cit.*).

[3] Καταλείπεται δὲ οὗτος, εἴτε ὑπὸ τύχης χρὴ λέγειν εἴτε ὑπὸ Θεοῦ προνοίας, σὺν ἑτέρῳ

The oldest European trace of the problem, aside from that of Hegesippus, is found in a manuscript of the beginning of the 10th century. It is also referred to in a manuscript of the 11th century and in one of the 12th century. It is given in



THE JOSEPHUS PROBLEM IN JAPAN

From Miyake Kenryū's *Shojutsu Sangaku Zuye* (1795 ed.), showing the problem of the stepmother, referred to on page 544

the *Ta'hbula* of Rabbi ben Ezra (*c.* 1140), and indeed it is to this writer that Elias Levita, who seems first to have given it in printed form (1518), attributes its authorship.

The problem, as it came to be stated, related that Josephus, at the time of the sack of the city of Jotapata by Vespasian, hid himself with forty other Jews in a cellar. It becoming necessary to sacrifice most of the number, a method analogous to the old Roman method of *decimatio* was adopted, but in such a way as to preserve himself and a special friend. It is

on this account that German writers still call the ancient puzzle by the name of Josephsspiel.

Chuquet (1484) mentions the problem, as does at least one other writer of the 15th century.[1] When, however, printed works on algebra and higher arithmetic began to appear, it became well known. The fact that such writers as Cardan[2] and Ramus[3] gave it prominence was enough to assure its coming to the attention of scholars.[4]

Like so many curious problems, this one found its way to the Far East, appearing in the Japanese books as relating to a stepmother's selection of the children to be disinherited. With characteristic Japanese humor, however, the woman was described as making an error in her calculations, so that her own children were disinherited and her stepchildren received the estate.

Testament Problem. There is a well-known problem which relates that a man about to die made a will bequeathing $\frac{1}{3}$ of his estate to his widow in case an expected child was a son, the son to have $\frac{2}{3}$; and $\frac{2}{3}$ to the widow if the child was a daughter, the daughter to have $\frac{1}{3}$. The issue was twins, one a boy and the other a girl, and the question arose as to the division of the estate.

The problem in itself is of no particular interest, being legal rather than mathematical, but it is worthy of mention because it is a type and has an extended history. Under both the Roman and the Oriental influence these inheritance problems played a very important rôle in such parts of analysis as the ancients had developed. In the year 40 B.C. the *lex Falcidia* required at least $\frac{1}{4}$ of an estate to go to the legal heir. If more than $\frac{3}{4}$ was otherwise disposed of, this had to be reduced by the rules of partnership. Problems involving this "Falcidian

[1] Anonymous MS. in Munich. See *Bibl. Math.*, VII (2), 32; Curtze, *ibid.*, VIII (2), 116; IX (2), 34; *Abhandlungen*, III, 123.

[2] In his *Practica* of 1539.

[3] See his edition of 1569, p. 125.

[4] It is also in Thierfelder's arithmetic (1587, p. 354), in Wynant van Westen's *Mathemat. Vermaecklyckh* (1644 ed., I, 16), in Wilkens's arithmetic of 1669 (p. 395), and in many other early works.

fourth" were therefore common under the Roman law, just as problems involving the widow's dower right were and are common under the English law.

The problem as stated above appears in the writings of Juventius Celsus (*c.* 75), a celebrated jurist who wrote on testamentary law; in those of Salvianus Julianus, a jurist in the reigns of Hadrian (117–138) and Antoninus Pius (138–161); and in those of Cæcilius Africanus (*c.* 100), a writer who was celebrated for his knotty legal puzzles.[1]

In the Middle Ages it was a favorite conundrum, and in the early printed arithmetics[2] it is often found in a chapter on inheritances which reminds one of the Hindu mathematical collections. It went through the same later development that characterizes most problems, and finally fell on account of its very absurdity. That is, Widman (1489) takes the case of triplets, one boy and two girls,[3] and in this he is followed by Albert (1534) and Rudolff (1526). Cardan (1539) complicates it by supposing 4 parts to go to the son and 1 part to the mother, or 1 part to the daughter and 2 parts to the mother, and in some way decides on an 8, 7, 1 division.[4] Texeda (1546) supposes 7 parts to go to the son and 5 to the mother, or 5 to the daughter and 6 to the mother, while other writers of the 16th century complicate the problem even more.[5] The final complications of the "swanghere Huysvrouwe" or "donna grauida" are found in some of the Dutch books, and these and

[1] Coutereels (Eversdyck edition of 1658, p. 382) traces the problem back to lib. 28, title 2, law 13, of the *Digest* of Julianus. He gives the usual 4, 2, 1 division as followed by Tartaglia, Rudolff, Ramus, Trenchant, Van der Schuere, and others. Coutereels, however, argues for the 4, 3, 2 division, and in this he has the support of various writers. Peletier gives 2, 2, 1, and others give 9, 6, 4. Brief historical notes appear in other books, as in the Schoner edition of Ramus (1586 ed., p. 186).

[2] Thus we have "Ein Testament" (Widman), "Erbteilung vnd vormundschaft" (Riese), "Erf-Deelinghe" (Van der Schuere), and "Erbtheilugs-Rechnung" (Starcken).

[3] Edition of 1558, fol. 97. He then divides the property in the proportion 4, 2, 1, 1.

[4] *Practica*, cap. 66, ex. 87.

[5] Ghaligai (1521), Köbel (1514), Riese (*Rechnung nach lenge*, 1550 ed.), Trenchant (1566), Van der Schuere (1600), Peletier (1607 ed., p. 244), Coutereels (1631 ed., p. 358), Starcken (1714 ed., p. 444), Tartaglia (*Tvtte l'opere d'aritmetica*, 1592 ed., II, 136).

the change in ideas of propriety account for the banishment of the problem from books of our day.[1] The most sensible remark about the problem to be found in any of the early books is given in the words of the "Scholer" in Robert Recorde's *Ground of Artes* (*c.* 1542): "If some cunning lawyers had this matter in scanning, they would determine this testament to be quite voyde, and so the man to die vntestate, because the testament was made vnsufficient."[2]

Problems of Pursuit. Problems of pursuit are among the most interesting elementary ones that have had any extended history. It would be difficult to conceive of problems that seem more real, since we commonly overtake a friend in walking, or are in turn overtaken. It would therefore seem certain that this problem is among the ancient ones in what was once looked upon as higher analysis. We have a striking proof that this must be the case in the famous paradox of Achilles and the Tortoise.[3]

It is a curious fact, however, that the simplest case, that of one person overtaking another, is not found in the Greek collections, although it appears in China[4] long before it does in the West. It is given, perhaps for the first time in Europe, among the *Propositiones ad acuendos juvenes* attributed to Alcuin, in the form of the hound pursuing the hare.[5] Thereafter it was looked upon as one of the necessary questions of European mathematics, appearing in various later medieval manuscripts. It is given in Petzensteiner's work of 1483, Calandri[6] used it in 1491, Pacioli has it in his *Sūma*[7] (1494),

[1] "Soo ontfangt sy ter tijdt haerder baringhe eenen Sone met een Dochter/ eñ een Hermaphroditus, dat is/half Man/half Vrouwe." Van der Schuere, 1600, fol. 98. In this case he divides 3175 guldens thus: d. 254, m. 508, s. 1524, h. 889. The same problem appears in Clausberg, *Demonstrative Rechen-Kunst*, 1772.

[2] 1558 ed., fol. X 8.

[3] For a study of this problem see F. Cajori, *Amer. Math. Month.*, XXII, 1 seq.

[4] For example, in the *Nine Sections* (*c.* 1105 B.C.?) and in Liu Hui's commentary (*c.* 263) on this classic. See also Volume I, page 32.


[5] "De cursu canis ac fuga leporis."

[6] "Una lepre e inanzi aun chane 3000 passi et ogni 5 passi delcane sono ꝑ 8 diquegli della lepre uosapere inquanti passi elcane ara giūto lalepre."

[7] "Vna lepre e dinançe a vn cane passa .60. e per ogni passa .5. che fa el cane la lepre ne fa .7. e finalmente el cane lagiongni [la giongi in the edition of 1523,

and most of the prominent writers on algebra or higher arithmetic inserted it in their books from that time on.[1]

In those centuries in which commercial communication was chiefly by means of couriers who traveled regularly from city to city (a custom still determining the name of *correo* for a postman in certain parts of the world) the problem of the hare



PROBLEM OF THE HARE AND HOUND

From a MS. of Benedetto da Firenze, *c.* 1460. It begins, "Vna lepre e înanzi a .1°. cane"

and hound naturally took on the form of, or perhaps paralleled, the one of the couriers. This problem was not, however, always one of pursuit, since the couriers might be traveling either in

from la giugnere, to overtake her] dimando in quanti passa el cane giõgera la lepre." Fol. 42, *v.* He says that the problem is not clear, because we do not know whether the "passa .60." are leaps of the dog or of the hare, showing that he felt bound to take the problem as it stood, without improving upon the phraseology.

[1] Thus Rudolff (*Kunstliche rechnung*, 1526; 1534 ed., fol. N vj); Köbel (*Rechenbuch*, 1514; 1549 ed., fol. 88, under the title "Von Wandern über Landt," with a picture in which the hare is quite as large as the hound); Cardan (*Practica*, 1539, cap. 66); Wentsel (1599, p. 51); Ciacchi (*Regole generali d'Abbaco*, p. 130, Florence, 1675); Coutereels (*Cyffer-Boeck*, 1690 ed., p. 584), and many others.

the same direction or in opposite directions.[1] This variant of the problem is Italian, for even the early German writers gave it with reference to Italian towns.[2] As a matter of course, it was also varied by substituting ships for couriers.[3]

It was natural to expect that the problem should have a further variant, namely, the one in which the couriers should not start simultaneously. In this form it first appeared in print in Germany in 1483,[4] in Italy in 1484,[5] and in England in 1522,[6] although doubtless known much earlier.

The invention of clocks with minute hands as well as hour hands gave the next variant, as to when both hands would be together,—a relatively modern form of the question, as is also the astronomical problem of the occurrence of the new moon. One of the latest forms has to do with the practical question of a railway time-table, but here graphic methods naturally take the place of analysis, so that of all the variants those of the couriers and the clock hands seem to be the only ones that will survive. Neither is valuable per se, but each is interesting, each is real within the range of easy imagination, and each involves a valuable mathematical principle,—a fairly refined idea of function.

[1] See Pacioli's *Sūma*, 1494, fol. 39, for various types.

[2] Thus Petzensteiner (1483, fol. 53), in his chapter "Von wandern," makes the couriers go to "rum" (Rome), thus: "Es sein zween gesellen die gand gen rum. Eyner get alle tag 6 meyl der ander geth an dem ersten tage 1 meyl an dem andern zwue etc. unde alle tag eyner meyl mer dan vor. Nu wildu wissen in wievil tagen eyner als vil hat gangen als der ander." Günther, *Math. Unterrichts*, p. 304.

[3] Thus Calandri (1491) says: "Una naue ua da Pisa a Genoua in 5 di: unaltra naue uiene da genoua a pisa in 3 di. uo sapere partendosi in nun medesimo tempo quella da Pisa per andare a Genoua et quella da Genoua p andare a pisa in quanti di siniscon terrano insieme."

[4] Petzensteiner's arithmetic, printed at Bamberg.

[5] Borghi's arithmetic.

[6] Tonstall's *De Arte Supputandi*, fol. 4, "Cvrsor ab Eboraco Londinvm proficiscens," etc. See also Cardan (*Practica*, 1539, cap. 66, with various types); Ghaligai (1521; 1552 ed., fol. 64); Albert (1540; 1561 ed., fol. P i); Baker (1568; 1580 ed., fol. 36); Coutereels (1631 ed., p. 371, and Eversdyck edition of 1658, p. 403); Trenchant (1566; 1578 ed., p. 280); Wentsel (1599, p. 51); Peletier (1549; 1607 ed., p. 290); Van der Schuere (1600, fol. 179); Schoner (notes on Ramus, 1586 ed., p. 174), and many others.

The Chessboard Problem. One of the best-known problems of the Middle Ages is that relating to the number of grains of wheat that can, theoretically speaking, be placed upon a chessboard, one grain being put on the first square, two on the second, four on the third, and so on in geometric progression, the total number being $2^{64} - 1$, or 18,446,744,073,709,551,615. The problem is Oriental. A chessboard problem of a different character appeared in the writings of one I Hang, a Chinese Buddhist of the T'ang Dynasty (620–907),[1] so that games on a checkered board had already begun to attract the attention of mathematicians in the East.



Zwen Bürger auß Oppenheym/ Einer Son Heynrich/der ander Contz vō Treber genant/ wolten mit einander gen Rom gehen/ Vnd Heynrich was alt/ vnd mocht einen tag nicht mehr denn zehen meilen gehen/ Aber Contz von Treber was jung vnd ſtarck/ der mocht einen tag 13. meilen gehen/ Deßhalben gieng Son Heynrich neun tag ehe auß Oppenheym denn Contz von Treber/ Alſo war Son Heynrich Contzen 90. meilen fürgangen/ ehe Contz angehabē hat außzugehen.

Nun iſt die frag/ in wie vil tagen Contz von Treber Son Heynrichen vbergangen/ vnd die zwen zuſamen kommen ſeind?

PROBLEM OF THE COURIERS

From Köbel's *Rechenbuch* (1514), the edition of 1564

Ibn Khallikân,[2] one of the best known of the Arab biographers (1256), relates[3] that when Sissah ibn Dahir invented the game of Chess, the king, Shihrâm, was filled with joy and commanded that chessboards should be placed in the temples. Furthermore, he commanded Sissah to ask for any reward he pleased. Thereupon Sissah asked for one grain of wheat for the first square, two for the next, and so on in geometric progression.


[1] G. Vacca, *Note Cinesi*, p. 135 (Rome, 1913). This problem is rather one of permutations.

[2] Or Challikân. Born September 22, 1211; died October 29, 1282.

[3] In his *Biographical Dictionary* (translation from the Arabic by Mac Guckin de Slane, 4 vols., Paris and London, 1843–1871), III, 69.

The result of the request is not recorded, but as an old German manuscript remarks, "Daz mecht kain kayser bezalen."[1]

The problem goes back at least as far as Mas'ûdî's *Meadows of Gold*[2] of the 10th century. It also appeared in the works of various other Arab writers,[3] and thence found its way into Europe through the *Liber Abaci*[4] (1202) of Fibonacci. It is found in numerous manuscripts of the 13th, 14th, and 15th centuries and in various early printed books.[5] The problem was much extended by later writers.[6] It found a variant in the problem of the horseshoe nails which appears in several manuscripts of the 15th and 16th centuries.



CHESSBOARD PROBLEM, *c.* 1400

From an Italian manuscript of *c.* 1400, now in the Columbia University Library

A Dutch arithmetician, Wilkens,[7] takes the ratio in the chessboard problem as three instead of two, and considers not only the number of grains but also the number of ships necessary to carry the total amount, the value of the cargoes, and the impossibility that all the countries of the world[8] should produce such an amount of wheat.[9]

[1] "No emperor could pay all that." Curtze, *Bibl. Math.*, IX (2), 113.

[2] Mas'ûdî died at Cairo in 956. A French translation in nine volumes, with Arabic text, appeared in Paris, 1861–1877. See also Boncompagni's *Bullettino*, XIII, 274.

[3] H. Suter, *Bibl. Math.*, II (3), 34.

[4] Boncompagni ed., I, 309.


[5] *E. g.*, Pacioli's *Sūma* (1494), fol. 43; Cardan's *Practica* (1539), cap. 66.

[6] As by Clavius, *Epitome* (1585), p. 297.

[7] 1669 ed., p. 112.

[8] "Al de Provintien van de gheheele werelt."

[9] For further historical notes see J. C. Heilbronner, *Historia Matheseos Universæ*, p. 440 (Leipzig, 1742).



THE PROBLEM OF THE HORSESHOE NAILS

From an anonymous MS. written in Italy *c.* 1535. As the problem is usually stated, the blacksmith receives one penny for the first nail, two pence for the second, four for the third, and so on, and there are twenty-four nails. This manuscript is in Mr. Plimpton's library.

The Mule and the Ass. Among the recreational problems that have come down to us there is one which appears in the form of an epigram with the name of Euclid attached. Rendered in English verse it is as follows:

A mule and an ass once went on their way with burdens of wine-skins;
Oppressed by the weight of her load, the ass was bitterly groaning.
The mule, observing her grievous complaints, addressed her this question:
"Mother, why do you murmur, with tears, for a maiden more fitting?
For give me one measure of wine, and twice your burden I carry;
But take one measure from me, and still you will keep our loads equal."
Tell me the measure they bore, good sir, geometry's master.[1]

3. Commercial Problems

Economic Problems. For the student of economics there is an interesting field in the problems of the 15th and 16th centuries, as may be seen from a few illustrations. The manuscripts and early printed books on arithmetic tell us that Venice was then the center of the silk trade, although Bologna, Genoa, and Florence were prominent. Florence was the chief Italian city engaged in the dyeing of cloth. "Nostra magnifica Città di Venetia," as Tartaglia so affectionately and appropriately called her, carried on her chief trade with Lyons, London, Antwerp, Paris, Bruges, Barcelona, Montpellier, and the Hansa towns, besides the cities of Italy. Chiarino (Florence, 1481) indicates the following as the most important cities with which Florence had extensive trade, his spelling being here preserved: Alessandria degypto, Marsilia, Mompolieri, Lisbona, Parigi, Bruggia, Barzalona, Londra, Gostatinopoli, and Dommasco, with the countries of Tunizi, Cypri, and Candia. Tartaglia gives Barcelona, Paris, and Bruges as the leading cities connected with Genoa in trade a half century later.

[1] *Euclidis Opera*, ed. Heiberg and Menge, VIII, 286 (Leipzig, 1916). The translation is by Professor Robbins, University of Michigan. See *The Classical Journal*, XV, 184.

[2] See the author's article in the *Amer. Math. Month.*, XXIV, 221, from which extracts are here made.

We also know from Chiarino the most important commodities of Florentine trade in the decade before America was discovered. These were rame (brass), stoppa (tow), zolphi (sulphur), smeriglio (emery), lana (wool), ghalla (gall), trementina (turpentine), sapone (soap), risi (rice), zucchari (sugar), cannella (cinnamon), piombo (lead), lini (flax), pece (pitch), acciai (thread), canapa (hemp), incenso (incense), indachi (indigo), mace (mace), cubeba (cubebs), borage (borax), and the ever-present saffron, the "king of plants," then everywhere used as a *sine qua non* in daily life, and now almost forgotten.



Ein Frow oder Haußmutter gehet auff den marckt/ kaufft vberhaupt ein Körblin mit Rebnerbyrn/darumb gibt sie achtzehen pfenning / so sie heim kompt / findet sie im Körblin hundert vnd achtzig byrn / Ist die frag/wie vil byren sie vmb ein pfenning habe? Thu/als obgelert/ so kompt dir zehen/ Also vil byren hat sie vmb einen pfenning / Vnd ist wolfeyl drumb.

THE PROBLEM OF THE MARKET WOMAN

From Köbel's *Rechenbüchlein* of 1514 (1564 edition)

The problems also tell us the cost of the luxuries and the necessities of life. Spanish linen was worth, for example, from 94 to 120 ducats per hundredweight, while Italian linen ran as high as 355 ducats and Saloniki linen as high as 380 ducats. French linen was much cheaper than the latter, selling for 140 ducats. The arithmetics tell us that the linen was baled and sent from Venice to towns like Brescia on muleback.

The problems "delle pigione" tell us that the houses of the bourgeoisie rented in Siena, in 1540, at about 25 to 30 lire per year, while a century later they rented in Florence for from 120 to 300 lire. We also have the prices of sugar, ginger, pepper, and other commodities, showing that these three, for example, were only within the reach of the wealthy.

Hotel life in a grand establishment is also revealed in various problems, of which this one, printed in 1561, is a fair type:

Item/Wenn in einem Gasthause weren 8 Kamern/in jglicher Kamer stünden 12 Bette/in jglichem Bette legen 3 Geste/vnd ein jglicher Gast gebe dem Hausgesinde 6 ß trinckgelt/Wie viel thuts in einer Summa?

That these conditions of 12 beds in a room and 3 guests in a bed are not exaggerated, many travelers in remote parts of the world today can testify.

Partnership. There are three historic stages in the conduct of mercantile business: (1) that of individual enterprise, (2) that of partnership,[1] and (3) that of corporations.[2] The first of these has always existed, but in extensive business affairs it early gave way to partnerships in which the profits were divided according to the money invested, the time that it was employed in the business, or both. As business operations became still more extensive the partnership generally gave place to the corporation. Although the corporation has only recently come into great prominence, there were *societates publicanorum*[3] in Rome, each directed by the *magister societatis* and made up of members who received shares of the profits in proportion to their investments. These societies were not formed for the conduct of general business, however, but only for collecting taxes for the censors.[4] The division of profits according to amounts invested goes back to the Babylonian merchants and is frequently mentioned in ancient records.[5]

Partnerships and Usury. Aside from the necessity of joining capital in large business enterprises there was another reason

[1] "Partner" is from the Latin *partionarius*, from *partitio*, a share or part. It comes through the Old French *parsonnier* and Middle English *parcener*.

[2] Latin *corporatio*, from *corpus*, a body. Compare "corporeal," "corps," "corse," and "corpse."

[3] That is, societies of the farmers-general of the revenues.

[4] From *censere*, to value or tax, whence our "census."

[5] A. H. Sayce, *Social Life among the Assyrians and Babylonians*, p. 63 (London, n. d.).

why partnerships flourished so extensively in the Middle Ages. The laws and the popular prejudice in Christendom against taking interest on money placed the "pope's merchants"[1] at a disadvantage with respect to the Jews. Merchants in need of money were generally helped by their guilds, ordinary borrowing being resorted to only in cases of emergency, as in the *Merchant of Venice*.[2] Hence, if a man had money lying idle for a time, it was natural that he should join with others in some temporary venture and take his share of the profit. He thus secured interest on his capital without incurring popular odium. A man might even be taken into partnership for a limited time only, or he might be compelled by his partners to withdraw;[3] in these cases it became necessary to divide the profits according to the amount invested and the time.

Various Names for Partnership. There is hardly a medieval writer on business arithmetic who does not give this subject an important place,[4] and nearly every printed commercial book for a period of four hundred years devoted a chapter to the topic. The Latin arithmeticians called it the *Regula de societate*[5] or *Regula consortii*,[6] while the Italian writers commonly used the plural term *compagnie*.[7] When the services of a

[1] W. Cunningham, *The Growth of English Industry and Commerce during the Early and Middle Ages*, pp. 329, 364 (London, 1896).

[2] The Christian laws had forced the business of money-lending into the hands of the Jews, as in the case of Shylock.

[3] An interesting case is told in the records of the famous business house of Kress, in Nürnberg: "Und do die rechnung geschah, do zalt man Paulus Forchtel sein gelt und wolten sein nit länger in unser gesellschaft haben." The records also relate: "Item wir haben gantze rechnung gemacht an sant barbara obent do man zelt von gotes gepurt 1395 yar und es westund [belonged to] yeden I^{c} XXXI gld. zu gewinn." G. von Kress, *Beiträge zur Nürnberger Handelsgeschichte aus den Jahren 1370 bis 1430*. See Günther, *Math. Unterrichts*, p. 291 n.

[4] *E.g.*, Fibonacci (1202), *Liber Abaci*, I, 139; Johannes Hispalensis (*c.* 1140), *Liber algorismi* (No. II of Boncompagni's *Trattati*), p. 111; and many others.

[5] Thus Huswirt (1501): "Regula de societate mercatorum et lucro" and "De societate et intercessione tpis" (temporis). Cardan (1539, cap. 52) has "De societatibus."

[6] Thus Gemma Frisius (1540): "Regula consortij, siue, vt dicunt, Societatis."

[7] Thus Feliciano (1526) has a chapter *Dele compagnie* (1545 ed., fol. 30). In Spanish the word appears as *compañia* and in French as *compagnie*, but the word *société* was also used, as by Peletier (1549).

partner were considered instead of any money contribution that he might have made, they used the term *soccite*.[1]

English writers, following the Italian practice, often used the word "company,"[2] although in general the word "fellowship"[3] was preferred.

538 Two marchants made a companie, A put in 300 pound for 2 monethes, and then putteth yet in 100 pound, and 6 monethes after that taketh out 200 poũd, and with the reſt remaineth vntill the yeares end. B put in 100 pound for one moneth, and then putteth yet in 700 pound, and 6 monethes after that taketh out a certaine ſumme of money, and with the reſt remaineth vntill the yeares end. and then finde to haue gained together 400 poũd, whereof B muſt haue 80 pound more then A, the queſtion is how much money B tooke out of the companie, without reckoning intereſt vpon intereſt.

300 . . 2 . . 600
100
400 . . 6 . . 2400
200
200 . . 4 . . 800
160 . . 3800 . . 240
19
5700
100 . . 1 . . 100
700
800 . . 6 . . 4800 *Facit* B tooke out 640 pound.
4900
800
160

400
80
320
160

If

PROBLEM IN PARTNERSHIP

From Masterson's *Arithmetike* (1592)

[1] Thus Cataneo (1546) follows his chapter *Delle Compagnie* by one *Delle Soccite*, saying: "lequali son simili alle compagnie," but that the latter consider "il capitale e non la persona & l'altro mette solo la persona senza altro capitale." Practically, he says, these problems have to do with the case in which some "gentil' huomo" puts in his cattle and some "uillano o soccio minore" puts in his time. The 16th century books also use the form *soccide*. The modern form is *soccio, soccita*.

[2] "Two men Company, and make a Stock of 700 l," in Hodder's *Arithmetick*, 10th ed., p. 152 (1672); but he calls the subject "The Rule of Fellowship."

[3] Thus Recorde (*c.* 1542) speaks of "the rule of Fellowshyppe . . . or Company" (1558 ed., fol. N 1), and Baker (1568) gives "the rule of Felowship." The term is used by the American Greenwood (1729) and in Pike's well-known arithmetic.

The Germans ordinarily preferred the term *Gesellschaft*[1] and the Dutch writers followed their lead.[2]

Pasturage Problems. Akin to partnership problems, and often classified with them, are pasturage problems.[3] These may have begun with the custom of the Roman *publicani* of renting to stock owners sections of the estates which the government had farmed out to them, payment being made in proportion to the number of cattle.[4] It is probable that the early use of commons by the shepherds was regulated according to the principles inherited from the early Roman conquerors.

The importance attached to the subject in the 16th and 17th centuries may be inferred from the fact that Clavius (1583) devotes thirty-two pages to it and Coutereels (1599) allows forty pages.[5]

Profit and Loss. The expression "profit and loss," still found in our arithmetics, although not always used in commercial parlance in quite the same sense, is an old Italian one. The books written in the vernacular used the term *guadagni e perdite*,[6] while those written in Latin called their chapter on the subject *De lucris & damnis*.[7] The term passed over into German as *Gewin und Verlust*,[8] into Dutch as *Winst ende*

[1] Thus Köbel (1514) has *Gesellschaft der Kaufleit*, and Albert (1534) has *Gesellschaft / oder der Kauffleut Regel / von eigelegtem Gelde.* Suevus (1593) gives the Latin form also, *Regvla societatis. Regel der Gesellschaft.*

[2] Thus Van der Schuere (1600) has *Reghel van Gheselschap.*

[3] Cardan (1539) speaks of them under the head *De societatibus bestiarum*, and Ortega (1512, the Rome edition of 1515) speaks of *compagnia pecoraria.* In the Dutch books of the 17th century the subject commonly went under the name *Vee-Weydinghe.*

[4] Ramsay and Lanciani, *Manual of Roman Antiquities*, 17th ed., p. 548 (London, 1901).

[5] Similarly, Pagani (1591), twenty-four pages; Werner (1561), twenty-six pages; Van der Schuere (1600), twenty-six pages; and Cardinael (1674 ed.), twenty-five pages.

[6] "Gains and losses," as in Sfortunati (1534), Cataneo (1546), and Pagani (1591).

[7] As in Cardan, 1539, cap. 59.

[8] Thus Rudolff (1526) gives an "Exẽpel von gewin vñ verlust," Riese (1522) has "Vom gewin vnd vorlust," and Köbel (1514) has "Regel vnd frag / von gewlñ der Kauffleut angekaufftet wahr / Regula Lucri."

verlies,[1] and into French as *gain & perte*.[2] The English writers used "loss and gain," Recorde (*c.* 1542) saying that "the fourth Chapter treateth of Losse and Gaine, in the trade of Marchandise."

The early American texts followed the English phraseology, speaking of "loss and gain." Thus Greenwood (1729) remarks that "the Intention of this *Rule* is, to discover what is *Lost*, or *Gained per Cent.* in the *Sale* or *Purchase* of any Quantity of *Goods*: in Order to raise, or fall the *Price* thereof accordingly."

The popularity of the subject in the 16th century may be inferred from the fact that Werner's *Rechenbuch* (1561) devotes forty-seven pages to it, and that other commercial arithmetics were similarly generous.

Commission and Brokerage. Although the subject of commission and brokerage is not new, these terms are relatively modern. The early printed books use such terms as "factorage"[3] and "factorie,"[4] from "factor,"[5] a middleman in the purchase and sale of products. The term "factor" was used in this sense in the Middle Ages, when the father of Fibonacci was (*c.* 1175) a factor in Bougie, and in the Renaissance period.[6] It is still used, although less commonly, in America and Great Britain, and warehouses for goods to be exported are still called "factories" in various parts of the world.

[1] As in Van der Schuere (1600). It also appears as *Winningh en verlies*, as in Bartjens (1633).

[2] So the Dutch-French work of Coutereels (1631) has "Comptes de gain & perte."

[3] Even as late as the 19th century Pike's arithmetic (8th ed., p. 204 (New York, 1816)) has the definition: "Factorage, Is an allowance of so much per cent. to a Factor or Correspondent, for buying and selling goods." It defines a broker as a merchant's assistant in buying or selling.

[4] Thus Rudolff (1526) and Werner (1561) have *Factorey*. Of the Dutch writers, Bartjens (1633) has *Factorie*; Raets (1580), *Rekeninghen van Facteurijen*; and Van der Schuere (1600), *Facteur-Rekeninghe*.

[5] That is, operator, from the Latin *facere*, to act or do. Compare the factor of a number.

[6] Thus Werner (1561): "Item ein Kauffman macht seinem Factor ein geding"; Trenchant (1566): "Aux compagnies d'entre marchans & facteurs"; Recorde (*c.* 1542): "A Merchant doth put in 800 pound into the hands of his Factor" (1646 ed., p. 519).

The word "broker" is not so common as "factor" in problems before the 19th century, although it appears in Middle English[1] to designate one who does business for another or acts as his agent.[2]

The term "commission," as now used to indicate a percentage, is relatively modern.

Equation of Payments. The absence of banking facilities to the extent now known in America, the difficulties in transmitting money, and the scarcity of currency before the great improvements in gold-mining in the 19th century rendered necessary until very recently an extensive credit system. Importing houses bought on credit and exported goods on credit to those from whom they bought, balancing their accounts from time to time. The process of finding the balance due, so that neither party should lose any interest, was the problem of the equation of payments. The subject is found in many manuscripts of the 14th and 15th centuries, and when textbooks began to appear in print it was looked upon as of great importance. Thus Recorde (*c.* 1542) says:

> Rules of Payment, which is a right necessarie Rule, and one of the chiefest handmaydes that attendeth vpon buying and selling.[3]

The subject went by various names,[4] but the later English and American writers generally used the expression "Equation of Payments."

Interest. The taking of interest is a very old custom, going back long before the invention of coins, to the period in which values were expressed by the weight of metal or by the quantity of produce. The custom of paying interest was well known

[1] *Brocour* or *broker.* It probably came from the Anglo-Saxon *brūcan*, to use or employ. The root is found in the Scandinavian and Teutonic languages, referring to business in general.

[2] A word coined in the 16th century from the Latin *agere*, to act or to do.

[3] Mellis ed., p. 478 (1594).

[4] Thus Hodder (1672 ed., p. 163) calls the chapter "Of Equation," and the Dutch works have such names as "Den Regel van Paeyement of Betalinghe" (Eversdyck's Coutereels, 1658 ed., p. 181) or simply "Reghel van Payementen" (Stockmans, 1609 ed., fol. Q 4; Houck, 1676 ed., p. 108).

in ancient Babylon. In Sumerian tablets of the period before 2000 B.C. the rate is often given as varying from the equivalent of 20 per cent to that of 30 per cent, according to whether it was paid on money, that is, on precious metals, or on produce. In general, in the later Babylonian records, the rate ran from $5\frac{1}{2}$ per cent to 20 per cent on money and from 20 per cent to $33\frac{1}{3}$ per cent on produce, although not expressed in per cents.[1] Even princes engaged in trade and insisted upon their interest, for one of the tablets relates the following:

> Twenty manehs of silver, the price of wool, the property of Belshazzar, the son of the king. . . . All the property of Nadin-Merodach in town and country shall be the security of Belshazzar, the son of the king, until Belshazzar shall receive in full the money as well as the interest upon it.[2]

Tablets of Nineveh as old as the 7th century B.C. have the following records:

> The interest [may be computed] by the year.
> The interest may be computed by the month.
>
> The interest on ten drachmas is two drachmas.
>
> Four manehs of silver . . . produce five drachmas of silver per month.[3]

Interest in Ancient India. The custom was also known in ancient India, appearing in the early legal writings of the *Śūtra* period, some centuries before the beginning of our era.[4] In the

[1] M. Jastrow, Jr., *The Civilization of Babylonia and Assyria*, pp. 323, 326, 338 (Philadelphia, 1915); A. H. Sayce, *Zeitschrift für Assyriologie*, V (1890), 276; T. G. Pinches, *ibid.*, I, 198, 202; A. H. Sayce, *Social Life among the Assyrians and Babylonians*, chap. v, p. 67 (London, n. d.) (hereafter referred to as Sayce, *Social Life*); G. Billeter, *Geschichte des Zinsfusses im griechisch-röm. Altertum* (Leipzig, 1898),—the leading authority. [2] Sayce, *Social Life*, p. 65.

[3] Since four manehs was about $180, and five drachmas was about $2, the interest on $180 was $24 a year, the rate being 13 per cent. See J. Menant, *La Bibliothèque du palais de Ninive*, p. 71 (Paris, 1880).

[4] Thus the *Dharma-sāstras* state that "5 *Māshās* for every 20 [*Kārshāpanas*] may be taken every month." Since 20 *māshās* were probably equal to a *kārshāpana*, the rate was $1\frac{1}{4}$ per cent per month, or 15 per cent annually. See R. C. Dutt, *A History of Civilization in Ancient India*, I, 174, 237 (London, 1893).

medieval period there are many evidences of the taking of interest. For example, Mahāvīra (*c.* 850) has various problems of the following type:

O friend, mention, after calculating the time, by what time 28 will obtain as interest on 80, lent out at the rate of $3\frac{1}{2}$ per cent [per month].

Bhāskara (*c.* 1150) also paid much attention to the subject, giving such problems as the following:

If the interest of a hundred[1] for a month be five, say what is the interest of sixteen for a year.

If the interest of a hundred for a month and one third be five and one fifth, say what is the interest of sixty-two and a half for three months and one fifth.

If the principal sum, with interest at the rate of five on the hundred by the month, amount in a year to one thousand, tell the principal and interest respectively.[2]

Interest Customs in Greece. The rate in Greece seems not to have been restricted by law and to have varied from 12 per cent to 18 per cent. In the time of Demosthenes 12 per cent was thought to be low. There were two general plans for computing interest: (1) at so much per month per mina, and (2) at such a part of the principal per year. Interest was usually paid at the end of each month.[3]

Interest in Rome. In Rome the rate of interest was at first unrestricted.[4] The Twelve Tables[5] limited the interest charged

[1] That is, the rate per cent.

[2] These extracts show that the rate of interest in India in the 12th century was about 60 per cent, and that interest was computed on a percentage basis. See Colebrooke's translation of the *Līlāvati*, pp. 36, 39.

[3] F. B. Jevons, *A Manual of Greek Antiquities*, p. 397 (London, 1895); Harper's *Dict. of Class. Lit. and Antiq.*, p. 665.

Interest was called *faenus*, or *fenus*, a later term being *usura* (from *uti*, to use), commonly expressed in the plural, *usurae*. So Cicero has "pecuniam pro usuris auferre." From this came the French *usure* and our "usury." Capital was *caput* (head, originally a head of cattle) or occasionally *sors* (lot or chance).

[5] *Duodecim Tabulae*, the first code of Roman law, 451–449 B.C., and the foundation of that law up to the time of the *Corpus Iuris* of Justinian, *c.* 530.

to Romans to one twelfth ($8\frac{1}{3}$ per cent) of the capital, and later (*c.* 100 B.C.) this limitation was extended to aliens as well. The *Lex Genucia* (342 B.C.) prohibited the taking of interest altogether,[1] but like the medieval canon law this seems not to have been enforced.

In later Roman times the Eastern custom of monthly interest came into use, the ordinary rate being 1 per cent per month, payable in advance, or 12 per cent per year. In Cicero's time 48 per cent per year was allowed, and under the first emperors 25 per cent was common. A little later 12 per cent per year was made the maximum; Justinian reduced this rate to $\frac{1}{2}$ per cent per month, which gave rise to the common rate of 6 per cent. In classical Latin works the rates of interest are usually mentioned either as *fenus unciarum*[2] or as *usurae centesimae*.[3]

Interest in the Middle Ages. In medieval Europe the canon law forbade the taking of usury, that is, the payment in advance for the use of money. The time had not come for borrowing money for such remunerative purposes as extensive manufacturing or as building railways and steamships, and so the principal was often consumed by usury instead of being increased. Usury would therefore have speedily resulted in the enslavement of the peasants, who were without money or financial ability. Hence the Church came to recognize a distinction between loans for production, which might reasonably have carried some remuneration, and those for consumption, which were contrary to public policy.[4]

[1] M. Cantor, *Politische Arithmetik*, p. 2 (Leipzig, 1898).

[2] Uncial interest, that is, interest by twelfths, $\frac{1}{12}$ being the common rate. This was $\frac{1}{120}$ per month when the ancient year consisted of ten months. When the year was later divided into twelve months the rate was still $\frac{1}{120}$ per month or $\frac{12}{120}$ per year. Since interest was paid by the month, this made the former rate $8\frac{1}{3}$ per cent and the latter 10 per cent per year. See Ramsay and Lanciani, *Manual of Roman Antiquities*, 17th ed., p. 472 (London, 1901).

[3] Hundredth interest, or 1 per cent a month. This was the ancient "per cent." If the security was poor, this was raised to *binae centesimae* (2 per cent per month) or even to *quaternae centesimae* (4 per cent per month).

[4] See the decree of the fifth Lateran Council (1512–1517) in Janet, *Le capital, la spéculation, et la finance, au XIX^e siècle*, p. 81 (Paris, 1892).

There was, however, another reason which was not so openly stated, namely, the desire of the Church and of the ruling classes to prevent the dangerous rivalry to authority which would have resulted from the accumulation of too large fortunes; in other words, to avoid the dangers of capitalism.

Origin of the Term "Interest." To overcome this restriction there accordingly developed a new economic custom. The borrower paid nothing for the use of the money if it was repaid at the time specified. If, however, he failed so to pay the principal, he was held to compensate his creditor by a sum which represented the difference, or "that which is between" ("id quod *interest*") the latter's position because of the delay and what his position would have been had he been paid promptly. *Id quod interest* was recognized by the Roman law, but as a certain per cent agreed to in advance it first appears in the 13th century, possibly suggested from the East.[1] Speaking of this method, Matthew Paris (1253) tells us that in his time 10 per cent was exacted every two months, and adds that in this way unscrupulous men "circumvented the needy in their necessities, cloking their usury under the show of trade."[2]

Among the economic movements of the Renaissance period was a serious questioning of the validity of the canon law against usury and a determination to recognize a new type of interest, namely, usury paid at the end of the term of borrowing.[3] As a result of this feeling the subject of interest found place in many of the early printed books, particularly in Italy,[4]

[1] Compare Bhāskara's *Lilāvati* (*c.* 1150), Colebrooke's translation, p. 39. Fibonacci (1202) gives problems involving 20 per cent interest, but the Hindu works give rates as high as 60 per cent.

[2] *Chronica Majora*, III, 329, published in the *Rolls Series*. See also W. Cunningham, *The Growth of English Industry and Commerce* . . . , p. 329 (London, 1896). One of the best historical sketches to be found in the early arithmetics is given by Sfortunati, *Nvovo Lvme*, 1534 (1544–1545 ed., fol. 60).

[3] See such works as F. de Platea, *Opus restitutionum usurarum et excommunicationum*, Venice, *c.* 1472 (de Platea lived *c.* 1300); J. Nider, *Tractatus de contractibus mercatorum* (s. l. a., but Cologne, with at least seven editions before 1501); and many other similar works of the period.

[4] So Calandri (1491), who uses thirty days to the month, sometimes using per cent ("per 3 anni a 10 per cento lanno") and sometimes stating the rate as the equivalent of so many pence in the pound.

although sometimes against the protest of the author.[1] This spirit of protest showed itself in the people's literature of England, as in Francis Thynne's (16th century) epigram:

> Stukelie the vsurer is dead, and bid vs all farwell,
> who hath a Iourney for to ride vnto the court of hell.

A similar testimony is found in Lauder (1568):

> Credit and frist [delay] is quyte away,
> No thing is let but for Usure;
> For euerie penny thay wyll haue tway:
> How long, Lord, will this warld indure?[2]

In spite of these protests the English parliament in 1545 sanctioned the taking of interest,[3] fixing the maximum rate at 10 per cent. The protest was such that the law was repealed in 1552, but it was reënacted in 1571,[4] and since that time all works on commercial mathematics have included the topic.

In Germany the opposition to interest was also very strong, and Martin Luther published a sermon on the subject in 1519.[5]

Compound Interest. The compounding of interest was known to the Romans and was not forbidden until rather late.[6] The late medieval and the Renaissance Italians, from whom we derive so much of our modern business arithmetic, used the word *merito*[7] for interest in general, and where it was computed "simply by the year"[8] it was called simple interest.[9]

[1] So Cataneo (*Le Pratiche*, 1546), under the title *De semplici meriti vsvreschi*, speaks of the practice as often "diabolical," and Pagani (1591) calls it "Cosa in vero molto biasmeuole, & diabolica."

[2] W. Lauder, *The Lamentationn of the Pure, twiching the Miserabill Estait of this present World*, published by the Early Eng. Text Soc., p. 28 (London, 1870).

[3] 37 Hen. VIII, c. 9.

[4] E. P. Cheyney, *Industrial and Social History of England*, p. 172 (New York, 1901).

[5] *Eyn Sermon von den Wucher*, Wittenberg, 1519.

[6] Harper's *Dict. of Class. Lit. and Antiq.*, p. 665.

[7] It passed into the French as *merite*, although the word *interest* was also used. Thus Trenchant (1566): "A calculer les merites ou interestz" (1578 ed., p. 299). The Italians also used the term *usura*. The 1515 Italian edition of Ortega (1512) has *Regula de lucro*.

[8] "Simplicemente all' anno," as Tartaglia (1556) says (1592 ed., II, fol. 95). See also Cardan's *Practica*, 1539, capp. 57 and 58.

[9] Ciacchi (1675, pp. 80, 228), a later writer, speaks *De' meriti semplici*.

Compound interest among the early Italians was computed from the beginning of each year[1] or period[2] and was called by the English writers of the 17th century "interest upon interest."[3] The taking of such interest was frequently charged against the Jews,[4] although unjustly so,[5] and is even characterized by their name.[6]

In the 15th and 16th centuries interest was usually computed either on a percentage basis or at so many pence to the pound.[7] The rate varied from the 60 per cent mentioned by Bhāskara (*c.* 1150) and his European contemporaries to smaller limits.[8]

The difficulty in computing interest gave rise in the 16th century to the use of tables. These were extended in the 17th century, a table of compound interest appearing in Richard Witt's *Arithmeticall questions* (London, 1613).

Discount. The computing of discount[9] for the payment of money due at a future time is relatively modern. It is found

[1] "À capo d' anno," as Tartaglia (1556) describes it (1592 ed., II, fol. 95).

[2] "À capo d' alcun tempo," as Cataneo (1546) describes it (1567 ed., fol. 53). Similarly Tartaglia (1592 ed., II, fol. 119): "Del meritar à capo d' anno, ò altro termine che d' alcuni è detto vsura." The expression passed over into French as "merite à chef de terme" (Trenchant, 1566; 1578 ed., p. 299).

[3] So in Hodder, 10th ed., 1672, p. 139. The Dutch commonly called it "interest op interest" or "Wins-ghewin (VVinsts-Gewin)."

[4] Thus Pagani (1591, p. 147): "e questo modo di meritare e communemente vsitato da gl' hebrei ne suoi Banchi."

[5] Günther, *Math. Unterrichts*, p. 290. Pagani also says that the Christians were equally to blame.

[6] "Ma d' altra sorte è la ragion dell' usura, che chiamano Guidaica" (*sic*, for "Giudaica"), as in the Italian edition (1567, fol. 32) of Gemma Frisius, but not in the Latin edition. Similarly Van der Schuere (1600, fol. 127) speaks of "een Ioodtsch profijt."

[7] "Meriter est bailler ses deniers pour profiter à raison d'vn tant pour ℔ ou pour 100: par an" (Trenchant, 1566; 1578 ed., p. 298).

[8] For example, Sfortunati (1534) gives the rate "a denari .2. la libra il mese," which is 10 per cent a year, and goes even as high as 4 pence per pound per month, or 20 per cent a year; Trenchant (1566) gives one problem at 12 per cent; Tartaglia (1556) gives 10 per cent, 16 per cent, 20 per cent, and other rates; one of the Dutch writers, Raets (1580) gives from 8 per cent to 14 per cent, 10 per cent being stated thus: "Soo 100 winnen in een iaer 10"; and Cardinael (1674 ed.) gives rates ranging from 10 per cent to 20 per cent.

[9] Formerly "discompt," from the Old French *descompter*, to reckon off, from *des-*, away, *compter*, to count.

in some of the 16th century arithmetics[1] but is more common in the century following, appearing under various names.[2]

Assize of Bread. One of the standard problems of the 16th century books related to the variation in size of a loaf of bread as the wheat varied in value. For example, if a 10-cent loaf weighs 14 oz. when wheat is worth $1.80 a bushel, how much should it weigh when wheat is worth $2.20 a bushel?[3]

The problem had its genesis in real conditions. Loaves were formerly of two kinds: (1) "assized bread," always sold at the same price but varying in weight according to the price of wheat, and (2) "prized bread," always of the same weight but varying in price.[4] The legal regulation for the assized bread goes back at least to 794, being found at that time in a Frankfort capitulary, and is probably of Roman origin. London regulations are found as early as the 12th century, and in the "assize of bread" of Henry II (1154–1189) these are worked out by inverse proportion.[5] As a result of these regulations, tables of the assize of bread were prepared and their use was

[1] Thus Cataneo (1546) has a chapter "Del semplice sconto" and one (corresponding to compound interest) "Dello sconto a capo d' alcvn tempo." Trenchant (1566) discounts an amount due in four years "à raison de 12 pour 100 par an."

[2] Thus in Coutereels's *Cyffer-Boeck* (1690 ed., p. 289) it appears as "Rabatteeren, Disconteeren, of af-korten," and in Hodder's *Arithmetick* (1672 ed., p. 175) as "The Rule of Rebate, or Discount."

[3] Thus Ortega (1512; 1515 ed., fol. 59): "℔ Si de vno misura de grano che costa 10 carlini·mi dano 4 vnze de pane per vno dinaro si voi sapere se de vna altra misura che costera 20 carlini quante vnze ne darano per uno medesimo dinaro." See also Gemma Frisius (1540; 1555 ed., fol. 66), Rudolff (*Kunstliche rechnung*, 1526; 1534 ed., fol. K 4), Albert (1534; 1561 ed., fol. N 1, under *Regula Detri Conuersa*), Suevus (1593, p. 320, with two pages "Vom Brodgewichte in thewren vnd wolfeihlen Jaren"), and many other writers of the 16th century and later.

[4] J. Nasmith, *An examination of statutes . . . the assize of bread*, Wisbech, 1800; S. Baker, *Artachthos Or a New Booke declaring the Assise or Weight of Bread*, London, 1621.

[5] "Quando *quartierium* frumenti se vendit pro sex sol.; tunc debet panis esse bonus et albus et ponderare sexdecim sol. de xx^{ti} lores [*i.e.*, 20 d. to 1 oz.]. . . . Quando pro quatuor solidis tunc debet ponderare triginta sex sol. et alius quadraginta sex sol. . . . ," and so on for different weights. W. Cunningham, *The Growth of English Industry and Commerce during the Early and Middle Ages*, p. 568 (London, 1896).

made obligatory.[1] This problem of the size of loaves was a common one in the early printed books and is often found as late as the second half of the 19th century.[2] The following, from the 1837 edition of Daboll's well-known American arithmetic, illustrates the type: "If when wheat is 7s. 6d. the bushel, the penny loaf will weigh 9 oz. what ought it to weigh when wheat is 6s. per bushel?"

Tare and Tret. Until the middle of the 19th century the subject of "tare and tret" was found in most of the English and American commercial arithmetics. "Tare" meant an allowance of a certain weight or quantity from the weight or quantity of a commodity sold in a box, cask, bag, or the like. The word came from the Arabic *ṭarḥa*[3] (what is thrown away) through the Spanish *tara* and the French *tare*, and shows the commercial influence exerted by the Arabs in Spain.

"Tret" meant about the same thing, but the word shows the Italian influence, meaning originally an allowance on things transported.[4] In England it was an allowance of 4 lb. in every 104 lb.

There was also a third term that was related to "tare" and "tret" and is commonly found in the English books of the 16th century. This term is "cloff," meaning an allowance of 2 lb. made on every 3 cwt. of certain goods in order that the weight might hold out in retailing. Thus Recorde (*c.* 1542) has problems of this type:

"Item at 3$\bar{s}$ 4d the pound weight, what shal 254$\frac{1}{2}$ be worth, in giuing 4 l weight vpon euery 100 for treate."

"Item if 100 l be worth 36$\bar{s}$ 8d, what shall 800 l be worth in rebating 4 pound upon euery 100 for tare and cloffe."[5]

[1] Such a table, from the Record Book of the city of Hull, is reproduced in facsimile in E. P. Cheyney, *Industrial and Social History of England*, p. 67 (New York, 1901).

[2] A rare and interesting tract on the subject is that of J. Powel, *Assize of Bread*, London, 1615, a guide for those who had to interpret the old law.

[3] From *ṭaraḥa*, he threw down.

[4] The word is from the Latin *trahere*, to draw or pull, whence *tractus*, Italian *tratto*, and French *trait*. From the same root we have "tract" and "traction."

[5] *Ground of Artes*, 1594 ed., p. 487. The origin of the term is uncertain.

In Baker's arithmetic (1568) "the eyght chapter treateth of Tares and allowances of Marchandise solde by weight," and other arithmetics of the period also presented the subject at considerable length.[1]

Cutting of Cloth. Problems relating to the cutting of cloth correlated so closely with the needs of merchants that the commercial schools seem generally to have included them in the 16th century. Thus Grammateus (1518) has problems on the cutting of cloth by tailors,[2] and Tartaglia (1556) also devoted considerable attention to them.[3] No attention was paid to the pattern, and the problems show that drapers had flexible consciences with respect to advising as to the amount needed for a garment.[4] Baker (1568) says that "the 5 Chapter treateth of lengthes and bredthes of Tapistrie, & other clothes,"[5] and John Mellis has a similar chapter in his addition (1582) to Recorde's *Ground of Artes*.

The custom of carpeting rooms, which reached its highest point in the 19th century, led to the inclusion of problems relating to this subject. The return to rugs in the 20th century is leading to a gradual elimination of the topic in America.

Barter. Of the applications of arithmetic none has had a more interesting history than barter,[6] a subject now very nearly obsolete in textbooks, although temporarily revived among nations as a result of the World War of 1914–1918, owing to conditions of exchange. There are three fairly well defined periods in the exchange of products. The first is that of pure barter, seen today in the exchange of guns and ammunition for a tusk of ivory in remote parts of Africa,—a period lasting throughout the era of savage life. This is also seen in the ancient method of paying taxes "in kind," so many fowls out

[1] Thus Ortega (1512; 1515 ed., fol. 53), "Regvla de tre de tara"; Stockmans (1589), "Reghel van Tara"; Ciacchi (1675), "Delle tare a vn tanto per cento"; Coutereels (1690 ed.), "Tara-Rekeningh."

[2] "Schneider regel" (1535 ed., fol. C6).

[3] 1592 ed., II, fol. 79. [4] Tartaglia, *loc. cit.*, fol. 81. [5] 1580 ed., fol. 126.

[6] Possibly from the Old French *barat, barate, barete*, whence *bareter*, to cheat or beguile. It appears in Italian as *baratto* (Ortega, 1512; 1515 Italian ed., fol. 78) or *baratti* (Feliciano da Lazesio, 1536).

of a dozen, or one cow out of a given number.[1] The second is that wherein a fixed value was assigned to certain products, such as grain or dates,[2] these products acting as media of exchange or as bases for determining values in bartering other products,—a period lasting until money was invented and indeed until currency became common. The third period is that of the adoption of money as a medium of exchange, this medium taking such forms as wampum, shells, coins, ingots, and government certificates.

Two influences perpetuated barter long after the first of these periods and indeed down to the present time, namely, the scarcity of currency[3] and the international fairs. In these fairs[4] the merchants found that barter was a necessity on account of the scarcity and diversity of money.[5]

Various Names for Barter. Barter also went by the name of "exchange," quite as we use the word "trade" at present. Thus an English writer of 1440 has the expression "Bartyrn or changyn or chafare[6] oone thynge for a othere, *cambio*,"[7] and

[1] For example, in Egypt. See H. Maspero, *Les Finances de l'Égypte sous les Lagides*, p. 29 (Paris, 1905).

[2] A. H. Sayce, *Social Life among the Assyrians and Babylonians*, chap. v (London, n. d.); W. Cunningham, *The Growth of English Industry and Commerce*, p. 114 (London, 1896).

[3] Ciacchi mentions this effect in his *Regole generali d'abbaco*, p. 114 (Florence 1675). It should be observed that the output of gold from 1850 to 1900 was greater than that of the preceding three hundred and fifty years, which accounts in part for the greater amount of currency now available.

[4] Compare the fair of Nijni Novgorod and the smaller fairs of Leipzig, Munich, and Lyons, all of which still continue, and the various international expositions which are modern relics of the ancient gatherings of merchants. In a MS. on arithmetic, written in Italy in 1684, nine pages are given to a list of great fairs, mostly European, which Italian merchants of that time were in the habit of attending.

[5] Thus Cataneo (1546), speaking "De baratti," says: "E Necessario al buon mercante non uolendo receuer danno esser molto experto nel barattere" (1567 ed., fol. 49).

[6] Middle English *chaffare*, *chepefare*, from the Anglo-Saxon *cēapian* (to buy) + *fare* (to go). There is the same root in the word "cheap," originally a bargain, and in "Cheapside," the well-known London street.

[7] Italian for "exchange." So Ghaligai (1521) speaks of "*Barattare*, ouer cambiare una Mercantia a un' altra"; and Pellos (1492) has "lo .xiij. capitol qui ensenha cābiar aut baratar vna causa per lautra."

Baker (1568) gives fifteen pages to his chapter which "treateth of the Rules of Barter: that is to say to change ware for ware." The early German writers had a similar usage.[1]

The French writers of the 16th century often used the interesting word *troquer*,[2] a word meaning to barter, the chapter being called *Des Troques*. From this word we have the English "truck," the material bartered, a word which came to mean the most common objects of exchange, such as garden truck, and the cart (truck) in which the dealer (truckman) carried it, and finally to mean worthless material in general.

Since merchandise was often bartered for future delivery, as in the case of goods from Damascus or China,[3] the question of interest, or its equivalent, often had to be considered. This gave rise to the distinction between barter without time and barter with time.[4]

Barter in America. In American colonial life the subject of barter played an important part. A diary of 1704, kept by one Madam Knight of Boston, gives an idea of the arithmetic involved, as the following extract will show:

> They give the title of merchant to every trader; who Rate their Goods according to the time and spetia they pay in: viz. Pay, mony, Pay as mony, and trusting. Pay is Grain, Pork, Beef, &c. at the prices sett by the General Court that Year; mony is pieces of Eight, Ryalls, or Boston or Bay shillings (as they call them,) or Good hard money, as sometimes silver coin is termed by them; also Wampom, vizt. Indian beads wch serves for change. Pay as mony is provisions, as aforesd one Third cheaper than as the Assembly or Genel Court sets it; and Trust as they and the mercht agree for time.

[1] Using their word *Stich*, meaning exchange. Thus Petzensteiner (1483): "Nu merck hübsch rechnung von stich." Rudolff (1526) tells how "zwen stechen mit einander," a phrase now used with respect to dice, and Albert (1534) relates that "Zween wöllen miteinander stechen."

[2] Thus Trenchant (1566): "Deux marchans veulent troquer leurs marchandises." Compare the Dutch *Mangelinge* (*Manghelinge*, *Mangelingh*) as a synonym for *troques* and *change* in the 16th and 17th century arithmetics.

[3] These two cases are mentioned in a MS. of 1684, written at Ancona, now in the library of Mr. Plimpton.

[4] Thus Ortega (1512; 1515 ed., fol. 78): " . . . baratto . . . p tempo como senza tẽpo."

Now, when the buyer comes to ask for a commodity, sometimes before the merchant answers that he has it, he sais, is Your pay redy? Perhaps the Chap Reply's Yes: what do You pay in? say's the merchant. The buyer having answered, then the price is set; as suppose he wants a sixpenny knife, in pay it is 12d—in pay as money eight pence, and hard money its own price; viz. 6d. It seems a very Intricate way of trade and what Lex Mercatoria had not thought of.

Another diary, kept by one Jeremiah Atwater, a New Haven (Connecticut) merchant, about 1800, had various entries of a similar nature, among which is the following:

To 5 yds Calico at 2s 6d per yard.
To be paid in turnips at 1s 6d and remainder in shoes. As far as the turnips pay, the calico is to be 2s 6d and the remainder toward shoes at 2s 8d.

Taxes. Of all the applications of arithmetic, taxation is one of the oldest. The tax collector is mentioned in the ancient papyri of Egypt,[1] in the records of Babylon,[2] in the Bible,[3] and, indeed, in the histories of all peoples. His methods reached the extreme of cruelty among the Saracens, and a decided trace of this cruelty is still seen among some of their descendants.

In Greece the tax[4] was levied directly on property or indirectly by tolls or customs. Resident aliens paid a poll[5] tax, and an indirect tax[6] of 2 per cent was levied at the customhouses.

Rome had an elaborate system of taxation,[7] and this was the source of our present systems. It included the tariff,[8] the

[1] See Volume I, page 45.
[2] Sayce, *Social Life*, p. 68.
[3] *E.g.*, 2 Kings, xxiii, 35; Luke, ii, 1.
[4] Τέλος (*tel'os*).
[5] Head, from the Danish *bol*, a ball, bowl, bulb, or head. Hence the "polls," where the heads of the electors are counted. The Greek term was μετοίκιον (*metoi'kion*).
[6] The πεντηκοστή (*pentecoste'*), fiftieth. This is the same word as our "pentecost," which refers to the fiftieth day after the Passover in the Jewish calendar. In Greek taxation the word referred to the tax of $\frac{1}{50}$, or 2 per cent, on exports and imports.
[7] Latin *taxatio*, from *taxare*, to estimate or evaluate.
[8] Spanish *tarifa*, a price list or book of rates; from the Arabic *ta'rif*, giving information, from the root *'arf*, knowing. The word shows the Arab influence, through the Spanish, upon modern business. The Spanish town Tariffa was

ground tax,[1] the poll tax,[2] the tithes (still familiar in certain parts of the world),[3] and, in later times, the tax on traders.[4]

It is a curious and interesting fact that the subject of taxation commanded but little attention in the early textbooks. It is possible that authors hesitated to touch upon such a sensitive spot because of the necessity for receiving an *imprimatur* from the taxing powers. Although it is occasionally found in the 16th century,[5] it was not common until textbook writers were more free in their offering.

Banking and Exchange. In the days when Europe was made up of a large number of small principalities, each with its own system of coinage, the subject of exchange was much more familiar to the average business man than it is today. How recently this was the case may be seen in a remark of Metternich's in 1845, that Italy "represents simply a group of independent states united under the same geographic term." The European traveler gets some idea of the early situation today, for at the railway stations on the borders he finds the exchange office,[6] where he may exchange the money of the country he is leaving for that of the country he is entering.

So important was the subject considered that the 1594 edition of Recorde's *Ground of Artes* devotes twenty-one pages to it, saying that it is of great value to the merchants dealing with Lyons, inasmuch as "there are 4 faires in a yeere, at which they do commonly exchange."[7]

named from the fact that it was the leading customhouse at one time. The Latin term was *portorium*, from *portare*, to carry, as in "import," "export," and "transport."

[1] *Tributum soli*, tribute of the land, *tributum* coming from *tribuere*, to bestow or pay. [2] *Tributum capitis*. [3] *Decimae*.

[4] *Collatio lustralis*. This was especially prominent in the 5th century, when the great social upheaval led to the aggrandizement of the aristocracy. See S. Dill, *Roman Society in the Last Century under the Western Empire*, p. 204 (London, 1898).

[5] As in Savonne (1563; 1571 ed., fol. 41), with the name "Reigles des impositions."

[6] *Bureau des changes, Wechselbureau, Wissel Bureau, Cambio.*

[7] The editor, Mellis, gives a long list of the leading fairs which an English merchant might attend.

Chain Rule. In the days when the value of coins varied greatly from city to city as well as from country to country, money changers employed a rule, probably of Eastern origin, which was known by various names[1] but was most commonly



RENAISSANCE DEALER IN EXCHANGE

From the 1500 edition of Widman's arithmetic (1489)


called the chain rule[2] or continued proportion. The following problem is adapted from Widman (1489) and illustrates the type and the solution:

A man went to a money changer in Vienna with 30 Nürnberg pence and asked that they be exchanged for Viennese money. Since the money changer was ignorant of their value, he proceeded thus: 7 pence of Vienna are worth 9 pence of Linz, and 8 of Linz are worth 11 of Passau, and 12 of Passau are worth

[1] For example, *Regula conjuncta, Regel conjoinct, Te Zamengevoegden Regel, Regel van Vergelykinge*, and *De Gemenghde Regel*, these terms being taken from various Dutch and Dutch-French books of the 17th and 18th centuries.

[2] *Den Kettingh-Regel, Den Ketting Reegel*, in the early Dutch books. "Gleichsam wie die Glieder einer 'Kette,'" as R. Just, a modern German writer, has it in his *Kaufmännisches Rechnen*, I, 81 (Leipzig, 1901).

13 of Wilsshof, and 15 of Wilsshof are worth 10 of Regensburg, and 8 of Regensburg are worth 18 of Neumarkt, and 5 of Neumarkt are worth 4 of Nürnberg. Then



so that $$\frac{7 \cdot 8 \cdot 12 \cdot 15 \cdot 8 \cdot 5 \cdot 30}{9 \cdot 11 \cdot 13 \cdot 10 \cdot 18 \cdot 4} = 13\tfrac{23}{29},$$

the value of the Nürnberg money in pence of Vienna.[1]

Like many Eastern problems it is found in the works of Fibonacci (1202),[2] and thereafter it was common until the latter part of the 19th century.

Early Banks. The early banks were established in places of greatest relative safety, and these were usually the temples. All kinds of valuables were thus protected from the depredations of thieves, both private and governmental, civil and military. This is seen, for example, in the great business interests carried on within the precincts of the temples in the Ur Dynasty (*c.* 2450–2330 B.C.) of Babylon.[3] On the tablets of this period may be found the records of loans, receipts, promissory notes, leases, mortgages, taxes, and other commercial activities. A little later, in the first millennium B.C., drafts appear[4] in quite the form used even today.

For the reason above stated, the priests in the Greek temples were frequently money lenders.[5] It was also on this account

[1] The rule closes: "Vñ multiplicir in kreucz durchauss auff 2 teyl vñ dividir" (fol. 152).

[2] "De baractis monetarum cum plures monete inter similes" (*Liber Abaci*, p. 126).

[3] M. Jastrow, Jr., *The Civilization of Babylonia and Assyria*, p. 318 (Philadelphia, 1915).

[4] A. T. Clay, *Babylonian Records in the Library of J. Pierpont Morgan*, Part I (New York, 1912).

[5] J. P. Mahaffy, *Old Greek Life*, p. 38 (London, 1885); hereafter referred to as Mahaffy, *Greek Life*.

that coins were struck[1] at the temple of Juno Moneta[2] in Rome.

The later Greek bankers and money changers were called τραπεζῖται,[3] a word derived from τράπεζα,[4] a table, just as "bank" comes from "bench."[5] Their tables were placed in the agoræ (public places) and the finding of "the tables of the money changers"[6] in the Temple at Jerusalem was not at all unusual. Indeed, one may see similar sights in the temples of Southern India today, or in the entrance to the great pagoda at Rangoon, Burma.

The business of the *trapezitai* included buying foreign money at a discount and selling it at a premium, paying interest on deposits, acting as pawnbrokers, and performing the duties of modern notaries.[7]

Banks in Rome. Bankers are mentioned by Livy (IX, 40, 16) as carrying on business as early as the 4th century B.C. At first a private banker was called an *argentarius* (silver dealer), an officer connected with the mint being a *nummularius*.[8] Somewhat later these terms were used along with *mensarius*[9] and *collectarius*[10] to represent any kind of banker.

[1] In early days they were stamped by the stroke of a hammer, and the word has remained in use.

[2] Juno the Admonisher (Adviser, Instructor). On this account the Romans used the word *moneta* to mean money, whence also our word "mint."

[3] *Trapezi'tai*, literally, "tablers." The Hebrew usage is the same.

[4] *Tra'peza*, whence our "trapezium," a figure representing a table (originally with two parallel sides), and "trapezoid," a figure shaped like a trapezium (originally with no parallel sides). The bankers were also called ἀργυραμοιβοί, *argyramoiboi'*, money changers. See Harper's *Dict. Class. Lit.*, p. 1597; Mahaffy, *Greek Life*, p. 38; F. B. Jevons, *Manual of Greek Antiq.*, p. 395 (London, 1895).

[5] Late Latin *bancus*, a bench; French *banc*, a long seat or table. So we have a bench of judges and the bank of a stream. "Banquet," simply a diminutive form, came to mean a feast instead of the table. In Italy *banca* came to mean a tradesman's stall, a counter, and a money changer's table, as well as a bank.

[6] As late as 1567 we find English writers telling of how "Christ overthrew the Exchaungers bankes."

[7] M. S. Koutorga, *Essai historique sur les trapézites ou banquiers d'Athènes*, Paris, 1859.

[8] Coin-man. Originally an officer of the mint who tested the silver before it was coined.

[9] Or *mensularius*, from *mensa*, table, influenced by the Greek name for banker.

[10] Late usage, found in Justinian's *Institutes*.

The exchange bureau was called the *permutatio*.[1] In the banking department the funds of the creditor were called the *depositum*,[2] whence our "deposit." This was subject to a *perscriptio*,[3] a check, quite as it is at present. The *depositum* drew no interest,[4] being like our common open accounts; but there developed also a kind of savings-bank department in which a deposit known as a *creditum*[5] drew interest.[6]

Letters of Credit. The ancient bankers issued letters of credit quite like those issued at present,[7] and also made drafts on one another.[8] The idea is therefore without foundation that it was the Jews who, driven from France to Lombardy in the 7th century, first made use of foreign drafts.[9]

Stockholders. Among the "seven greater arts" recognized in medieval Florence was that of the money changers.[10] As early as 1344 the city government, finding itself unable to pay some $300,000 that it owed, formed a bank[11] and issued shares of stock[12] which were transferable as in modern corporations.[13]

[1] Compare our "permutation." J. Marquardt, *La vie privée des Romains*, French translation, p. 15 (Paris, 1893).

[2] *De* + *ponere*, to place. A *depositum* is that which is placed down.

[3] *Per* + *scribere*, to write; a written order.

[4] It was *vacua pecunia*, unproductive money.

[5] From *credere*, to trust, to have confidence in, to believe, whence our "creed"; in banking business a sum held in trust.

[6] Because of this interest feature, the claims of the *depositarii* were legally preferred to those of the *creditores* in case of the failure of a bank. See the *Digest*, XVI, 3, 7, 2.

[7] Mahaffy, *Greek Life*, p. 38. They were called by the Greeks συστατικαὶ ἐπιστολαί (*systatikai' epistolai'*), letters of introduction.

[8] Cicero uses "permutare Athenas" to mean "to draw on Athens"; and "ab Egnatio solvat" to mean "to pay by draft on Egnatius."

[9] Unger, *Die Methodik*, p. 90.

[10] *Arte del cambio* or *dei cambiatori*. The other six *arti maggiori* were those of (1) dressers of foreign cloth, (2) dealers in wool, (3) judges and notaries, (4) physicians and apothecaries, (5) dealers in silk, (6) furriers. E. G. Gardner, *The Story of Florence*, p. 28 (London, 1900).

[11] *Monte*, mount, bank, money; compare the French *mont-de-piété*.

[12] Originally the word meant a thing that was stuck or fixed, and hence a post, the Anglo-Saxon *stocc*, as in "stockade." The same root is found in "etiquette" (Old French *estiquet*), a little note "stuck up" on the gate of a court. From the same source we have "stack" and "ticket." See also page 194.

[13] C. A. Conant, *History of Modern Banks of Issue*, p. 21 (New York, 1896).

The idea of issuing notes payable in coin but only partly covered by a reserve was a development of the 17th century, beginning in Amsterdam (1609) and developing into more modern form in Stockholm (1661).[1]

Bills of Exchange. Probably the first bill of exchange to appear in a printed work on mathematics is the one given by Pacioli (1494), the form being substantially the same as the one now in use.[2]

Four Kinds of Exchange. In the early printed arithmetics there were four kinds of exchange, of which we have preserved two. The four types were as follows:

1. Common exchange, the mere interchange of coins, the work of the "money changers."[3]

2. Real exchange, by means of drafts.[4]

3. Dry exchange,[5] a method of evading usury laws by means of fictitious bills[6] of exchange,—drafts that bore no fruit.[7]

[1] *Ibid.*, p. 24.

[2] 1494 adi 9 agosto ī v^{a}.

Pagate per questa prima nostra a Lodouico de francesco da fabriano e compagni once cento doro napolitane insu la proxima fiera de fuligni per la valuta daltretanti receuuti qui dal Magnifico homo miser Donato da legge quondā miser Priamo. E ponete $\overline{p}$ noi. Idio da mal ve guardi.

vostro Paganino de paganini da Brescia ss. . . .
Domino Alphano de Alphanis e cōpagni in peroscia. . . .
Sūma, 1494 ed., fol. 167, *v.*

For later examples, see Cardan's *Practica*, 1539, cap. 56; Trenchant's arithmetic of 1566 (1578 ed., p. 350), and other commercial works.

[3] The Italians called it "cambio menuto, ouer commune," as in Tartaglia's work (1592 ed., II, fol. 174). In Spanish it appeared as "cambio por menudo" (Saravia, 1544), and in French as "change menu ou commun" (Trenchant, 1566). It is this form that is referred to in 1335, in the English act of 9 Edw. III, stat. 2, c. 7: "Et que table deschange soit a Dovorri & aillours, ou & q^{i}nt il semblera a nos & notre consail per faire eschange,"—our "bureau des changes."

[4] Saravia (1544) makes two divisions of this type. See his Italian translation of 1561, fols. 108, 110.

[5] Italian, *cambio secco*; French, *change sec*; German, *trockener Wechsel*, as the terms appear in the 16th century.

[6] From the Latin *bulla*, a bubble, a leaden seal that looked like a bubble, and hence the sealed document, like a papal bull. From the same root we have "bullet," "bulletin," "bowl," and "bullion" (a mass of sealed or stamped metal).

[7] So called, as Saravia (Italian translation, 1561) says, from their resemblance to an "albero secco, il quale non ha humore, ne foglie ne frutto."

Such exchange was placed under the ban by an English statute of 1485/6: "eny bargayne . . . by the name of drye exchaunge . . . be utterly voide." English writers sometimes spoke of it as "sick" exchange,[1] confusing the French *sec* (dry) with an English word of different meaning.



FROM THE MARGARITA PHYLOSOPHICA (1503)

Showing geometry as largely concerned with gaging and similar practical work

4. Fictitious exchange,[2] the plan of collecting a debt by drawing on the debtor.

As coinage came to be better settled a definite par of exchange was recognized, and so, beginning in the arithmetics of the latter part of the 16th century, we find various rules relating to this subject. Thus Recorde (*c.* 1542) says that "as touching the exchange, it is necessary to vnderstand or know the Pair, which the Italians call Pari."[3]

Days of Grace. The Italian cities had fixed rules as to the number of days after sight or after date at which drafts should be paid. Drafts between Venice and Rome were payable ten days after sight; between Venice and London, three months after date; of Venice on Lyons, at the next succeeding quarterly fair, and so on, thus giving the payer time to obtain money. In these customs is to be found the origin of the

[1] So T. Wilson, writing on usury in 1584, speaks of "sicke and drie exchange."

[2] The *cambio fittitio* of the Italian writers of the 16th century, and the *change fict* of the French.

[3] 1594 ed., p. 557. We also have such Dutch terms as *Rekeninghe vander Pary* (Raets, 1580) and *Den Reghel Parij* (Stockmans, 1676 ed.), and such French terms as *le per* (Savonne, 1563).

"days of grace," formerly allowed in England and America, but generally, owing to improved banking facilities, abandoned in the latter country about the opening of the 20th century.



THE GAGER (GAUGER)

From Köbel's *Vysirbůch*, 1515, showing the tools of the art

Meaning of a Check. At present the check (in England, cheque) is extensively used instead of a bank draft. The word has an interesting history, coming from the Persian *shah*, a king. In the game of chess the player called out "*shah*"

when the king was in danger, and "*shah-mat*" ("the king is dead") when the king could no longer move. From this we have our "check" and "checkmate" in chess, "check"



THE GAGER AT WORK

From Johann Frey's *Ein new Visier büchlein*, Nürnberg [1543]

being thought to mean simply "stop." Hence we have the Middle English *chek*, French *échec* (a check or defeat), Italian *scacco* (a chess board), "checkers," and "check" (a stop in one's account at a bank).

Gaging. Before the size of casks was standardized as the result of manufacturing in quantities or of general laws affecting large territories, the subject of gaging[1] (gauging) played

[1] The Middle English was *gagen* or *gawgen*, to gage. The *u* came in through the Old French *gauger*. In medieval Latin *gaugia* meant a standard wine cask, but the origin of the word is uncertain.

an important rôle in applied mathematics. The word relates to the finding of the capacity of casks and barrels. In Germany a gager was called a Visierer,[1] and in the 15th century there appeared numerous manuals with such titles as *Vysirbůch*, *Vysierbuch*, and *Visyrbuechleynn*.[2] The custom was carried over to England, and even in our early American arithmetics there were chapters on gaging. The first notable German book on the subject was Köbel's work (see page 579) of 1515, although the *Margarita phylosophica* of 1503 pictured geometry (see page 578) as chiefly concerned with such measurements.

Other Applications. It would be difficult to give a satisfactory list of all the applications of elementary mathematics to the manifold interests of man that have developed in the centuries past. These applications include, besides those already mentioned, such topics as the adulteration of goods,[3] accountancy and exchange,[4] small commerce from town to town,[5] the leather trade,[6] grazing, and baking,—a list that might be extended to include many other topics and that illustrates the way in which arithmetic has met human needs.

[1] From *visieren*, to visé, to indorse a standard, to show that it has been seen (subjected to vision) and approved.

[2] *E.g.*, Grammateus, 1523.

[3] Thus Köbel: "Die Regel Fusci mit jhrer Erklårung/folget hernach. . . . Das wort Fusci/bedeut nicht anders/dann ein zerbrochen gůt gemülb/oder andere vnreynigkeit/so in der Specerei funden wirt/als under den Negelin/Imber/Saffran/&c. Auch Silber vnderm Golt/Kupffer vnderm Silber." *Rechenbüchlein*, 1514; 1549 ed., fol. 77. See also other German works of the period, such as Albert (1534; 1561 ed., fol. D viii) and Thierfelder (1587, p. 116). The word also appears as *Fusti*.

[4] Frequently found in the early Dutch books as "Rekeninghe voor cassiers" (Van der Schuere, 1600, fol. 65), "Reductio, ofte Cassiers Rekeninge" (Cock, 1696 ed., p. 96), "Cassiers Rekeninghe" (Mots, 1640, fol. 1), "Den Regel vander Muntē Oft den Reghel gheheeten Regula Cassiers" (Stockmans, 1676 ed., p. 205).

[5] A common topic in the French, Dutch, and German books of the 16th and 17th centuries under such titles as "Rechnung vberland" (Rudolff, *Kunstliche rechnung* (1526; 1534 ed., fol. k1)); "Overlantsche Rekeninghe" and "Comptes de Voyage" (Coutereels, Dutch and French arithmetic, 1631, p. 283); "Uytlandtsche Rekeningen" and "Voyages" (Eversdyck's edition of Coutereels, 1658, p. 299); and "Voyagien" (Houck, 1676 ed., p. 148).

[6] Particularly in such centers as Leipzig, where we find arithmetics of about 1700 with such chapters as "Leder und Rauchwahren-Rechnung."

4. Applications of Algebra

General Nature. The first applications of algebra were in the nature of number puzzles. Such was the first algebraic problem[1] of Ahmes (*c.* 1550 B.C.), already mentioned,—"Mass, its whole, its seventh, it makes 19." Such are, in general, the problems of Diophantus (*c.* 275), and the problems which still form the large majority of those given in current textbooks. When a pupil is called upon today to solve the equation $x + \frac{1}{7}x = 19$, he is really solving the first problem of Ahmes, and all our abstract work in equations is a development of this type.

The second general application is geometric, and this characterizes the works of the Greek writers, with the exception of Diophantus.

The third general application is to fanciful problems relating to human affairs, and this is essentially Oriental in spirit.

The fourth type is characterized by the attempt to relate algebra actually to the affairs of life. The first steps in this direction were taken when algebra was more or less a part of arithmetic, problems often being given that were essentially algebraic but were solved without any further symbolism than that afforded by the medieval algorism. It is with this type that we are working at present and are making some advance.

We shall now consider the first three of these stages in detail.

The Number Puzzle. It was with the number puzzle that algebra seems to have taken its start. The desire of the early philosophers to unravel some simple numerical enigma was similar to the child's desire to find the answer to some question in the puzzle column of a newspaper. A few types will be selected involving linear equations, the quadratic being considered later.

Ahmes (*c.* 1550 B.C.) gave numerous problems like "Mass, its $\frac{2}{3}$, its $\frac{1}{2}$, its $\frac{1}{7}$, its whole, it makes 33."[2]

[1] No. 24 in Peet, *Rhind Papyrus*, p. 61, with slightly different translation.

[2] *Ibid.*, No. 31. In modern symbols,

$$x + \tfrac{2}{3}x + \tfrac{1}{2}x + \tfrac{1}{7}x = 33.$$

The problems of Diophantus (*c.* 275) were often of this type, as in the first one given in Book I:[1] "To divide a given number into two having a given difference," with the particular case of $2x + 40 = 100$.

In the Middle Ages in Europe the standard algebraic problem was of the same general nature. This is seen, for example, in the *De Numeris Datis* of Jordanus Nemorarius (*c.* 1225), where all the problems are abstract, as in the following case:

"If there should be four numbers in proportion and three of them should be given, the fourth would be given."[2]

From early times to the present the number puzzle has played a leading part in algebra, and under current conditions, when algebra is required as a school subject, this is not altogether fortunate.

Simultaneous Linear Equations. Problems involving simultaneous linear equations were more numerous in the Orient in early times than they were in Europe. Thus we find in India a considerable number of such problems, together with rules that amount to directions for solving various types of simultaneous equations. For example, Mahāvīra (*c.* 850) has the following problem:

> The mixed price of 9 citrons and 7 fragrant wood-apples is 107; again, the mixed price of 7 citrons and 9 fragrant wood-apples is 101. O you arithmetician, tell me quickly the price of a citron and of a wood-apple here, having distinctly separated those prices well.

His rule for the solution is similar to the one used today for eliminating one unknown.[3]

Another of Mahāvīra's problems, evidently suggested by one that appeared in the Greek epigrams and was there attributed to Euclid, is as follows:

> Three merchants saw [dropped] on the way a purse [containing money]. One [of them] said [to the others], "If I secure this purse,

[1] Heath, *Diophantus*, 2d ed., p. 131.

[2] Problem 30: "Si fuerit [*sic*] IIII numeri proporcionales et tres eorum dati fuerint et quartus datus erit" (*Abhandlungen*, II, 143).

[3] See his work, p. 130.

I shall become twice as rich as both of you with your moneys on hand." Then the second [of them] said, "I shall become three times as rich." Then the other [, the third,] said, "I shall become five times as rich." What is the value of the money in the purse, as also the money on hand [with each of the three merchants]?[1]

Indeterminate Problems. The indeterminate problem, leading to what is now called the indeterminate equation, is very old. It is probable that it formed a type of recreation long before the time of Archimedes, since in the problems assigned to him there appears one of very great difficulty. This problem has already been discussed on page 453.[2]

Although Diophantus (*c.* 275) proposed many indeterminate equations, they were not in the form of applied problems. Cases of the latter kind seem to have come chiefly from the Orient, at least in early times.

In the *Greek Anthology* (*c.* 500?) there are two problems involving indeterminate linear equations. The first (XIV, 48) is as follows:

The three Graces were carrying baskets of apples and in each was the same number. The nine Muses met them and asked them for apples, and they gave the same number to each Muse, and the nine and three had each of them the same number. Tell me how many they gave and how they all had the same number.[3]

The second (XIV, 144) is a dialogue between two statues:

A. How heavy is the base on which I stand, together with myself!

B. My base together with myself weighs the same number of talents.

A. I alone weigh twice as much as your base.

B. I alone weigh three times the weight of yours.[4]

[1] See his work, p. 155.

[2] The problem was discovered by the German dramatist G. E. Lessing, in 1773, while he was serving as court librarian at Wolfenbüttel. On the history of the problem see B. Lefebvre, *Notes d'Histoire des Mathématiques*, p. 33 n. (Louvain, 1920). On a few indeterminate problems due to Heron, see Heath, *History*, II, 444.

[3] The equations reduce to $x = 4y$. There were $12n$ apples in all.

[4] The equations are $x + y = u + v$, $x = 2v$, $u = 3y$.

With their usual desire to give a fanciful but realistic touch to algebra, the Chinese applied the indeterminate equation to a problem commonly known as that of the Hundred Fowls. This problem goes back at least to the 6th century[1] and differs so greatly in its nature from those of the Greeks that it seems to have originated in the East. The problem is as follows:

If a cock is worth 5 *sapeks*; a hen, 3 *sapeks*; and 3 chickens together, 1 *sapek*, how many cocks, hens, and chickens, 100 in all, will together be worth 100 *sapeks*?

From China the problem apparently found its way to India, for it appears in Mahāvīra's work (*c.* 850) in the following form:

Pigeons are sold at the rate of 5 for 3 [*paṇas*], *sārasa* birds at the rate of 7 for 5 [*paṇas*], and peacocks at the rate of 3 for 9 [*paṇas*]. A certain man was told to bring at these rates 100 birds for 100 *paṇas* for the amusement of the king's son, and was sent to do so. What [amount] does he give for each [of the various kinds of birds that he buys]?

Mahāvīra gave a method for solving such problems that was sufficient to satisfy those who were interested in puzzles, but which had little merit otherwise.[2]

This fanciful type of problem was probably made known in Europe at the time of the general penetration of Oriental ideas, and here it developed into a form somewhat like this:

20 persons, men, women, and girls, have drunk 20 pence worth of wine; each man pays 3 pence, each woman 2 pence, and each girl $\frac{1}{2}$ penny; required the number of each.

Algebraically, $$m + w + g = 20$$

and $$3m + 2w + \tfrac{1}{2}g = 20,$$

[1] Kaye, *Indian Math.*, p. 40; L. Vanhée, "Les cents volailles ou l'analyse indéterminée en Chine," *T'oung-pao*, XIV, and reprint; L. Matthiessen, "Vergleichung der indischen Cuttaca- und der chinesischen Tayen-Regel, unbestimmte Gleichungen und Congruenzen ersten Grades aufzulösen," *Sitzungsberichte der math.-naturwiss. Section in der Verhandl. der Philol. Vers. zu Rostock*, 1875.

[2] See his work, pp. 133–135.

which would be indeterminate if it were not for the fact that there were some of each and that the result must be in positive integers, thus admitting of only one set of answers, namely, 1, 5, and 14. In general, all problems of this type are of the form

$$x + y + z = m,$$

$$ax + by + cz = n,$$

reducing to

$$px + qy = r,$$

thus being indeterminate unless the physical conditions are such as to exclude all but one set of values.

The Regula Coecis. Such problems are known in European works as early as the 9th century, and thereafter they become common.[1] In the 15th century they begin to mention the persons as being at a *cecha*,[2] and hence the rule for solving such problems became known to 16th century writers as the *Regula Coecis*. There has been much dispute as to the origin of the term *coecis*,[3] but from such historic evidence as we now have it seems to relate to the fact above stated, namely, that the persons were at a *cecha*.[4] The problem seems to relate to drinking where each paid his own share,[5] very likely from the fact that *zecche* meant originally the money[6] paid for the drinks.

[1] Thus a 14th century MS. in Munich (cod. lat. Monac. 14684): "Sint hic milites, pedites et puelle, et sint in universo 12, et habeant 12 panes, parciendos, et quilibet miles accipiat duos, quilibet pedes quartam partem panis, quilibet puella medietatem panis: queritur, quot erunt milites, pedites et puelle." *Bibl. Math.*, IX (2), 79.

[2] Curtze, for example, found, in a MS. of 1460, a problem beginning: "ponam casus, quod sint 20 persone in una cecha" (*Abhandlungen*, VII, 35).

[3] Which also appears in such forms as *zekis*, *zeches*, *cekis*, *ceci*, *coeci*, *caeci*, and *caecis*. *Bibl. Math.*, XIII (2), 54, and VI (3), 112.

[4] We find similar expressions in the 16th century. Thus Rudolff (*Kunstliche rechnung*, 1526) has a topic "Von mancherley person an einer zech," and says further: "Es sitzen 20 person an einer zech/man/frawen/vnd jungfrawen . . .," adding that this is the "regel/welche sie nennen Cecis oder Virginū/" (1534 ed., fol. N vij).

[5] L. Diefenbach, in his *Novum Glossarium* (Frankfort a. M., 1867, p. 339), says: "symbolum . . . vulgo zecha quo quisque suam portionem confert." In American slang, a *zecha* was a "Dutch treat."

[6] Italian *zecca*; compare *zecchiere*, a mint master, as in Ciacchi, *Regole generali d' Abbaco* (Florence, 1675, p. 247).

There have been various other speculations as to the word. One of these is that it comes from an Arabic expression signifying not content with one but demanding many, referring to the many possible solutions.[1] It has also been thought to come from the Latin *caecus* (blind), with such fanciful explanations as that a problem of this kind was solved by the blind Homer, or that the solver went blindly to work to find the solution. Indeed, it was often called the Blind Rule in the 16th and 17th centuries.[2]

Another name for the rule was *Regula potatorum* (rule of the drinkers), and this gives added reason for the interpretation of *zecha* as a drinking bout.[3] A still more common name was *Regula virginum*, rule of the girls, usually explained by the fact that the solutions show more girls than men or women.[4]

In spite of the fact that this style of problem is interesting, the arithmeticians often discouraged its use because of its indeterminateness,[5] although there were found others who increased the difficulty by using more than three unknowns.

Alligation. Beginning apparently in the Renaissance period as an application of indeterminate equations, the *Regula Alli-*

[1] "Cïntu Sekîs, hoc est adulteram indigetarunt: propterea, ut opinor, quod uno ac legitimo quaestionis enodatu non contenta, plures plerumque admittat solutiones." J. W. Lauremberg, *Arithmetica* (Sorö, 1643), quoted by Zeuthen in *L'Intermédiaire des Mathématiciens*, p. 152 (Paris, 1896) (hereafter referred to as *L'Intermédiaire*); *Bibl. Math.*, X (2), 96. Carra de Vaux (*Bibl. Math.*, XI (2), 32) says that Lauremberg's expression should have the word *sikkir*, which means toper, and this is more reasonable.

[2] Thus Thierfelder asks: "Warumb wirdt dise Regel Cecis genannt?" and explains that the problem is indeterminate and that "ein Ungeübter nicht bald finden kan/darumb ist es jm ein blinde Regel" (1587, p. 211). Cardinael speaks of it as "Den Blinden-Reghel" (1674 ed., p. 88).

[3] The Dutch writer, Bartjens, also speaks of it as "Bachus-rekeninge" (1752 ed., p. 213). See also Unger, *Die Methodik*, p. 101.

[4] This origin is stated by Jacobus Micyllus (1555): "regula quam ab eo, ut videtur, appellarunt, quod virginum personae ac nomen inter exempla illius subinde repetuntur." So we find "Coecis oft Virginum" (Van der Schuere, 1600, fol. 174), "Rekeninge Coecis, ofte Virginum" (Bartjens, 1633), and similar forms, expecially among the early Dutch writers.

[5] Thus Stockmans: "Desen regel niet soo seker en is in zijn werckinge als de ander voorgaēde" (1676 ed., p. 380); and Coutereels remarks that it is "meer vermakelijkheyd als sekerheyd" (1690 edition of the *Cyffer-Boeck*, p. 559).

gationis,[1] or Rule of Alligation, attracted considerable attention for nearly three hundred years. Problems in alligation were sometimes indeterminate and sometimes not. The following, for example, is indeterminate:

How many Raisins of the Sun, at *7d. per lb.* and Malaga Raisins at *4d. per lb.* may be mixed together for *6d. per lb.*?[2]

Such a problem becomes determinate by the addition of some further appropriate condition, as in the following case:

A tobacconist mixed 36 lb. of tobacco, at 1s. 6d. per lb. 12 lb. at 2s. a pound, with 12 lb. at 1s. 10d. per lb.; what is the price of a pound of this mixture?[3]

In general, such problems were simply ingenious efforts to make algebra seem real, but they were usually solved without the aid of algebraic symbols, and hence they found place in higher arithmetics until the close of the 19th century, when they generally disappeared except as a few remained in the form of mixture problems[4] in the elementary algebras. Indeed, Recorde (*c.* 1542) remarked that the rule of alligation "might be well called the rule of Myxture."[5] He was the first English writer to suggest other applications than those referring to alloys, saying: "it hath great vse in composition of medicines,

[1] From *ad*, to, + *ligare*, to bind. From the same roots come the French *alliage*, and our words *alloy* and *ally*. So we have in French "La regle des aliages" (Trenchant, 1566; 1578 ed., p. 191) and "La Reigle d'Alligation" (Peletier, 1549; 1607 ed., p. 247).

[2] T. Dilworth, *The Schoolmasters Assistant*, new ed., p. 97 (London, 1793). This was one of the most celebrated English arithmetics of the period; it had great influence on American textbooks.

[3] N. Daboll, *Schoolmaster's Assistant*, 1837 ed., p. 177. This was one of the most celebrated of the early American arithmetics.

[4] The Dutch writers called them problems solved by "Den Reghel van Menginghe" (Cardinael, 1674 ed., p. 66, with 23 pages to the subject). So we find "Allegatio, Menginghe" (Van der Schuere, 1600), "Rekeninghen van Mengelingen" (Raets, 1580 ed., fol. K 3), "Alligationis, ofte Menginghe" (Bartjens, 1676 ed., p. 165), and "Alligatio, Alliage, of Mengingh" (Coutereels, *Cyffer-Boeck*, 1690 ed., p. 484). The Italian writers sometimes had a chapter *De' mescoli* (on mixtures), and this may have suggested the German *Mischungsrechnung*. For a recent use of the topic in Germany, see R. Just, *Kaufmännisches Rechnen*, I, p. 86 (Leipzig, 1901).

[5] *Ground of Artes*, 1558 ed., fol. Y 3.

and also in myxtures of metalles, and some vse it hath in myxtures of wines, but I wshe it were lesse vsed therin than it is now a daies." These practices rendered the subject so popular that Baker (1568) gave forty-eight pages to it in his 1580 edition.

Mint Problems. One of the leading applications of alligation was found in the general need for the mixing of chemicals and metals by the alchemists of the Renaissance,[1] by bell founders, and by mint masters. As to the coining of money, it should be remembered that this was not in general a government monopoly in the Middle Ages, so that it was looked upon as something unheard of that Ferdinand and Isabella should assert this right in 1496.[2] The privilege belonged rather to cities or districts and even in a single small country was often claimed by several people,[3] often as an inherited right.[4] Add to this fact the great awakening in the mining industry of Germany in the latter part of the 15th century and the extensive importations of gold and silver from the Americas in the first half of the 16th century, and it will be seen that the subject of alligation naturally had at that time a new and popular field in the mixing of alloys for purposes of coinage.[5] This explains the interest in the subject of coinage in the 16th century.[6]

[1] "Il consolare oro, ed argento non è altro, che vn' allegazione di que' due metalli, per li quali la maggior parte degli Alchimisti son diuenuti miseri, e mendichi, per volere inuestigare la congelazione del Mercurio in vera, ed ottima Luna, o Sole, la quale senza il diuino aiuto in vano da gli Alchimisti vien tentata." Ciacchi, *Regole generali d' Abbaco*, p. 244 (Florence, 1675).

[2] H. B. Clarke, in the *Cambridge Mod. Hist.*, I, chap. xi.

[3] W. Cunningham, in the *Cambridge Mod. Hist.*, I, chap. xv.

[4] E. P. Cheyney, *Documents Illustrative of Feudalism*, p. 34 (Philadelphia, 1898).

[5] Of course alloys for this purpose had been known and used to some extent ever since the early coinage from natural electron in Asia Minor and from bronze in Rome. On the latter, see F. Gnecchi, *Monete Romane*, 2d ed., p. 86 (Milan, 1900).

[6] So Rudolff (1526; 1534 ed., fol. M 6) and Grammateus (1518; 1535 ed., fols. C 8, D 4, etc.) have chapters on *Muntzschlag*, and various writers speak of the problems of the *müntzmeister* (Rudolff), *Munt-meester* (Van der Schuere, 1600), and *mint-master* (Hodder, 1672 ed.). The Germans also had chapters on *Silber Rechnung*, *Goldt Rechnung*, and *Kupfer Rechnung* (Riese, 1522; 1550 ed., fols. 37–40); the Dutch, on *Rekeninghe van Goudt end Silber*, with *Comptes d'or & d'argent* (Coutereels's Dutch and French editions of 1631, p. 298, and other dates); and the Italians, on *Del consolare dell' oro et dell' argento* (Cataneo, 1546; Tartaglia, 1556).

Problem of Hiero's Crown. Closely related to this subject is the problem of Hiero's crown, which Robert Recorde (*c.* 1542) states in quaint language as follows:

Hiero kynge of the Syracusans in Sicilia hadde caused to bee made a croune of golde of a wonderfull weight, to be offered for his good successe in warres: in makynge wherof, the goldsmyth fraudulētly toke out a certayne portion of gold, and put in syluer for it. [Recorde then relates the usual story of Archimedes and the bath, telling how the idea of specific gravity came] as he chaunced to entre into a bayne full of water to washe hym, [and] reioycing excedingly more then if he had gotten the crown it self, forgat that he was naked, and so ranne home, crying as he ranne ἔυρηκα, ἔυρηκα,[1] I haue foūd, I have found.

Cardan[2] asserts that the story is due to Vitruvius, who simply transmitted the legend. It appears in various books of the 16th century and is still found in collections of algebraic problems.

First Problems in the New World. To those especially who live in the New World some local interest attaches to the problems that appear in the first mathematical work (1556) to be printed there. Among these problems are the following:[3]

I bought 10 varas of velvet at 20 pesos less than cost, for 34 pesos plus a vara of velvet. How much did it cost a vara? Add 20 pesos to 34 pesos, making 54 pesos, which will be your dividend. Subtract one from 10 varas, leaving 9. Divide this into 54, giving 6, the price per vara.

I bought 12 varas of velvet at 30 pesos less than cost, for 98 pesos minus 4 varas. How much was the cost per vara? The following is a short method: add the 30 pesos and the 98 pesos, making 128; add the number of varas, 12 and 4, making 16; divide 16 into 128, giving 8, the price per vara.

[1] *Eu'reka, eu'reka*; more precisely, *heu'reka*.

[2] *Practica*, 1539, cap. 66, ex. 45. He relates that Archimedes "nudus e balneo exultās domū reuertebatur," adding, "nescio an ob amorē veritatis potius laudandus quā ob importunā & impudicā nuditatē vituperādus."

[3] See also pages 385, 392, and Volume I, pages 353–356.

I bought 9 varas of velvet for as much more than 40 pesos as 13 varas at the same price is less than 70 pesos. How much did a vara cost? Add the number of pesos, 40 and 70, making 110. Add the number of varas, 9 and 13, making 22. Dividing 110 by 22, the quotient is 5, the price of each vara.

A man traveling on a road asks another how many leagues it is to a certain place. The other replies: "There are so many leagues that, squaring the number and dividing the product by 5, the quotient will be 80." Required to know the number of leagues.

A man is selling goats. The number is unknown except that it is given that a merchant asked how many there were and the seller replied: "There are so many that, the number being squared and the product quadrupled, the result will be 90,000." Required to know how many goats he had.


The work contains other and more difficult problems in algebra and the theory of numbers, but the above are types of the ordinary puzzles which the author places before his readers.

5. Magic Squares

Oriental Origin. The magic square seems unquestionably to be of Chinese origin. The first definite trace that we have of it is in the *I-king*, where it appears as one of the two mystic arrangements of numbers of remote times. This particular one, the *lo-shu*, is commonly said to have come down to us from the time of the great emperor Yu, *c.* 2200 B.C. The tradition is that when this ruler was standing by the Yellow River a divine tortoise appeared, and on its back were two mystic symbols, one being the *lo-shu* already described in Volume I, Chapter II. As may be seen from the illustration there given, it is merely the magic square here shown.


4	9	2
3	5	7
8	1	6

This particular square is found in many recent Chinese works, and every fortune teller of the East makes use of it in his trade. Little by little the general knowledge of magic squares seems to have been extended, and when Ch'êng Tai-wei



MAGIC SQUARES IN JAPAN

Half of a magic square as given in Hoshino Sanenobu's *Kō-ko-gen Shō*, 1673




MAGIC CIRCLE FROM SEKI'S WORKS

From the reprint of the works of the great Japanese mathematician, Seki Kōwa (*c.* 1661)

wrote his *Systematized Treatise on Arithmetic*[1] in 1593, he included not only a discussion of magic squares but also one of magic circles.

Japan. The Japanese became particularly interested in the subject in the 17th century, as the illustrations from the works



MAGIC CIRCLE OF 129 NUMBERS

From Muramatsu Kudayū Mosei's *Mantoku Jinkō-ki*, 1665. The numbers in each radius add to 524, or 525 with the center 1

of some of their leading writers show.[2] Among the prominent scholars who gave attention to these forms were Muramatsu Kudayū Mosei, who wrote several works on arithmetic and

[1] *Suan-fa Tong-tsung*. See Volume I, page 352.

[2] For their treatment of the subject see Smith-Mikami, pp. 57, 69, 71, 73, 79, 116, 120, 177.

geometry, beginning in 1663; Hoshino Sanenobu, whose *Triangular Extract*[1] appeared in 1673; Isomura Kittoku, whose *Ketsugi-shō* appeared in 1660; and the great Seki Kōwa, who devoted one of his *Seven Books*[2] to the theory of magic squares and magic circles.

7	12	1	14
2	13	8	11
16	3	10	5
9	6	15	4

India. From China the magic square seems to have found its way into India and the adjacent southern countries, but whether this was direct or through the Arab influence we can only conjecture. It appears in a Jaina inscription in the ancient town of Khajurāho, India, where various ruins bear records of the Chandel dynasty (870–1200), and is probably not older than the 11th or 12th century. This Indian square, shown above, displays a somewhat advanced knowledge of the subject, for not only has it an even number of squares on a side, but each of the four minor squares has a relation to the others, as may be seen by the illustration at the right.

19 9 25 15	15 9 25 19
19 25 9 15	15 25 9 19

This is perhaps the earliest trace of such fantastic elaborations of the magic square,[3] although no careful study of the history has yet been made.[4] It is probable that the astrologers carried such ideas to the West, where their influence upon the medieval mathematics of Europe is apparent.

Today the magic square is used as a charm all through India, being found in fortune bowls, in medicine cups, and in amulets. In Thibet it is particularly in evidence, being found in the

[1] *Kō-ko-gen Shō.*

[2] Or *Shichibusho.* The particular book is the *Hōjin Yensan*, revised in MS. in 1683.

[3] F. Schilling, *Jahresbericht der deutschen Math. Verein.*, XIII, 383.

[4] But see such works as W. S. Andrews, *Magic Squares* (Chicago, 1907), and subsequent articles in *The Open Court.* For Roman and Egyptian claims see E. Falkener, *Games, Ancient and Oriental*, p. 277 (London, 1892).

"wheel of life"[1] and worn as an amulet to ward off evil. It is also seen in Sumatra, in the Malay Peninsula, and in the other countries which have had close relations with India and China.

The squares are not always of the pure type, however; that is, the sums of the rows, columns, and diagonals are not always constant. In some of them, for example, the columns add successively to 300, 200, 100, and the like,[2] and in many of them the numbers are repeated whenever it was necessary to make the sums come as desired.



THIBETAN TALISMAN

With signs of the zodiac, the ancient *pa-kua* or trigrams, and the *lo-shu* in Thibetan numerals

Connection with Alchemy. It seems that the numbers must often have been connected with the old alchemistic idea of the planets and the metals,—an idea that permeated the doctrines of many of the medieval mystics. Of the three triads made up of the nine digits, the first had the following relations:

1 = gold = the sun, ☉
2 = silver = the moon, ☽
3 = tin = Jupiter, ♃ (the hand grasping the thunderbolt)

[1] See the illustration in Volume I, page 27.

[2] S. S. Stitt, "Notes on some Maldivian Talismans," *Journ. Royal Asiatic Soc.*, pp. 121, 130 (London, 1906).

The second triad was as follows:

4 = gold again = the sun, ☉
5 = mercury = Mercury, ☿
6 = copper = Venus, ♀

The third triad was as follows:

7 = silver again = the moon, ☽
8 = lead = Saturn, ♄
9 = iron = Mars, ♂

With such a relationship it is possible to understand such talismans as one found in the Maldivian Islands, in the Indian Ocean, where the charm to protect a virgin sums to 18, whose digits sum to 9, the number of Mars, the protector. In these numbers there also enters the idea of congruence, particularly to the modulus 9. Thus a talisman to keep out Satan has its rows $80 \equiv 8$, $1600 \equiv 7$, $180 \equiv 9$, $\cdots$, which have for their sum $69 \equiv 15 \equiv 6$ = Venus,—a vagary that can be explained only by conjecture.[1]

Hebrews. The magic square played an important part in the cabalistic writings of the Hebrews. Rabbi ben Ezra (*c.* 1140) mentions it, although it had been used by Hebrew writers long before his time.[2] The Eastern Jews early found in the ancient Chinese *lo-shu*, with its constant sum 15, a religious symbol, inasmuch as 15 in Hebrew is naturally יה (10 + 5), which is made up of the first two letters of Jahveh (Jehovah), that is, יהוה, although, lest they should be guilty of profanity, they always wrote 9 + 6 for 15. If the corner numbers of the common form of square are suppressed, the even (feminine) elements are eliminated and there remain only the odd numbers, this cruciform arrangement serving as a charm among various Oriental peoples.[3]

	9	
3	5	7
	1	

[1] Stitt, *loc. cit.*, p. 144. [2] M. Steinschneider, *Abhandlungen*, III, 98.
[3] D. Martines, *Origine e progressi dell' Aritmetica*, p. 39 (Messina, 1865), with considerable information upon this subject.

Arabs. Whether the magic square reached the Arabs from India, from China, or from Persia, we do not know. It might readily have come from any one of these countries through either of the others, but at any rate it was well known to various late Arab writers, appearing, for example, in the works of the philosopher Ġazzâlî[1] about 1100.

Christians. In general the magic square found no recognition as a Christian symbol, although the occult writers naturally made use of it. When it appears in medieval mathematical works it is usually in the form of a problem whose solution requires the arrangement of the ordinary 9-celled square shown below.[2]

The textbook writers of the 16th century paid considerable attention to the subject. Cardan, for example, gives seven different squares bearing respectively the names of the sun, the moon, and the five planets then known, and gives some directions for the formation of such squares.[3]

In art the first instance of the use of these forms is probably the one in the well-known *Melancholia* by Dürer.

One of the most elaborate examples found in architectural decoration is cut in the wall in the Villa Albani at Rome.[4] It contains eighty-one cells and is dated 1766.

2	7	6
9	5	1
4	3	8

[1] Mohammed ibn Mohammed ibn Mohammed, Abû Ḥâmid, al-Ġazzâlî, born at Ṭûs, 1058/59; died at Ṭûs, 1111. The Arabic name of the magic square is *shakal turâbi*. See E. Rehatsek, "Explanations and Facsimiles of eight Arabic Talismanic Medicine-cups," *Journ. of the Bombay Branch of the Royal Asiatic Soc.*, X, 150.

[2] For example, Günther reports a 13th century MS. with this problem: "Tres erant fratres in Colonia, habentes 9 vasa vini. Primum vas continet 1 amam, secundum 2, tertium 3, · · · nonum 9. Divide vinum illud aequaliter, inter illos tres, vassis inconfractis."

The division is 1 + 5 + 9, 3 + 4 + 8, 2 + 6 + 7, but that this is indeterminate is seen by the magic square here represented.

[3] *Practica*, 1539, capp. 42, 66.

[4] The inscription is "Caetanus Gilardonus Romanus philotechnos inventor. A.D. MDCCLXVI." On the recent mathematical investigations the reader will do well to consult such works on mathematical recreations as those of Lucas, Ball, Schubert, and Ahrens; the work by W. S. Andrews, mentioned on page 594; and the references in G. A. Miller, *Historical Introduction to Mathematical Literature*, p. 20 (New York, 1916).



DÜRER'S "MELANCHOLIA," WITH MAGIC SQUARE

One of the first magic squares to appear in print. Chiefly interesting because it is an even-celled square. Dürer (1471–1528) also wrote on higher plane curves in connection with art

TOPICS FOR DISCUSSION

1. The general progress of mathematical recreations, with a consideration of the leading works upon the subject.

2. The influence of mathematical recreations upon the development of mathematics.

3. Traces of the ancient puzzle problems in the elementary mathematical literature of the present time, with a consideration of the value of these problems.

4. Types of puzzle problems dependent upon indeterminate equations, with a study of their history.

5. Mathematics as an aid to the study of the history of economics and commerce.

6. The historical development of the corporation, as traced through the problems of arithmetic.

7. Racial influences upon business customs, as seen in commercial problems.

8. The standard applications of arithmetic from remote times to the present, with a consideration of their important changes.

9. Commercial problems at one time important and now nearly obsolete, with a consideration of the causes of their rise and fall.

10. Certain commercial problems and customs, at one time important but now nearly or quite obsolete, left their impress upon our language. Consider a few such cases.

11. Ancient problems of the bank compared with those of the present time.

12. The need for barter in ancient times and the reasons for its retention, even in highly civilized countries, until recently.

13. The history of various types of taxation in ancient, medieval, and modern times.

14. The reason why gaging was considered as an important branch of mathematics in the early printed arithmetics and geometries, and why it lost its standing in the 19th century.

15. Reasons for the study of magic squares in ancient times, in the Middle Ages, and at the present time, with typical illustrations.

16. Instances of the use of the magic square in the East as a talisman or amulet.


17. The relation of magic squares to alchemy, and the relation of alchemy to modern science.

CHAPTER VIII

TRIGONOMETRY

1. General Development of Trigonometry

Meaning of the Term. If we take trigonometry to mean the analytic science now studied under this name, we might properly place its origin in the 17th century, after the development of a satisfactory algebraic symbolism. If we take it to mean the geometric adjunct to astronomy in which certain functions of an angle are used, we might look for its real origin in the works of Hipparchus (*c.* 140 B.C.), although there are earlier traces of its use. If we take it to mean literally "triangle measurement,"[1] the origin would naturally be placed much earlier, say in the second or third millennium B.C. Since this third phase is considered under geometry, we may properly confine our work to the development of the idea of the functions of an angle, giving first a brief sketch of the rise of the science and then the history of certain of its details.



Egypt. In the Ahmes Papyrus (*c.* 1550 B.C.) there are five problems[2] relating to the mensuration of pyramids, and four of these make mention of the *seqt*[3] of an angle. Ahmes is not at all clear in expressing the meaning of this word, but from the context it is thought that the *seqt* of

[1] Τρίγωνον (*trí'gonon*, triangle) + μέτρον (*met'ron*, measure).

[2] Nos. 56–60 in the list. See Peet, *Rhind Papyrus*, p. 97; Eisenlohr, *Ahmes Papyrus*, pp. 134–148. Throughout this chapter much use has been made of A. Braunmühl, *Geschichte der Trigonometrie*, 2 vols., Leipzig, 1900, 1903; hereafter referred to as Braunmühl, *Geschichte*.


[3] Eisenlohr takes the word to mean ratio number. It is also transliterated *śḳd* and *seqet*. It may be significant that the Hebrew *sgd* means "bowing."

the regular pyramid shown on page 600 is probably equivalent to cot $\angle OMV$.[1] The Egyptian pyramids were generally constructed so that $\angle OMV$ was approximately constant (about 52°) and $\angle OAV$ was about 42°. At present we are without means of knowing what use was made of this function.

Babylon. The relation between the mathematical knowledge of the Egyptians and that of the Babylonians in the third millennium B.C., as seen in the unit fraction (p. 210), leads us to suppose that the latter people may have known of the primitive Egyptian trigonometry. We have, however, no direct knowledge that this was the case. There are evidences of angle measure at a very early date, as witness fragments of circles which seem to have been used for this purpose and which have come down to us. These fragments seem to have been parts of primitive astrolabes, as stated on page 348.

There is also extant an astrological calendar of King Sargon, of the 28th century B.C., and a table of lunar eclipses beginning 747 B.C., so that evidence of an interest in astronomy is not lacking throughout a long period of Babylonian history. All this involved a certain amount of angle measure, but there is no direct evidence of any progress in what we commonly understand as trigonometry.

The Gnomon. Herodotus (*c.* 450 B.C.) tells us that the Greeks obtained their sundial from Babylon. This is very likely true, for we know that the Egyptians used a sun clock as early as 1500 B.C., and the Babylonians could hardly have been behind them in the knowledge of such a device. The relation of the sundial to trigonometry is seen in the fact that it is an instrument for a form of astronomical observation. A staff *GN*, called by the Greeks a *gnomon* (p. 16), is erected and the shadow *AN* observed. It is longest at noon when *S*, the sun, is farthest south, this being at the winter



[1] For discussion see Braunmühl, *Geschichte*, I, 2; Eisenlohr, *Ahmes Papyrus*, p. 137. Peet, *Rhind Papyrus*, p. 98, gives cot $\angle OMV$.

solstice, and shortest when it is farthest north, at the summer solstice; and hence an examination of its limits enables the observer to measure the length of the year. The daily lateral motion of the point A allows for the measure of diurnal time, quite as the motion of noon along AN allows for the measure of annual time. The gnomon being constant, the length of AN at noon varies with $\angle A$, and to us this means a recognition that AN, or $AN : GN$, is a function of $\angle A$, namely, the cotangent. We have no trace, however, of any name (except the *seqt*) for such a relation in the period of which we are speaking.

China. In the *Chóu-peï Suan-king* (*c.* 1105 B.C.)[1] the right-angled triangle is frequently used in the measure of distances, heights, and depths, and it is quite probable that the ratios of the sides were recognized. One passage reads, "The knowledge comes from the shadow, and the shadow comes from the gnomon," so that possibly a primitive plane trigonometry was known in China in the second millennium B.C. Aside from this there is no evidence that the early Chinese had names for any functions of an angle. The early astronomical interests of the Chinese, however, like those of other ancient peoples, necessitated some kind of angle measure.

Greece. When Thales measured the height of a pyramid by means of its shadow, he used what was already known, probably in various parts of the world, as "shadow reckoning."[2] In his "Banquet of the Seven Wise Men" Plutarch speaks of Nilax, one of the guests, as saying to Thales:

> Whereas he[3] honors you, he particularly admires you for divers great accomplishments and particularly for the invention whereby, with little effort and by the aid of no mathematical instruments, you found so accurately the height of the pyramids. For, having fixed your staff erect at the point of the shadow cast by the pyramid, two triangles


[1] But see Volume I, page 30; also Mikami, *China*, p. 4.

[2] We have this term, substantially at least, in the *Chóu-peï Suan-king*; in the works of Brahmagupta (*c.* 628), under "Measure by Shadow," p. 317; in Mahāvīra (*c.* 850), under "Calculations relating to Shadows," p. 275; and in Bhāskara, under "Ch'háyá-vyavahára" (determination of shadow), p. 106.

[3] The king of Egypt, called by the Greeks Amasis, *c.* 570 B.C.

were formed by the tangent rays of the sun, and from this you showed that the ratio of one shadow to the other was equal to the ratio of the [height of the] pyramid to the staff.

Essentially the measure of heights by means of shadows involves the knowledge that, in this figure, $BC : AB = B'C' : AB'$. To us it seems as if tan A would be suggested by such a relation, but we have no evidence that this was the case in the time of Thales. We only know that, centuries later, AB was called the *umbra recta* (right shadow), showing that the relation of AB to BC entered trigonometry through shadow reckoning.



It is said that Anaximander (*c.* 575 B.C.) erected near Sparta the first gnomon in Greece. It was probably in the form of an obelisk, a mere post placed perpendicular to the apparent plane of the earth's surface, and not the triangular form later in use. It could have been used for determining the meridian line, and tradition says that this was done; but besides this it served, as it probably did in Egypt and Babylon, to measure the year, the seasons, and the time of day.

Relation to Astronomy. In this early work of Anaximander, as in similar cases among the Babylonians and Egyptians, it is evident that the real purpose in view was the study of astronomy, the unraveling of the mysteries of the universe. This led to the study of the celestial sphere, the triangles being, therefore, spherical figures. This accounts for the fact that the study of spherical triangles kept pace with that of plane triangles in the Greek trigonometry. We find, however, no tangible evidence of the definition or even of the idea of a spherical triangle before the appearance of the work of Menelaus on spherics (*c.* 100).[1]

Early Works on Spherics. The oldest extant works on spherics, and indeed the oldest Greek mathematical texts that have come down to us, are two astronomical treatises by Autol'ycus[2]

[1] Heath, *History*, II, 262.

[2] Αὐτόλυκος. See Heath, *History*, I, 348.

of Pitane (*c.* 330 B.C.). The first is on a moving sphere[1] and consists of twelve elementary propositions relating to the principal circles. The second work was on the risings and settings of the fixed stars,[2] in two books. Neither of these works shows, however, any knowledge of spherical trigonometry.

Aristarchus. The next important step in the development of trigonometry was taken by the astronomer Aristarchus of Samos (*c.* 260 B.C.).[3] He attempted to find the distances from the earth to the sun and the moon, and also the diameters of these bodies. His geometric reasoning was accurate, but his instruments were so crude that he could come no nearer the ratio of the distance of the moon to that of the sun than to say that it was between $\frac{1}{18}$ and $\frac{1}{20}$. In his proof he makes use of ratios which are suggestive of the tangent of an angle.[4]

Hipparchus. In his commentary on the *Almagest*,[5] Theon of Alexandria (*c.* 390) asserts that Hipparchus (*c.* 140 B.C.), the greatest of the Greek astronomers, wrote twelve books on the computation of chords[6] of angles, but of these books we have no further trace. Hipparchus himself, in the fragment of his work that has come down to us, leads us to believe that he was engaged in such computations and in the graphic solution of spherical triangles.[7] It therefore seems reasonable to assert,

[1] Περὶ κινουμένης σφαίρας.

[2] Περὶ ἐπιτολῶν καὶ δύσεων. The two works were edited by Hultsch in 1885.

[3] P. Tannery thinks that this step, which is usually attributed to Aristarchus, was taken by Eudoxus (*c.* 370 B.C.). See his "Aristarque de Samos," *Mém. de la Soc. des sciences de Bordeaux*, V (2), 241; *Mémoires Scientifiques*, I, 371; Heath, *History*, II, 1.

[4] This proof is given in Braunmühl, *Geschichte*, I, 8, and by Tannery, *Mémoires Scientifiques*, I, 376. See also the Commandino edition of Aristarchus, 1572; R. Wolf, *Geschichte der Astronomie*, p. 172 (Munich, 1877). The work of Aristarchus, Περὶ μεγεθῶν καὶ ἀποστημάτων ἡλίου καὶ σελήνης, was translated by A. Nokk and published, with a commentary, in a Programm, Freiburg, 1854. See also Heath, *History*, II, 4.

[5] Halma's French translation, p. 110 (Paris, 1821).

[6] The Greeks called the chord εὐθεῖα (*eutheĭa*), the Latin *chorda* being from the Greek χορδή (*chordē*, intestine), whence it meant a string made of dried intestine used in a lyre, and hence a straight chord of a bow (arc).

[7] Braunmühl, *Geschichte*, I, 10; Heath, *History*, II, 257.

from the evidence that we have, that the science of trigonometry begins with Hipparchus.[1] It has been asserted, but the proof is unsatisfactory, that the formulas for sin $(A \pm B)$ and cos $(A \pm B)$, and for the radius of the circumscribed circle $\left(R = \frac{abc}{4\Delta}\right)$, were essentially known to him.[2] In order to solve a triangle Hipparchus and other early writers always supposed it inscribed in a circle. The sides were then considered as chords, and these were computed as functions of the radius. In this way the table of chords was of special value. Triangles on a sphere were always decomposed into right-angled triangles, and these were solved separately. Although not mentioning the subject of spherical triangles in any of his works now extant, Hipparchus solves a certain problem in which he must have used the equivalent of the formula $\tan b = \cos A \tan c$, where $C = 90°$, and both he and Ptolemy (*c.* 150) knew the relation which we express by the equation $\sin^2 A + \cos^2 A = 1$.[3]

The treatise of Theodosius of Tripoli (*c.* 100) on the sphere[4] may be passed with mere mention, since it contains no work on trigonometry.

Heron of Alexandria. Although Heron (*c.* 50?)[5] showed much ingenuity in his mensuration of the triangle, and was thoroughly conversant with the art of surveying as practiced in Egypt, it cannot be said that he gave any evidence of appreciating the significance of trigonometry. He made use of certain rules which we should express in formulas for finding the area of regular polygons, giving in each case the product of the square of a side by a certain number, and these rules afford some evidence of a kind of prognosis of trigonometric functions. That

[1] On his astronomical work, see J. B. J. Delambre, *Histoire de l'astronomie ancienne*, I, 106 (Paris, 1817); P. Tannery, *Recherches sur l'histoire de l'astronomie ancienne*, Paris, 1893.

[2] That they are essentially involved in Euclid's *Elements*, VI, 16, is shown in the Simson additions, Props. C and D. See Heath's *Euclid*, Vol. II, pp. 224, 225.

[3] Heath, *History*, II, 259.

[4] Latin ed., Paris, 1529; Greek ed., Beauvais, 1558. See Volume I, page 125.

[5] Or possibly as late as *c.* 200. See Volume I, page 125.

is, taking A as the area of a regular n-gon and s_n as the side, he stated that the following relations exist:

$$A_3 = \tfrac{1}{3}\tfrac{3}{0} s_3^2, \qquad A_8 = \tfrac{2}{6}\tfrac{9}{0} s_8^2,$$


$$A_5 = \tfrac{5}{3} s_5^2, \quad \text{or} \quad \tfrac{1}{7}\tfrac{2}{0} s_5^2, \qquad A_9 = \tfrac{5}{8}\tfrac{1}{0} s_9^2, \quad \text{or} \quad \tfrac{3}{6}\tfrac{8}{0} s_9^2,$$

$$A_6 = \tfrac{1}{5}\tfrac{3}{0} s_6^2, \qquad A_{10} = \tfrac{1}{2}\tfrac{5}{0} s_{10}^2,$$

$$A_7 = \tfrac{4}{1}\tfrac{3}{0} s_7^2, \qquad A_{12} = \tfrac{4}{1}\tfrac{5}{0} s_{12}^2.$$

Since $A_n = \frac{1}{4} n s_n^2 \cot \frac{180^\circ}{n}$, it might be inferred that Heron had some knowledge of $\cot \frac{180^\circ}{n}$, but there is nothing in the coefficients to indicate this knowledge.[1]

Menelaus. About 100 A.D. the astronomer Menelaus of Alexandria, then living in Rome, took up the study of spherical triangles, a subject which, as we have seen, may have occupied the attention of Hipparchus. He wrote a work in six



books on chords, and although this is lost we have his treatise on spherics,[2] which not only forms the oldest known work on spherical trigonometry but reveals a remarkable knowledge of geometry and trigonometry in general.


[1] P. Tannery, "Arithmétique des Grecs dans l'Héron d'Alexandrie," *Mémoires de la Soc. des sciences de Bordeaux*, IV; *Mémoires Scientifiques*, I, 189; Heath, *History*, II, 326.

[2] *Menelai Sphaericorum Libri III*, translated by Maurolycus from Arabic and Hebrew sources and published at Messina in 1558. Mersenne (*c.* 1630) published an edition in 1644, and Halley's edition appeared posthumously at Oxford in 1758. For other editions see A. A. Björnbo, "Studien über Menelaos' Sphärik," *Abhandlungen*, XIV, 1, especially p. 17. See Heath, *History*, II, 261.

In the plane and spherical triangles shown on page 606 he proved the following relations:

Plane Triangle	*Spherical Triangle*
$\frac{CE}{AE} = \frac{CF}{DF} \cdot \frac{DB}{AB}$	$\frac{cd\,2\,CE}{cd\,2\,AE} = \frac{cd\,2\,CF}{cd\,2\,DF} \cdot \frac{cd\,2\,DB}{cd\,2\,AB}$
$\frac{CA}{AE} = \frac{CD}{DF} \cdot \frac{FB}{BE}$	$\frac{cd\,2\,CA}{cd\,2\,AE} = \frac{cd\,2\,CD}{cd\,2\,DF} \cdot \frac{cd\,2\,FB}{cd\,2\,BE}$

where *cd* 2 *CE* stands for the chord of twice the arc *CE*, that is, for what we call 2 sin *CE*. Since six quantities are involved in each equation, this was known in the Middle Ages as the *regula sex quantitatum* and was looked upon as the fundamental theorem of the Greek trigonometry. Whether it is due to Menelaus, to Hipparchus, or possibly to Euclid is a matter of dispute, but it is found first in definite form in the *Spherics* of Menelaus,[1] the proposition on the plane triangle being a lemma for the other one.



Menelaus also gave a *regula quatuor quantitatum*, as follows:[2] If the two triangles *ABC* and *DEF* have $\angle A = \angle D$ and $\angle C = \angle F$ then

$$\frac{cd\,2\,AB}{cd\,2\,BC} = \frac{cd\,2\,DE}{cd\,2\,EF}.$$

Ptolemy. The original contributions of Ptolemy (*c.* 150) to trigonometry are few, if any; but we are greatly indebted to him for his summary, in the *Almagest*, of the theorems known to Hipparchus.[3] Like other Greek writers, he used chords of angles instead of sines, but the idea of the sine seems to have been in his mind.[4] He extended the table of chords begun by Hipparchus, and it is quite probable that this is the source of the table of sines used by the early Hindu writers.

[1] On this point consult M. Chasles, *Aperçu historique*, 2d ed., 291; Delambre, *Histoire de l'astronomie ancienne*, I, 245 (Paris, 1817); A. A. Björnbo, *Abhandlungen*, XIV, 96, 99; Heath, *History*, II, 266.

[2] For other features see Braunmühl, *Geschichte*, I, 17; A. A. Björnbo, *loc. cit.*, p. 124. [3] Heath, *History*, II, 276. [4] *Ibid.*, II, 283.


Hindu Trigonometry. Although the Hindus had already produced the *Sūrya Siddhānta* (*c.* 400), and although this work treated of the ancient astronomy, gave a table of half chords apparently based, as stated above, upon Ptolemy's work, and showed some knowledge of trigonometric relations,[1] it was not until Āryabhaṭa wrote his *Āryabhaṭīyam* (*c.* 510) that we had in Oriental literature a purely mathematical treatise containing definite traces of the functions of an angle. In this work he speaks of the half chord, as the *Sūrya Siddhānta* had done before him.[2]

The subsequent work of the Hindus was concerned chiefly with the construction of tables, and this will be mentioned later.[3]

Arab and Persian Trigonometry. The chief interest that the Arab and Persian writers had in trigonometry lay, as with their predecessors, in its application to astronomy. On this account we find a growing appreciation of the science, beginning with the founding of the Bagdad School and extending to the close of the Mohammedan supremacy in scientific matters.

The chief Arab writer on astronomy was Albategnius[4] (*c.* 920), who ranked as the Ptolemy of Bagdad. Like the Hindus, he used half chords instead of chords. He also gave the rule for finding the altitude of the sun, which we express by the formula

$$x = \frac{l \sin(90° - \phi)}{\sin \phi},$$



which is simply equivalent to saying that

$$x = l \cot \phi,$$

but there is no evidence to show that he had any real knowledge of spherical trigonometry.

[1] See Volume I, pages 34, 145. There is a translation by Burgess in the *Journal of the American Oriental Society*, VI.

[2] L. Rodet, *Leçons de Calcul d'Âryabhata*, pp. 11, 24 (Paris, 1879).

[3] Colebrooke, *Aryabhaṭa*, pp. 90 n., 309 n.

[4] Al-Battânî. His work on the movements of the stars was translated by Plato of Tivoli (*c.* 1120) under the title *De motu stellarum*. It is known to us through the writings of Regiomontanus, and was published at Nürnberg in 1537. There was also a Bologna edition of 1645.

Abû'l-Wefâ (*c.* 980) did much to make the *Almagest* known, computed tables with greater care than his predecessors, and began a systematic arrangement of the theorems and proofs of trigonometry. With him the subject took on the character of an independent science.

It was, however, Naṣîr ed-dîn al-Ṭûsî (*c.* 1250), a Persian astronomer, who wrote the first work[1] in which plane trigonometry appears as a science by itself.

Ulugh Beg (*c.* 1435) of Samarkand was better known as an astronomer than as a writer on trigonometry, but the tables of sines and tangents computed under his direction helped to advance the science.

Arab Influence in Europe. With the decline of Bagdad the study of trigonometry assumed greater importance in Spain, particularly as related to those spherical triangles needed in the work in astronomy. The most important writers were the astronomers Ibn al-Zarqâla[2] (*c.* 1050), who constructed a set of tables, and Jabir ibn Aflah[3] (*c.* 1145). In the 13th century Alfonso X (*c.* 1250) directed certain scholars at Toledo to compute a new set of tables, chiefly for astronomical purposes; these Alfonsine Tables were completed *c.* 1254[4] and were long held in high esteem by later astronomers.

Fibonacci (1220) was acquainted with the trigonometry of the Arabs and, in his *Practica Geometriae*, applied the subject to surveying.

Peurbach and Regiomontanus. By the 14th century England knew the Arab trigonometry, and in the 15th century, thanks largely to Peurbach (*c.* 1460), who computed a new table of sines, and to his pupil Regiomontanus (*c.* 1464), European scholars in general became well acquainted with it. The work

[1] *Shakl al-qaṭṭâ'* (*Theory of Transversals*). There was a French translation published at Constantinople in 1891.

[2] Ibrâhîm ibn Yaḥyâ al-Naqqâsh, Abû Isḥâq, known as Ibn al-Zarqâla, or, in the translations, as Arzachel. He lived in Cordova.

[3] Or Jeber (Geber) ibn Aphla, of Seville. The German transliteration is Dschabir ibn Aflah. His astronomical work was published at Nürnberg in 1543

[4] See page 232 and also Volume I, page 228.

of Regiomontanus[1] had great influence in establishing the science as independent of astronomy. He computed new tables and may be said to have laid the foundation for the later works on plane and spherical trigonometry. In this general period there were also various minor writers, like Leonardo of Cremona (*c.* 1425), but they contributed little of value.[2]

Copernicus (*c.* 1520) completed some of the work left unfinished by Regiomontanus and embodied it in a chapter *De Lateribus et Angulis Triangulorum*, later (1542) published separately by his pupil Rhæticus.

Influence of Printed Books. The first printed work on the subject may be said to be the *Tabula directionum* of Regiomontanus, published at Nürnberg before 1485.[3]

The first book in which the six trigonometric functions were defined as functions of an angle instead of an arc, and substantially as ratios, was the *Canon doctrinae triangulorum* of Rhæticus (Leipzig, 1551), although it gives no names for $\sin\phi$, $\cos\phi$, and $\csc\phi$ except *perpendiculum*, *basis*, and *hypotenusa*.[4] Rhæticus was the first to adopt the semiquadrantal arrangement of the tables, giving the functions to 45° and then using the cofunctions. He found $\sin n\phi$ in terms of $\sin\phi$, $\sin(n-1)\phi$, and $\cos(n-2)\phi$, a subject elaborated by Jacques Bernoulli[5] (1702).

Vieta and his Contemporaries. Vieta (*c.* 1580) added materially to the analytic treatment of trigonometry. He also computed $\sin 1'$ to thirteen figures and made this the basis for the rest of the table. With him begins the first systematic development of the calculation of plane and spherical triangles by the

[1] *De triangulis omnimodis Libri V*, written *c.* 1464, first printed at Nürnberg in 1533. He also edited Ptolemy's *Almagest*, the first edition appearing at Venice in 1496. On his indebtedness to Naṣîr ed-dîn and others see A. von Braunmühl, "Nassîr Eddîn Tûsi und Regiomontan," *Abh. der Kaiserl. Leop.-Carol. Deutschen Akad. der Naturforscher*, LXXI, p. 33 (Halle, 1897).

[2] See J. D. Bond, *Isis*, IV, 295.

[3] Second ed., Venice, 1485; 3d ed., Augsburg, 1490. See Hain, 13,799.

[4] On his double use of this term for secant and cosecant, see Braunmühl, *Geschichte*, I, 147.

[5] See page 629; Tropfke, *Geschichte*, II (1), 229.

aid of all six functions. In one of his tracts there appears the important formula

$$\frac{a+b}{a-b}=\frac{\tan\frac{1}{2}(A+B)}{\tan\frac{1}{2}(A-B)},$$

which had already been discovered by Fincke, as mentioned below.

Albert Girard published at The Hague in 1626 a small but noteworthy work on trigonometry, and in this he made use of the spherical excess in finding the area of a spherical triangle. This was also given in his algebra of 1629. It also appeared at about the same time in Cavalieri's *Directorium generale* (Bologna, 1632) and, a little later, in his *Trigonometria plana et spherica*[1] (Bologna, 1643).

Thomas Fincke,[2] a Danish mathematician, published an important work, the *Geometria Rotundi*, in Basel in 1583 (2d ed., 1591). He gave the law relating to $a+b:a-b$, expressing it as

$$\frac{\frac{1}{2}(a+b)}{\frac{1}{2}(a+b)-b}=\frac{\tan\frac{1}{2}(180^\circ-C)}{\tan\left[\frac{1}{2}(180^\circ-C)-B\right]}.$$

The equivalent of our present form is due to Vieta, as already stated.

Pitiscus (1595) published an important trigonometry in which he corrected the tables of Rhæticus and modernized the treatment of the subject. In this work the word "trigonometry" appears for the first time as the title of a book on the subject.

British Writers. Besides his invention of logarithms, which has already been considered, Napier replaced the rules for spherical triangles by one clearly stated rule, the Napier Analogies,[3] published posthumously in his *Constructio* (Edinburgh, 1619).

Oughtred's trigonometry appeared in 1657. In this work he attempted to found a symbolic trigonometry; and although

[1] "In omnibus verò triangulis sphaericis tres eorum anguli simul sumpti superant duos rectos. Et excessus eorum est . . ." (p.29). He adds: "quod ego probaui in meo Directorio P. 3, Cap. 8."

[2] See N. Nielsen, *Matematiken i Danmark, 1528–1800* (Copenhagen, 1912).

[3] For a discussion of this subject see Braunmühl, *Geschichte*, II, 26.

algebraic symbolism was now so advanced as to make this possible, the idea was not generally accepted until Euler's influence was exerted in this direction in the 18th century.

John Newton (1622-1678) published in 1658 a treatise on trigonometry[1] which, while based largely on the works of Gellibrand and other writers, was the most complete book of the kind that had appeared up to that time. Newton and Gellibrand even went so far as to anticipate our present tendency by giving tables with centesimal divisions of the angle.

The greatest contribution to trigonometry made by John Wallis (1616–1703) was probably his encouragement of the statement of formulas by equations instead of by proportions, and his work on infinite series. The former advanced the analytic feature and the latter made possible the calculation of functions by better methods.

Sir Isaac Newton[2] (1642–1727) made many improvements in trigonometry, as in all other branches of mathematics. He expanded $\sin^{-1}x$, or arc sin x, in series, and by reversion he then deduced a series for $\sin x$. He also communicated to Leibniz the general formulas for $\sin nx$ and $\cos nx$.

The first to derive general formulas for $\tan nx$ and $\sec nx$ directly from the right-angled triangle was a French writer, Thomas-Fantet de Lagny (*c.* 1710). He was also the first to set forth in any clear form the periodicity of the functions. The word "goniometry" was first used by him (1724), although more in the etymological sense of mere angle measure than is now the case.

The Imaginary recognized in Trigonometry. The use of the imaginary in trigonometry is due to several writers of the first half of the 18th century. Jean Bernoulli discovered (1702) the relation between the arc functions[3] and the logarithm of an imaginary number. In his posthumous work of 1722 Cotes showed that

$$\phi i = \log(\cos\phi + i\sin\phi),$$

[1] *Trigonometria Britannica, or the doctrine of triangles in two books*, London. 1658.

[2] Braunmühl, *Geschichte*, II, chap. iii.

[3] Such as arc sin x, or $\sin^{-1}x$.

although no writers at that time used this particular symbolism. As early as 1707 De Moivre knew the relation

$$\cos\phi = \tfrac{1}{2}(\cos n\phi + i\sin n\phi)^{\frac{1}{n}} + \tfrac{1}{2}(\cos n\phi - i\sin n\phi)^{\frac{1}{n}},$$

which is obviously related to the theorem

$$(\cos\phi + i\sin\phi)^n = \cos n\phi + i\sin n\phi,$$

published in 1722 and usually called by his name.[1]

Euler gave (1748) the equivalent of the formula

$$e^{\phi i} = \cos\phi + i\sin\phi,$$

but this was no longer new. His use of i for $\sqrt{-1}$ (1777) was, however, a welcome contribution. Lambert (1728–1777) extended this phase of trigonometry and developed the theory of hyperbolic functions which Vincenzo Riccati had already (*c.* 1757) suggested and which Wallace[2] elaborated later.

Functions as Pure Number. The first writer to define the functions expressly as pure number was Kästner[3] (1759), although they had already been used as such by various writers.[4]

Trigonometry becomes Analytic. As already stated, through the improvements in algebraic symbolism European trigonometry became, in the 17th century, largely an analytic science, and as such it entered the field of higher mathematics.


In the Orient, however, the science continued in its primitive form, largely that of shadow reckoning, until the Jesuits carried European methods to China, beginning about the year 1600. From that time on the Western influence generally prevailed, not merely in such centers as Nanking and Peking but also, somewhat later, in Japan.

[1] Braunmühl, *Geschichte*, II, 76. [2] See Volume I, page 458.

[3] A. G. Kästner, *Anfangsgründe der Arithmetik Geometrie ebenen und sphärischen Trigonometrie und Perspectiv*, Göttingen, 1st ed., 1759; 2d ed., 1764; 3d ed., 1774. He remarks: "Bedeutet also nun x den Winkel in Graden ausgedruckt, so sind die Ausdrückungen sin x; cos x; tang x u. s. w. Zahlen, die für jeden Winkel gehören" (3d ed., p. 380).

[4] Thus Regiomontanus (c. 1463) speaks of the tangents as *numeri*. This occurs in his *tabula foecunda*, prepared for astronomical purposes, and so called "quod multifariam ac mirandam utilitatem instar foecundae arboris parare soleat."

With this brief summary of the development of the science we may proceed to a consideration of a few of the special features which the teacher will meet in elementary trigonometry.



JAPANESE TRIGONOMETRY OF *c.* 1700

From Murai Masahiro's *Riochi Shinan*, early in the 18th century, showing European influence

2. Trigonometric Functions

Sine. The most natural function for the early astronomer to consider was the chord of an arc of a circle having some arbitrary radius. Without any good notation for fractions it was not convenient to take a radius which would give difficult fractional values for the approximate lengths of the chords. A convenient radius, such as 60, being taken, the chord of the arc, being considered purely as a line, was the function first studied by the astronomers.

The first table of chords of which we have any record was computed by Hipparchus (*c.* 140 B.C.), but this table is lost and we have no knowledge as to its extent or its degree of accuracy. The next table of chords of which we have good evidence

was that of Menelaus (*c.* 100), but this is also lost, although his work on spherics shows his use of the function. The third important table of chords is that of Ptolemy (*c.* 150). He divided the circle into 360° and the diameter into 120 equal parts,[1] a relation doubtless suggested both by the numerous factors of 120 and, since $3 \times 120 = 360$, by the ancient use of 3 for π. Influenced like Hipparchus by Babylonian precedents, he used sexagesimal fractions, the radius consisting of 60 *moirai*, each *moira* of 60 minutes, and so on.[2]

Origin of the Sine. A special name for the function which we call the sine is first found in the works of Āryabhaṭa (*c.* 510). Although he speaks of the half chord,[3] he also calls it the chord half[4] and then abbreviates the term by simply using the name *jyā* or *jīva* (chord). He follows Ptolemy in dividing the circle into 360°, and gives a table of sines, of which a portion is shown on page 626.

It is further probable, from the efforts made to develop simple tables, that the Hindus were acquainted with the principles which we represent by the formulas

$$\sin^2\phi + \cos^2\phi = 1,$$

$$\sin^2\phi + \operatorname{versin}^2\phi = 4\sin^2\frac{\phi}{2},$$

and

$$\sin\frac{\phi}{2} = \sqrt{\frac{1-\cos\phi}{2}},$$

the last two of these appearing in the *Pañca Siddhāntikā* of Varāhamihira (*c.* 505).

The table of sines given by Āryabhaṭa was reproduced by Brahmagupta (*c.* 628), but he did nothing further with trigonometry. Bhāskara (*c.* 1150), however, in his *Siddhānta Śiromāni*, gave a method of constructing a table of sines for every degree.[5]

[1] Τρήματα (*tre'mata*, literally "holes," and hence the holes, or pips, of dice). These parts were also called μοῖραι (*moi'rai*, parts), usually translated as degrees.

[2] The minutes were ἑξηκοστὰ πρῶτα (*hexekosta' pro'ta*), first sixtieths; the seconds were ἑξηκοστὰ δεύτερα (*hexekosta' deu'tera*), second sixtieths.

[3] *Ardhā-jyā*, *ardhajyā*, or *ardhā-djyā*.

[4] *Jyā-ardhā*.

[5] Tropfke, *Geschichte*, II (1), 192.

Name for Sine. The *jyā*[1] of Āryabhaṭa found its way into the works of Brahmagupta as *kramajyā*, that is, straight sine, or *sinus rectus*, as distinguished from the *sinus versus*, the versed sine. This was changed to *karaja* when it went over into Arabic, and as such appears in the Bagdad School of the 9th century. In particular, al-Khowârizmî used it in the extracts which he made from the *Brahmasiddhāntā* of Brahmagupta, probably the work known as the *Sindhind*. It is also found, with natural variants in form, in the writings of the Spanish Arab Ibn al-Zarqâla[2] (*c.* 1050).

The sine also appears in the *Pañca Siddhāntikā* of Varāhamihira (*c.* 505), where a table is computed with the Greek diameter of 120. Indeed, the probability of Greek influence upon the methods used by the Hindus is very strong.

The Arabs used the meaningless word *jîba*, phonetically derived from the Hindu *jyā*. The consonants of the word permitted the reading *jaib*, which means bosom, and so this was adopted by later Arabic writers.

Sine in Latin Works. When Gherardo of Cremona (*c.* 1150) made his translations from the Arabic[3] he used *sinus* for *jaib*, each word meaning a fold,[4] and this usage, possibly begun even earlier, was followed by other European scholars. The word "chord" was also used for the same purpose.[5]

[1] It has such forms as *djyā*, *dschyā* (German transliteration), *jīva* and *jība*.

[2] Or al-Zarkâla, the Latin Arzachel. In the Latin translation there is a chapter "De inventione sinus et declinationis per Kardagas." On his use of *kardagas* see Braunmühl, *Geschichte*, I, 78; on such variants as *gardaga* and *cada*, see *ibid.*, page 102; and on such special uses as *cardaga* for arc 15°, see *ibid.*, pages 110, 120.

[3] *E.g.*, the *Canones sive regulae super tabulas Toletanas* of al-Zarqâla: "Sinus cuius libet portionis circuli est dimidium corde duplicis portionis illius." See also the *Astronomia Gebri filii Affla Hispalensis*, which Apianus edited and published at Nürnberg in 1533.

[4] *Jaib* means bosom, breast, bay; and *sinus* means bosom, bay, a curve, the fold of the toga about the breast, the land about a gulf, a fold in land.

[5] Thus Plato of Tivoli: ". . . sive mentionē cordarū de medietatis cordis opportere intelligi, nisi aliquo proprio nomine signauerimus, quod & cordā integram appellabimus." On this subject, which has caused much controversy, see Braunmühl, *Geschichte*, I, 49, with bibliography. For the absurd suggestion that sinus = s. ins. = semissis inscriptae [chordae], with bibliography, see Tropfke, *Geschichte*, II (1), 212.

Abû'l-Wefâ (*c.* 980) defined clearly the chord, sine, and versed sine (sinus versus). He showed that

$$\sin\phi = \tfrac{1}{2}\,\mathrm{cd}\,2\phi,$$

$$\frac{2r - \mathrm{cd}\,(180^\circ - \phi)}{\mathrm{cd}\,\frac{\phi}{2}} = \frac{\mathrm{cd}\,\frac{\phi}{2}}{r}, \quad \text{our} \quad 2\sin^2\frac{\phi}{2} = 1 - \cos\phi,$$

$$\frac{\mathrm{cd}\,\phi}{\mathrm{cd}\,\frac{\phi}{2}} = \frac{\mathrm{cd}\left(180^\circ - \frac{\phi}{2}\right)}{r}, \quad \text{our} \quad \sin\phi = 2\sin\frac{\phi}{2}\cos\frac{\phi}{2},$$

and $\sin(\phi \pm \phi') = \sqrt{\sin^2\phi - \sin^2\phi\sin^2\phi'} \pm \sqrt{\sin^2\phi' - \sin^2\phi\sin^2\phi'}$.

He also constructed a table of sines for every 15′.

Ibn al-Zarqâla, mentioned on page 616, computed a table[1] of sines and versed sines, using as his arbitrary radius 150′ and also, following Ptolemy, 60$^\mu$, where μ stands for μοῖραι (*moi'rai*)[2] and is here used so as not to confuse units of line measure with degrees of angle measure. It is probable that such tables were known to Rabbi ben Ezra (*c.* 1140).

In his *Practica Geometriae* (1220) Fibonacci defines the *sinus rectus arcus* and *sinus versus arcus*,[3] and from that time on the terms were generally recognized in the Middle Ages. Tables of sines were given in various works thereafter,[4] so that their use became common.

Other Names for Sine. The term "sine" was not, however, universally recognized, for Rhæticus (*c.* 1560) preferred *perpendiculum*. Of the special terms which appeared from time to time there may be mentioned the *sinus totus* and *sinus perfectus*, both of which were used for sin 90°.[5]

[1] See page 616, n. 2. [2] See page 232, n. 3.

[3] ". . . .be. uocatur sinus rectus utriusque arcus .ab. et .bc.; et recta .ae. uocatur sinus uersus arcus .ab." (*Scritti*, II, 94).

[4] *E.g.*, Johannes de Lineriis (*c.* 1340). Ulugh Beg's tables (*c.* 1435) were computed for every minute of arc.

[5] *E.g.*, by Johann von Gmünden (*c.* 1430). Regiomontanus (1463) used *sinus totus rectus*. Rhæticus (*c.* 1550) used *sinus totus*, as did most other writers of the time.

Abbreviations for Sine. The first writer to make any general use of a satisfactory abbreviation for sine was Girard (1626). He designated the sine of A by A, and the cosine of A by a.[1] As early as 1624 the contraction *sin* appears on a drawing representing Gunter's scale, but it does not appear in Gunter's work published in that year.[2] In a trigonometry published by Richard Norwood (London, 1631) the author states that "in these examples s stands for *sine*: t for *tangent*: sc for *sine complement*: tc for *tangent complement*: sec for *secant*." The first writer to use the symbol *sin* for *sine* in a book seems to have been the French mathematician Hérigone (1634). Cavalieri (1643) suggested *Si*, and in the 1647 edition of Oughtred the symbol S is used. In 1654, Seth Ward, Savilian professor of astronomy at Oxford, himself a pupil of Oughtred's, used s, taking S' for the *sinus complementi*. Oughtred's symbol was adopted by various English writers of the 17th century.[3] The symbols $\sin^{-1}x$, $\cos^{-1}x$, $\cdots$, for arc $\sin x$, arc $\cos x$, $\cdots$, were suggested by the astronomer Sir John F. W. Herschel (1813).

Versed Sine. The next function to interest the astronomer was neither the cosine nor the tangent, but, strange as it may seem to us, the versed sine. This function, already occasionally mentioned in speaking of the sine, is first found in the *Sūrya Siddhānta* (*c.* 400) and, immediately following that work, in the writings of Āryabhaṭa, who computed a table of these functions. A sine was called the *jyā*; when it was turned through 90° and was still limited by the arc, it became the turned (versed) sine, *utkramajyā*,[4] so that the versin $\phi = 1 - \cos\phi$.

From India it passed over to the Arab writers, and Albategnius (al-Battânî, *c.* 920), for example, expressly states that he uses the expression "turned chord"[5] for the versed sine.

Since the early writers were given to fanciful resemblances and spoke of the bow (ACB) and string (AB), or the *arcus*


[1] Cantor, *Geschichte*, II (2), 709; Tropfke, *Geschichte*, II (1), 217.

[2] F. Cajori, "Oughtred's Mathematical Symbols," *Univ. of Calif. Pub. in Math.*, I, 185. Consult this article also for the rest of this topic.

[3] *E.g.*, Sir Charles Scarburgh (1616–*c.* 1696), a name also given as Scarborough.

[4] Or *utramadjyā*. [5] In some of the Latin translations, *chorda versa*.

and *chorda*, it was natural for them to speak of the versed sine as the arrow. So the Arabs spoke of the *saḥem*, or arrow, and the word passed over into Latin as *sagitta*, a term used by Fibonacci (1220)[1] to mean versed sine and commonly found in the works of other medieval writers.[2] Among the Renaissance writers there was little uniformity. Maurolico (1558) used *sinus versus major* of ϕ to designate versin $(180° - \phi)$, but others preferred the briefer term *sagitta*.



Cosine. Since the Greeks used the chord of an arc as their function, they had no special use for the chord of the complement. When, however, the right-angled triangle was taken as the basis of the science, it became convenient to speak of the sine of a complement angle. Thus there came into use the *koṭijyā* of Āryabhaṭa (*c.* 510),[3] although the sine of $90° - \phi$ commonly served the purpose then as it did later with the Arabs.[4] Even when a special name became necessary it was developed slowly. Plato of Tivoli (*c.* 1120) used *chorda residui* or spoke of the complement angle.[5] Regiomontanus (*c.* 1463) used *sinus rectus complementi.* Rhæticus (1551) preferred *basis*, Vieta (1579) used *sinus residuae*, Magini (1609) used *sinus secundus*, while Edmund Gunter (1620) suggested *co.sinus*, a term soon modified by John Newton (1658) into *cosinus*, a word which was thereafter received with general favor. Cavalieri (1643) used the abbreviation *Si.2*; Oughtred, *s co arc*; Scarburgh, *c.s.*; Wallis,[6] Σ; William Jones (1706), *ś*; and Jonas Moore[7] (1674), *Cos.*, the symbol generally adopted by later writers.

[1] *Scritti*, II, 94, ll. 10, 16, 18.

[2] *E.g.*, Levi ben Gerson (*c.* 1330), in his *De sinibus, chordis, et arcubus*, where his translator also uses *sinus versus*; and Johann von Gmünden (*c.* 1430).

[3] The possible *seqt*, *śḳd*, or *seqet* of the Egyptians (p. 600). Also transliterated *koṭidjyā*.

[4] As with Albategnius (al-Battânî) and others.

[5] "Quod ad perficiendum 90 deficit."


[6] "Σ, co-sinus, seu sinus complementi" (*Opera*, 1693, II, 591).

[7] *Bibl. Math.*, I (3), 69.

Tangent and Cotangent. While the astronomers found the chord and sine the functions most useful in their early work, and so developed them first, the more practical measurements of heights and distances first required the tangent and cotangent,—the gnomon and shadow respectively. It is possible that Ahmes (*c.* 1550 B.C.) knew the tangent, but in any case we know that shadow reckoning was an early device for finding heights, and that it was related to the sundial which Anaximander (*c.* 575 B.C.) introduced into Greece. Unlike the sine and cosine, the tangent and cotangent developed side by side, the reason being that the gnomon and shadow were equally important, the complementary feature playing no part at first. The Greeks, however, made no use of these functions of an angle, so far as we know, except as Thales measured the heights of pyramids by means of shadows and similar triangles.

The Umbra Recta and Umbra Versa. The *Sūrya Siddhānta* (*c.* 400) and other Hindu works speak of the shadow, particularly in connection with astronomical rules, but it was the Arabs who first made any real use of it as a function.

It was Ahmed ibn ʿAbdallâh,[1] commonly known as Habash al-Ḥâsib, "the computer" (*c.* 860), who constructed the first table of tangents and cotangents,[2] but it exists only in manuscript. The Arab writers distinguished the straight shadow, translated by the later medieval Latin writers as *umbra*, *umbra recta*, or *umbra extensa*, and the turned shadow, the *umbra versa* or *umbra stans*, the terms varying according as the gnomon was perpendicular to a horizontal plane, as in ordinary dials, or to a vertical wall, as in sundials on a building. They were occasionally called the horizontal and vertical shadows.[3] The shadow names were also used by most of the later Latin authors and by writers in general until relatively modern times, being frequently found as late as the 18th century.



[1] Or al-Mervazî. See Volume I, page 174.

[2] These are given in a MS. of his astronomical tables preserved at Berlin. See Suter, *Abhandlungen*, X, 209.

[3] *E.g.*, by Abû'l-Ḥaṣan Ali ibn 'Omar al-Marrâkoshî, of Morocco (*c.* 1260).

These functions do not seem to have interested the western Arab writers, no trace of either *umbra* being found in the works of Jabir ibn Aflah (*c.* 1145).

The terms *umbra recta* and *umbra versa* were not used by Gerbert, but Robertus Anglicus (*c.* 1231) speaks of the *umbra*, so that by his time it had come to be somewhat recognized. Thereafter the names *umbra recta* and *umbra versa* were in fairly common use.

Table of Shadows. The first writer whose table of shadows is generally known is Albategnius (al-Battânî, *c.* 920), the table giving the cotangents for each degree of the quadrant.[1]

Abû'l-Wefâ (*c.* 980) constructed a table of tangents for every 15′, the first table of tangents that is known to us; and about this time there was computed a table of cotangents for every 10′. Under the direction of Ulugh Beg (*c.* 1435) there was prepared a table of tangents for every 1′ from 0° to 45° and of every 5′ from 45° to 90°, but his table of cotangents was constructed only for every 1°.[2]

Names and Symbols. Although Rhæticus (1551) did not use these common names for tangent and cotangent, he defined each as a ratio and gave the most complete table that had appeared up to that time.

Vieta (*c.* 1593) called the tangent the *sinus foecundarum* (abridged to *foecundus*[3]) and also the *amsinus* and *prosinus*. It was not until Thomas Fincke wrote his *Geometria Rotundi* (1583) that the term "tangent" appeared as the equivalent of *umbra versa*.[4] The name was adopted by Pitiscus (1595), and the reputation of this great writer gave it permanent standing. Magini (1609) used *tangens secunda* for cotangent. The term *cotangens* was first used for this function by Edmund Gunter (1620).

[1] That is, he gave the value of $u = l \cdot \frac{\cos\phi}{\sin\phi}$ for $\phi = 1°, 2°, \cdots, l$ being the length of the gnomon.

[2] Braunmühl, *Geschichte*, I, 75.

[3] On the origin of the term see page 613 and Braunmühl, *Geschichte*, I, 161 n.; Tropfke, *Geschichte*, II (1), 210.

[4] "Recta sinibus connexa est tangens peripheriae, aut eam secans" (*Geometria Rotundi*, p. 73 (Basel, 1583)).

As abbreviations for tangent and cotangent, Cavalieri (1643) used *Ta* and *Ta.2*; Oughtred (1657), *t arc* and *t co arc*; Sir Charles Scarburgh, *t.* and *ct.*; and Wallis (1693), T and τ. The abbreviation *tan*, as in $\overset{\text{tan}}{\text{A}}$, was first used by Girard (1626), and *Cot.* was suggested by Jonas Moore (1674), but even yet we have no generally accepted universal symbols for tangent and cotangent.

Secant and Cosecant. Since neither the astronomer nor the surveyor of early times had any need for the secant and cosecant, except as the hypotenuse of a right-angled triangle, these functions were developed much later than the others. The secant seems first to have been considered by al-Mervazî (Habash, *c.* 860), although the two functions first appear in definite form in the works of Abû'l Wefâ (*c.* 980), but without special names. Little was done with them by the Arabs, however, and it was not until tables for navigators were prepared in the 15th century that secants and cosecants appeared in this form.[1] Although Copernicus (1542) knew the secant, speaking of it as the *hypotenusa* and computing a set of values of these functions, it was his pupil Rhæticus who first included secants in a printed table. The secant and cosecant appear with the other four functions in his *Canon doctrinae triangulorum* (Leipzig, 1551), although Rhæticus speaks of each in that work as a hypotenuse. The name "cosecant" seems to have appeared first in his posthumous *Opus Palatinum* (1596). Maurolico (1558) included in his tables[2] the secants from 0° to 45°.

Names for Secant and Cosecant. The name "secant" was first used by Fincke (1583) and, although Vieta (1593) called this function the *transsinuosa*, the more convenient and suggestive name soon came into general use. The cosecant was called the *secans secunda* by Magini (1592) and Cavalieri (1643). Pitiscus (1613) gave the secants and cosecants in his tables, and since then they have been commonly found in similar publications.

[1] Braunmühl, *Geschichte*, I, 114; 115 n.
[2] *Tabula benefica*, in his work on spherics.

By way of abbreviations for secant and cosecant, Cavalieri (1643) used *Se* and *Se.2*; Oughtred (1657), *sc arc* and *sec co arc*; Wallis (1693), *s* and σ; but the more convenient symbol *sec*, suggested by Girard (1626) in the form $\overset{\text{sec}}{\text{A}}$ soon came into general use. There is as yet no international symbol for cosecant, *cosec* and *csc* both being used.

Relation between Functions. Although the functions themselves were not specifically named, various early writers make statements which involve in substance many of the relations that we now recognize. Thus the formula

$$\sin\phi = \sqrt{1 - \cos^2\phi}$$

or

$$\sin^2\phi + \cos^2\phi = 1$$

is essentially the Pythagorean Theorem and as such was known to the Greeks.

Abu'l-Wefâ (*c.* 980) knew substantially the formulas

$$\tan\phi : 1 = \sin\phi : \cos\phi,$$
$$\cot\phi : 1 = \cos\phi : \sin\phi,$$
$$\sec\phi = \sqrt{1 + \tan^2\phi},$$

and

$$\csc\phi = \sqrt{1 + \cot^2\phi}.$$

Rhæticus (1551) knew the relations

$$\sec\phi : 1 = 1 : \cos\phi$$

and

$$\csc\phi : 1 = 1 : \sin\phi.$$

Vieta (1579) gave the following proportions:[1]

$$1 : \sec\phi = \cos\phi : 1 = \sin\phi : \tan\phi,$$
$$\csc\phi : \sec\phi = \cot\phi : 1 = 1 : \tan\phi,$$

and

$$1 : \csc\phi = \cos\phi : \cot\phi = \sin\phi : 1.$$

3. Trigonometric Tables


Early Methods of Computing. The more important of the earliest trigonometric tables have been mentioned in connection with the several functions. A brief statement will now be

[1] Tropfke, *Geschichte*, II (1), 225.

made as to the general methods of computing these tables and as to the early printed tables themselves.

The first methods of which we have definite knowledge are those of Ptolemy (*c.* 150).[1] His computation of chords depends on four principles:

I. From the sides of the regular inscribed polygons of 3, 4, 5, 6, and 10 sides he obtained the following:



$$\text{cd } 36^\circ = 37^\mu\ 4'\ 55'',$$
$$\text{cd } 72^\circ = 70^\mu\ 32'\ 3'',$$
$$\text{cd } 60^\circ = 60^\mu,$$
$$\text{cd } 90^\circ = 84^\mu\ 51'\ 10'',$$

and

$$\text{cd } 120^\circ = 103^\mu\ 55'\ 23''.$$

In a semicircle as here shown, $\overline{BC}^2 + \overline{CA}^2 = \overline{AB}^2$, and so

$$\text{cd}(180^\circ - 36^\circ) = \text{cd } 144^\circ = \sqrt{120^2 - \text{cd}^2\ 36^\circ} = 114^\mu\ 7'\ 37''.$$

II. In an inscribed quadrilateral the sum of the rectangles of the two pairs of opposite sides is equal to the rectangle of the two diagonals. This is known as Ptolemy's Theorem and is found in the *Almagest*.[2]

III. The chord of a half arc can be found from the chord of the arc; that is, from cd 12° it is possible to find cd 6°, and then cd 3°, and so on.[3]

IV. By a scheme of interpolation it is possible to approximate the chord of $\frac{1}{3}\phi$, when cd ϕ is known.[4]

With the help of these principles Ptolemy was able to find the chords of all angles to a fair degree of approximation. Thus he found that

$$\text{cd } 1^\circ\ 0' = 1^\mu\ 2'\ 50'',$$

which would make

$$\sin 30' = \tfrac{1}{2}\,\text{cd } 1^\circ\ 0' = 0^\mu\ 31'\ 25'' = 0.0087268,$$

[1] The tables are given at the end of Lib. I, cap. ix, of the *Almagest*.

[2] Halma ed., I, 29; Heiberg ed., p. 36; Braunmühl, *Geschichte*, I, 19.

[3] Essentially he has $\sin\frac{\phi}{2} = \sqrt{\frac{1 - \cos\phi}{2}}$. For the mathematical discussion, see Braunmühl, *Geschichte*, I, 20.

[4] For mathematical discussion, see Braunmühl, *Geschichte*, I, 21; Tropfke, *Geschichte*, II (1), 296.

whereas our seven-place tables give it as 0.0087265. Abû'l-Wefâ (*c.* 980) computed this result as $0^{u}\ 31'\ 24''\ 55'''\ 54^{iv}\ 55^{v}$, —a value which is correct as far as the tenth decimal place.

द्वितीयाऽध्याय ३३

घटाने पर शेष २२५ होगा। इस २२५ को प्रथम ज्याई २२५ के साथ जोड़ देने से योगफल ४५८ होगा। यही द्वितीय ज्याई है ॥१५॥ उस द्वितीय ज्याई ४५८ को प्रथम ज्याई से भाग करके भागफल २ लेकर यह २ इस के साथ पूर्व द्वितीय ज्याई निष्कासन भाग फल से जो १ मिला है, जोड़ने से ३ होगा। इस ३ को उस भाजक २२५ से घटाने पर २२२ बचेगा, इसी २२२ को द्वितीय ज्याई ४५८ के साथ जोड़ने से ६७९ होगा, यही तृतीय ज्याई है। इसी प्रकार क्रमशः २४ ज्याई गणना करनी होगी ॥१६॥ किसी वृत्त के चतुर्थांश जिस का व्यासार्द्ध ३४३८ उस के ३४ अंश की ज्याई निम्नलिखित होंगी ॥

	अंश	वा	कला	ज्या		अंश	वा	कला	ज्या
प्रथम कोण	$1\frac{1}{2}$		२२५	२२५	१३ वां कोण	४८	॥	२४७५	२२६७
द्वितीय ,,	$3\frac{1}{2}$		४५०	४४८	१४ वां ,,	$52\frac{1}{2}$		२७००	२४४१
तृतीय ,,	$11\frac{1}{2}$		६७५	६७१	१५ वां ,,	$56\frac{1}{2}$		२८२५	२५८२
चतुर्थ ,,	१५		९००	८९०	१६ वां ,,	६०		३१५०	२७२८
पञ्चम ,,	$18\frac{3}{4}$		११२५	११०५	१७ वां ,,	६३	॥	३३७५	२८५९
षष्ठ ,,	$22\frac{1}{2}$		१३५०	१३१५	१८ वां ,,	$67\frac{1}{2}$		३६००	२९७८
सप्तम ,,	$26\frac{1}{4}$		१५७५	१५२०	१९ वां ,,	$71\frac{1}{2}$		३८२५	३०८४
अष्टम ,,	३०		१८००	१७१९	२० वां ,,	७५		४०५०	३१७७
नवम ,,	$33\frac{3}{4}$		२०२५	१९१५	२१ वां ,,	७८	॥	४७२५	३३७२
दशम ,,	$37\frac{1}{2}$		२२५०	२०८३	२२ वां ,,	$82\frac{1}{2}$		४९५०	३४०९
एकादश ,,	$41\frac{1}{2}$		२४७५	२२६७	२३ वां ,,	$86\frac{3}{4}$		५१७५	३४३१
द्वादश ,,	४५		२७००	२४३१	२४ वां ,,	९०		५४००	३४३८

पूर्वोक्त ज्याई परिमाण सब को उलटे प्रकार से ३४३८ व्यासार्द्ध से पृथक् पृथक् घटाने पर जो अङ्क घटाने से बचेंगे उन को उत्क्रमज्या कहते हैं। प्रति ३४ अंश में इस प्रकार उत्क्रमज्या हो जाती हैं। १६–२२ श्लोक तक ॥

मुनयोरन्ध्रयमला रसषट्कामुनीश्वराः। द्व्यष्टैकारूप-
षड्द्वस्राः सागरार्थहुताशनाः ॥२३॥ खर्तुवेदा नवाद्र्यर्धा
दिङ्नगास्त्र्यर्धकुञ्जराः। नगाम्बरवियच्चन्द्ररूपभूधरश-

FROM THE FIRST PRINTED EDITION OF THE SŪRYA SIDDHĀNTA

Printed at Meerut, India, *c.* 1867. This is the oldest Hindu work on astronomy. It shows the table of which a portion is given on page 626

A table of sines is given in the *Sūrya Siddhānta* (*c.* 400), and Āryabhaṭa (*c.* 510) gives a table of sines and versed sines. The following portion of the table of sines, substantially as in

some of the manuscripts of the *Sūrya Siddhānta* and as given also in Āryabhaṭa's work, will serve to show the degree of accuracy:

Arc	Sine	Modern Value
$3^\circ\ 45'$	$225'$	$224.84'$
$7^\circ\ 30'$	$449'$	$448.72'$
$11^\circ\ 15'$	$671'$	$670.67'$
15°	$890'$	$889.76'$

Āryabhaṭa's method of working out his table was to take $\sin 3^\circ\ 45'$[1] as equal to arc $3^\circ\ 45'$, and from this to find the sines of multiples of this angle by the rule already given in the *Sūrya Siddhānta*,[2]

$$\sin(n+1)\phi = \sin n\phi + \sin n\phi - \sin(n-1)\phi - \frac{\sin n\phi}{\sin \phi},$$

which is correct except for the last term.

Arab Methods. The early Arabs used the Hindu results, but later scholars developed original methods of attack. Of these, one of the best known is given by Mîram Chelebî (*c.* 1520)[3] in his commentary on Ulugh Beg (*c.* 1435).[4] He gives two methods, the first being somewhat similar to the one used by Ptolemy. The second method is interesting because it involves the approximate solution of a cubic equation of the form

$$ax - b = x^3.$$

European Computers. Of the later computers of the Middle Ages and early Renaissance, Regiomontanus (1546) stands at the head, but his methods were not new. Indeed, there was no particular originality shown in the computations from the time of Ptolemy to the invention of the modern methods based on series.

[1] Known by the special name *kramajyā*. [2] Braunmühl, *Geschichte*, I, 35.

[3] Mûsâ ibn Mohammed ibn Maḥmûd ibn Qâḍizâdeh al-Rûmî, a teacher in Gallipoli, Adrianople, and Brusa; died 1524/25.

[4] M. Woepcke, "Discussion de deux méthodes arabes pour déterminer une value approchée de sin 1°," *Journal de math. pures et appliquées*, XIX (1854), 153; A. Sédillot, *Prolégomènes des Tables astronomiques d'Ouloug-Beg*, Paris, 1853; Braunmühl, *Geschichte*, I, 72, with incorrect date.

In general, all ancient tables were constructed with Ptolemy's radius of 60; that is, the *sinus totus*, or sin 90°, was 60. This was due to the necessity of avoiding fractions in the period before the invention of decimals. The first to adopt the simpler form, sin 90° = 1, was Jobst Bürgi (*c.* 1600), but his tables computed on this basis are not extant. Although the invention of decimal fractions had now made the use of unity possible for the *sinus totus*, this idea was not fully appreciated until a memoir by de Lagny was written in 1719.[1] It was nearly thirty years later that the plan received its first great support at the hands of Euler.[2]

Early Printed Tables. Of the early printed tables there may be mentioned as among the more important the table of sines with the radius divided decimally, published by Apianus in 1533; the table of all six functions based on a semiquadrantal arrangement, published by Rhæticus in 1551, calculated to every 10′ and to seven places; Vieta's extension of the tables of Rhæticus to every minute (1579, but the printing began in 1571); the table of tangents by Reinhold (1511–1553) to every minute, printed in 1554; the table of all six functions, published in England by Blundeville in 1594; the *Opus Palatinum*, with the functions for every 10″ to ten decimal places, with tables of differences, compiled by Rhæticus and published by Valentin Otto (or Otho) in 1596. Dr. Glaisher, referring to the work of Rhæticus, speaks of him as "by far the greatest computer of pure trigonometrical tables" and as one "whose work has never been superseded." The *Opus Palatinum* was so named in honor of the elector palatine, Friedrich IV, who paid for its publication.[3]


The serious use of tables based upon the centesimal division of the angle was a result of the movement that led to the metric system. An elaborate set of such tables was prepared in Paris at about the close of the 18th century, and little by little the plan found favor. Such a set of tables appeared even in Japan

[1] *Histoire et Mémoires de l'Acad. d. sci.*, Paris, 1721, p. 144; 1726, p. 292; 1727, p. 284; 1729, p. 121.

[2] *Introductio in analysin infinitorum*, I, § 127. Lausanne, 1748.

[3] For a summary of such tables see the *Encyc. Brit.*, 11th ed., XXVI, 325.

as early as 1815, but it was not until the close of the 19th century that the idea took any firm hold upon the mathematics of Europe, and then with the French schools still in the lead.



CENTESIMAL TABLES OF JAPAN

From a manuscript of a work on trigonometry, by Miju Rakusai, written in 1815, showing a table of natural functions on the decimal division of the angle. This page shows the cotangents and cosines, beginning at the top with 0°

4. Typical Theorems

Addition Theorem of Sines. It is impossible, in the space allowed, to mention more than a few of the important theorems of trigonometry, and these will now be considered.

The Greeks knew essentially that

$$\sin(\phi \pm \phi') = \sin\phi \cos\phi' \pm \cos\phi \sin\phi'.$$

Stated as a proposition involving chords, it is probable that Hipparchus (*c.* 140 B.C.) knew it. It was certainly known to Ptolemy (*c.* 150), and it often bears his name. Bhāskara (*c.* 1150) also gives the theorem. As already stated on page 617, Abû'l-Wefâ (*c.* 980) gave it essentially under the form

$$\sin(\phi \pm \phi') = \sqrt{\sin^2\phi - \sin^2\phi \sin^2\phi'} \pm \sqrt{\sin^2\phi' - \sin^2\phi \sin^2\phi'}.$$

Functions of Multiple Angles. The formula

$$\sin 2\phi = 2 \sin\phi \cos\phi$$

is a corollary of the general case of $\sin(\phi + \phi')$. It is first expressly given as a rule by Abû'l-Wefâ, the form being, as already stated, chord ϕ : chord $\frac{1}{2}\phi$ = chord $(180° - \frac{1}{2}\phi) : r$.

Vieta[1] (1591) first gave the formulas

$$\sin 3\phi = 3\cos^2\phi \sin\phi - \sin^3\phi,$$
$$\cos 3\phi = \cos^3\phi - 3\sin^2\phi \cos\phi,$$

and connected $\sin n\phi$ with $\sin\phi$ and $\cos\phi$.

Rhæticus (1569) found the relation

$$\cos n\phi = \cos(n-2)\phi - 2\sin\phi \sin(n-1)\phi.$$

Newton (1676) gave the well-known relation

$$\sin n\phi = n\sin\phi + \frac{(1-n^2)n}{3!}\sin^3\phi + \cdots,$$

and Jacques Bernoulli (1702) showed that

$$\sin n\phi = \cos^n\phi - \frac{n(n-1)}{2!}\cos^{n-2}\phi \sin^2\phi + \cdots,$$
$$\cos n\phi = \frac{n}{1}\cos^{n-1}\phi \sin\phi - \frac{n(n-1)(n-2)}{3!}\cos^{n-3}\phi \sin^3\phi + \cdots.$$

Functions of Half an Angle. Ptolemy[2] (*c.* 150) knew substantially the sine of half an angle, expressed as half a chord, and it is probable that Hipparchus (*c.* 140 B.C.) and certain that Varāhamihira (*c.* 505) knew the relation which we express as

$$\sin\frac{\phi}{2} = \sqrt{\frac{1-\cos\phi}{2}}.$$

[1] Tropfke, *Geschichte*, II (2), 57–61, with bibliography.
[2] See the Heiberg edition of Ptolemy, p. 39.

After the development of analytic trigonometry in the 17th century, these relations were greatly extended. Four others may be mentioned as typical, the first two, due to Euler (1748), being

$$\tan 2\phi = \frac{2\tan\phi}{1-\tan^2\phi},$$

$$\cot 2\phi = \frac{\cot\phi - \tan\phi}{2};$$

and the others, due to Lambert (1765), being

$$\sin 2\phi = \frac{2\tan\phi}{1+\tan^2\phi},$$

$$\cos 2\phi = \frac{1-\tan^2\phi}{1+\tan^2\phi}.$$

Theorem of Sines. The important relation now expressed as

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

was known to Ptolemy (*c.* 150) in substance, although he expressed it by means of chords.[1]

While recognized by Albêrûnî and other Oriental writers, it was Naṣîr ed-dîn (*c.* 1250) who first set it forth with any clearness. A little later Levi ben Gerson (*c.* 1330) stated the law in his work *De sinibus, chordis, et arcubus*;[2] but the first of the Renaissance writers to express it with precision was Regiomontanus[3] (writing *c.* 1464).

[1] Thus, if C is a right angle in triangle ABC, then

$$a = \frac{c \text{ chord } 2A}{120^{\mu}},$$

where 120 μοῖραι is the diameter of the circumcircle.

[2] "... omnium triangulorum rectilineorum talem proportionem una linea habet ad aliam, qualem proportionem unus sinus angulorum, quibus dictae lineae sunt subtensae, habet ad alium." See Braunmühl, *Geschichte*, I, 106.

[3] "In omni triangulo rectilineo proportio lateris ad latus est, tamquam sinus recti anguli alterum eorum respicientis, ad sinum recti anguli reliquum latus respicientis" (Lib. II, prop. 1). See Tropfke, *Geschichte*, V (2), 74.

Theorem of Cosines. The fact that

$$c^2 = a^2 + b^2 - 2\,ab\cos C$$

is essentially a geometric theorem of Euclid.[1] In that form it was known to all medieval mathematicians. In the early printed books it appears in various forms, Vieta (1593) giving it substantially as

$$2\,ab/(a^2 + b^2 - c^2) = 1/\sin(90° - C),$$

and W. Snell (1627) as

$$2\,ab/[c^2 - (a - b)^2] = 1/(1 - \cos C).$$

Theorem of Tangents. The essential principle of the law of tangents, which was given by Vieta[2] (p. 611) and improved by Fincke (1583), was known to Ptolemy (*c.* 150). Regiomontanus (*c.* 1464) expressed it by a rule which we should state as

$$\frac{\sin A + \sin B}{\sin A - \sin B} = \frac{\tan \frac{1}{2}(A + B)}{\tan \frac{1}{2}(A - B)}.$$

Areas. The first evidence of the rule which resulted in the formula for the area of a triangle, which we know as

$$\Delta = \tfrac{1}{2}\,ab\sin C,$$

is found in the trigonometry of Regiomontanus (*c.* 1464),[3] but the theorem is not explicitly stated by him. Snell (1627) gave it in the form

$$1 : \sin A = bc : 2\,\Delta.$$

Right-angled Spherical Triangle. The Greek mathematicians made use of the right-angled spherical triangle in their computations, but nowhere do we find a systematic treatment of the subject. Taking the hypotenuse as c, we have the following six cases:

1. $\cos c = \cos a \cos b$.
2. $\cos c = \cot A \cot B$.
3. $\cos A = \cos a \sin B$.
4. $\cos A = \tan b \cot c$.
5. $\sin b = \sin c \sin B$.
6. $\sin b = \tan a \cot A$.

[1] *Elements*, II, 12, 13.

[2] "Ut aggregatum crurum ad differentiam eorundem, ita prosinus dimidiae angulorum ad basin ad prosinum dimidiae differentiae" (*Opera*, Schooten ed., p. 402). See Tropfke, *Geschichte*, V (2), 80.

[3] *De triangulis omnimodis*, Nürnberg, 1533.

In his astronomical problems Ptolemy (*c.* 150) makes use essentially of the first, fourth, fifth, and sixth of these cases,[1] although without the functions he could not give the rules.

The third case is essentially given by Jabir ibn Aflah (*c.* 1145), and so it was commonly known as Jabir's Theorem.[2]

The first writer to set forth essentially all six cases was Naṣîr ed-dîn (*c.* 1250). Napier's Rules for the right-angled spherical triangle appeared in his tables of 1614.[3]

Oblique-angled Spherical Triangle. The oblique-angled spherical triangle was not seriously studied by itself until the Arabs began to consider it in the 10th century.

The Theorem of Sines,

$$\frac{\sin a}{\sin A} = \frac{\sin b}{\sin B} = \frac{\sin c}{\sin C},$$

and the Theorem of Cosines of Sides,

$$\cos a = \cos b \cos c + \sin b \sin c \cos A,$$

may have been known to them, but they are first found in print in the *De triangulis* written by Regiomontanus *c.* 1464.[4]

The Theorem of Cosines of Angles,

$$\cos A = -\cos B \cos C + \sin B \sin C \cos a,$$

was given in substance by Vieta in 1593, although he had used it before this date.[5] It was first proved by Pitiscus in 1595.

The Theorem of Cotangents,

$$\frac{1}{\sin b \csc A} = \frac{\cot c + \cos A \cot b}{\cot C},$$

was also given in substance by Vieta, but was modified by Adriaen van Roomen (1609) and proved by Snell (1627).[6]

[1] Tropfke, *Geschichte*, V (2), 131, with references. Consult this work and Braunmühl's *Geschichte* for further details.

[2] It was possibly known to Ṯâbit ibn Qorra (*c.* 870).

[3] *Mirifici logarithmorum canonis descriptio*, 1614, Lib. II, cap. iv.

[4] On the Theorem of Sines see Tropfke, V (2), 133; on the Theorem of Cosines see *ibid.*, p. 139.

[5] For the priority question see Tropfke, *Geschichte*, V (2), 139.

[6] For the general literature on this subject see Braunmühl, *Geschichte*, I (1), 25; Tropfke, *Geschichte*, V (2), 137 seq., especially p. 143.

TOPICS FOR DISCUSSION

1. The etymology of the words "trigonometry," "geometry," "mensuration," "agrimensor," "survey," "geodesy," and other terms having a related meaning.

2. Primitive needs that would naturally tend to the development of trigonometry.

3. The relation of shadow reckoning to plane trigonometry in various countries and at various times.

4. The influence of astronomy upon the development of the science of trigonometry.

5. The Greek astronomers who contributed most to the study of trigonometry, the function which they developed, and the reason why this function was selected.

6. The contributions of Menelaus to the study of trigonometry.

7. The Hindu contributions to the science.

8. The assertion that the chief contribution to mathematics made by the Arab scholars was to the science of trigonometry, and that this contribution was important.

9. The Arab and Persian writers on trigonometry, and the important features of their work.

10. Influence of Peurbach and Regiomontanus.

11. The change of trigonometry from being essentially geometric to being largely analytic, and the influence of this change upon the later development of the science.

12. Development of the concept of the sine of an angle, and the origin of the name.

13. Development of the concept and name of the cosine.

14. Development of the concepts of the versed sine and the coversed sine, with reasons for their gradual disappearance.

15. The favorite functions in astronomy and those used in practical mensuration.

16. Development of the tangent and cotangent.

17. Development of the secant and cosecant, and the causes leading to their gradual disuse in modern times.

18. The relation between trigonometric functions.

19. Development of the leading methods of computing trigonometric tables.


20. The history of typical and important theorems of trigonometry.

CHAPTER IX

MEASURES

1. Weight

Measures in General. The subject of metrology is so extensive that it is impossible, in a work like this, to do more than give a few notes relating to the measures in common use. The purpose of this chapter, therefore, is simply to lay before the student some of the points of interest in the history of the most familiar of our several units of measure, to suggest the significance of the names of these units, and to indicate some of the works on the subject to which he may go for further information.[1]



EGYPTIAN WEIGHT

A porphyry weight found near the North Pyramid at Lisht and now in the Metropolitan Museum. The inscription reads, "Senusert, giving life eternally, 70 gold debens." It was used for weighing gold

Egypt. The use of the balance for purposes of weighing is doubtless prehistoric, for weights are found in remains of the first dynasty of Egypt (*c.* 3400 B.C.).[2] The first inscribed weight that has been found is of the 4th dynasty, the time of the Great

[1] An excellent summary of the history is given under "Weights and Measures" in the *Encyc. Brit.*, 11th ed. The measure of angles and arcs has been already considered in Chapter VIII.

[2] See W. M. Flinders Petrie, *Proceedings of the Soc. of Biblical Archæol.*, London, XXIII, 385. See also *Bulletin of the Metrop. Mus. of Art*, New York, XII, 85.

Pyramid.[1] The earliest Egyptian scales were simple balances, either held in the hand or supported on a standard. These are frequently illustrated in the temple wall pictures. The steelyard with its sliding weight and fixed fulcrum was used as early as 1350 B.C.

The Egyptian weights of which the names and values are known with certainty were the *deben* (*dbn*, formerly read *uten*, about 13.6 grams, but commonly taken as 15 grams) and the *kidet* (*kdt*, *kite*, 0.1 of a *deben*).[2]



TABLE OF BABYLONIAN MEASURES OF CAPACITY AND WEIGHT

A fragment of a clay tablet found at Nippur and dating from *c.* 2200 B.C. The reverse side contains tables of weight, length, and area. Courtesy of the University of Pennsylvania

Babylonia. The Babylonians used a cubic foot of rain water to establish their unit of weight, the standard talent.[3] The chief subdivision of the talent was called a *maneh* and was $\frac{1}{60}$ of a talent. They also had a unit of weight known as the *she*, about 45 mg. Our knowledge of the Babylonian measures in general is derived from a number of inscribed tablets such as the one here shown.

Hebrews. The Hebrew standards were kept in the temple, as was also the case in other nations. Thus we read[4] of "the shekel of the Sanctuary," that is, the standard shekel, about a quarter of an ounce in early times, or a half ounce after the

[1] *Proceedings of the Soc. of Biblical Archæol.*, XIV, 442.

[2] Peet, *Rhind Papyrus*, p. 26; *Encyc. Brit.*, 11th ed., XXVIII, 480.

[3] Mahaffy, *Greek Life*, p. 67. See also J. Brandis, *Das Münz- Mass- und Gewichtswesen in Vorderasien*, pp. 1, 41 (Berlin, 1866); G. A. Barton, *Haverford Library Collection of Cuneiform Tablets*, Philadelphia, 1905, 1909.

[4] Exodus, xxx, 13.

time of the Maccabees (1st century B.C.). This shekel of the Hebrews was the *sicilicus* of the Romans.[1] The Hebrew *maneh* was 100 shekels, or $\frac{1}{30}$ of a talent. The shekel was also used as a unit of capacity, and with the Babylonians it was equivalent to 0.07 liter.[2]

Greece. The Greek unit of weight in Homer's time was the talent,[3] a standard that varied from country to country. The ancient Greek talent weighed about 57 pounds, but the Hebrews used the term for a unit of about $93\frac{3}{4}$ pounds. It was also used as a unit of value, generally the value of a talent of silver, this being about $1180 in Greece and from about $1650 to about $1900 among the Hebrews, according to present standards.

For a smaller weight the Greeks used the drachma,[4] originally "a handful" but used by the ancients to designate both a weight and, as in the case of the talent, a unit of value. In modern Greece it is a coin identical in value with the franc at the normal rate of exchange.

The later Greek weights may be thus summarized: 1 *talanton* = 60 *mnai* = 6000 *drachmai* = 36,000 *oboloi* = 288,000 *chalkoi*.[5]

Rome. The Roman unit of weight was the pound. This was divided into twelfths (*unciae*).[6] The usual Roman weights may be summarized as follows:

1 *libra* = 12 *unciae* = 48 *sicilici* = 288 *scripula* = 576 *oboli* = 1728 *siliquae*.

The ounce was about 1.09 oz. avoirdupois, or 412 grains.

[1] Or *siclus*; Greek *σίγλος*. [2] Barton, *loc. cit.*, II, 18.

[3] Τάλαντον (*tal'anton*). It was originally smaller than the later talent. See F. Hultsch, *Griechische und römische Metrologie*, p. 104 (Berlin, 1882); Harper's *Dict. Class. Lit.*; Pauly-Wissowa; A. Böckh, *Metrologische Untersuchungen*, Berlin, 1838.

[4] Δραχμή (*drachme'*). The Lydian drachma of the 7th century B.C. was $\frac{1}{4}$ of a shekel.

[5] In Greek, τάλαντον; μνᾶ, μναῖ; δραχμή, δραχμαί; ὀβολός, ὀβολοί; χαλκοῦς, χαλκοῖ. The *chalkous* was about 0.091 g. As a measure of value it was a copper coin worth $\frac{1}{8}$ of an obol, somewhat less than $\frac{1}{2}$ of an American cent. The *talanton* was about 26,196 g.

[6] Hultsch, *loc. cit.*, p. 144; Ramsay and Lanciani, *Manual of Roman Antiqs.*, 17th ed., p. 461 (London, 1901). There are many works on the subject. Among

The Romans had a table known as the *mensa ponderaria*, in the stone top of which were cavities like washbasins, with a plug in the bottom of each cavity. These were standards of capacity, or of capacity with respect to weight.[1]

Far East. In India and other parts of the Far East the weights and currencies were commonly based upon the weights of certain seeds. The favorites were the *abrus precatorius*, a creeper having a small, bright-red seed with a black spot on it,[2] and the *adenanthera pavonina*, a large pod-bearing tree with a bright-red seed which is conventionally taken as weighing twice an abrus seed.[3]

England. In England the grain was originally the weight of a barleycorn, a barley grain. The Latin *granum* has the same root (*gar*) as our word "corn."[4]

the earlier ones are A. Alciatus, *Libellvs. De Ponderibvs et mensuris*, Copenhagen, 1530; L. Portius, *De sestertio pecvniis ponderibvs et mensvris antiqvis libri duo*, s. l. a. (Venice?, *c.* 1500), with editions at Florence (1514?) and Basel (1520 and 1530); G. Budaeus, *De asse, et partibvs eivs, libri V*, Paris, 1514 (title as in Lyons edition, 1551); G. Agricola, *Libri quinque de Mensuris & Ponderibus*, Paris, 1533; Venice, 1533 and 1535; Basel, 1549 and 1550; an epitome, Lyons, 1552; H. Uranius, *De re nvmaria, mensvris et ponderibus Epitome ex Budaeo, Portio, . . .*, Solingen, 1540; M. Neander, ΣΥΝΟΨΙΣ *mensvrarvm et pondervm, . . .* Basel, 1555. These show the interest taken in the subject in the 16th century. Of the 18th century works one of the best is J. Arbuthnot, *Tables of Antient Coins, Weights, and Measures*, 2d. ed., London, 1754.

[1] A Naples specimen is illustrated in Mau's *Pompeii*, Kelsey's second edition, New York, 1902.

[2] Often seen for sale in European and American shops. The name *precatorius* (from *precator*, one who prays) comes from the fact that certain Buddhists use these as beads for their rosaries.

[3] R. C. Temple, "Notes on the Development of Currency in the Far East," *Indian Antiquary*, 1899, p. 102. Other seeds were used, as is shown by H. T. Colebrooke, "On Indian Weights and Measures," *Asiatic Researches*, V (1799), 91, with tables. See also his *Lilāvati*, p. 1, § 2.

[4] Whence also "garner," to gather grain; "pomegranate," from the French *pomme* (apple) and *grenate* (seeded); "granite," a grained or spotted stone; "garnet"; "grange"; and the Spanish *granada*. The Scotch statute required that the inch be "iii bear cornys gud and chosyn but tayllis" (tailless). The Latin statute of England read: "Tria grana ordei sicca et rotunda faciunt pollicem." On the history of British measures in general see F. W. Maitland, *Domesday Book and Beyond*, Cambridge, 1897, p. 368; J. H. Ramsay, *The Foundations of England*, London, 1898, I, 533; F. Seebohm, *The English Village Community*, 4th ed., p. 383 (London, 1896); R. Potts, *Elementary Arithmetic*, London, 1886.

The word "pound" comes from the Latin *pondo* (by weight),[1] and the ounce, as already stated, from the Latin *uncia,* a twelfth of the Roman pound.[2]

England had developed a system of weights before the Troy weight was introduced from the French town of Troyes, one of the many places in which fairs were held in the Middle Ages. This introduction seems to have taken place as early as the second half of the 13th century, for Grafton's *Chronicles*[3] has this to say of the matter:

About this tyme[4] was made the statute of weightes and measures, that is to say, that a sterlyng penny should waye .xxxij. graynes of wheate drie and round, and taken in the middes of the eare,[5] and .xx. of those pence shoulde make an ounce, and .xij. ounces make a pound Troy: And .viij. pound Troy weight make a gallon of Wine, and .viij. wyne galons to make a London bushell, which is the .viij. part of a quarter. Also three barly Cornes dry and round should make an ynch, & .xij. ynches a foote, and thre foote a yard, a fiue yards, halfe a perch, or poll, & .xl. pol in length & thre in bredth an acre of land.

And these standardes of weight and measures were confirmed in the .xv. yere of king Edward the thirde, and also in the tyme of Henry the sixt and of Edward the fourth, and lastly confirmed in the last yere of Henry the seuenth. But in the time of king Henry the sixt it was ordeyned that the same ounce should be deuided into .xxx. pence, and in the tyme of king Edward the fourth, into .xl. pence, and in the tyme of king Henry the eight into .xliiij. pence: But the weight of the ounce Troy, and the measure of the foote continued alwayes one.

In due time the Troy weight was replaced by the avoirdupois for general purposes and was thenceforth limited chiefly to

[1] From *pendere*, to weigh. From the same root we have such words as "depend," "spend," and "pendulum," and the French *poids* and our "poise."

[2] Lack of space precludes any discussion of the relation of the apothecaries' weight to the ancient Greek and Roman systems and symbols. There is an extensive literature on the subject.

[3] 1569 ed.; 1809 reprint, p. 277. [4] "The LIJ Yere of Henry III," *i.e.*, 1268.

[5] So Recorde (*c.* 1542) says: "Graine, meaninge a grayn of corne or wheat drye, and gathered out of the myddle of the eare." *Ground of Artes*, 1558 ed., fol. L 4.

the use of goldsmiths.[1] These goldsmiths also used in this connection the carat, a weight consisting of 12 grains.[2] The word had a variety of meanings, being commonly used to express the purity of gold, "22 carats fine" meaning an alloy that is $\frac{22}{24}$ "fine gold." It appears in various forms,[3] and its meaning in this sense comes from the fact that a gold mark was 24 carats, so that a mark that had only 18 carats of gold was only $\frac{18}{24}$ pure. So Recorde (*c.* 1542) says: "The proofe of gold is made by Caracts, whereof 24 maketh a Marke of fine gold: the Caract is 24 graines."[4]

Avoirdupois Weight. The word "avoirdupois" is more properly spelled "averdepois," and it so appears in some of the early books. It comes from the Middle English *aver de poiz*,[5] meaning "goods of weight." In the 16th century it was commonly called "Haberdepoise," as in most of the editions of Recorde's (*c.* 1542) *Ground of Artes*. Thus in the Mellis edition of 1594 we have:

> At London & so all England through are vsed two kinds of waights and measures, as the Troy waight & the Haberdepoise.

[1] So the Dutch arithmetics of the 17th century speak of it as *Assay-gewicht*. *E.g.*, Coutereels's *Cyfer-Boeck*, 1690 ed., p. 16. The Dutch writers also called it *Trois gewicht*, as in Bartjens's arithmetic, 1676 ed., p. 155.

[2] So the Dutch arithmetics of Petrus (1567), Van der Schuere (1600), and others give 12 grains = 1 karat, 24 karats = 1 marck (for gold), and 20 angels = 1 ounce, 8 ounces = 1 marck (for silver). Trenchant (1566) says: "Per ansi le marc d'or sans tare est à 24 kar. de fin aloy." In this sense it comes from the Arabic *qīrāt*, a weight of 4 barleycorns; but the Arabs derived it from the Greek κεράτιον (*kera'tion*), the fruit of the locust tree, L. Latin *cerates*. Perhaps the Arabic use is responsible for the carat weight's being 4 diamond grains, now taken as 200 milligrams.

[3] Italian *carato*, French *carat*, and Spanish *quilate*. So Texada (1546): "24. quilates son de puro oro"; Sfortunati (1534): "Io mi trouo oro di .24. charatti"; Trenchant (1566): "18 karats de fin"; and Rudolff (1526): "fein 18 karat."

[4] Compare "4 marcx d'Or a 14 Carats de fin," in Coutereels's Dutch-French arithmetic, 1631 ed., p. 309.

[5] *Aver de pois*, *peis*, etc. The English *aver*, from the Old French *aveir* or *avoir*, meant goods, and *poiz* was the French *peis* or *pois*, Latin *pensum*, from *pendere*, to weigh. About the year 1500 the old Norman *peis* was superseded by the modern *pois*. The incorrect *du*, for *de*, came in about 1650. Even as late as 1729 the American Greenwood used "averdupois."

The system was introduced into England from Bayonne *c.* 1300, but is essentially Spanish. The name is limited to the English-speaking countries, the pound of 16 ounces being called on the Continent by various names, such as the pound merchant.[1] Troy weight was the more popular until the 16th century, when, as Digges (1572) tells us, "Haberdepoyse" became the "more vsuall weight." Even a century later, however, the Troy weight was given first and was used for weighing such commodities as figs and tobacco and even lead and iron.[2] There was also the Tower pound of 11.25 Troy ounces, but this was abandoned about the year 1500. In the latter part of the 18th century a popular writer[3] thus refers to the matter:

When *Averdupois Weight* became first in Use, or by what *Law* it was at first settled, I cannot find out in Statute Books; but on the contrary, I find that there should be but one Weight (and one Measure) used throughout this Realm, viz. that of Troy, (Vide 14 Ed. III, and 17 Ed. III). So that it seems (to me) to be first introduced by Chance, and settled by Custom, viz. from giving good or large weight to those Commodities usually weighed by it, which are such as are either very Coarse and Drossy, or very subject to waste; as all kinds of Grocery Wares.

2. Length

Babylonia and Egypt. The Babylonian measures, like those of most early peoples, were derived to a considerable extent from the human body. For example, one of the world's primitive measures was the cubit,[4] the length of the *ulna*, or forearm, whence the English *ell* and French *aune*, but applied to various lengths. This standard is found among the Babylonians, the

[1] Thus Trenchant (1566) says, "La liure marchande vaut 16 onces." The Dutch writers sometimes called it "Holland weight," as in Coutereels's *Cyffer-Boeck*, 1690 ed., p. 17, where 1 pound = "2 Marck" or "16 once" or "32 loot."

[2] Hodder's arithmetic, 1672 ed., pp. 15, 66, 68.

[3] J. Ward, *The Young Mathematician's Guide*, 12th ed., p. 32. London, 1771.

[4] Latin *cubitum*, elbow. Sir Charles Warren, *The Ancient Cubit*, London, 1903; a scholarly and extended treatment of the subject.

length varying from 525 mm. to 530 mm.[1] It was known somewhat earlier in Egypt and numerous specimens are still extant.[2]

Greece and Rome. The method of fixing standards by measurement of the human body naturally led to many variations. Thus the Attic foot[3] averaged 295.7 mm.; the Olympic, 320.5 mm.; and the Æginetan, 330 mm. A similar variation is found in Western Europe, the Italian foot being 275 mm.; the Roman, 296 mm. (substantially the same as the Attic); and the *pes Drusianus,* 333 mm.[4] The foot was not a common measure until *c.* 280 B.C., when it was adopted as a standard in Pergamum.

The fingerbreadth[5] was used by both Greeks and Romans, as was also the palm[6] of four digits. The cubit was six palms, or twenty-four digits, the Roman foot was $13\frac{1}{2}$ digits, and the fathom[7] was the length of the extended arms. The mile[8] was, as the name indicates, a thousand units, the unit being a double step.

In general, the most common Roman measures of length may be summarized as follows: the *pes* (foot) was 0.296 m. long, and 5 *pedes* (feet) made 1 *passus*; 125 *passus* made 1 *stadium*, about 185 m.; and 8 *stadia* made 1 mile, about 1480 m.[9]

[1] J. Brandis, *Das Münz- Mass- und Gewichtswesen in Vorderasien* (Berlin, 1866), p. 21; Hilprecht, *Tablets*, p. 35. The Babylonian name was *ammatu*, and this unit was divided into 30 *ubânu* (*ubânê*). To use our common measures, we may say that the average Roman cubit was 17.4 inches; the Egyptian, 20.64 inches; and the Babylonian, 20.6 inches. See Peet, *Rhind Papyrus*, p. 24.

[2] As the *mahi*, three of which made the *xylon*, the usual length of a walking staff, about 61.5 inches, and 40 of which made the *khet*. Other measures are also known, such as the foot, which was equivalent to about 12.4 of our inches.

[3] Πούς (*pous*). The general average as given by Hultsch (*loc. cit.*, p. 697) is 308 mm. [4] K. R. Lepsius, *Längenmasse der Alten*, Berlin, 1884.

[5] Δάκτυλος (*dak'tylos*); Latin, *digitus*.

[6] Δοχμή (*dochme'*); Latin, *palmus*. This is our "hand," used in measuring the height of a horse's shoulders. Homer speaks of handbreadths (δῶρον, *do'ron*) and cubits (πυγών, *pygon'*).

[7] Anglo-Saxon *fæthm*, embrace. The Greek word was ὄργυια (*or'guia*), the length of the outstretched arms; Latin, *tensum*, stretched.

[8] *Mille passuum* (colloquially *passum*). The pace was a double step, and hence a little over 5 Anglo-American feet.

[9] The Greek stadium (στάδιον, *sta'dion*) varied considerably in different cities. The Athenian stadium was about 603–610 Anglo-American feet.

Far East. The finger appears in India as "eight breadths of a *yava*" (barleycorn), four times six fingers making a cubit,[1] as in Greece. The other Oriental units have less immediate interest.

England. In England there was little uniformity in standards before the Norman Conquest. The smaller units were determined roughly by the thumb,[2] span,[3] cubit, ell,[4] foot, and pace.

A relic of this primitive method is seen in the way in which a woman measures cloth, taking eight fingers to the yard, or the distance from the mouth to the end of the outstretched arm.

For longer distances and for farm areas it was the custom to use time-labor units, as in a day's journey or a morning's plowing, such terms being still in use in various parts of the world. The furlong (40 rods, or an eighth of a mile) probably came from the Anglo-Saxon *furlang*, meaning "furrow long."

The word "yard" is from the Middle English *yerd* and the Anglo-Saxon *gyrd*, meaning a stick or a rod, whence also a yardarm on a ship's mast. That the standard was fixed in England by taking the length of the arm of Henry I (1068–1135) is not improbable. Thus an old chronicle relates: "That there might be no Abuse in Measures, he ordained a Measure made by the Length of his own Arm, which is called a *Yard*."

The words "rod" and "rood" may have had a common origin. The rod was used for linear measure and the rood came to be used for a fourth of an acre.[5]


[1] *Hasta*, *cara*, the forearm. Colebrooke's *Lilāvati*, p. 2, § 35. The word *cubit* appears in India and Siam as *covid*, in Arabia as *covido*, and in Portugal as *covado*.

[2] Latin *pollex*, whence the French *pouce*, an inch. The word "inch," like "ounce," is (as already stated) from the Latin *uncia*, the twelfth of a foot or the twelfth of a pound. Originally the word meant a small weight and is allied to the Greek ὄγκος (*on'kos*), bulk, weight. The old Scotch inch was averaged from the thumbs of three men, "þat is to say, a mekill man and a man of messurabill statur and of a lytill man." See Maitland, *loc. cit.*, p. 369.

[3] The distance spanned by the open hand, from thumb to little finger; finally taken as 9 inches.

[4] The ell has varied greatly. In England it is 45 inches, that is, $1\frac{1}{4}$ yards. The old Scotch ell was 37.2 inches, and the Flemish ell was 27 inches.

[5] For a bibliography of the subject of measures of length consult the encyclopedias. Among the most ingenious studies of the subject is W. M. F. Petrie, *Inductive Metrology*, London, 1877.



MEASURES OF LENGTH

European of the 18th century, showing the general appearance of the common measuring sticks of the period. The three shortest pieces are ells

3. Areas

Acre. The common unit of land measure known as the acre[1] has varied greatly in different countries and at different periods.[2] It was commonly taken to mean a morning's plowing,[3] a strip of land 4 rods wide and one furrow long, that is, 4 rods by 1 furlong, 4 rods by 40 rods, or 160 square rods.

The rood was a fourth of an acre and was also called a perch.[4] It is thus described by Recorde (*c.* 1542):

> 5 Yardes and a halfe make a Perche . . . [and] 1 Perche in bredth & 40 in length, do make a Rodde of land, which some cal a Rood, some a Yarde lande, and some a Forthendale.[5]

4. Capacity

Modern Times. The subject of measures of capacity is so extensive as to make it impossible to mention more than a few facts concerning our modern British and American units.

The gill was the Old French *gelle*, a sort of wine measure, from the Middle Latin *gillo* or *gellus*, a wine vessel.[6]

The quart[7] is, of course, simply the quarter of a gallon.[8]

[1] Anglo-Saxon *æcer*. The ancient units of area have no particular significance at the present time. It is, therefore, sufficient, merely to mention the Greek *plethron*, about 0.235 of an acre, and the Roman *jugerum*, about 0.623 of an acre.

[2] For some of these variations see F. W. Maitland, *Domesday Book and Beyond*, p. 374 (Cambridge, 1897).

[3] The cattle used in plowing in the morning were put out to pasture in the afternoon. Compare the German *Morgen*.

[4] Latin *pertica*, a pole, staff, or rod. The word has various other uses, as in the case of a perch of stone or masonry, the contents of a wall 18 inches thick, 1 foot high, and 1 rod long, or $24\frac{3}{4}$ cubic feet. The perch as a unit of length was the same as the rod.

[5] 1558 ed., fol. L 6.

[6] It may come from the same root as "gallon." The United States gill contains 7.217 cu. in., or 118.35 cu. cm. The British gill contains 142 cu. cm.


[7] French *quarte*, from the Latin *quartus*, fourth, which is related to *quattuor*, four, and to such words as "quadrilateral" (four-sided), "quarry" (a place where stones are squared), "quarantine" (originally a detention of forty days), "quarto," "quire" (Low Latin *quaternum*, a collection of four leaves), "square" (probably Low Latin *ex* (intensive) + *quadrare*, to square), "squad," and "squadron."

[8] The United States gallon contains 231 cu. in., like the old English wine gallon. The imperial (British) gallon contains 277.274 cu. in.

The pint may possibly receive its name from the Spanish *pinta*,[1] a mark, referring to a marked part of a larger vessel.[2] The word "bushel"[3] means a small box, but the origin of the word "peck," as applied to a measure, is obscure.

5. VALUE

Early Units. In the measure of value it became necessary at an early period to develop media of exchange of one kind or another. The primitive pastoral people naturally used cattle of some kind, whence the Latin noun *pecunia*[4] (money) and the English adjective "pecuniary."



EARLY ROMAN MONEY (PECUNIA)
Showing how a coin was stamped to represent the value of an ox, 4th century B.C. From Breasted's *Survey of the Ancient World*

For media of exchange the Greeks often used copper utensils, and ingots of silver and gold. The Babylonians and Egyptians also made use of ingots and rings of the precious metals, selling these by weight, whence came the *aes infectum*[5] of the Latins. From this relation of money to value came the double use of such measures as the talent and the pound. Even

[1] Latin *picta*, marked or painted, whence "picture." The Middle English form is *pynte*. The origin is, however, uncertain.

[2] Of the British measures whose names are still heard in the colonies and in America, kilderkin was the Dutch *kindeken* (German *Kinderchen*), a babekin, that is, a mere baby in bulk as compared with a tun or vat. "Tun" and "ton" are the same word (Middle English *tonne*, Low Latin *tunna*), meaning a large barrel and hence also a great weight.

[3] Middle English *buschel* or *boischel*; Low Latin *bussellus*, or *bustellus*, diminutive related to Old French *boiste*, a box. Compare "pyx," Greek πυξίς (*pyxis'*), a box, particularly one made of πύξος (*pyx'os*, Lat. *buxus*), boxwood. The imperial (British) bushel contains 2218.192 cu. in., and the Winchester bushel (which became the legal standard in the United States) contains 2150.42 cu. in.

[4] From Latin *pecus*, sheep, cattle. For discussion, see T. Gomperz, *Les penseurs de la Grèce*, French translation, p. 8 (Lausanne, 1904); F. Hultsch, *Griechische und römische Metrologie*, p. 162 (Berlin, 1882).

[5] *Aes*, bronze, copper, money; *infectum*, *in* + *facere*, to make; that is, uncoined money. See F. Gnecchi, *Monete Romane*, 2d ed., p. 86 (Milan, 1900).

at present, in certain mining districts, the ounce of gold is commonly spoken of as a unit of value. The pound[1] became the *libra*[2] in most Latin countries.

From the *aes infectum* as pieces of metal came the large and heavy metal disks of the early Romans. When these were stamped they became the *aes signatum*.[3]

Coins. The earliest stamped coins found in the Mediterranean countries were probably struck in Lydia in the 7th century B.C.,[4] or possibly in Ægina in the 8th century. They seem to have appeared in China at about the same time.

The first silver money coined in Rome (268 B.C.) was based upon the relation of 10 *asses* to the *denarius*, but the number was changed at a later date. Pliny tells us that the first gold money coined in Rome appeared in 217 B.C.[5] The *aureus*, or gold *denarius*, was first coined under Augustus (31 B.C.–14 A.D.) as $\frac{1}{40}$ of a pound, but it underwent gradual changes until, under Constantine (306–337), it became $\frac{1}{72}$ of a pound, then taking the name *solidus*.[6]

Great Britain. When Cæsar went to Britain (*c.* 55 B.C.) he found the natives using certain weights of metal as media of exchange. Coinage was introduced soon thereafter, based on Roman values. The figure of Britannia, which is still seen

[1] Latin, *pondo libra*, "a pound by weight." From the same root as *pondus*, a weight, we also have such words as "ponder," and such units as the Spanish *peso*; see also page 638, note 1.

[2] Latin *libra*, a balance, a pound weight, from the Greek λίτρα (*lī'tra*), a pound, whence *litre*, liter. The constellation Libra has for its symbol ♎, the scale beam. From the same root we have such words as "deliberate," to weigh our thoughts. The *libra* appears in French as *livre* and in Italian as *lira*, the old pound in weight as well as in value. The French *livre* was also called a *franc*, as in Trenchant (1566): "la liure autremẽt appellee *frãc*."

[3] Gnecchi, *loc. cit.*, p. 89.

[4] Herodotus, I, 94. Judging from the museum pieces the early coins seem to have been both of gold and of silver, and both circular and oblong. See an illustration in Volume I, page 56.

[5] Gnecchi, *loc. cit.*, chap. xiv and p. 145.

[6] *I.e.*, a solid piece of money. From this we have the word "soldier," a man who fought for money, and such words as the Italian *soldo* and the French *sol* and *sou*. The English symbol for shilling (/) comes from the old form of *s* (ſ) and was the initial for *solidus*, just as £ is the initial of *libra* (pound) and as *d* (for penny) is the initial of *denarius*.

on British coins as mistress of the seas, is not at all modern. It appears on one of the pieces of Hadrian's time (*c.* 130).[1]

The most ancient coin of the Saxon period (*c.* 600) is the *sceat*, a silver coin weighing about 1 gram. The word *scilling*, for shilling, appears at this time, the word possibly meaning a little scar.[2] The origin is doubtful, and the word may mean the "clinking coin."[3]

The word "penny" may have come from the Latin *pannus*, a cloth, and hence is the value of a certain piece; but, as in many similar cases, the origin is obscure. Since a piece of cloth was a convenient pledge for money borrowed, the word "pawn" may have come from the same source.[4]

The word "farthing," the Anglo-Saxon *feorthling*, is simply the "little fourth" of a penny.[5]

The Saxon coins were regulated by the pound weight. This pound was commonly known as the Cologne pound, having been brought from that city, and was probably the same as the weight known after the Conquest as the Tower pound, so called because the mint of London was in the Tower. A pound sterling was this weight of silver coins.

United States. The word "dollar" comes from the Low German *Daler*, German *Thaler*, from *Joachimsthaler*, since these large silver pieces first appeared in the Thal[6] of St. Joachim,

[1] For a brief résumé of the history of British coins, see R. Potts, *Elementary Arithmetic with brief notices of its history*, § II (London, 1886).

[2] *Skil* means to divide and comes from *skal* or *skar*, to cut; so that *scar-ling* means a little cut on a tally stick, to distinguish the mark from the larger scar (score) which indicated 20 shillings, or a pound. *Skar* is the root of Greek κείρειν (*kei'rein*), to cut close, and is allied to the Anglo-Saxon *scær* and the German *scheren*, whence our "shear." The English "jeer" may be from the same source through the Dutch phrase *den gheck sheeren*, "to shear the fool." A "share" of stock is from the same root.

[3] Compare the German *schellen*, to sound or tinkle. See Greenough and Kittredge, *Words and their Ways*, p. 140 (New York, 1901).

[4] Similarly "panel," a piece of anything marked off. The Middle English of "penny" is *peni*, plural *penies* and *pens*. Compare the Anglo-Saxon *pening* and the German *Pfennig*.

[5] The word is substantially the same as "firkin," from the Dutch *vier* (four) + *kin* (diminutive, as in "lambkin"), once a fourth of a barrel.

[6] English "dale" and "dell." Coined there because of the silver mines in the valley.

Bohemia, in the 16th century. "Dime" is from the French *disme* and Latin *decem* (ten), "cent" is from *centum* (hundred), and "mill" is from *mille* (thousand). It took over fifty years to replace the English system by the "Federal" in the United States. The origin of the symbol $ is uncertain. It seems to have first appeared in print in Chauncey Lee's *The American Accomptant* (Lansingburgh, 1797) (although in a form very different from the one familiar at the present time), but it was used in manuscripts before that date. The Italian and British merchants had long used ℔ for pounds, writing it quite like our dollar sign (see Volume I, page 233), and it is possible that our merchants in the closing years of the 18th century simply adopted this symbol, just as we have adopted the English word "penny" to mean a cent, which is only a halfpenny.[1]

6. Metric System

Need for the System. The ancient systems of measures were open to two serious objections: (1) they were planned on a varying scale instead of the scale of ten by which the civilized world always counts, and (2) they were not uniform even in any single country. Before the metric system was adopted there were, in northern France alone, eighteen different *aunes*,[2] and in the entire country there were nearly four hundred ways of expressing the area of land.

This condition was not unique in France; it was found in all European countries. Before the days of good roads and easy communication from place to place the difference in standards was not very troublesome, but by the end of the 18th century it became evident that some uniformity was essential.

Early Attempts at Reform. As early as 650 there was an effort made at uniformity in France, a standard of measure being kept in the king's palace. Under Charlemagne (*c.* 800) there

[1] See *Rara Arithmetica*, p. 470. There are various hypotheses as to the origin of the symbol $, most of them obviously fanciful.

[2] The cloth measure, the old English ell, as already given. On the metric system as a whole, see the excellent historical work of G. Bigourdan, *Le système métrique*, Paris, 1901.

was nominal uniformity throughout the kingdom, the standards of the royal court being reproduced for use in all leading cities. After Charlemagne's death, however, the numerous feudal lords adopted scales to suit their own interests. Attempts were made at various other times, as in 864, 1307, and 1558, to unify the systems in France, but none of these were successful.

Rise of the Metric System. In 1670, Gabriel Mouton, vicar of the church of St. Paul, at Lyons, proposed a system which should use the scale of 10, and which took for its basal length an arc 1′ long on a great circle of the earth. This unit he called a *milliare* or *mille*, 0.001 of a *mille* being called a *virga* and 0.1 of a *virga* being called a *virgula*. It will be seen that this was, in general plan, not unlike the metric system.

In England, Sir Christopher Wren (*c.* 1670) proposed as the linear unit the length of a pendulum beating half seconds. In France, Picard suggested (1671) the length of a pendulum beating seconds, and Huygens (1673) approved of this unit.

In order to avoid the difficulty involved in the varying length of the second pendulum in different latitudes, La Condamine suggested (1747) the use of a pendulum beating seconds at the equator, a proposal which would, if adopted, have given a standard approximating the present meter. In 1775 Messier determined with great care the length of the second pendulum for 45° of latitude, and an effort was made to adopt this as the linear unit, but it met with no success.

France works out the Metric System. In 1789 the French Académie des Sciences appointed a committee to work out a plan for a new system of measures, and the following year Sir John Miller proposed in the House of Commons a uniform system for Great Britain. About the same time Thomas Jefferson proposed to adopt a new system in the United States, taking for a basal unit the length of the second pendulum at 38° of latitude, this being the mean for this country. In 1790 the French National Assembly took part in the movement, and as a result of the widespread agitation it was decided to proceed at once with the project of unification. The second pen-

dulum was given up and an arc of one ten-millionth of a quarter of a meridian was selected as the basal unit. A careful survey was made of the length of the meridian from Barcelona to Dunkirk, but troubles with the revolutionists (1793) delayed the work. The committees which began and carried on the enterprise were changed from time to time, but they included some of the greatest scientists of France, such as Borda, Lagrange, Lavoisier, Tillet, Condorcet, Laplace, Monge, Cassini, Meusnier, Coulomb, Haüy, Brisson, Vandermonde, Legendre, Delambre, Berthollet, and Méchain. Owing to a slight error in finding the latitude of Barcelona, the original idea of the unit was not carried out, but a standard meter was fixed, and from this copies were made for use in all civilized countries.

The system was merely permissive in France until 1840, when it was made the only legal one. The expositions held in London (1851) and Paris (1855, 1867) aided greatly in making the system known outside of France. In 1919 the system was the only legal one in thirty-four countries, with a population of about 450,000,000; was optional in eleven countries, with a population of about 730,000,000; and was recognized by twenty-six countries, with a population of about 690,000,000, this recognition taking the form of assisting in the support of the International Bureau of Weights and Measures at Paris.

The modern plan of determining such standards adopts as the unit the length of a light wave of a defined type.[1]

[1] Owing to the importance of the subject and the necessity for condensing the treatment in this chapter, the following bibliographical references are added: W. S. B. Woolhouse, *Measures, Weights, and Moneys of all Nations*, 6th ed., London, 1881; F. Hultsch, *Griechische und römische Metrologie*, Berlin, 1882; A. Böckh, *Metrologische Untersuchungen*, Berlin, 1838; F. W. Clarke, *Weights, Measures, and Money of all Nations*, New York, 1875; William Harkness, "The Progress of Science as exemplified in the Art of Weighing and Measuring," *Bulletin of the Philosophical Society of Washington*, X, p. xxxix; E. Noel, *Science of Metrology*, London, 1889; C. E. Guillaume, *Les récents progrès du système métrique*, Paris, 1913. See also the various *Procès-verbaux des séances* of the *Comité international des poids et mesures*, Paris, 1876 to date; W. Cunningham, *The Growth of English Industry and Commerce*, p. 118 (London, 1896); A. De Morgan, *Arithmetical Books*, p. 5 (London, 1847); Albêrûnî's *India*, translated by E. C. Sachau, 2 vols., London, 1910, for the metrology of India *c.* 1000 (see particularly Volume I, chapter xv).

7. Time

Problem Stated. Before the time of printed calendars, when astronomical instruments were crude affairs and the astronomer was merely a court astrologer, and when the celebration on a movable feast day of the anniversary of a fixed religious event did not seem to arouse adverse criticism, even among the enemies of the various faiths, the regulation of the calendar[1] naturally ranked as one of the chief problems of mathematics.

The Computus. Accordingly there arose in all ecclesiastical schools of any standing in the Christian church the necessity for instructing some member or group of the priestly order in the process of computing the dates of Easter and the other holy days. For this purpose there were prepared short treatises on the subject. A book of this kind was generally known as a *Computus Paschalis*, *Computus Ecclesiasticus*, or, more commonly, a *Computus* or *Compotus*.[2]

General Nature of the Computi. Briefly stated, the modern form of the ancient *computus* begins with the assertion that Easter day, on which the other movable feasts of the Church depend, is the first Sunday after the full moon which happens upon or next after March 21. If the full moon happens upon a Sunday, Easter day is the Sunday following. The full moon

[1] From the Latin *kalendarium*, a list of interest payments due upon the first days of the months, the *kalendae*. The word was not used in its present sense by the Romans. They used *fasti* to indicate a list of days in which the holidays were designated.

Of the many works on the calendar, one of the latest and most extensive is F. K. Ginzel, *Handbuch der math. und techn. Chronologie. Das Zeitrechnungswesen der Völker*, 3 vols., Leipzig, 1906–1914. See also J. T. Shotwell, "The Discovery of Time," *Journal of Philosophy, Psychology, and Scientific Methods*, XII, Nos. 8, 10, 12, and *Records of Civilization, Sources and Studies*, chapter iv (New York, 1922). For a popular essay see M. B. Cotsworth, *The Evolution of Calendars*, Washington, 1922.

[2] Hieronymus Vitalis (Girolamo Vitali), *Lexicon Mathematicvm*, Paris, 1688 (Rome edition of 1690, p. 173), thus defines *computus*: "Significatio pressiùs accepta est, atque antonomasticè haesit annorum ratiocinio, & temporum distributioni, quod proprie Chronologos, & Astronomos spectat." The spelling *compotus* was at one time the more common,—possibly a kind of pun upon the convivial habits of the computers, a *compotatio*, the Greek *symposium* (συμπόσιον, *sympos'ion*), meaning a "drinking together."

is taken as the 14th day of a lunar month, "reckoned according to an ancient ecclesiastical computation and not the real or astronomical full moon."

In order to use this ancient computation it becomes necessary to be able to find the Golden Number of the year. This is done by adding 1 to the number of the year and dividing by 19, the remainder being the number sought. If the remainder is 0, the Golden Number is 19. Thus the Golden Number of 1930 is found by taking the remainder of $(1930 + 1) \div 19$, which is 12.

Taking the seven letters A, B, C, D, E, F, G, the letter A belongs to January 1, B to January 2, and so on to G, which belongs to January 7, after which A belongs to January 8, and so on. If January 2 is Sunday, the Dominical letter of the year is B. By means of the Dominical letter it is possible to find the day of the week of any given date. The finding of this letter depends upon a few simple calculations connected with tables given in the *computi*.[1]

Universality of the Problem. The problem was not confined to the Christian church. Since most early religions were connected with sun worship or with astrology, work somewhat similar to that of preparing the *computus* was needed in all religious organizations. Hence we find a problem analogous to the Christian one in the routine work of the Hebrew, Mohammedan, Brahman, and ancient Roman priests, all of whom acted as guardians of the calendar.[2] We shall now consider some of the astronomical difficulties in the way of making a scientific calendar.

[1] For a full discussion see A. De Morgan, "On the Ecclesiastical Calendar," *Companion to the Almanac for 1845*, p. 1 (London, n.d.), and "On the Earliest Printed Almanacs," *Companion to the Almanac for 1846*, p. 1.

One of the best studies of the *computus* is C. Wordsworth, *The Ancient Kalendar of the University of Oxford*, Oxford, 1904. This represents the calendar as it stood *c.* 1340.

[2] There is extant an Egyptian papyrus of about the beginning of the Christian Era that evidently was intended to serve the same purpose as the later *computi*. See W. M. Flinders Petrie, "The geographical papyrus (an almanack)," in *Two Hieroglyphic Papyri from Tanis*, published by the Egypt Exploration Fund, London, 1889.

The Day. Of the chief divisions of time the most obvious one is the day. This was, therefore, the primitive unit in the measure of time and the one which for many generations must have been looked upon as unvarying. As the race developed, however, various kinds of day were distinguished. First from the standpoint of invariability is the sidereal day, the length of time of a revolution of the earth as shown by observations on the fixed stars, namely, 23 hours 56 minutes 4.09 seconds[1] of our common time. First from the standpoint of the casual observer, however, is the true solar day, the length of time between one passage of the sun's center across the meridian and the next passage. This varies with the season, the difference between the longest and shortest days being 51 seconds; but for common purposes the solar day sufficed for thousands of years, the sundial being the means by which it was most frequently measured. As clocks became perfected a third kind of day came into use, the artificial mean solar day, the average of the variable solar days of the year, equal to 24 hours 3 minutes 56.56 seconds of sidereal time. In addition to these general and obvious kinds of day, writers on chronology distinguish others which do not concern the present discussion.[2]

The day began with the Babylonians at sunrise; with the Athenians, Jews, and various other ancient peoples, and with certain Christian sects, at sunset; with the Umbrians, at noon; and with the Roman and Egyptian priests, at midnight.[3]

The Month. The next obvious division of time was the month, originally the length of time from one new moon to the next, and one that served as the greater unit for many thousands of years. As science developed, however, it became apparent, as in the case of the day, that there are several kinds of months. There is the sidereal month, the time required for a passage of the moon about the earth as observed with reference to the

[1] All such figures are approximations, varying slightly with different authorities.

[2] See, for example, A. Drechsler, *Kalenderbüchlein*, p. 19 (Leipzig, 1881).

[3] Thus Pliny: "Ipsum diem alii aliter observavere. Babylonii inter duos solis exortus; Athenienses inter duos occasus; Umbri a meridie ad meridiem; . . . Sacerdotes Romani, et qui diem diffiniere civilem, item Aegyptii et Hipparchus, a media nocte in mediam" (*Hist. Nat.*, II, cap. 79).

fixed stars, namely, 27 days 7 hours 43 minutes 11.5 seconds. There is also the synodical month, from one conjunction of the sun and moon to the next one, averaging 29 days 12 hours 44 minutes 3 seconds, or 2 days 5 hours 0 minutes 51.5 seconds more than the sidereal month. This is the month of those who use a lunar calendar, and is the basis of the artificial month, twelve of which make our common year.[1]

The Year. Less obvious than the day or the month was the year, a period observable only about one three hundred sixty-fifth as often as the day and about one twelfth as often as the month. It took the world a long time to fix the length of the year with any degree of accuracy, and the attempt to harmonize time-reckoning by days, months, and years has given rise to as many different calendars as there have been leading races. First of all there has to be considered what constitutes a year. The sidereal year is the period of revolution of the earth about the sun, namely, 365 days 6 hours 9 minutes 9.5 seconds (365.256358 days). The tropical year is the period of apparent revolution of the sun about the earth from the instant of one vernal equinox to the next, and would be the same as the sidereal year if it were not for the slight precession of the equinox, amounting to about 50″ a year. This precession makes the length of the tropical year 365 days 5 hours 48 minutes 46.43 seconds (365.242204 days).[2] There is also the anomalistic year of 365 days 6 hours 14 minutes 23 seconds, measured from the time when the earth is nearest the sun to the next time[3] that they are in the same relative position,—a year that is slightly longer than the sidereal. There is also the lunar year of twelve synodical months, probably the first one recognized by the primitive observers of nature, and in addition to this there are various other periods which have gone by the same general name.[4]

[1] For other types of month see Drechsler, *loc. cit.*, p. 24.

[2] This was the length in the year 1800. It varies about 0.59 seconds a century. B. Peter, *Kalenderkunde*, 2d ed., p. 20 (Leipzig, 1901).

[3] From perihelion to perihelion.

[4] For the list, see Drechsler, *loc. cit.*, p. 26. On the cycle, see *ibid.*, p. 30; on the era, p. 44.

The Week. The week was less obvious than the day, the month, or even the year, having no astronomical events by which to mark its limits. It seems very likely that it arose from the need for a longer period than the day and a shorter period than the month. Hence we have the half month, known as the fortnight (fourteen nights), and the quarter month, known as the week.

Early Attempts in making a Calendar. Of the various attempts to perfect a calendar only a few will be mentioned, and in general those that had some bearing upon the Christian system.[1]

The Babylonians, whose relation to the invading Chaldeans was such as to make their later calendars substantially identical, seem to have been the first of the world's noteworthy astronomers. Aristotle relates that before 2200 B.C. they attempted scientific observations of the heavens, and Porphyrius (*c.* 275) tells us that Callisthenes (*c.* 330 B.C.) took to the Greek sage the results of a series of Chaldean observations extending over 1903 years. The Chaldeans knew the length of the year as 365 days 6 hours 11 minutes, but used both the lunar month and lunar year for civil purposes. They divided both the natural day and the natural night into twelve hours each, and in quite early times the sundial and water clock were known, the latter for use at night. For astronomical purposes the day was divided into twenty-four equal hours. They probably very early used a fourth of a month as a convenient division of time, or rather the half of the half, as was the customary way of thinking in the ancient world.

The Chinese Calendar. We are ignorant of the nature of the primitive Chinese calendars.[2] Certainly earlier than 2000 B.C.

[1] A good treatment of the subject is given by L. Ideler, *Handbuch der math. und techn. Chronologie*, 2 vols., 1825; 2d ed., Breslau, 1883, a facsimile of the first edition. Good résumés are given by Drechsler, *loc. cit.*, p. 56; Peter, *loc. cit.*, p. 5 (chiefly on the Christian calendar). From all these sources information has freely been taken. On the general question of chronology see also J. B. Biot, "Résumé de Chronologie Astronomique," *Mémoires de l'Académie des Sciences*, XXII, 209–476 (Paris). See especially Ginzel's work mentioned on page 651 n.

[2] Mikami, *China*, pp. 5, 45.

the subject occupied the attention of the astrologers. It is, however, very difficult to unravel a system which changed with each emperor, and only a few general principles can be set forth. Under the emperor Yau (*c.* 2357–*c.* 2258 B.C.) an effort was made to establish a scientific calendar for the whole country, and possibly this was done even earlier, under the emperor Huang-ti (*c.* 2700 B.C.).[1] In accordance with a decree of Wu-wang (1122 B.C.)[2] the day seems to have begun with midnight, although before this time, under the Shang dynasty (1766–1122 B.C.), it began at noon. The civil day has twelve hours, and the middle of the first hour is midnight.[3] Each hour is divided into eight parts (*khe*), each being our quarter, and each of these into fifteen *fen*, each therefore being our minute. In modern times the *fen* has been divided into sixty *miao* (seconds) under European influence. At present also the American clock is becoming common in China, so that the ancient system seems destined soon to pass away. Nevertheless the connection between the old Chinese calendar and that of Europe seems apparent. The Chinese days were named in such a way as to give seven-day periods corresponding to our weeks, and the month began, as was so often the case in early times, with the new moon. The twelve lunar months were supplemented in such a way as to harmonize the lunar and solar years, the Chang Cycle[4] being used by the Chinese before the Greeks knew of the same system under the name of the Metonic Cycle.

The Egyptian Calendar. In the ancient Egyptian calendar, which influenced all the Mediterranean countries beginning with Crete and the mainland of Greece, the business day included the night, the natural day and night being each divided into twelve hours, these hours varying in length with the

[1] Or Hoang-ti. His reign seems to have begun in the year 2704 B.C. See Volume I, page 24; Mikami, *China*, p. 2.

[2] Founder of the Chau (Cheu, Choi) dynasty, 1122–225 B.C.

[3] These hours (*shi*) are each 120 European minutes in length.

[4] For details of the complex system, and for the influence of China on Japan, see Drechsler, *loc. cit.*, pp. 71, 88.

season. The civil day seems to have commonly begun at sunset, although the priests are said by Pliny to have begun theirs at midnight. In later times, for astronomical purposes, the day began at noon[1] and was divided into twenty-four equal hours. The Romans considered the planets as ruling one hour of each day, in the following order, beginning with the first hour of Saturday: Saturn, Jupiter, Mars, Sun, Venus, Mercury, Moon, the sun and moon being placed among the "wanderers."[2] From these planets they named the days by the following plan: Taking Saturn for the first hour of Saturday and counting the hours forward, it will be seen that the second hour is ruled by Jupiter, and so on to the twenty-fourth, which is ruled by Mars. Then the next hour, the first of Sunday, is ruled by the Sun, the first hour of the next day by the Moon, and so on. Thus the days of the week were named by the ruling planets of their first hours, and we have Saturn's day, Sun's day, Moon's day, Mars's day,[3] Mercury's day,[4] Jupiter's day,[5] Venus's day,[6] a system that has come down to our time and seems destined to continue indefinitely, in spite of the fact that we are using names of heathen gods in the calendars of various religions.

Each month except the last (*Mesori*) in the native Egyptian calendar contained thirty days, five days being added to *Mesori* so as to make the year one of three hundred sixty-five days. Since this gave an error of about one fourth of a day, the year was a changing one, coming back to its original position with respect to the heavenly bodies once in 4×365 common years, or 1460 years (1461 Egyptian years). The year began with the first day of the month of Thoth, the god who, according to Plato's *Phædrus*, introduced the calendar and numbers into Egypt. As early as the 14th century B.C. the Egyptians recognized the value of a fixed year, but the changing one was too strongly implanted in the religious canons of the people to be given up. The fixed year was used to the extent of a division

[1] So with Ptolemy the astronomer, *c.* 150.

[2] "Planet" is from the Greek πλανήτης (*plane'tes*), originally a wanderer.

[3] French, *Mardi*.

[4] French, *Mercredi*.

[5] In the Northern lands, Thor's day.

[6] In the Northern lands, Frigg's day, Frigg being the goddess of marriage.

into three seasons, regulated by the river,—the Water Season, the Garden Season, and the Fruit Season,[1]—these being easily determined by the temple observers. From the temple, too, came the announcements of the turn in the rise or fall of the river, the nilometers being under the observation of the priests. These early nilometers may be seen in the temples today; they were concealed from the observation of the common people, the water being admitted by subterranean channels.

The Alexandrian Calendar. After Egypt became a Roman province (*c.* 30 B.C.) the Alexandrian calendar, including the fixed year, was introduced, although the varying year of the ancients remained in popular use until the 4th century. The Alexandrian system was used until the first half of the 7th century, when the country yielded (638) to the Mohammedan conqueror, with an attendant change of the calendar except in Upper Egypt, where the Coptic, Abyssinian, or Ethiopian churches maintained their supremacy. Since 1798, when the French obtained brief control of the country, the European system has been used side by side with the Mohammedan.

The Athenian Calendar. The Athenian calendar followed the Egyptian in beginning the new day at sunset and in dividing both day and night into twelve hours. The seven-day week was not used, however, the lunar month being divided into three parts. Of these the first consisted of ten days, numbered in order, the "5th day of the beginning of the month" being the fifth. Then followed nine days, numbered as before, but with the designation "over ten." From there to the end of the month the numbers were 20, 1 over 20, and so on, these days also being numbered backwards from the end of the month. In the popular calendar the month began with the new moon, and twelve of these months made three hundred fifty-four days, requiring the insertion of a new month[2] every three years.[3] Meton (432 B.C.) constructed a nineteen-year cycle

[1] Namely, June 21 to October 20; October 21 to February 20; February 21 to June 20.

[2] A second month of Poseideon, known as Poseideon II.

[3] The detailed variations of this plan need not be considered here. See any work on the calendar.

in which the third, fifth, eighth, eleventh, thirteenth, sixteenth, and nineteenth years should contain the extra month,[1] a plan which Callippus, a century later (325 B.C.), modified to include four nineteen-year cycles.[2] Still later Hipparchus (150 B.C.) suggested the use of four of the cycles of Callippus, less a day, or 110,036 days in all, but neither of the last two calendars came into popular use.

Roman Calendar. The oldest of the Roman calendars seems to have been the one attributed to Romulus. The year probably consisted of ten months of varying length, or of 304 days, beginning with March. Numa Pompilius (715–672 B.C.) is said to have added two other months, January and February, and his year was probably lunar. The Decemvirs (5th century B.C.) decreed a solar year, the regulation of which was left in the hands of the priests. The calendar was so mismanaged, however, that by the time of Julius Cæsar each day was eighty days out of its astronomical place, and radical measures were necessary for its reform. Cæsar therefore decreed that the year 46 B.C. should have four hundred forty-five days[3] and that thereafter the year should consist of three hundred sixty-five days, with a leap year every fourth year.[4]

Names of the Months. Following the older custom as to the beginning of the year in March, Caesar at first used the following plan for the calendar:

1. Martius,	31 days.	7. Septembris,	30 days.
2. Aprilis,	30 days.	8. Octobris,	31 days.
3. Maius,	31 days.	9. Novembris,	30 days.
4. Junius,	30 days.	10. Decembris,	31 days.
5. Quintilis,	31 days.	11. Januarius,	31 days.
6. Sextilis,	31 days.	12. Februarius,	28 days.

[1] 19 years = 235 months = 6939$\frac{3}{4}$ days. The months as arranged, however, contained 6940$\frac{2}{3}$ days.

[2] 4 × 19 years = 76 years = 940 months. The months were 29 or 30 days and totaled 27,759 days.

[3] Hence called *annus confusionis*.

[4] A calendar of *c.* 100 B.C. was recently found at Anzio, in the Campagna, based upon a lunar year of 353 days with an intercalary month of 27 days on alternate years.

This accounts for our names "September" (7th month), "October," "November," and "December." On his original plan every alternate month, beginning with March, had thirty-one days, the others having thirty days, except that February received its thirtieth day only once in four years. Cæsar later decreed that the year should begin with January, and finally, but during his life, the name of Quintilis, the month in which he was born, was changed to Julius. He also changed the number of days in certain months, and the result appears in our present calendar. After his death, in the second year of his calendar, a further confusion arose, apparently through a misunderstanding on the part of the priests as to the proper date for leap year. This was corrected by Augustus, and in his honor the name of Sextilis was changed to bear his name. From that time on the Julian calendar remained in use until its reformation under Gregory XIII in 1582, and it was used by the Greek Catholics, including the Russians, until the World War of 1914–1918, the dates until that time differing by thirteen days from those of the calendar of Western Europe.

Christian Calendar. The indebtedness of the present European calendar to those already described is apparent, and it is also evident that our calendar has had an extensive history. The beginning of the year, for example, has not been uniform from time to time and in different countries. In the early centuries the year usually began with April in the East[1] and with March in the West, although sometimes with the Feast of the Conception, Christmas day,[2] Easter, or Ascension day, or at other times according to the fancy of the popes. Finally Innocent XII again decreed that the year should begin on

[1] Although the Byzantine calendar began with September 1.

[2] As in Spain until the 16th century and in Germany from the 11th century. March 1 and March 25 (the Annunciation) were favorite dates, although Advent Sunday (the fourth Sunday before Christmas) has generally been recognized as the beginning of the ecclesiastical year. March 1 was used generally in medieval France, in Oriental Christendom, and (until 1797) in Venice. March 25 was used by the medieval Pisans and Florentines. Most of the Italian states adopted January 1 in 1750. For further details consult Ginzel (see page 651 n.) or such works as A. Drechsler, *Kalenderbüchlein*, p. 77 (Leipzig, 1881).

January 1, beginning with 1691, as Philip II had done for the Netherlands in 1575,[1] and as Julius Cæsar had done before the Christian Era.

Numbering the Years. Following the Roman custom, the years in the early centuries of Christianity were dated from the accession of the emperor or consul. We have a relic of this in the dating of acts of parliament in England and of presidential proclamations in America.[2] It was not until the abbot Dionysius Exiguus[3] (533) arranged the Christian calendar that the supposed[4] date of the birth of Christ was generally taken for the beginning of our era, Christmas day being therefore appropriately selected as the first day of the year 1. This calendar was adopted in Rome in the 6th century,[5] in various other Christian countries in the 7th century, and generally throughout Europe in the 8th century.[6]

Changes in Easter. Not only the beginning of the year but the determination of Easter has been the subject of much change. We now consider Easter as the first Sunday after the first full moon following the vernal equinox,[7] as decreed in 325 by the Council of Nice (Nicæa). Formerly it fell on the date

[1] England adopted January 1 in 1752.

[2] As "in the 150th year of our independence."

[3] Dionysius the Little. He went to Rome *c.* 500 and died there in 540.

[4] He considered the birth of Christ as taking place in the year 754 of the founding of Rome, although early Christians placed it in the year 750.

[5] There are, however, no extant inscriptions of the 6th century which bear dates in the Christian Era. See M. Armellini, *Archeologia Cristiana*, p. 479 (Rome, 1898). Sporadic efforts had been made before the 6th century to use a Christian calendar. The oldest known specimen of such a calendar dates from 354. See B. Peter, *Kalenderkunde*, 2d ed., p. 4 (Leipzig, 1901).

[6] The exceptions were the Spanish peninsula and Southern France. Charlemagne was the first great ruler to use (783) the Dionysian calendar.

[7] In Rome. It is possible to have a difference of a week between this Sunday in Rome and in (say) Honolulu, the full moon occurring on Sunday in Rome when it is still Saturday in some places to the west. This has occasionally occurred as an astronomical fact although not as an ecclesiastical one. It should be understood that, for Church purposes, March 21 is taken as the date of the vernal equinox, and that the full moon is not determined by modern astronomy but by certain rules as laid down, say, in the Book of Common Prayer. Easter, therefore, now varies from March 22 to April 25. For a good résumé of the Easter problem see Peter, *loc. cit.*, p. 58.

of the Jewish Passover, but in order to avoid this coincidence the Church readjusted its calendar. Justinian, with this in view, decreed (547) that Easter should be 21 days (instead of 14 days) after the first new moon after March 7. In this way, in general, the Passover and Easter do not come together, although occasionally they synchronize.[1]

The Gregorian Calendar. The present calendar of Western Europe and the Americas, the so-called Gregorian calendar, was necessitated by the fact that the year is not $365\frac{1}{4}$ days long, as recognized by Cæsar, but is about 11 minutes 14 seconds shorter than this. Therefore once in 128 years the Julian calendar receded one day from the astronomical norm, and by the close of the 14th century the departure of Easter day from its traditional position became so noticeable that it was the subject of much comment. It was not until Gregory XIII, however, consulting with such scientists as Aloysius Lilius[2] and Christopher Clavius (*c.* 1575), determined on a reform, that anything was really accomplished. He decreed that October 4, 1582, should be called October 15, and that from the total number of leap years there should be dropped three in every four centuries. In particular he decreed that only such centennial years as are divisible by 400 (1600, 2000, 2400, etc.) should be leap years. This requires no further adjusting of the calendar for over 3000 years. Italy, Spain, Portugal, Poland, France, and a part of the Netherlands adopted this calendar in 1582. In 1583 it was recognized in part of Germany, the old style being also used until 1700. Part of Switzerland adopted it in 1584, and the other part, together with Denmark and the rest of the Netherlands, in 1700. It was also adopted in Hungary in 1587, in Prussia in 1610, in England in 1752, and in Sweden in 1753. So fixed had the Julian calendar become in the minds of the people, however, that even as late as the opening of the 19th century O. S. (old style) and N. S. (new style)

[1] As in 1805, 1825, 1828, 1832, and early in the present century, on April 12, 1903. Among the many suggestions for Easter is that of Jean Bernoulli that it should be the first Sunday after March 21, without reference to the moon.

[2] Ludovico Lilio, Luigi Lilio Ghiraldi (1510–1576).

were used in dating letters in America, while writers on arithmetic felt it necessary to include a description of the Julian and Gregorian calendars as late as the close of that century. The changes brought about by the World War of 1914–1918 led to a more general adoption of the Gregorian calendar in the few countries which had continued to use the Julian or other types.

The Calendar in the French Revolution. In the early days of the French Revolution an attempt was made to impose a new calendar upon the country, partly as a protest against the Christian church. It was hoped that this reform, like that which resulted in the metric system, would receive international recognition. The new era was to begin with the autumnal equinox which occurred on September 22, 1792. There were twelve months of thirty days each, and these months were divided into decades in which the days were named numerically,—Primidi, Duodi, and so on. The extra five or six days of the year were grouped at the end as holidays. The months were named according to natural conditions, thus: In autumn, Vendémiaire (vintage), September 22–October 21; Brumaire (fog), October 22–November 20; Frimaire (sleet), November 21–December 20. In winter, Nivôse (snow), December 21–January 19; Pluviôse (rain), January 20–February 18; Ventôse (wind), February 19–March 20. In spring, Germinal (seed), March 21–April 19; Floréal (blossom), April 20–May 19; Prairial (pasture), May 20–June 18. In summer, Messidor (harvest), June 19–July 18; Fervidor or Thermidor (heat), July 19–August 17; Fructidor (fruit), August 18–September 16. As might have been known, the scheme failed, and on August 30, 1805, a decree was signed reëstablishing the Gregorian calendar, beginning January 1, 1806.[1]

Other Calendars. The other calendars are of no special interest in the history of mathematics. The Hindus began their year with the day of the first new moon after the vernal equinox. The Jews begin their day at sunset, their week on Saturday night (*i.e.*, when their holy day ends and Sunday be-

[1] This decree may still be seen in the Musée des Archives Nationales, in Paris.

gins), and their year with *Tishri* 1.[1] Their calendar, more lunar than ours, is quite complicated.

The Maya civilization[2] had a curious system, the year beginning with the winter solstice and being divided into eighteen months, entirely independent of astronomical considerations. Scholars have recently asserted that their calendar goes back as far as the 34th century B.C.

The Mohammedans begin their day with sunset, and, like many other Eastern peoples, divide both daytime and nighttime into twelve hours, the length of the hour varying with the season. The week begins on Sunday, and Friday is the day of rest. Their month begins with the new moon, and the year is purely lunar, of 354 or 355 days. The year 1343 A.H. began on August 2, 1924, of the Christian calendar. The era began with the Hejira, the flight of Mohammed from Mecca on July 15 or 16, 622. On account of frequent references to the Mohammedan calendar in literature, it may be added that a simple rule, accurate enough for practical purposes, for translating a year of the Hejira into a Christian year is as follows: To 97 per cent of the number of the year add 622; the result is the Christian year. Thus 1326 A.H. = 97 per cent × 1326 + 622 = 1908 A.D.

Early Christian Computi. The first noteworthy Christian work[3] on the calendar was that of Victorius of Aquitania (457). About a century later a second *Computus Paschalis* appeared, probably written by Cassiodorus (562). In the next century the question of Easter had become so complicated as to cause (664) a dispute between the church in England and the authorities in Rome. The best of the early works on the

[1] *I.e.*, the first new moon after the autumnal equinox. Their year formerly began with *Nisan*, their seventh month, thus using, like that of the Hindus, the vernal equinox. In 1908 *Tishri* 1 was September 26 of the Christian calendar. See S. B. Burnaby, *Elements of the Jewish and Muhammedan Calendars*, London, 1901.

[2] S. G. Morley, *An Introduction to the Study of the Maya Hieroglyphs*, Washington, 1915; C. P. Bowditch, *The Numeration, Calendar Systems, and Astronomical Knowledge of the Mayas*, Cambridge, Massachusetts, 1910.

[3] On this work and the works of later scholars on the same subject see B. Lefebvre, *Notes d'Histoire des Mathématiques*, p. 39 (Louvain, 1920).

computus is the one written by Bede[1] in the 8th century. This contains a precise statement as to the method of finding the date of Easter in any year.

In the 9th century both Hrabanus Maurus (*c.* 820) and Alcuin (*c.* 775) wrote upon the problem, and Charlemagne thought the subject so important that he urged that it be considered in every monastery.[2]

Medieval Works. Lectures were held upon the subject in the 13th and 14th centuries in the various European universities. Sacrobosco (*c.* 1250) wrote a work on it,[3] such a practical calculator as Paolo Dagomari (*c.* 1340) did not hesitate to do the same, and even the Jewish scholars contributed treatises on the Christian calendar as well as their own.[4] It is to a commentary by Andalò di Negro (*c.* 1300) on a work by Jacob ben Machir (d. 1307) that we owe the first prominent use in Europe of the Arabic word *almanac*,[5] later brought into general use by such writers as Peurbach (*c.* 1460) and Regiomontanus (*c.* 1470).

Printed Works. The first printed *computus* was that of Anianus.[6] In this work there appears the original of the familiar rime beginning "Thirty days hath September."[7]

[1] *De temporum ratione.*

[2] D. C. Munro, *Selections from the Laws of Charles the Great*, p. 15 (Philadelphia, 1900), "Admonitio generalis," 789. See also T. Ziegler, *Geschichte der Pädagogik*, p. 28 (Munich, 1895); Günther, *Math. Unterrichts*, p. 66.

[3] *Libellus de anni ratione, seu ut vocatur vulgo computus ecclesiasticus.*

[4] There is a MS. now in Petrograd, written by Jechiel ben Josef (1302), under the title *Injan Sod ha-Ibbur*, with a chapter on the Christian *computus*. See M. Steinschneider, "Die Mathematik bei den Juden," *Bibl. Math.*, XI (2), 16, 38, 74; XII (2), 5, 33.

[5] Heb. אלמנך, from the Arab. *al-manākh*. The word is not pure Arabic, however, and the real origin is unknown. See Boncompagni's *Bullettino*, IX, 595; Günther, *Math. Unterrichts*, p. 190 n.


[6] *Cōputus manualis magri aniani. metricus cū?mēto* (Strasburg, 1488). There is said to have been an edition printed at Rome in 1486. For bibliography of Anianus see C. Wordsworth, *The Ancient Kalendar of the University of Oxford*, p. 113 (Oxford, 1904). See also the facsimile on page 668.

[7] The Latin form as given by Anianus is as follows:

> Junius aprils september et ipse nouember
> Dant triginta dies reliquis supadditur vnus,
> De quorum numero februarius excipiatur.—Fol. B 8

See *Rara Arithmetica*, p. 33.

This was not original with Anianus, however, for it is found in various medieval manuscripts.[1] It first appeared in English verse in 1590.




A COMPUTUS OF 1393

In this MS. there appears, in Italian, the verse "Thirty days hath September"

[1] *E.g.*, in the above anonymous Italian MS. of 1393, beginning:

"Trenta di a nouembre apile giugno & settembre."

The MS. is in Mr. Plimpton's library. See *Rara Arithmetica*, p. 443.


Sunt aries thaurus gemini
cancer leo virgo libra sc
scorpius sagittarius caper
amphora pisces

FROM A COMPUTUS OF 1476

This page shows the usual verses beginning "Sunt aries, thaurus, gemini, cancer." The work also contains the verses, in Latin, beginning "Thirty days hath September," which are found as early as the 13th century. From Mr. Plimpton's library

Liber qui Computus inscribitur: vna cum figuris et ma/
nibus necessariis tam in suis locis q̄ in fine libri positis.
Incipit feliciter .

Lux orta est iusto Psalmista Ista ver
ba possunt dupliciter considerari. Primo pñt
dici de deo q̄ est lux vera. ideo dicebat dauid.
lux orta est iusto. Et de ista luce dicit Johā. i.
Erat lux vera q̄ illuminat omnē hominē venie/
tē in hunc mundū. Secūdo potest de scientia.
Et dicit lux quasi scientē reddētē lucidū. quia facit hominē scie/
tem esse lucidū. In quibus verbis ad cōmendationē scie duo bre
uiter tangūtur. Primo enī tangit scientie altitudo preciosa p hoc
quod dicit lux. Secundo largitudo gloriosa per hoc q̄ dicit ori/
ta est. Primū probatur auctoritate et ratiōe. auctoritate Isidori
sic dicentis / Scientia est fons indeficiens. bonitatis via. sui sal/
uatoris cognitio. Ratiōe sic. illud est validū et preciosum quod
de inualido et imperfecto facit validum et pfectum. scientia ē huius
modi. ergo ꝛc. maior est manifesta. minor declaratur p phīm ter/
cio de aīa sic dicentē: Aīa in principio sue creationis ē tanq̄ tabu
la rasa in qua nihil depictū est. depingibilis nī scientiis et virtuti/
bus. Primum probat auctoritate boetii et ratiōe in prologo arit/
metice / Scientia ē eor̄ que vera sunt et impmutabilis essentie no
straq̄ comphensione veritatis. Ratiōe sic: illud est tanq̄ summū
bonū quod habet largitionē gloriosam scia ē hmōi. ergo ꝛc. ma
ior est vera. minor pbatur p diffinitionē scientie q̄ talis ē. Scia est
ddā habitus aie rōnalis nō innatus sed acqsitus oīm hūanar̄ re
rum indagatrix et totius humane vite gubernatrix / q̄ scia sit ha/
bitus ptz. q̄ scia est aliquod existens in aīa. sed omne illud quod
est in aīa aut ē habitus aut potētia aut passio. Et hoc testat Aristo.
in secūdo ethicor̄. q̄ scia nō sit passio ptz q̄ passiones sunt in vo
luntate scia nō ē hmōi. ergo ꝛc. q̄ nō sit potētia ptz. q̄ ēl:bz po/
tentia sit a natura. sicut irascibilis. et cōcupiscabilis. et sic relinqui
tur q̄ nō sit potentia. q̄ sit habitus aie rōnalis ptz p ddicta. q̄ aūt

a ii

THE COMPUTUS OF ANIANUS (1488)

First page of the edition of *c.* 1495. The work of Anianus was the first one printed on the *computus*. See page 665

Among the prominent *computi* printed in the 16th century is that of Arnaldo de Villa Nova (*c.* 1275).[1] Many of the early arithmetics also gave a brief treatment of the *computus*.[2]

The *computus* finally found a place in various liturgical works, and at present can be conveniently studied in pages prefixed to most editions of the Book of Common Prayer.

Early Timepieces. One of the general problems in connection with the calendar has to do with the finding of the hours of the day. For this purpose the shadow cast by some obstruction to the sun's rays was probably used by all primitive peoples. At first it is probable that a prominent tree, a rock, or a hill was selected, but in due time an artificial gnomon[3] was erected and lines were drawn on the earth to mark off the shadows. Since the hour shadow is longer when the sun is near the horizon, either concave surfaces or curve lines on a plane were placed at the foot of the gnomon.

Hours. The ancients usually had twelve hours in the day and the same number in the night. There have been various speculations as to why twelve was selected for this purpose, among them being one which referred the custom to the Babylonian knowledge of the inscribed hexagon.[4] It is probable, however, that twelve was used in measuring time for the same reason that it was used for measuring length and weight,—because the common fractional parts (halves, thirds, and fourths) were easily obtained. The day hours were longer than the night hours in the summer and shorter in the winter, a fact referred to by several ancient writers.[5]

[1] *Computus Ecclesiasticus & Astronomicus*, Venice, 1501. There was another work by this name printed at Venice in 1519.

[2] *E.g.*, the Treviso arithmetic (1478), fol. 57. Köbel's *Rechenbüchlin* (1531 ed.) devotes ten pages to the subject.

[3] Herodotus uses this term γνώμων (*gno'mon*), and it is common in Greek literature. Later writers sometimes called it a horologe (ὡρολόγιον, *horolog'ion*) when used for the sundial specifically. In still later times it was called the pole (πόλος, *pol'os*).

[4] G. H. Martini, *Abhandlung von den Sonnenuhren der Alten*, p. 18 (Leipzig, 1777).

[5] So Vitruvius: "Brumalis horae brevitates"; St. Augustine: "Hora brumalis aestivae horae comparata, minor est."

Although Herodotus (II, 109) speaks of the "12 parts" of the day among the Babylonians and the Greeks, the word "hour"[1] was not used either by him or by Plato or Aristotle. It was apparently a later idea to give these divisions a special name.

Early Dials. The sundial seems to have been used first in Egypt, but it is found also at an early date in Babylonia. Herodotus (II, 109) says that it was introduced from Babylon into Greece, and tradition says that this was done by Anaximander (*c.* 575 B.C.), the gnomon being placed at the center of three concentric circles. The early Egyptian dial has already been mentioned in Volume I, page 50. The first concave dial to be used in Greece is said to have been erected on the island of Cos by Berosus.[2] Several such dials have been found in the Roman remains,[3] and the early ones have no numerals on the hour lines, these lines being easily distinguished without such aids.[4]

Besides the plane dials and the concave spherical dials there were both concave and convex cylindric forms. Vitruvius (*c.* 20 B.C.) tells us that one Dionysodorus[5] invented the cylindric

[1] Ὥρα (*ho'ra*). There may be some relation between the word and the name of the Egyptian Horus, god of the rising sun, and the Hebrew *or* (light).

[2] A priest of Belus at Babylon. The name was probably Bar (Ber) Oseas or Barosus, that is, son of Oseas. Fl. *c.* 250 B.C. Vitruvius (IX, 4; X, 7, 9) says that Berosus went to Cos in his later years, founded a school of astrology, and invented what seems to have been a hemispherical sundial. For a general description of early dials see G. H. Martini, *loc. cit.*, pp. 24, 70. On Anaximander's connection with the gnomon, as recorded by Diogenes Laertius, Favorinus, and Herodotus, see W. A. Heidel, "Anaximander's Book, the earliest known geographical treatise," *Proceedings of the American Academy of Arts and Sciences*, LVI, 239.

[3] The first one was discovered (1741) in a Tusculan villa. For early descriptions see G. L. Zuzzeri, *D'una antica villa scoperta sul dosso del Tusculo, e d'un antico orologio a sole*, Venice, 1746. Boscovich also described it in the *Giornale de' Letterati per . . . 1746*, art. 14. The second one was found (1751) at Castelnuovo, near Rome; the third, also near Rome; and the fourth (1764), at Pompeii, although apparently made in Egypt.

[4] Thus Persius (*Satires*, III, 4): ". . . quinta dum linea tangitur umbra," the shadow resting on the fifth line of the day, an hour before noon.

[5] Also spelled Dionysiodorus. He is said by Pliny to have found the radius of the earth to be *c.* 5000 miles, but nothing is known of his life. He lived *c.* 50 B.C.

form of dial, but we do not know whether it was convex or concave. We also know from Vitruvius that there were various other forms in use by the Romans.[1]

Difficulties with the Gnomon. One difficulty that was experienced with the large sundials of the ancients was that the gnomon did not cast a distinct shadow. The size of the sun is such as to have the shadow terminate in a penumbra which rendered the determination of the solstice, for example, a difficult matter. This is one reason why it is not probable that the Egyptian obelisks were used by scientific observers as gnomons. To overcome the difficulty the Greeks often used a column with a sphere on top, the center of the sphere corresponding to the center of the shadow and the center of the sun. Such gnomons are found on medals of the time of Philip of Macedon, and it is possible that this is the explanation of the column on the coin of Pythagoras shown on page 70 of Volume I. Dials of this type were introduced in Rome by Menelaus (*c.* 100), or at least were improved by him.[2]

As would naturally be expected, there were many special forms of dials. The "dial of Ahaz" (Isaiah, xxxviii), for example, was probably a flight of stairs, very likely curved, upon which a ray of sunlight fell. This dates from about the 8th century B.C.

It is impossible to do more at this time than to refer thus briefly to the use of dials among the ancients. The literature of the subject is very extensive.

Hourglasses and Clepsydræ. The need was early felt for some kind of device to tell the hours at night as well as during the day, and in cloudy weather as well as when the sun's direct rays gave their aid. Various methods were employed, such as burning tapers, hourglasses, and water clocks. The hourglass was known probably as early as 250 B.C. Plato (*c.* 380 B.C.) gave much thought to the matter, and his conclusions may have

[1] "Aliaque genera et qui supra scripti sunt, et alii plures inventa reliquerunt."
[2] On this type see Bigourdan, *L'Astronomie*, p. 91.

suggested to Ctesib'ius (*c.* 150 B.C.)[1] the idea of a water clock, the clepsydra,[2] which the second P. Cornelius Scipio Nasica ("Scipio with the pointed nose") is said to have introduced into Rome (*c.* 159 B.C.). In the early forms of these clocks the water trickled from one receptacle to another in a given time, much as the sand flows in an hourglass, but the later forms were more complicated. It was such an instrument that Harun al-Rashid sent to Charlemagne in 807. A clepsydra of the primitive type was until recently in operation in one of the ancient towers of Canton, China.

Influence on Later Timepieces. Since the priesthood, which composed the learned class, kept account of the official time in the early days of civilization, the Church continued to undertake this task until the modern period. The priest tolled the hour as determined by the dial or hourglass. The dial was put in a conspicuous place in the town, on the church tower, and hence in modern times the clock is often seen in the church tower and the hours are struck on the church bell. Because of this fact we have our name "clock," a word probably derived from the Celtic and meaning bell, whence the French *cloche*, a bell.

When the hour lines were marked on the dial, Roman numerals were used, always with IIII instead of IV for four, and hence we see the same markings upon modern clocks. The ancient gnomon was under the care of the priests, and brass plates are still to be seen in the floors of some of the churches in the Mediterranean countries, the sun shining through a certain window and telling the seasons as marked upon them.[3]

[1] Κτησίβιος, a native of Alexandria. None of his works are extant, but he is said to have invented not only a water clock but also a hydraulic organ and other machines.

[2] Κλεψύδρα (*klepsy'dra*). See Volume I, page 69 n.; Vitruvius, *De Architectura*, IX, cap. 9; Pliny, *Hist. Nat.*, VII.

[3] Those who wish to obtain further information upon the subject should consult such encyclopedias as the *Britannica* and such works as the following: A. Fraenkel, "Die Berechnung des Osterfestes," Crelle's *Journal*, CXXXVIII, 133; E. M. Plunket, *Ancient Calendars and Constellations*, London, 1903; C. P. Bowditch, "Memoranda on the Maya Calendars," *American Anthropologist*, III (N.S.), 129.

Clocks. It should not be thought that clocks of the general form known at the present time date only from Galileo's discovery of the isochronal property of the pendulum. As a matter of fact, wheel clocks go back to Roman times, and Boethius is said to have invented one (*c.* 510). Such clocks are known to have been used in churches as early as 612. The invention of those driven by weights is ascribed to Pacificus, archdeacon of Verona, in the 9th century, although a similar claim is made on behalf of various others. Clocks involving an assemblage of wheels are medieval in origin, and one was set up in St. Paul's, London, as early as 1286. Small portable clocks were in use in the 15th century, as witness a letter of 1469, written by Sir John Paston and containing the following admonition:

> I praye you speke wt Harcourt off the Abbeye ffor a lytell clokke whyche I sent him by James Gressham to amend and yt ye woll get it off him an it be redy.

The oldest mechanical clock of which we have any complete description was made by a German named Heinrich De Vick and was set up in the tower of the palace of Charles V of France in 1379.[1] The principle employed was that of a weight suspended by a cord which was wound about a cylinder. This cylinder communicated power to a train of geared wheels which, in turn, transformed by means of a "scape wheel" the rotary motion to a backward-and-forward motion controlling the hands. The tendency of the weight to descend too rapidly was checked by a device for regulating the action of the wheels.

The pendulum clock was introduced about 1657 and seems chiefly due to Huygens. The principle of the pendulum, properly attributed to Galileo, had been observed as early as the 12th century by Ibn Yûnis (*c.* 1200), and had been employed

[1] On the general topic see W. I. Milham, *Time and Timekeepers*, New York, 1923, with full bibliography; E. von Bassermann-Jordan, *Die Geschichte der Zeitmessung und der Uhren* (Berlin, 1920–); R. T. Gunther, *Early Science in Oxford*, Vol. II (Oxford, 1923); H. T. Wade, "Clocks," *New International Cyclopædia*, V, 470 (2d ed., New York, 1914).

by astronomers to estimate intervals of time elapsing during an observation, but it had not been applied to a clock. It was made known in England through Ahasuerus Fromanteel, a Dutch clockmaker, about 1662.

John Harrison's[1] great construction of a ship's chronometer with a high degree of precision was made in the second half of the 18th century and finally secured for him the prize of £20,000 offered by the British government in 1714 for a method of ascertaining within specified limits the longitude of a ship at sea.[2] At the present time the noon of Greenwich mean time (G. M. T.) is communicated to ships by wireless, and so, in the case of the larger vessels, the finding of longitude no longer depends upon the chronometer alone.

Harrison's contribution to practical navigation was so important as to warrant a brief statement about the nature of his work. Although he was by trade a carpenter, his mechanical tastes led him to experiment with clocks. Having observed the need for a pendulum of constant length, he devised (1726) the "gridiron pendulum," in which the downward expansion of the steel rods compensated for the upward expansion of the brass ones. After the British government (1714)[3] had offered the prize already mentioned, Harrison gave his attention to the perfection of a watch that should serve to assure Greenwich time to a ship at sea. By 1761 he had constructed one that, after a voyage of several months, had lost only 1 min. 54½ sec. and assured the longitude within 18 miles. The government paid him £10,000 in 1765, and a like sum in 1767,—a modest reward for an invention of such great value to the world, even though the degree of accuracy would now be considered very unsatisfactory.

[1] Born at Foulby, parish of Wragby, Yorkshire, early in 1693; died in London, March 24, 1776.

[2] For a list of such prizes see Bigourdan, *L'Astronomie*, p. 166.

[3] The original act reads, "At the Parliament to be Held at *Westminster*, the Twelfth Day of *November*, *Anno Dom.* 1713," but was printed (and doubtless enacted) in 1714. It is entitled "An Act for Providing a Publick Reward for such Person or Persons as shall Discover the Longitude at Sea," there being nothing "so much wanted and desired at Sea, as the Discovery of the Longitude." On the entire topic see R. T. Gould, *The Marine Chronometer, its History and Development*, London, 1923.

TOPICS FOR DISCUSSION

1. Additional information concerning weight, length, area, and volume as found in the various encyclopedias.

2. The cubit, shekel, talent, and various other measures referred to in Biblical literature.

3. The development of measures, including weights, in accord with human needs.

4. An etymological study of such words as *metric*, *groschen*, *doubloon*, *measure*, *watch*, *day*, *month*, and *year*.

5. Influence of the Roman system of measures, including weights and values, upon other European systems.

6. The universality of certain primitive units of measure such as the cubit and inch.

7. Primitive customs as related to units of measure, such as "a day's journey," "a watch in the night," and a "Morgen."

8. Primitive measures still found in various parts of the country, having been transmitted from generation to generation like the folklore of the people.

9. Supplementary information on the calendar, as found in the various encyclopedias.

10. Meaning and use of the "golden number" and "dominical letter" as set forth in various encyclopedias or in works on the Church calendar.

11. Methods of finding the date of Easter as given in the books referred to in the preceding topic.

12. The reform of the calendar under Gregory XIII, including a study of earlier attempts at reform.

13. Significance of such technical terms as *computus Paschalis*, *calendar*, *a red-letter day*, and *almanac*.

14. History of the discovery of the approximate length of the year, and the method of ascertaining it.

15. Relation among calendars used by various peoples of ancient and medieval times.

16. Influence of Rome upon the calendars of European countries, and the effort of France, during the Revolution, to break away from tradition with respect to the divisions of the year and the names of the days and months.

17. The mathematics of the sundial.

CHAPTER X

THE CALCULUS

1. Greek Ideas of a Calculus

General Steps Described. There have been four general steps in the development of what we commonly call the calculus, and these will be mentioned briefly in this chapter. The first is found among the Greeks.[1] In passing from commensurable to incommensurable magnitudes their mathematicians had recourse to the method of exhaustion, whereby, for example, they "exhausted" the area between a circle and an inscribed regular polygon, as in the work of Antiphon (*c.* 430 B.C.).

The second general step in the development, taken two thousand years later, may be briefly called the method of infinitesimals. This method began to attract attention in the first half of the 17th century, particularly in the works of Kepler (1616) and Cavalieri (1635), and was used to some extent by Newton and Leibniz.

The third method is that of fluxions and is the one due chiefly to Newton (*c.* 1665). It is this form of the calculus that is usually understood when the invention of the science is referred to him.

The fourth method, that of limits, is also due to Newton, and is the one now generally followed.

Contributions of the Greeks. As stated above, the Greeks developed the method of exhaustion about the 5th century B.C. The chief names connected with this method have already been

[1] Sir T. L. Heath, "Greek Geometry with special reference to infinitesimals," *Mathematical Gazette*, March, 1923; D. E. Smith, *Mathematics*, in the series "Our Debt to Greece and Rome," Boston, 1923; G. H. Graves, "Development of the Fundamental Ideas of the Differential Calculus," *The Mathematics Teacher*, III, 82.

mentioned, but a few details of their work and that of their contemporaries will now be given.

Zeno of Elea (*c.* 450 B.C.) was one of the first to introduce problems that led to a consideration of infinitesimal magnitudes. He argued that motion was impossible, for this reason:

> Before a moving body can arrive at its destination it must have arrived at the middle of its path; before getting there it must have accomplished the half of that distance, and so on *ad infinitum*: in short, every body, in order to move from one place to another, must pass through an infinite number of spaces, which is impossible.[1]

Leucippus (*c.* 440 B.C.) may possibly have been a pupil of Zeno's. Very little is known of his life and we are not at all certain of the time in which he lived, but Diogenes Laertius (2d century) speaks of him as the teacher of Democritus (*c.* 400 B.C.). He and Democritus are generally considered as the founders of the atomistic school, which taught that magnitudes are composed of indivisible elements[2] in finite numbers. It was this philosophy that led Aristotle (*c.* 340 B.C.) to write a book on indivisible lines.[3]

Democritus is said to have written on incommensurable lines and solids, but his works are lost, except for fragments, and we are ignorant of his method of using the atomic theory.

Method of Exhaustion. Antiphon (*c.* 430) is one of the earliest writers whose use of the method of exhaustion is fairly well known to us. In a fragment of Eudemus (*c.* 335 B.C.), conjecturally restored by Dr. Allman,[4] we have the following description:

> Antiphon, having drawn a circle, inscribed in it one of those polygons[5] that can be inscribed: let it be a square. Then he bisected each side of this square, and through the points of section drew straight lines at right angles to them, producing them to meet the cir-

[1] Allman, *Greek Geom.*, p. 55.

[2] Ἄτομοι (*a'tomoi*). Allman, *Greek Geom.*, p. 56.

[3] Περὶ ἀτόμων γραμμῶν (first edition, Paris, 1557). The work is also attributed to Theophrastus.

[4] Allman, *Greek Geom.*, p. 65.

[5] That is, according to the usage of the time, regular polygons.

cumference; these lines evidently bisect the corresponding segments of the circle. He then joined the new points of section to the ends of the sides of the square, so that four triangles were formed, and the whole inscribed figure became an octagon. And again, in the same way, he bisected each of the sides of the octagon, and drew from the points of bisection perpendiculars; he then joined the points where these perpendiculars met the circumference with the extremities of the octagon, and thus formed an inscribed figure of sixteen sides. Again, in the same manner, bisecting the sides of the inscribed polygon of sixteen sides, and drawing straight lines, he formed a polygon of twice as many sides; and doing the same again and again, until he had exhausted the surface, he concluded that in this manner a polygon would be inscribed in the circle, the sides of which, on account of their minuteness, would coincide with the circumference of the circle.

We have in this method a crude approach to the integration of the 17th century.

Bryson (*c.* 450 B.C.), who seems to have lived just before Antiphon's period of greatest activity, was at one time thought to have used a method that had the merit of circumscribing as well as inscribing regular polygons and exhausting the area between them. This was probably not the case (Vol. I, p. 84), although the method was used by some of his successors. There is also no reliable evidence to prove the assertion that Bryson assumed that the area of the circle is the arithmetic mean between the areas of two similar polygons, one circumscribed and the other inscribed.

The Contribution of Eudoxus. Eudoxus of Cnidus (*c.* 370 B.C.) is probably the one who placed the theory of exhaustion on a scientific basis. It is uncertain just how much reliance is to be placed upon the tradition which asserts that Book V of Euclid's *Elements* (the book on proportion) is due to him, but it is thought that the fundamental principles there laid down are his. The fourth definition in Book V is: "Magnitudes are said to have a ratio to one another which are capable, when multiplied, of exceeding one another," and this excludes the relation of a finite magnitude to a magnitude of the same kind

which is either infinitely great or infinitely small.[1] It is in this definition and the related axiom that Dr. Allman finds a basis for the scientific method of exhaustion and discerns the probable influence of Eudoxus. According to Archimedes, this method had already been applied by Democritus (*c.* 400 B.C.) to the mensuration of both the cone and the cylinder.

It is known that Hippocrates of Chios (*c.* 460 B.C.) proved that circles are to one another as the squares on their diameters, and it seems probable that he also used the method of exhaustion,—a subject which was evidently much discussed about that time. Archimedes tells us that the "earlier geometers" had proved that spheres have to one another the triplicate ratio of their diameters, so that the method was probably used by others as well.

Archimedes and Integration. It is to Archimedes himself (*c.* 225 B.C.) that we owe the nearest approach to actual integration to be found among the Greeks.[2] His first noteworthy advance in this direction was concerned with his proof that the area of a parabolic segment is four thirds of the triangle with the same base and vertex, or two thirds of the circumscribed parallelogram. This was shown by continually inscribing in each segment between the parabola and the inscribed figure a triangle with the same base and the same height as the segment. If A is the area of the original inscribed triangle, the process adopted by him leads to the summation of the series

$$A + \tfrac{1}{4}A + (\tfrac{1}{4})^2 A + \cdots,$$

or to finding the value of

$$A[1 + \tfrac{1}{4} + (\tfrac{1}{4})^2 + (\tfrac{1}{4})^3 + \cdots],$$

so that he really finds the area by integration and recognizes, but does not assert, that

$$(\tfrac{1}{4})^n \to 0 \text{ as } n \to \infty,$$

this being the earliest example that has come down to us of the summation of an infinite series.

[1] Heath, *Euclid*, Vol. II, p. 120; see also his *Archimedes*, p. xlvii.
[2] Heath, *Archimedes*, p. cxlii.


Area of the Parabola. In his proof relating to the quadrature of the parabola Archimedes first proves two propositions numbered 14 and 15 in his treatise on this curve. These assert that, with respect to the figure here shown,[1]

$$\triangle EqQ < 3(FO_1 + F_1O_2 + \cdots + F_{n-1}O_n + \triangle E_nO_nQ,$$

and $$\triangle EqQ > 3(R_1O_2 + R_2O_3 + \cdots + R_{n-1}O_n + \triangle R_nO_nQ.$$

He then states (Prop. 16) that the area of the segment of the parabola is equal to $\frac{1}{3}\triangle EqQ$. The proof is by a *reductio ad absurdum* and is given by Heath substantially as follows:

I. Suppose that the area of the segment is greater than $\frac{1}{3}\triangle EqQ$. Then the excess can, if continually added to itself, be made to exceed $\triangle EqQ$. And it is possible to find a submultiple of the triangle EqQ less than the said excess of the segment over $\frac{1}{3}\triangle EqQ$.



Let the triangle FqQ be such a multiple of the triangle EqQ. Divide Eq into equal parts each equal to qF, and let all the points of division including F be joined to Q meeting the parabola in $R_1, R_2, \cdots, R_n$ respectively. Through $R_1, R_2, \cdots, R_n$ draw diameters of the parabola meeting qQ in $O_1, O_2, \cdots, O_n$ respectively.

Let O_1R_1 meet QR_2 in F_1.

Let O_2R_2 meet QR_1 in D_1 and QR_3 in F_2.

Let O_3R_3 meet QR_2 in D_2 and QR_4 in F_3, and so on.

We have, by hypothesis,

$$\triangle FqQ < \text{area of segment} - \tfrac{1}{3}\triangle EqQ,$$

or $$\text{area of segment} - \triangle FqQ > \tfrac{1}{3}\triangle EqQ. \qquad (1)$$

Now, since all the parts of qE, such as qF and the rest, are equal, we have $O_1R_1 = R_1F_1$, $O_2D_1 = D_1R_2 = R_2F_2$, and so on; therefore

$$\begin{aligned}\triangle FqQ &= FO_1 + R_1O_2 + D_1O_3 + \cdots \\ &= FO_1 + F_1D_1 + F_2D_2 + \cdots + F_{n-1}D_{n-1} + \triangle E_nR_nQ. \qquad (2)\end{aligned}$$

But area of segment $< FO_1 + F_1O_2 + \cdots + F_{n-1}O_n + \triangle E_nO_nQ.$ (3)

[1] For the proof, see Heath, *Archimedes*, p. 241, preferably F. Kliem's German translation, pp. 361–365 (Berlin, 1914). The proof of the next proposition is taken from the same work, p. 244.

Subtracting the equation (2) from the inequality (3), we have

area of segment $-\triangle FqQ < R_1O_2 + R_2O_3 + \cdots + R_{n-1}O_n + \triangle R_nO_nQ$;

whence, *a fortiori*, by (1),

$$\tfrac{1}{3}\triangle EqQ < R_1O_2 + R_2O_3 + \cdots + R_{n-1}O_n + \triangle R_nO_nQ.$$

But this is impossible, since [Props. 14, 15]

$$\tfrac{1}{3}\triangle EqQ > R_1O_2 + R_2O_3 + \cdots + R_{n-1}O_n + \triangle R_nO_nQ.$$

Therefore area of segment $\not> \tfrac{1}{3}\triangle EqQ$.

II. If possible, suppose the area of the segment less than $\tfrac{1}{3}\triangle EqQ$.

Take a submultiple of the triangle EqQ (as the triangle FqQ), less than the excess of $\tfrac{1}{3}\triangle EqQ$ over the area of the segment, and make the same construction as before.

Since $\triangle FqQ < \tfrac{1}{3}\triangle EqQ$ — area of segment, it follows that

$$\triangle FqQ + \text{area of segment} < \tfrac{1}{3}\triangle EqQ$$
$$< FO_1 + F_1O_2 + \cdots + F_{n-1}O_n + \triangle E_nO_nQ.$$

[Props. 14, 15]

Subtracting from each side the area of the segment, we have

$$\triangle FqQ < \text{sum of spaces } qFR_1,\ R_1F_1R_2,\ \cdots,\ E_nR_nQ,$$
$$< FO_1 + F_1D_1 + \cdots + F_{n-1}D_{n-1} + \triangle E_nR_nQ, \textit{ a fortiori};$$

which is impossible, because, by (2) above,

$$\triangle FqQ = FO_1 + F_1D_1 + \cdots + F_{n-1}D_{n-1} + \triangle E_nR_nQ.$$

Hence area of segment $\not< \tfrac{1}{3}\triangle EqQ$.

Since, then, the area of the segment is neither greater than nor less than $\tfrac{1}{3}\triangle EqQ$, it is equal to it.

The Method of Archimedes. As to the working of the mind of Archimedes in arriving at the conclusion in regard to the area of the parabola (a conclusion which led to the above proof) we have some interesting evidence. In a manuscript discovered in Constantinople in 1906 by Professor Heiberg, the editor of the works of Archimedes, the latter's method of approach to certain propositions is set forth. In particular the first proposition relates to the steps taken in arriving at the conclusion with respect to the quadrature of the parabola.[1]

[1]The Heiberg edition was translated by Lydia G. Robinson, Chicago, 1909, and by Sir Thomas L. Heath, Cambridge, 1912.


The following is the translation as given in Heath's edition:

Let ABC be a segment of a parabola bounded by the straight line AC and the parabola ABC, and let D be the middle point of AC. Draw the straight line DBE parallel to the axis of the parabola and join AB, BC.

Then shall the segment ABC be $\frac{4}{3}$ of the triangle ABC.

From A draw AKF parallel to DE, and let the tangent to the parabola at C meet DBE in E and AKF in F. Produce CB to meet AF in K, and again produce CK to H, making KH equal to CK.

Consider CH as the bar of a balance, K being its middle point.



Let MO be any straight line parallel to ED, and let it meet CF, CK, AC in M, N, O, and the curve in P.

Now, since CE is a tangent to the parabola and CD the semiordinate,

$$EB = BD;$$

for this is proved in the *Elements* [*of Conics*].[1]

Since FA, MO are parallel to ED, it follows that

$$FK = KA,$$

and

$$MN = NO.$$

[1] *I.e.*, the works on conics by Aristæus and Euclid. See the similar expression in *On Conoids and Spheroids*, Prop. 3, and *Quadrature of Parabola*, Prop. 3.

Now, by the property of the parabola, which is proved in a lemma,

$$MO : OP = CA : AO \qquad [\textit{Quadrature of Parabola}, \text{Prop. } 5]$$
$$= CK : KN = HK : KN. \qquad [\text{Eucl. VI. } 2]$$

Take a straight line TG equal to OP, and place it with its center of gravity at H, so that $TH = HG$; then, since N is the center of gravity of the straight line MO, and $MO : TG = HK : KN$, it follows that TG at H and MO at N will be in equilibrium about K.

[*On the Equilibrium of Planes*, I, 6, 7]

Similarly, for all other straight lines parallel to DE and meeting the arc of the parabola, (1) the portion intercepted between FC, AC with its middle point on KC and (2) a length equal to the intercept between the curve and AC placed with the center of gravity at H will be in equilibrium about K.

Therefore K is the center of gravity of the whole system consisting (1) of all the straight lines as MO intercepted between FC, AC and placed as they actually are in the figure and (2) of all the straight lines placed at H equal to the straight lines as PO intercepted between the curve and AC.

And, since the triangle CFA is made up of all the parallel lines like MO, and the segment CBA is made up of all the straight lines like PO within the curve, it follows that the triangle, placed where it is in the figure, is in equilibrium about K with the segment CBA placed with its center of gravity at H.

Divide KC at W so that $CK = 3\,KW$; then W is the center of gravity of the triangle ACF: for this is proved in the books on equilibrium.

[*On the Equilibrium of Planes*, I, 15]

Therefore $\triangle ACF$: segment $ABC = HK : KW$

$$= 3 : 1.$$

Therefore segment $ABC = \frac{1}{3} \triangle ACF$.

But $\triangle ACF = 4 \triangle ABC$.

Therefore segment $ABC = \frac{4}{3} \triangle ABC$.

Now the fact here stated is not actually demonstrated by the argument used, but that argument has given a sort of indication that the conclusion is true. Seeing, then, that the theorem is not demonstrated, but at the same time suspecting that the conclusion is true, we shall have recourse to the geometrical demonstration which I myself discovered and have already published.

Archimedes anticipates Modern Formulas. In his treatment of solids bounded by curved surfaces he arrives at conclusions which we should now describe by the following formulas:[1]

Surface of a sphere, $4\pi a^2 \cdot \frac{1}{2}\int_0^{\pi} \sin\theta\, d\theta = 4\pi a^2$.

Surface of a spherical segment,

$$\pi a^2 \int_0^{\alpha} 2\sin\theta\, d\theta = 2\pi a^2(1-\cos\alpha).$$

Volume of a segment of a hyperboloid of revolution,

$$\int_0^{b} (ax + x^2)\, dx = b^2(\tfrac{1}{2}a + \tfrac{1}{3}b).$$

Volume of a segment of a spheroid,

$$\int_0^{b} x^2 dx = \tfrac{1}{3}b^3.$$

Area of a spiral, $\frac{\pi}{a}\int_0^{a} x^2 dx = \frac{1}{3}\pi a^2$.

Area of a parabolic segment,

$$\frac{1}{A^2}\int_0^{A} \Delta^2 d\Delta = \tfrac{1}{3}A.$$

2. Medieval Ideas of the Calculus

Relation to Mensuration. The only traces that we have of an approach to the calculus in the Middle Ages are those relating to mensuration and to graphs. The idea of breaking up a plane surface into infinitesimal rectangles was probably present in the minds of many mathematicians at that time in the West as well as in the East, but it was never elaborated into a theory that seemed worth considering. For example, a Jewish writer, Jehudah Barzilai, living in Barcelona in the 13th century,[2] as-

[1] Heath, *Archimedes*, p. cxlvi seq.; G. Loria, *Le scienze esatte nell' antica Grecia*, 2d ed., p. 108 (Milan, 1914); Heath, *Method of Archimedes*, p. 8 (1912).

[2] *Sefer Jezira*, commentary by Judah ben Barzilai, p. 255 (Berlin, 1884).

serts that "it has been said that there is no form in the world except the rectangle, for every triangle or rectangle is composed of rectangles too small to be perceived by the senses."

The next important step in the preparation for the calculus taken in the Middle Ages is the one already described in connection with the geometric work of Oresme (*c.* 1360). His method of latitudes and longitudes gave rise to what we should now call a distribution curve or graph,—a step that is fundamental to the modern method of finding the area included between a curve and certain straight lines.

3. Modern Forerunners of the Calculus

Early Writers. As is usual in such cases, it is impossible to determine with certainty to whom credit belongs, in modern times, for first making any noteworthy move in the calculus, but it is safe to say that Stevin is entitled to serious consideration. His contribution is seen particularly in his treatment of the subject of the center of gravity of various geometric figures, anticipating as it did the work of several later writers.[1] Other writers, even in medieval times, had solved various problems in mensuration by methods which showed the influence of the Greek theory of exhaustion and which anticipated in some slight degree the process of integration. Among them may be mentioned the name of Ṯâbit ibn Qorra (*c.* 870), who found the volume of a paraboloid. Soon after Stevin wrote, Luca Valerio[2] published his *De quadratura parabolae* (Rome, 1606), using a method of attack that was essentially Greek in its spirit.

Kepler. Among the more noteworthy attempts at integration in modern times were those of Kepler (1609). In his notable work on planetary motion he asserted that a planet describes equal focal sectors of ellipses in equal times. This naturally demands some method for finding the areas of such sectors,

[1] *De Beghinselen der Weeghconst*, Leyden, 1686. For a summary see H. Bosmans, "Le calcul infinitésimal chez Simon Stevin," *Mathesis*, XXXVII (1923).

[2] Born *c.* 1552; died in 1618. He was professor of mathematics and physics at Rome.

and the one invented by Kepler was called by him the method of the "sum of the radii," a rude kind of integration. He also became interested in the problem of gaging, and published a work on this subject and on general mensuration as set forth by Archimedes.[1] Far from being an elementary treatment of gaging, this was a scientific study of the measurement of solids in general. Kepler considers solids as composed "as it were" (*veluti*) of infinitely many infinitely small cones or infinitely thin disks, the summation of which becomes the problem of the later integration.

Cavalieri. It was Kepler's attempts at integration that led Cavalieri to develop his method of indivisibles,[2] a subject which may also have been suggested to him by Aristotle's tract *De lineis insecabilibus*, to take the common Latin title.[3] It may also have been suggested by one of the fragments of Xenocrates (*c.* 350 B.C.), an Athenian, who wrote upon indivisible lines.[4]

Cavalieri's Lack of Clearness. Cavalieri was not always clear in his statements respecting the nature of an indivisible magnitude. In general, however, he seems to have looked upon a solid as made up practically of superposed surfaces, a surface as made up of lines, and a line as made up of points, these component parts being the ultimate possible elements in the decom-

[1] *Ausszug auss der vralten Messe-Kunst Archimedes, . . . Erklärung vnd Bestättigung der Oesterreichischen Weinvisier-Ruthen*, Linz, 1616; ed. Frisch, V, 497, 614 (Frankfort a. M., 1864). Kepler's letters (*ibid.*, p. 626) show that he was working on the subject as early as 1605. On this entire period see C. R. Wallner, *Bibl. Math.*, V (3), 113.

[2] *Geometria indivisibilibus continuorum nova quadam ratione promota*, Bologna, 1635; 2d ed., *ibid.*, 1653; *Exercitationes geometricae sex*, Bologna, 1647.

[3] H. Vogt referred to this tract in the *Bibl. Math.*, X (3), 146, and F. Cajori called attention to it more prominently in *Science* (U.S.), XLVIII (N.S.), 577. See also Heath, *History*, I, 346.

[4] " . . . in infinitum vero dividi non posse, sed in atomos quasdam desinere: has porro atomos non esse partium expertes et minimas, sed pro quantitate et materia dividi posse et partes habere: caeteroqui specie atomos et prima naturae statuens esse primas quasdam lineas insecabiles et ex his facta plana et solida prima."

This Latin translation by Simplicius is given, with the original Greek, in the "Xenocratis Fragmenta," F. W. A. Mullach, *Fragmenta Philosophorum Graecorum*, III, 118, § 21 (Paris, 1881).

position of the magnitude. He then proceeded to find lengths, areas, and volumes by the summation of these "indivisibles," that is, by the summation of an infinite number of infinitesimals.

Such a conception of magnitude cannot be satisfactory to any scientific mind, but it formed a kind of intuitive step in the development of the method of integration and undoubtedly stimulated men like Leibniz to exert their powers to place the theory upon a scientific foundation.[1]

Illustration of Cavalieri's Method. Some idea of Cavalieri's method may be obtained by considering his comparison of a triangle with a parallelogram having the same base and the same altitude. Calling the smallest element of the triangle 1, the next will be 2, the next 3, and so on to n, the base. The area is therefore $1 + 2 + 3 + \cdots + n$, or $\frac{1}{2}n(n+1)$. But each element of the parallelogram is n, and there are n of them, as in the triangle, and so the area is n^2. Then the ratio of the area of the triangle to the area of the parallelogram is

$$\frac{1}{2}n(n+1) : n^2 \quad \text{or} \quad \frac{1}{2}\left(1 + \frac{1}{n}\right).$$

But

$$\frac{1}{2}\left(1 + \frac{1}{n}\right) \to \frac{1}{2} \quad \text{as} \quad n \to \infty,$$

and so the triangle is half the parallelogram.

By means of his method Cavalieri was able to solve various elementary problems in the mensuration of lengths, areas, and volumes, and also to give a fairly satisfactory proof of the theorem of Pappus with respect to the volume generated by the revolution of a plane figure about an axis.[2]

[1] For a discussion of Cavalieri's work and its relation to the calculus, see H. Bosmans, "Sur une contradiction reprochée à la théorie des 'indivisibles' chez Cavalieri," *Annales de la Société scientifique de Bruxelles*, XLII (1922), 82. For his work on the center of gravity see E. Bortolotti, "Le prime applicazioni del calcolo integrale alla determinazione del centro di gravità di figure geometriche," *Rendiconto . . . della R. Accad. delle Scienze*, Bologna, 1922, reprint.

[2] For a translation of Cavalieri's Theorem relating to the volumes of solids, see G. W. Evans, in *Amer. Math. Month.*, XXIV, 447. On the method in general see also H. Bosmans, "Un chapitre de l'œuvre de Cavalieri," *Mathesis*, XXXVI, 365.

Roberval. At the same time that Cavalieri was working on the problem of indivisibles Roberval[1] was proceeding upon a similar hypothesis. He considered the area between a curve and a straight line as made up of an infinite number of infinitely narrow rectangular strips, the sum of which gave him the required area. In the same way he attacked the problems of rectification and of cubature. He also found the approximate value of $\int_0^1 x^m dx$, m being a positive integer, by finding the value of

$$\frac{0^m + 1^m + 2^m + \cdots + (n-1)^m}{n^{m+1}},$$

asserting that this approaches $1/(m+1)$ as $n \to \infty$.

Fermat. Fermat (1636) reached the same conclusion, basing his treatment upon a method set forth by Archimedes, and also extended the proof to include substantially the cases in which m is fractional (1644) or negative (1659), although not using either the fractional or the negative exponent in his work. He also attacked (1636) the problem of maxima and minima, that is, of finding the points on a curve at which the tangent is parallel to the x-axis. It was probably because of this step that Lagrange expressed himself as follows:[2]

> One may regard Fermat as the first inventor of the new calculus. In his method *De maximis et minimis* he equates the quantity of which one seeks the maximum or the minimum to the expression of the same quantity in which the unknown is increased by the indeterminate quantity. In this equation . . . he divides . . . by the indeterminate quantity which occurs in them as a factor; then he takes this quantity as zero and he has an equation which serves to determine the unknown sought. . . . His method of tangents depends upon the same principle.[3]

[1] *Traité des indivisibles*, mémoire, Paris, 1634. See A. E. H. Love, "Infinitesimal Calculus," *Encyc. Brit.*, 11th ed.

[2] *Œuvres de Lagrange*, ed. Serret, X, 294. See Cajori in *Amer. Math. Month.*, XXVI, 16.

[3] For confirmation of this opinion by Laplace and Tannery, see Cajori, *loc. cit.*, p. 17. On the work *De maximis et minimis* consult the *Supplément* to Volumes I–IV of the *Œuvres de Fermat*, edited by C. de Waard, Paris, 1922, and the review by H. Bosmans, *Revue des Questions Scientifiques*, Brussels, April, 1923.

With his name should be joined that of a later writer, Antonio di Monforte[1] (1644–1717), a Neapolitan mathematician who worked along similar lines.

Problem of Tangents. The problem of tangents, the basic principle of the theory of maxima and minima, may be said to go back to Pappus (*c.* 300).[2] It appears indirectly in the Middle Ages, for Oresme (*c.* 1360) knew that the point of maximum or minimum ordinate of a curve is the point at which the ordinate is changing most slowly. It was Fermat, however, who first stated substantially the law as we recognize it today, communicating (1638)[3] to Descartes a method which is essentially the same as the one used at present, that of equating $f'(y)$ to zero. Similar methods were suggested by René de Sluze[4] (1652) for tangents, and by Hudde[5] (1658) for maxima and minima.

Other Writers. From then until Newton finally brought the work to a climax various efforts were made in the same direction by such writers as Huygens, Torricelli, Pascal,[6] and Mersenne. The fact that the area of the hyperbola $xy = 1$, found by Grégoire de Saint-Vincent[7] (1647), is related to logarithms was recognized by Fermat, and Nicolaus Mercator[8] made use of the principle in his calculation of these functions.

Wallis. The first British publication of great significance bearing upon the calculus is that of John Wallis, issued in 1655.

[1] F. Amodeo, "La Regola di Fermat-Monforte per la ricerca dei massimi e minimi," *Periodico di Matematica*, XXIV, fasc. VI.

[2] There is a good summary of the history of tangents as related to the calculus in a work by Anibal Scipião Gomes de Carvalho, *A Teoria das Tangentes antes da Invenção do Cálculo Diferencial*, Coimbra, 1919.

[3] *Opera varia*, Toulouse, 1679.

[4] See also "A short and easy method of drawing tangents to all geometrical curves," *Phil. Trans.*, 1672.

[5] *De reductione aequationum et de maximis et minimis*, in a letter published in 1713.


[6] H. Bosmans, *Archivio di Storia della Scienza*, IV, 369.

[7] *Opus geometricum quadraturae circuli et sectionum coni*, 2 vols., Antwerp, 1647.

[8] *Logarithmotechnia*, London, 1668 and 1674.

It is entitled *Arithmetica Infinitorum, sive Nova Methodus Inquirendi in Curvilineorum Quadraturam, aliaque difficiliora Matheseos Problemata*, and is dedicated to Oughtred. By a method similar to that of Cavalieri the author effects the quadrature of certain surfaces, the cubature of certain solids, and the rectification of certain curves. He speaks of a triangle, for example, "as if" (*quasi*) made up of an infinite number of parallel lines in arithmetic proportion, of a paraboloid "as if" made up of an infinite number of parallel planes, and of a spiral as an aggregate of an infinite number of arcs of similar sectors, applying to each the theory of the summation of an infinite series. In all this he expresses his indebtedness to such writers as Torricelli and Cavalieri. He speaks of the work of such British contemporaries as Seth Ward and Christopher Wren, who were interested in this relatively new method, and, indeed, his dedication to Oughtred is the best contemporary specimen that we have of the history of the movement just before Newton's period of activity.[1] All this, however, was still in the field of integration, the first steps dating, as we have seen, from the time of the Greeks.

Barrow. What is considered by us as the process of differentiating was known to quite an extent to Barrow (1663). In his *Lectiones opticae et geometricae*[2] he gave a method of tangents in which, in the annexed figure, Q approaches P as in our present theory, the result being an indefinitely small (*indefinite parvum*) arc. The triangle PRQ was long known as "Barrow's differential triangle,"[3] a name which, however, was not due to him. It is evident that this method, and the figure as well, must have had a notable influence upon the mathematics of his time.



[1] The work also appears in his *Opera Mathematica*, I, 255–470 (Oxford, 1695).

[2] London, 1669.. The work seems to have been written in 1663 and 1664. Love, *loc. cit.*; J. M. Child, *Geometrical Lectures of Isaac Barrow*, Chicago, 1916.

[3] On this close approach to the later calculus see Whewell's edition of Barrow's *Mathematical Works*, p. xii (Cambridge, 1860), and Child, *loc. cit.*

It is quite probable that Barrow had advised Newton of his work on this figure as early as 1664.[1] Pascal had already published a figure of somewhat the same shape,[2] so that the study of triangles of the general nature illustrated above was being undertaken and discussed at this time in both England and France. The triangles given by both Barrow and Pascal were apparently known to Leibniz, and they assisted him in developing his own theory.[3]

Barrow also recognized the fact that integration is the inverse of differentiation, but he did not use this relation to aid him in solving the quadrature problem.

Period of the Invention of the Newtonian Method. We now approach the period which is popularly thought to be the one in which the calculus was invented. It is evident, however, that a crude integral calculus was already in use and that some approach had been made to the process of differentiation. It is also evident that the lines of approach to the calculus in general have been two in number, one representing the static phase as seen in the measurement of fixed lengths, areas, or volumes, and in the making use of such ideas as those of infinitesimals and indivisibles; the other representing the dynamic phase as seen in the motion of a point. To the former belong such names as Kepler and Cavalieri and, in general, Archimedes; to the latter belong the great leaders in the mathematics of the time of Newton and Leibniz.[4]

[1] J. M. Child, *The Early Mathematical Manuscripts of Leibniz*, p. 11 (Chicago, 1920), a work which students of the history of the calculus should consult, not merely for its translations but for its notes. On the figures used by Barrow, Pascal, and Leibniz, see *ibid.*, p. 15. This work is hereafter referred to as Child, *Leibniz Manuscripts*.

[2] In his *Lettres de A. Dettonville* (Paris, 1659), the part relating to the triangle having been written in 1658.

[3] Child, *Leibniz Manuscripts*, p. 16. See Leibniz's admission as to Barrow in his letter to Jacques Bernoulli (1703), *ibid.*, p. 20.

[4] On the general history of the development of the calculus in the 17th century the following works may be consulted with profit:

W. W. R. Ball, *History of Mathematics*, 10th ed., London, 1922, the treatment of the calculus being particularly complete; A. De Morgan, *On a point connected with the dispute between Keill and Leibniz about the Invention of Fluxions*, London, 1846; also the *Companion to the British Almanack*, 1852, and *Philo-*

4. Newton and Leibniz

Newton. Newton's great contribution to the theory consists in part in his extension of the method to include the other functions then in common use, in his recognition of the fact that the inverse problem of differentiation could be used in solving the problem of quadrature, in his introduction of a suitable notation, and in his wide range of applications of the subject. Starting with the knowledge already acquired by Barrow, he developed, beginning in 1665, his method of "fluxions." This he afterward set forth in three tracts,[1] which, in accordance with his unfortunate plan of avoiding publicity in his discoveries, were not printed until many years later.

Newton's Three Types. Newton recognized three types of the calculus. In his *Principia* (1687) he made some use of

sophical Magazine, June and November, 1852; G. J. Gerhardt, *Die Entdeckung der Differenzialrechnung durch Leibniz*, Halle, 1848; J. Raphson, *The History of Fluxions*, London, 1715; Latin edition the same year; H. Sloman, *Leibnizens Ansprüche auf die Erfindung der Differenzialrechnung*, Leipzig, 1857; English translation with additions, Cambridge, 1860; M. Cantor, *Geschichte der Mathematik*, III, chap. 97; W. T. Sedgwick and H. W. Tyler, *A Short History of Science*, chap. xiii (New York, 1917); H. Weissenborn, *Die Principien der höheren Analysis, als historisch-kritischer Beitrag zur Geschichte der Mathematik*, Halle, 1856; D. Brewster, *Memoirs of the Life, Writings, and Discoveries of Sir Isaac Newton*, 2 vols., Edinburgh, 1885; 2d ed., 1860; J. Collins, *Commercium Epistolicum de Varia Re Mathematica*, London, 1712; 2d ed., London, 1722 (two editions); 3d ed., London, 1725; French ed., Paris, 1856; G. Vivanti, *Il concetto d' infinitesimo e la sua applicazione alla matematica*, Mantua, 1894; 2d ed., Naples, 1901; J. M. Child, *The Early Mathematical Manuscripts of Leibniz*, Chicago, 1920.

On the history of fluxions in Great Britain, beginning with Newton, the best work is F. Cajori, *A History of the Conceptions of Limits and Fluxions in Great Britain from Newton to Woodhouse*, Chicago, 1919. On the general history of the later development of the calculus, see J. A. Serret and G. Scheffers, *Lehrbuch der Differential- und Integralrechnung*, II, 581–626 (5th ed., Leipzig, 1911), and III, 694–720 (Leipzig, 1914); Sedgwick and Tyler, *loc. cit.*, chap. xv; E. W. Brown, "Mathematics," in *The Development of the Sciences*, New Haven, 1923.

[1] (1) *De Analysi per Equationes numero terminorum infinitas*, written in 1666 and sent to Barrow, who made it known to John Collins, who allowed Lord Brouncker to copy it; it was not published, however, until (London) 1711. (2) *Method of Fluxions and Infinite Series*, written in 1671 but not printed until (London) 1736, and then in John Colson's translation with this title. (3) *Tractatus de Quadratura Curvarum*, apparently written in 1676, but not published until (London) 1704 (appendix to his *Opticks*).

infinitely small quantities,[1] but he apparently recognized that this was not scientific, for it is not the basis of his work in this field.

Method of Fluxions. His second method was that of fluxions. For example, he considered a curve as described by a flowing point, calling the infinitely short path traced in an infinitely short time the moment of the flowing quantity, and designated the ratio of the moment to the corresponding time as the "fluxion" of the variable, that is, as the velocity. This fluxion of x he denoted by the symbol $\dot{x}$. In his *Method of Fluxions*[2] he states that "the moments of flowing quantities are as the velocities of their flowing or increasing,"—a statement which may be expressed in the Leibnizian symbolism as

$$\frac{dy}{dx} = \frac{dy}{dt} : \frac{dx}{dt}.$$

His treatment of fluxions may be illustrated by the following extract from his work:[3]

If the moment of x be represented by the product of its celerity $\dot{x}$ into an indefinitely small quantity o (that is $\dot{x}o$), the moment of y will be $\dot{y}o$, since $\dot{x}o$ and $\dot{y}o$ are to each other as $\dot{x}$ and $\dot{y}$. Now since the moments as $\dot{x}o$ and $\dot{y}o$ are the indefinitely little accessions of the flowing quantities, x and y, by which these quantities are increased through the several indefinitely little intervals of time, it follows that these quantities, x and y, after any indefinitely small interval of time, become $x + \dot{x}o$ and $y + \dot{y}o$. And therefore the equation which at all times indifferently expresses the relation of the flowing quantities will as well express the relation between $x + \dot{x}o$ and $y + \dot{y}o$ as between x and y; so that $x + \dot{x}o$ and $y + \dot{y}o$ may be substituted in the same equation for those quantities instead of x and y.

Therefore let any equation

$$x^3 - ax^2 + axy - y^3 = 0$$

[1] F. Cajori, in *Amer. Math. Month.*, XXIV, 145; *ibid.*, XXVI, 15.
[2] Colson translation, p. 24 (London, 1736).
[3] Pp. 24, 25. See also G. H. Graves, *loc. cit.*, III, 82.

be given, and substitute $x + \dot{x}o$ for x and $y + \dot{y}o$ for y, and there will arise

$$\begin{aligned} & x^3 + 3x^2\dot{x}o + 3x\dot{x}\dot{x}oo + \dot{x}^3o^3 \\ & - ax^2 - 2ax\dot{x}o - a\dot{x}^2oo \\ & + axy + ax\dot{y}o + a\dot{x}oy + a\dot{x}o\dot{y}o \\ & - y^3 - 3\dot{y}oy^2 - 3\dot{y}^2ooy - \dot{y}^3o^3 = 0. \end{aligned}$$

Now, by supposition,

$$x^3 - ax^2 + axy - y^3 = 0,$$

which therefore being expunged and the remaining terms being divided by o, there will remain

$$3\dot{x}x^2 + 3\dot{x}^2ox + \dot{x}^3oo - 2a\dot{x}x - a\dot{x}^2o + a\dot{y}x + a\dot{x}y + a\dot{x}\dot{y}o - 3\dot{y}y^2 - 3\dot{y}^2oy - \dot{y}^3oo = 0.$$

But whereas o is supposed to be infinitely little that it may represent the moments of quantities, the terms which are multiplied by it will be nothing in respect to the rest. Therefore I reject them and there remains:

$$3\dot{x}x^2 - 2a\dot{x}x + a\dot{y}x + a\dot{x}y - 3\dot{y}y^2 = 0.$$

Method of Limits. Newton's third method, that of limits, appears in his *Tractatus de Quadratura Curvarum* (1704). In the introduction he says:

Let a quantity x flow uniformly and let it be required to find the fluxion of x^n. In the time in which x by flowing becomes $x + o$, the quantity x^n becomes $\overline{x + o}|^n$; *i.e.*, by the method of infinite series,

$$x^n + nox^{n-1} + \frac{nn - n}{2} oox^{n-2} +, \text{ etc.},$$

and the increment o and

$$nox^{n-1} + \frac{nn - n}{2} oox^{n-2} +, \text{ etc.},$$

are to each other as 1 and

$$nx^{n-1} + \frac{nn - n}{2} ox^{n-2} +, \text{ etc.}$$

Now let the increment vanish and their last ratio will be 1 to nx^{n-1}.

He also gives the interpretation of these ratios as the slopes of a secant through two points on a curve and of the tangent which is the limiting position of this secant.[1] He adds:

If the points are distant from each other by an interval however small, the secant will be distant from the tangent by a small interval. That it may coincide with the tangent and the last ratio be found, the two points must unite and coincide altogether. In mathematics, errors, however small, must not be neglected.

In the *Principia* (Section I) Newton set forth his idea of these ultimate ratios as follows:

Ultimate ratios in which quantities vanish, are not, strictly speaking, ratios of ultimate quantities, but limits to which the ratio of these quantities, decreasing without limit, approach, and which, though they can come nearer than any given difference whatever, they can neither pass over nor attain before the quantities have diminished indefinitely.

In the fluxional notation Newton represented the fluent of x by $\dot{x}o$, or simply by $\dot{x}$. The fluent of $\dot{x}$ he represented by $\ddot{x}$, and so on,[2] a notation first published in the *Algebra* of John Wallis (1693).

Summary of Newton's Method. Ball has clearly summarized Newton's general method of treatment as follows:

There are two kinds of problems. The object of the first is to find the fluxion of a given quantity, or, more generally, "the relation of the fluents being given, to find the relations of their fluxions." This is equivalent to differentiation. The object of the second, or inverse, method of fluxions is, from the fluxion or some relations involving it, to determine the fluent; or, more generally, "an equation being proposed exhibiting the relation of the fluxions of quantities, to find the relations of those quantities, or fluents, to one another." This is

[1] Graves, *loc. cit.*

[2] "Sint v, x, y, z fluentes quantitates, & earum fluxiones his notis $\dot{v}$, $\dot{x}$, $\dot{y}$, $\dot{z}$, designabuntur respective. . . . Qua ratione $\dot{v}$ est fluxio quantitatis v, & $\ddot{v}$ fluxio ipsius $\dot{v}$, & $\dddot{v}$ fluxio ipsius $\ddot{v}$" (*Opera*, II, 392).

equivalent either to integration, which Newton termed the method of quadrature, or to the solution of a differential equation which was called by Newton the inverse method of tangents.[1]

Leibniz. Leibniz (1684) was well aware of the work of men like Barrow, Huygens, Grégoire de Saint-Vincent, Pascal, and Cavalieri. He was in London in 1673, and there he probably met with scholars who were perfectly familiar with the discoveries of Barrow and Newton, and with Barrow himself he had extended correspondence. After leaving England he set to work upon the problems of tangents and quadratures and invented a notation which was original and at the same time was generally more usable than that of Newton,—the "differential notation." He proposed to represent the sum of Cavalieri's indivisibles by the symbol $\int$, the old form of *s*, the initial of *summa*, using this together with Cavalieri's *omn.* (for *omnia*), and to represent the inverse operation by *d*. By 1675 he had settled this notation,[2] writing $\int y dy = \frac{1}{2}y^2$ as it is written at present.

Leibniz published his method in 1684[3] and 1686,[4] speaking of the integral calculus as the *calculus summatorius*, a name connected with the *summa* ($\int$) sign. In 1696 he adopted the term *calculus integralis*, already suggested by Jacques Bernoulli in 1690.

His Conception of the Differential. Some idea of his conception of the differential may be obtained from a statement in a letter written by him to Wallis on March 30, 1699:

It is useful to consider quantities infinitely small such that when their ratio is sought, they may not be considered zero, but which are rejected as often as they occur with quantities incomparably greater.

[1] W. W. R. Ball, *Hist. of Math.*, 6th ed., p. 344 (London, 1915), to which the reader is referred for further details, Mr. Ball having given special attention to the work of Newton. See also A. von Braunmühl, *Bibl. Math.*, V (3), 355.

[2] But published in 1686. Love, *loc. cit.*

[3] "Nova methodus pro maximis et minimis, itemque tangentibus . . . ," in the *Acta Eruditorum.*

[4] "De geometria recondita et analysi indivisibilium atque infinitorum," also in the *Acta Eruditorum.* On an early case of integration (1599) before the symbolism appeared, see F. Cajori, in *Bibl. Math.*, XIV (3), 312.

Thus if we have $x + dx$, dx is rejected. But it is different if we seek the difference between $x + dx$ and x, for then the finite quantities disappear. Similarly we cannot have xdx and $dxdx$ standing together. Hence if we are to differentiate xy we write:

$$(x + dx)(y + dy) - xy = xdy + ydx + dxdy.$$

But here $dxdy$ is to be rejected as incomparably less than $xdy + ydx$. Thus in any particular case the error is less than any finite quantity.[1]

As to the approximate period at which he began to arrive at his laws for the differentiation of algebraic functions, we have a manuscript of his which was written in November, 1676, and in which he gives the following statements:[2]

$$\overline{dx} = 1, \quad \overline{dx^2} = 2x, \quad \overline{dx^3} = 3x^2, \text{ etc.};$$

$$\overline{d\frac{1}{x}} = -\frac{1}{x^2}, \quad \overline{d\frac{1}{x^2}} = -\frac{2}{x^3}, \quad \overline{d\frac{1}{x^3}} = \frac{3}{x^4}, \text{ etc.};$$

$$\overline{d\sqrt{x}} = \sqrt{\frac{1}{x}}, \text{ etc.};$$

$$dx^e = e\,x^{e-1}, \text{ and conversely } \int \overline{x^e} = \frac{x^{e+1}}{e+1}.$$

Hence $\overline{d\frac{1}{x^2}} = \overline{dx^{-2}}$ will be $-2x^{-3}$ or $-\frac{2}{x^3}$, and $\overline{d\sqrt{x}}$ or $dx^{\frac{1}{2}}$ will be $-\frac{1}{2}x^{-\frac{1}{2}}$ or $-\frac{1}{2}\sqrt{\frac{1}{x}}$.

Some of these results are incorrect, probably because of careless writing, and some appear in his earlier manuscripts, but they all serve to show how the mind of Leibniz was working in this period. By the end of the year 1676 he had developed the rule for differentiating a product, and by July, 1677, he had the differentiation of algebraic functions well in hand.[3]

[1] *Leibnitzens Mathematische Schriften*, Gerhardt ed., IV, 63 (Series III, in *Leibnitzens Gesammelte Werke*, Pertz ed., Halle, 1859) (this portion translated by Mr. Graves).

[2] J. M. Child, *The Early Mathematical Manuscripts of Leibniz*, p. 124 (Chicago, 1920), the results being as there stated, including errors.

[3] Child, *loc. cit.*, p. 116.

His notation for differentiation was used in England by John Craig as early as 1693,[1] and the same writer used his sign for integration ten years later.[2] Both symbols were somewhat familiar to English mathematicians throughout the 18th century, although it was not until the 19th century that their use in Great Britain became general.

Priority Dispute. The dispute between the friends of Newton and those of Leibniz as to the priority of discovery was bitter and rather profitless. It was the subject of many articles[3] and of a report by a special committee of the Royal Society.[4]

English readers of the 18th century were so filled with the arguments respecting the controversy as set forth in the *Commercium Epistolicum* (1712) and Raphson's *History of Fluxions* (1715), that they gave Leibniz little credit for his work. It was not until De Morgan (1846) reviewed the case that they began generally to recognize that they had not shown their usual spirit of fairness. On the other hand, Leibniz was so stung by the accusations of his English critics that he too showed a spirit that cannot always be commended.

Leibniz states his Case. It is interesting to read the words of Leibniz in his own defense, as presented in his *Historia et Origo Calculi Differentialis*:[5]

Since therefore his[6] opponents, neither from the *Commercium Epistolicum* that they have published, nor from any other source, brought forward the slightest bit of evidence whereby it might be established that his rival used the differential calculus before it was published by our friend;[7] therefore all the accusations that were brought against him by these persons may be treated with contempt

[1] *Methodus Figurarum* (London, 1693). [2] *Tractatus Mathematicus.*

[3] Beginning with a publication by a Swiss scholar, Nicolas Fatio de Duillier (1664–1753), whose *Lineae brevissimi descensus investigatio geometrica duplex* appeared in London in 1699. See Child, *Leibniz Manuscripts*, pp. 22, 23.

[4] The report appeared in 1712. See Collins, *Commercium Epistolicum.* It was also edited by Biot and Lefort and published at Paris in 1856.

[5] Found in MS. by Dr. C. I. Gerhardt in the Royal Library at Hannover and published in Latin in 1846; English translation by Child, *Leibniz Manuscripts*, pp. 22, 57.

[6] *I.e.*, Leibniz's, the work being written in the third person.

[7] *I.e.*, himself.

as beside the question. They have used the dodge of the pettifogging advocate to divert the attention of the judges from the matter on trial to other things, namely to infinite series. But even in these they could bring forward nothing that could impugn the honesty of our friend, for he plainly acknowledged the manner in which he made progress in them; and in truth in these also, he finally attained to something higher and more general.

Brief Summary of the Dispute. The facts are that Leibniz knew of Barrow's work on the "differential triangle" before he began his own investigations, or could have known of it, and that he was also in a position to know something of Newton's work. The evidence is also clear that Newton's discovery was made before Leibniz entered the field; that Leibniz saw some of Newton's papers on the subject as early as 1677; that he proceeded on different lines from Newton and invented an original symbolism; and that he published his results before Newton's appeared in print. With these facts before us, it should be possible to award to each his approximate share in the development of the theory.[1]

Successors of Newton and Leibniz. Most of the British writers of the period 1693–1734, failing to comprehend Newton's position, considered a fluxion as an infinitely small quantity.[2] The first noteworthy improvement in England is due to Bishop Berkeley, who, in his *Analyst* (1734), showed the fallacy of this method of approach and attempted to prove that even Newton was at fault in his logic. Berkeley provoked great discussion in England, and the result was salutary, not that it affected Newton's standing, but that it put an end to much of the lax reasoning of his followers.[3]

[1] On the general controversy see the summary given in Ball, *Hist. of Math.*, 6th ed., pp. 356–362; H. Sloman, *The Claim of Leibniz to the Invention of the Differential Calculus*, English translation, London, 1860.

[2] F. Cajori, *Amer. Math. Month.*, XXIV, 145; XXVI, 15; to these articles the reader is referred for valuable details relating to this period.

[3] On the gradual improvement of the Leibniz theory through the laying of a scientific foundation for the doctrine of limits, see F. Cajori, "Grafting of the theory of limits on the calculus of Leibniz," *Amer. Math. Month.*, XXX, 223, with excellent bibliography.

Cauchy's Contribution. Perhaps the one to whom the greatest credit is due for placing the fundamental principle of the calculus on a satisfactory foundation is Cauchy.[1] He makes the transition from

$$\frac{dy}{dx} = \frac{f(x+i)-f(x)}{i} = f'(x)$$

to

$$dy = f'(x)\,dx$$

as follows:

Let $y = f(x)$ be a function of the independent variable x; i, an infinitesimal, and h, a finite quantity. If we put $i = \alpha h$, α will be an infinitesimal and we shall have the identity

$$\frac{f(x+i)-f(x)}{i} = \frac{f(x+\alpha h)-f(x)}{\alpha h},$$

whence we derive

$$\frac{f(x+\alpha h)-f(x)}{\alpha} = \frac{f(x+i)-f(x)}{i}\,h. \tag{1}$$

The limit toward which the first member of this equation converges when the variable α approaches zero, h remaining constant, is what we call the "differential" of the function $y = f(x)$. We indicate this differential by the characteristic, d, as follows:

$$dy \text{ or } df(x).$$

It is easy to obtain its value when we know that of the derived function, y' or $f'(x)$. In fact, taking the limits of both members of equation (1), we have in general:

$$df(x) = hf'(x). \tag{2}$$

In the particular case where $f(x) = x$, equation (2) reduces to

$$dx = h.$$

Thus the differential of the independent variable, x, is simply the finite constant, h. Substituting, equation (2) will become

$$df(x) = f'(x)dx,$$


or, what amounts to the same thing,

$$dy = y'dx.$$

[1] *Résumé des Leçons sur le Calcul Infinitésimal, Quatrième Leçon*, Paris, 1823; *Œuvres Complètes, Sér. II, Tome IV*, Paris, 1899.

5. JAPAN

The Yenri. There developed in Japan in the 17th century a native calculus which may have been the invention of the great Seki Kōwa (1642–1708), as tradition asserts, although we have no positive knowledge that he ever wrote upon the subject. This form of the calculus is known as the *yenri*, a word meaning "circle principle" or "theory of the circle" and possibly



EARLY STEPS IN THE CALCULUS IN JAPAN


Crude integration, from Sawaguchi Kazuyuki's *Kokon Sampō-ki*, 1670. Sawaguchi was a pupil of Seki Kōwa, the Newton of Japan

suggested by an earlier Chinese title or by the fact that the method was primarily used in the measurement of this figure.

The mensuration of the circle by crude forms of integration is found in various works of the 18th century, such as the one illustrated above and the one shown on page 702, published by Mochinaga and Ōhashi in 1687. A similar use of the theory is found in connection with the mensuration of the sphere in

Isomura's work of 1684, and thereafter it appeared in numerous works in the closing years of the 17th century and the early part of the century following.

In a general way it may be said that the *yenri* was an application of series to the ancient method of exhaustion. For example, Takebe Kenkō (1722) found the approximate value of π by inscribing regular polygons up to 1024 sides, and probably more, giving the value to upwards of forty decimal places. In this work Takebe states that his method of approximation



EARLY STEPS IN THE CALCULUS IN JAPAN

From the *Kaisan-ki Kōmoku*, by Mochinaga and Ōhashi, representatives of the Seki School. The work was published in 1687. The method is essentially that of Sawaguchi

was not the one used by Seki Kōwa. In fact we know that the latter found an approximate value of π by computing successive perimeters, whereas Takebe based his work upon the squares of the perimeters, π^2 being taken as the square of the perimeter of a regular polygon of 512 sides. The value of π is expressed as a continued fraction, a plan which he states was due to his brother, Takebe Kemmei. Some of the formulas and series used by Takebe were very ingenious.[1]

[1] Smith-Mikami, p. 143.

TOPICS FOR DISCUSSION

1. General steps in the development of the calculus from the time of the Greeks to the present.

2. Zeno's paradoxes, their purpose, their fallacies, and their relation to the calculus.

3. The study of indivisible elements among the Greek philosophers, and its influence upon mathematics.

4. The atomistic philosophy of the Greeks, its founder and advocates; its bearing upon ancient mathematics and its relation to the modern calculus.

5. The relation of the method of exhaustion, especially as developed by Archimedes, to the integral calculus.

6. The various Greek writers on the method of exhaustion, together with a consideration of its results.

7. The contributions of Archimedes to the making of the calculus. His methods of proof.

8. The method employed by Archimedes in discovering his geometric propositions.

9. Formulas of the modern calculus anticipated by Archimedes and any other Greek writers.

10. The contributions of the Greeks to the subject of mechanics, and especially those of Aristotle and Archimedes.

11. Influence of Oresme with respect to the calculus.

12. Causes leading to Kepler's study of the problem of the calculus, together with a statement of the results of his work.

13. General nature of Cavalieri's contribution; the problems studied; the weakness of his method; the special results that he accomplished; and his influence upon Leibniz.

14. Fermat's contributions to the calculus compared with those of Cavalieri, Barrow, and Roberval.

15. The contributions of Roberval, Barrow, and other immediate predecessors of Newton.

16. Newton's discoveries in the calculus, with particular reference to the fundamental principles employed by him.

17. Leibniz's discoveries and the question of priority.

18. General nature of the developments in the calculus after Newton and Leibniz.

19. General nature of the early Japanese calculus.

INDEX

Since certain proper names are mentioned many times in this volume, only such page references have been given as are likely to be of considerable value to the reader, the first reference being to the biographical note in case one is given. In general, the biographies and bibliographies are to be found in Volume I. As a rule, the bibliographical references give only the page on which some important book or reference is first mentioned. Except for special reasons (such as a quotation, a discovery, or a contribution to which a reader may be likely to refer), no references are given to elementary textbooks or to the names of authors which are already given in Volume I and are mentioned only incidentally in Volume II. Obsolete terms are usually indexed only under modern forms. For further information consult the index to Volume I.

Aāhmesu. *See* Ahmes
Abacus, 7, 86, 156, 177; arc, 181; Armenian, 174; Babylonian, 160; Chinese, 168; dust, 157; Egyptian, 160; in France, 191; Gerbert's, 180; in Germany, 183, 190; Greek, 161; grooved, 166; Japanese, 170; Korean, 171, 174; line, 181, 186; Mohammedan, 174; Polish, 176; Pythagorean table, 177; Roman, 165; Russian, 175, 176; Turkish, 174; Western European, 177
Abbreviations of fractions, 221
Abel, N. H. (*c.* 1825), 469
Abhandlungen, 15
Abraham ben Ezra (*c.* 1140), 353, 437, 442, 543
Abscissa, 318, 324
Absolute number, 12
Absolute term, 394
Absolute value, 267
Abstract, 11
Abû Bekr Mohammed. *See* al-Karkhî
Abû Ja'far al-Khâzin (*c.* 960), 455
Abû'l-Faradsh (*c.* 987), 466
Abû'l-Ḥaṣan (*c.* 1260), 620
Abû'l-Weîâ (*c.* 980), 467, 609, 617, 622, 623
Abundant number, 20
Achilles problem, 546
Acre, 644
Acts (operations), 36
Adams, G. (*c.* 1748), 206
Addend, 88
Addition, 88, 184; of fractions, 223; symbols of, 395
Adelard of Bath (*c.* 1120), 12, 382
Adriaan. *See* Adriaen
Adriaen Anthoniszoon (*c.* 1600), 310
Adrianen Metius (*c.* 1600), 310
Adriaenszoon, J. M. (*c.* 1608), 373
Æbutius Faustus, L., 361
Æbutius Macedo, M., 357
Aethelhard. *See* Adelard
Affected quadratics, 450
Afghanistan, 72
Aggregation symbols, 416
Agnesi, M. G. (*c.* 1748), 331
Agricola, G., 637
Agrimensor, 361
Aguillon, F. (1613), 344
Aḥmed ibn 'Abdallâh al-Mervazî (*c.* 860), 620
Ahmes (*c.* 1650–1550 B.C.), 210, 386, 498, 500
Aḥmōse. *See* Ahmes
Ahrens, W., 536, 542
Akhmim, 212
Albategnius (*c.* 920), 608
al-Battânî. *See* Albategnius
Albêrûnî (*c.* 1000), 73, 308
Alchemy, 595
Alciatus, A. (1530), 637
Alcobatiensis, Codex, 198
Alcuin (*c.* 775), 535
Alessandro (*c.* 1714), 116
Alexandre de Villedieu (*c.* 1240), 14, 80
Alexandrian calendar, 658
al-Fazârî (*c.* 773), 72
Alfonsine Tables (*c.* 1250), 609
Alfonso X, el Sabio, 609

Algebra, 378; applications, 582; Arabic, 382; Chinese, 380; Egyptian, 379; Greek, 381; Hindu, 379; medieval, 382; name, 386; Persian, 382; powers, 393; related to geometry, 320; symbols, 382, 395; unknown quantity, 393
Algorism (algorithm), 9, 78, 88
Algus, 9, 78
al-Ḥaitam of Baṣra (*c.* 1000), 455
al-Ḥasan (Alhazen). *See* al-Ḥaitam
al-Ḥaṣṣâr (*c.* 12th century), 118
'Alî ibn Velî (*c.* 1590), 393
al-Karkhî (*c.* 1020), 388, 504
al-Kashî (*c.* 1430), 310, 505
al-Khayyâmî. *See* Omar Khayyam
al-Khowârizmî (*c.* 825), 9, 72, 382, 388, 446
Alliage, 588
Alligation, 587
Allman, G. J. (*c.* 1880), 677
Allotte de la Fuye, 38
Alloy, 588
Almagest. *See* Ptolemy
Almâhânî (al-Mâhânî) (*c.* 860), 382, 455
al-Mâmûn (*c.* 820), 372
Almanac, 665
al-Mervazî (Ḥabash al-Ḥâsib) (*c.* 860), 622
al-Rashid (*c.* 800), 672
al-Rûmî (*c.* 1520), 626
al-Zarqâla (Zarkâla) (*c.* 1050), 609, 616
Ambrose of Milan (*c.* 370), 542
Amicable (amiable) numbers, 23
Amodeo, F., 689
Analytic geometry, 316, 322, 324; solid, 325
Anatolius (*c.* 280), 5
Anaximander (*c.* 575 B.C.), 603
Andalò di Negro (*c.* 1300), 665
Anderson, G., 80
Andrés, M. J. (*c.* 1515), 200
Andrews, W. S., 594
Angle, 277; sum of angles in a triangle, 287; trisection, 297, 298
Anharmonic ratio, 333, 334
Anianus (1488), 665, 668
Annotio, Perito. *See* Cataldi
Antecedent, 483
Anthology, Greek, 532
Antilogarithms, 523
Antiphon (*c.* 430 B.C.), 677
Antonio de Dominis (1611), 343
Apianus, P. (*c.* 1527), 341, 441, 508, 509
Apices, 75
Apollonius (*c.* 225 B.C.), 318
Approximate roots, 253
Arabic numerals, 69, 70. *See* Hindu-Arabic numerals
Arabs, achievements, 272, 455, 467; algebra, 382; computation of tables, 626; in Europe, 609; geometry, 272; magic squares, 597; measure of the earth, 372; trigonometry, 608. *See* Arabic numerals, Hindu-Arabic numerals
Arbalète, 346
Arbuthnot, J., 637
Arcerianus, Codex, 504
Archibald, R. C., 21, 30, 287, 293, 302
Archimedes (*c.* 225 B.C.), 5, 454, 679, 681, 684; cattle problem, 453, 584; on the circle, 307; cubic of, 80
Arcus Pythagoreus, 177
Area, of a circle, 298, 302; of a polygon, 606; of a triangle, 631
Areas, 286, 644
Arenarius, 5
Argand, J. R. (*c.* 1810), 266
Argus, 9, 10
Aristarchus (*c.* 260 B.C.), 604
Aristotle (*c.* 340 B.C.), 2
Arithmetic, 7, 8. *See* Calculate, Calculating machines, Logistic, Numerals, Problems, Series, and the various operations and rules
Arithmetica, 1, 7
Arithmetics, American, 86
Armillary sphere, 350, 370
Arnaldo de Villa Nova (*c.* 1275), 669
Arnauld, A. (*c.* 1650), 28
Arnauld de Villeneuve (*c.* 1275), 669
Arnold, Sir E., 80
Ars Magna, 461–464
Ars supputandi, 11
Articles, 12, 14
Artificial numbers, 208
Āryabhaṭa the Elder (*c.* 510), 379, 387, 444, 608, 615, 626
Āryabhaṭa the Younger, 379
Arzachel. *See* al-Zarqâla
As, 208
Aśoka (3d century B.C.), 65–68
Assize of bread, 566
Astrolabe, 348, 601
Astrology, 73
Astronomical fractions, 229
Astronomical instruments, 348, 364
Astronomical progression, 495
Astronomy, 601–607

Asymptote, 318
Athelhard. *See* Adelard
Athenæus (*c.* 300?), 289
Athenian calendar, 650
Atomic theory, 677
Augrim. *See* Algorism
August, E. F. (*c.* 1850), 270
Augustine of Hippo (*c.* 400), 200
Aurelius Clemens Prudentius (*c.* 400), 166
Ausdehnungslehre, 268
Autolycus (*c.* 330 B.C.), 603
Aventinus, J. (*c.* 1552), 200
Avoirdupois weight, 639
Axioms, 280, 281
Ayer Papyrus, 396
Ayutâs, 308

Baba Nobutake (*c.* 1700), 367
Babbage, C. (*c.* 1840), 204
Babylonians, calendar, 655; geometry, 270; measures, 635, 640; numerals, 36; trigonometry, 601
Bachet, C. G. (*c.* 1612), 535
Backer Rule, 490
Backgammon, 166
Bacon, Roger (*c.* 1250), 282, 340, 372
Bagdad, 72
Baily, F. (1843), 376
Baker, H. (1568), 493, 502
Baker, S., 566
Bakhshâlî manuscript, 71
Ball, W. W. Rouse, 324, 691, 695, 696
Bamboo rods, 169–171, 432
Banerjee, G. N., 64
Bank, 187
Banking, 572, 574
Baraniecki, M. A., 176
Barbaro, E. (*c.* 1490), 186
Barbieri, M., 197
Barcelona, 75
Barclay, A. (*c.* 1500), 187
Barnard, F. P., 157
Bar Oseas (*c.* 250 B.C.), 670
Barozzi, F. (*c.* 1580), 82
Barrow, I. (*c.* 1670), 413, 690
Bartels, J. M. C. (*c.* 1800), 336
Barter, 568
Bartoli, C. (*c.* 1550), 349, 356
Barton, G. A., 38, 635
Base line, 376
Bassermann-Jordan, E. von, 673
Bastard Rule, 490
Battânî. *See* Albategnius
Baumeister, A., 162
Bayley, Sir E. C., 157, 158, 197
Bechtel, E. A., 200
Bede the Venerable (*c.* 710), 200
Beer, R., 75
Behâ Eddîn (*c.* 1600), 388
Beldamandi (*c.* 1410), 502
Bell, J. D., 168
Belli, S. (*c.* 1570), 285, 286, 355
Beman, W. W., 261, 267, 407
Ben. *See* Ibn
Benedetto da Firenze (*c.* 1460), 547
Benedict. *See* Benedetto
Benedict, S. R., 32
Ben Ezra. *See* Abraham ben Ezra
Ben Mûsâ. *See* al-Khowârizmî
Bentham, 327
Benvenuto d' Imola, 528
Berkeley, G. (*c.* 1740), 699
Berlin Papyrus, 432, 443
Bernoulli, Jacques (I) (*c.* 1690), 505, 528, 629
Bernoulli, Jean (I) (*c.* 1700), 612, 662; on complex numbers, 264
Bernoulli numbers, 505
Berosus (*c.* 250 B.C.), 670
Beyer, J. H. (1616), 245
Bézout, É. (*c.* 1775), 450
Bhandarkar, 70
Bhâskara (*c.* 1150), 380, 425, 426, 446, 484, 501, 525, 615
Bianco, F. J. von, 319
Bierens de Haan, D. (*c.* 1870), 518, 528
Biering, C. H., 298
Bigourdan, G., 347, 648
Bija Ganita, 380, 426
Bill of exchange, 577
Billeter, G., 560
Billion, 84
Binet, J. P. M. (*c.* 1812), 477
Binomial Theorem, 507, 511
Bion, N. (*c.* 1713), 359
Biot, J. B. (*c.* 1840), 655
Biquadratic equation, 466
Birkenmajer, A., 341
Bissaker, R. (1654), 205
Björnbo, A. A., 606
Bloomfield, M., 71
Blundeville, T. (*c.* 1594), 627
Bobynin, V., 34, 213, 219
Boccardini, G., 282
Böckh, A., 636
Böklen, E., 17
Boethius (Boetius) (*c.* 510), 6, 73, 524
Bolyai, F. (*c.* 1825), 337
Bolyai, J. (*c.* 1825), 335, 337
Bombelli, R. (1572), 19, 386, 428

Boncompagni, B. (*c.* 1870), 15, 27, 34, 71, 104, 108, 114, 153, 254
Bond, J. D., 610
Bonola, R., 282
Borghi (Borgi), P. (1484), 81
Borrowing process, 99
Bortolotti, E., 26, 459, 687
Bosanquet, R. H. M., 230
Bosmans, H., 240, 252, 328, 430, 435, 459, 465, 508, 685, 687, 688, 689
Bouelles, Charles de (*c.* 1500), 22, 327
Bouguer, P. (1734), 327, 376
Bouvelles. *See* Bouelles
Bowditch, C. P., 664, 672
Bowring, J., 166
Brachistochrone, 326
Brahe, Tycho, 310
Brahmagupta (*c.* 628), 380, 387; on quadratics, 445
Brāhmī forms, 67, 69
Brandis, J., 635
Braunmühl, A. von (*c.* 1900), 600, 610
Breasted, J. H., 645
Brewster, D., 692
Bridges, J. H., 340
Briggs, H. (*c.* 1615), 516
Bring (*c.* 1786), 470
Brocard, H. (*c.* 1900), 326
Broken numbers, 217
Broker, 558
Brouncker, W. (*c.* 1660), 420, 452, 692
Brown, E. W., 692
Brown, R., 58
Brugsch, H. K., 34
Bryson (*c.* 450 B.C.), 678
Buckley, W. (*c.* 1550), 236
Budé (Budaeus), G. (*c.* 1516), 209, 637
Budge, E. A. W., 316
Buée, Abbé (1805), 266
Bühler, G., 71
Bürgi, J. (*c.* 1600), 431, 523, 627
Bürk, A., 288
Burgess, E., 608
Burnaby, S. B., 664
Burnam, J. M., 75
Burnell, A. C., 70
Burnside, W. S., 473
Bushel, 645
Buteo (Butéon), J. (*c.* 1525), 428, 434, 541
Butler, R., 246
Butler, W., 11

Cabala, 525
Cabul, 72
Cæcilius Africanus (*c.* 100), 545
Cæsar, Julius (*c.* 46 B.C.), 659, 660
Cæsar, J. (1864), 542
Cajori, F., 5, 64, 88, 205, 231, 246, 283, 397, 404, 507, 546, 618, 686, 688, 692, 693, 699
Calandri, P. (1491), 142
Calculate, 166
Calculating machines, 202
Calculatores, 166
Calculi, 166. *See also* Counters
Calculones, 166
Calculus, 676
Calendar, 651, 655–664; Athenian, 658; Babylonian, 655; Chinese, 655; Christian, 660; Egyptian, 656; French Revolution, 663; Gregorian, 662; Roman, 659
Callippus (Calippus) (*c.* 325 B.C.), 659
Cambien, 183
Cambio, 569
Canacci, R. (*c.* 1380), 391
Canon Paschalis. *See* Calendar
Cantor, M. (*c.* 1900), 10, 160, 177, 345, 562
Capacity, 644
Capella (*c.* 460), 3, 200
Cappelli, A., 62
Capra, B. (1655), 246
Caracteres, 75
Carat, 639
Cardan, H. (or J.) (*c.* 1545), 384, 428, 459–464, 467, 530
Cardinal numbers, 26
Cardioid, 326
Cardo, 317
Carlini, F., 376
Carlini, L., 529
Carmen de Algorismo, 78
Carnot, L. N. M. (*c.* 1800), 333
Carpeting problems, 568
Carra de Vaux, 64, 229, 587
Carrying process, 93, 183
Carslaw, H. S., 282
Cartesian geometry, 318. *See also* Analytic geometry
Carvalho, A. S. G. de, 689
Casati, P. (1685), 246
Cassini's oval, 329
Cassiodorus (*c.* 502), 3
Castellum nucum, 26
Castillon (Castiglione), G. F. M. M. Salvemini, de (*c.* 1750), 326, 511
Casting accounts, 98
Casting out nines, 151
Cataldi, P. A. (*c.* 1590), 419

Catenary, 327
Catoptrics, 339
Cauchy, A. L. (*c.* 1830), 477, 700
Cavalieri, B. (*c.* 1635), 686
Cayley, A. (*c.* 1870), 322, 477
Celestial sphere, 364, 365, 367
Cellini, B., 342
Celsius, A. (*c.* 1740), 375
Celsus, J. (*c.* 75), 545
Censo, 394, 427
Census, 554. *See also* Censo
Cent, 648
Centesima rerum venalium, 247
Centesimal angle division, 627
Ceulen, L. van (*c.* 1580), 310
Chace, A. B., 436, 498, 500
Chain rule, 573
Chaldea. *See* Babylonians
Chalfant, F. H., 40
Challikân, Ibn (1256), 549
Chamberlain, B. H., 171
Champlain's astrolabe, 350
Chances, doctrine of, 529
Characteristic, 514
Charlemagne (*c.* 780), 648, 672
Chasles, M. (*c.* 1850), 9, 13, 80, 109, 157, 175, 177, 322, 378, 607
Chassant, L. A., 57
Chaucer, 9, 188
Check, 579; of elevens, 154; of nines, 151. *See also* Checks
Checkered board, 187
Checks on operations, 151
Chelebî (*c.* 1520), 626
Chessboard problem, 549
Cheyney, E. P., 564
Child, J. M., 690, 691
Chinese, algebra, 425, 432, 457, 475; calendar, 655; determinants, 475; geometry, 271; numerals, 39, 67, 68; series, 499; trigonometry, 602; values of π, 309
Ch'in Kiu-shao (*c.* 1250), 42, 381
Chords, table of, 604, 607, 614, 624
Choreb, 174
Chóu-peï Suan-king, 215, 602
Christian of Prag (*c.* 1400), 77, 95
Christian calendar, 660
Chrysippus (*c.* 240 B.C.), 524
Chrystal, G., 253
Chuquet, N. (1484), 84, 414, 502, 519
Chu Shï-kié (*c.* 1299), 257, 381
Chûzen. *See* Murai
Ciacchi, 29
Ciermans, J. (*c.* 1640), 203
Ciphering, 11
Circle, 278; quadrature of, 298, 302. *See also* π
Circumference, 278; of the earth, 369
Cissoid, 314, 327
Cistern problem, 536
Clairaut, A. C. (*c.* 1760), 325, 464
Clairaut, J. B. (*c.* 1740), 206
Clarke, F. W., 650
Clarke, H. B., 589
Classification, of numbers, 11; of equations, 442
Clavius, C. (*c.* 1583), 430, 662
Clay, A. T., 574
Clepsydra, 538, 671
Cloche, 672
Clock, 671, 672, 674
Clock problem, 548
Clodd, E., 195
Cloff, 567
Cloth, cutting of, 568
Cochlioid, 327
Cocked hat (curve), 327
Codex Alcobatiensis, 198
Codex Arcerianus, 504
Coefficient, 393
Coins, 646
Colebrooke, H. T., 91, 380, 637
Collins, J. (*c.* 1700), 415, 692
Colmar, T. de (*c.* 1820), 204
Colson, J., 693
Columna rostrata, 60
Combination lock, 527
Combinations, 524
Commercial problems, 552
Commercium Epistolicum, 511
Commission and brokerage, 558
Commutative law, 395
Company. *See* Partnership
Compasses, sector, 347
Complement, 98
Complex numbers, 261, 267
Composites, 12, 14
Compotus Reinheri, 62
Compound interest, 564
Compound numbers, 14
Compound proportion, 491
Comptroller, 186
Computing table. *See* Abacus
Computus, 651, 664. *See also* Calendar, Compotus
Conant, C. A., 576
Conant, L. L., 59
Conchoid, 298, 327
Concrete, 11
Condamine, La, 649

Cone, frustum of, 294
Congruence theorems, 285
Congruent figures, 285
Congruent numbers, 30
Conic sections, 317, 454, 679
Conjoint rule, 492
Conjugate numbers, 267
Consequent, 483
Contenau, G., 37
Continued fractions, 311, 418
Continued products, 311
Continuous magnitude, 26
Convergence, 507
Coordinate paper, 320
Coordinates, 316, 324
Copernicus, 610, 622
Cordovero, M. (*c.* 1560), 526
Cornelius de Judeis (1594), 354
Corporation, 576
Corpus Inscriptionum Etruscarum, 58
Corpus Inscriptionum Latinarum, 57
Corssen, W., 58
Cortese, G. (*c.* 1716), 115
Cosa. *See* Coss
Cosecant, 622
Cosine, 619, 631, 632
Cosmic figures, 295
Coss, 392
Cossali, P., 108, 384
Cotangent, 620–622, 632
Cotes, R., 265, 613
Cotsworth, M. B., 651
Coulba, 174
Counter, 181, 188
Counters, 158, 165, 166, 186, 190. *See also* Abacus
Counting. *See* Numerals
Counting rods, 169
Counting table, 174
Court of the Exchequer, 188
Cowry shells, 501
Crafte of Nombryng (*c.* 1300), 32, 78, 92, 98, 102, 104
Craig, J. (1693), 698
Cramer, G. (*c.* 1740), 328
Credit (creditum), 576
Credit, letter of, 576
Cretan numerals, 48
Cross ratio, 334
Cross staff, 346
Cruma, 361
Ctesibius of Alexandria (2d century B.C.), 538, 672
Cube, 292; duplication of, 298, 313
Cube numbers, 19
Cube root, 144, 148
Cubes, sum of, 504
Cubic curves, 324
Cubic equation, 454–467
Cubit, 640
Cuboid, 292
Cubus, 427
Cuento, 82, 88
Cuneiform numerals, 36, 68
Cunningham, W., 555
Currency, 569
Curtze, E. L. W. M., 123, 256, 345, 393, 544, 550, 586
Curve surfaces, 325
Curves, algebraic, 324; characteristics of, 326; cubic, 324, 325; of descent, 326; of double curvature, 325; plane, 324; of pursuit, 327; tautochronous, 328; transcendental, 324; well-known, 326. *See also* Cissoid and other names
Cusa, Nicholas (*c.* 1450), 327
Cushing, F. H., 59
Cycloid, 327
Cypriote numerals, 48, 49

Daboll, N., 588
Dagomari, Paolo (*c.* 1340), 81, 665
D'Alembert's Theorem, 474
Damianus, 340
Danfrie, 363
Danti, E. (*c.* 1573), 339, 340
Danzel, T. W., 59
Darboux, J. G., 333
Daremberg, C., 209
Darius vase, 161
Dase, Z. (*c.* 1860), 311
Day, 653
Days, names of, 657
Days of grace, 578
Decagon, 290
Decimal point, 238. *See also* Fractions
Decimanus, 317
Decimatio, 541
Decker, E. de (1626), 518
De Colmar. *See* Colmar
Decourdemanche, J. A., 64
Decussare principle, 56
Deficient number, 20
Degree, 232, 374, 443
De Haan. *See* Bierens de Haan
Delamain, R. (*c.* 1630), 205
Delambre, J. B. J. (*c.* 1800), 605
De latitudinibus formarum, 319
Democritus (*c.* 400 B.C.), 677
De Moivre, A. (*c.* 1720), 265, 529, 613
Demonstrative geometry, 271

De Morgan, A. (*c.* 1850), 148, 652, 691, 698
Demotic writing, 47, 68
Denarius, 208
Denominate number, 12
Denominator, 220
Density of the earth, 376
Deparcieux, A. de (1746), 530
Deposit (depositum), 576
Desargues, G. (*c.* 1640), 332
Descartes, R. (*c.* 1637), 322, 328, 343, 431, 443, 469, 471, 689
Descriptive geometry, 331
Determinant, 433, 475
Develey, E. (*c.* 1800), 85
De Vick, H. (*c.* 1379), 673
Devil's curve, 328
De Witt, J. (1658), 324
Dhruva, H. H., 70
Dial of Ahaz, 671. *See also* Sundial
Diameter, in geometry, 278; as number, 6
Diaz. *See* Diez
Dickson, L. E., 2, 29, 301
Dickstein, S., 11, 176
Dieck, W., 322
Diefenbach, L., 586
Diego de Landa, 43
Diez, J. (*c.* 1550), 385, 392; problems by, 590
Difference, 97
Differences, finite, 512
Differential, 696
Differential notation, 696–698
Differential triangle, 690
Differentiation, 691
Digges, L. and T. (*c.* 1572), 488
Digits, 12, 13, 15
Dill, S., 572
Dilworth, T., 588
Dime, 647
Dimension of an equation, 443
Dionysodorus (*c.* 50 B.C.), 670
Diophantus (*c.* 275), 422–424, 450, 452, 455
Dioptrics, 339
Discount, 565
Discrete magnitude, 26
Distances, 285
Distributive law, 395
Dividend, 131
Divine proportion, 291
Divisibility, 221
Division, 128; *a danda*, 141; *batello*, 136; of common fractions, 226; complementary, 134; definition, 128; *ferrea*, *aurea*, and *permixta*, 135; galley method, 136; Gerbert's method, 134; Greek, 133; long, 140, 142; *repiego*, 135; *scapezzo*, 136; sexagesimal, 233; short, 133; symbol, 406
Divisor, 131, 222; advancing, 139
D'Ocagne, M., 203
Dodson, J. (1742), 523
Dollar, 647
Dominical letter, 652
Dominis, A. de (1611), 343
Double False, 438
Drachma, 636
Dragoni, A., 198
Draughts, 166
Drechsler, A., 653
Drieberg, F. von, 16
Drumhead trigonometry, 357
Duality, 325
Duchesne, S. (*c.* 1583), 310
Dürer, A. (*c.* 1510), 296, 328, 342, 597, 598
Duhem, P. (*c.* 1900), 342
Duillier. *See* Fatio
Duplation, 33
Duplication of the cube, 298, 313
Duranis, 72
Dust table, 177
Dutt, R. C., 213
Dynamis, 394

e, logarithmic base, 517
Earth, density, 376; form, 369, 374; measure, 368–376
Easter, 651, 659. *See also* Calendar
Eastlake, F. W., 197
Economic problems, 552
Egypt, 45, 68, 270, 379, 600
Egyptians, calendar, 656; equations, 431, 435; fractions, 210; geometry, 270; measures, 634; numerals, 45; symbols, 410
Elastic curve, 328
Elchatayn. *See* False Position
Elefuga proposition, 284
Elevens, check of, 154
Ell, 640, 643
Ellipse, 317, 454
Elworthy, F. T., 200
Encyklopädie, 257
Eneström, G., 13, 96, 120, 139, 263, 431, 437, 461, 466, 511, 526
Engel, F., 267, 335
English measures, 640, 642
Enriques, F., 282
Epanthema of Thymaridas, 432

Epicycloid, 326, 328
Equality, symbol of, 395
Equating to zero, 431
Equation, 394; dimension of, 443; of payments, 559
Equations, Arab forms, 424, 434, 436; biquadratic, 384, 466; Chinese forms, 425, 432, 457; classification of, 442; cubic, 384, 454–467; Egyptian forms, 431, 435; factoring, 448; fifth degree, 469; fundamental theorem, 473; Hindu forms, 425, 434, 458; indeterminate, 451–453, 584; Japanese forms, 433; linear, 432, 435, 583; literal, 435; number of roots, 473; numerical higher, 471; in printed form, 426–432; quadratic, 443; simultaneous, 431, 432, 583; solution of, 435
Equiangular spiral, 329
Eratosthenes, 5, 370
Erlangen, *Sitzungsberichte*, 293
Erman, J. P. A., 130
Etruscans, 58, 64
Etten, H. van (1624), 535
Euclid (*c.* 300 B.C.), 4, 338; first editions, 272, 273; geometric terms, 274; on quadratics, 444
Eudoxus (*c.* 370 B.C.), 678
Euler, L. (*c.* 1750), 265, 311, 431, 450, 453, 464, 469, 613, 627, 629
Euler's Theorem, 296
Evans, A. J., 50
Evans, G. W., 687
Even numbers, 16, 18
Exchange, 569, 572, 577
Exchequer, 188
Exhaustion, method of, 303, 677–679
Exponents, 414
Eyssenhardt, 22

Fabri, O. (1752), 355
Factor (broker), 558
Factors, 30; of equations, 448
Fairs, 569
Fakhrî, 382, 388
Falkener, E., 594
False Position, 437–4
Famous problems, three, 297
Farthing, 647
Fathom, 641
Fatio de Duillier, N. (*c.* 1700), 698
Faulhaber, J. (*c.* 1620), 518
Faustus, L. Æbutius, 361
Favaro, A., 126, 206, 427, 436
Fegencz, H. G., 195
Fellowship. *See* Partnership
Fenn, J. (1769), 283
Fermat, P. de (*c.* 1635), 452, 453, 688, 689
Fermat, S. (1679), 322
Fermat's Numbers, 30
Fermat's Theorem, 30
Fernel (Fernelius), J. (*c.* 1535), 374
Ferramentum, 361
Ferrari, L. (*c.* 1545), 467
Ferro, Scipio del (*c.* 1500), 459
Fibonacci, Leonardo (*c.* 1202), 6, 310, 382, 384, 437, 457, 471, 505, 609
Figurate numbers, 24, 170
Fihrist, Kitâb al- (*Book of Lists*), 466
Finæus. *See* Fine
Fincke (Fink, Finke, Finchius), T. (*c.* 1583), 611, 621
Fine, Oronce (*c.* 1525), 345, 347
Finger notation, 196
Finger reckoning, 12, 120, 196
Finite differences, 512
Fink, E., 336
Fior, Antonio Maria (*c.* 1506), 459
Fischer, 475
Fitz-Neal (*c.* 1178), 188
Fleet, J. F., 70
Fleur de jasmin (curve), 328
Floridus. *See* Fior
Fluxions, 693. *See also* Calculus
Foecundus, 613, 621
Folium of Descartes, 328
Fontana, G. (*c.* 1775), 324
Foot, 641
Fortunatae Insulae, 317
Fracastorius, H. (Fracastoro, G.) (*c.* 1540), 373
Fraction, definition, 219; name, 217; terms, 220
Fractional exponent, 414
Fractions, addition of, 223; astronomical, 229; bar in writing, 215; Chinese, 215; common, 215, 219; complex, 219; continued, 311, 418; decimal, 235; division of, 226; Egyptian, 210; general, 213; Greek, 214, 231; multiplication of, 224, 232; operations with, 222; periodic, 30; physical, 229; Roman, 208, 214; sexagesimal, 228; subtraction of, 223; unit, 210, 212; vulgar, 219
Francesca. *See* Franceschi
Franceschi, Pietro (*c.* 1475), 296, 342
Franco of Liège (*c.* 1066), 310
Frank, J., 350
Frankland, W. B., 335

Freigius, J. T., 61, 179
Frey, J., 580
Friedlein, G., 16, 19, 50, 126, 127
Frisius. *See* Gemma
Frizzo, G., 123
Fromanteel, A. (*c.* 1662), 673
Frustum, of a cone, 294; of a pyramid, 293
Fujita Sadasuke (*c.* 1780), 41
Fundamental operations, 32, 35, 416
Furlong, 642
Fuss, P. H. von, 454

Gaging, 580
Galileo (*c.* 1600), 347, 373, 673
Gallon, 644
Galois, E. (*c.* 1830), 469
Gambling and probability, 529
Games, number, 16
Garbe, R., 71
Gardner, E. G., 392
Gardthausen, V., 52
Gauging. *See* Gaging
Gauss, C. F. (*c.* 1800), 337, 469, 474, 476, 507
Geber, 390. *See also* Jabir
Gebhardt, M., 427
Geet, M., 354
Geiler of Kaiserberg (*c.* 1500), 190
Gelcich, E., 321
Gellibrand, H. (*c.* 1630), 612
Gematria, 54, 152
Gemma Frisius (*c.* 1540), 520
Gemowe lines, 411
Geometric cross, 346
Geometric series. *See* Series
Geometric square, 345
Geometry, 270; analytic, 316; Babylonian, 270; Chinese, 271; demonstrative, 271; descriptive, 331; Egyptian, 270; elliptic, 338; Greek, 271; hyperbolic, 338; instruments of, 344; intuitive, 270; modern, 331; name for, 273; non-Euclidean, 331, 336; parabolic, 338; projective, 272, 332; Roman, 271
Gerbert (*c.* 1000), 74
Gergonne, J. D. (*c.* 1810), 334
Gerhardt, C. I., 10, 92, 396, 692, 698
Gernardus (13th century ?), 34, 100
Gersten, C. L. (1735), 204
Ghaligai, F. (*c.* 1520), 427
Gherardo Cremonense (*c.* 1150), 382, 616
Ghetaldi, M. (*c.* 1600), 321
Giambattista della Porta (1558), 373
Gibson, G. A., 515
Gill, 644
Ginsburg, J., 525
Ginzel, F. K., 651
Girard, A. (*c.* 1630), 415, 430, 474, 618, 622, 623
Glaisher, J. W. L., 515, 518
Gnecchi, F., 589
Gnomon, 16, 601, 603, 669, 671
Ġobâr numerals, 73, 175
Golden Number, 652
Golden Rule, 484, 486, 491
Golden Section, 291
Gomperz, T., 504
Goniometry, 612
Goschkewitsch, J., 168
Gosselin, G. (*c.* 1577), 430, 392, 435
Gould, R. T., 674
Gouraud, C. (1848), 528
Gow, J., 50, 145
Grafton's *Chronicles*, 638
Grain (weight), 637
Gramma symbol, 428
Grassmann, H. G. (*c.* 1850), 268
Graunt, J. (1662), 530
Gravelaar, 240
Graves, G. H., 676
Great Britain, 640, 642, 646
Greatest common divisor, 222
Greek Anthology, 532, 584
Greeks, algebra, 381; astronomy, 603; geometry, 271; measures, 636, 641; numerals, 47; trigonometry, 602
Green, J. R., 189
Greenough, J. B., 193, 647
Greenwood, I. (1729), 86, 494
Gregorian calendar, 662
Gregory, D. (*c.* 1700), 339
Gregory XIII, 662
Griffith, F. L., 432
Groma, 361
Gromaticus, 361
Group of an equation, 470
Gruma, 361
Günther, S., 126, 319, 419, 478
Guillaume, C. E., 650
Gunter, E. (*c.* 1620), 619, 621
Gunther, R. T., 362, 673
Gupta forms, 67
Gwalior inscription, 69
Gyula v. Sebestyén, 194

Haan. *See* Bierens de Haan
Ḥabash al-Ḥâsib (*c.* 860), 620
Hager, J., 501
Haldane. E. S., 323

Half-angle functions, 629
Hall, H., 188
Halley, E. (*c.* 1690), 530
Halliman, P. (1688), 255
Halliwell, J. O., 100
Halsted, G. B., 282, 335
Hamilton, W. R. (*c.* 1850), 267
Hanai Kenkichi (*c.* 1850), 203
Hankel, H. (*c.* 1870), 118, 261
Hare-and-hound problem, 546
Harkness, W., 650
Harley, R., 470
Harmonic points, 332
Harmonic series, 503
Harpedonaptae, 288
Harper's *Dictionary of Classical Literature and Antiquities*, 162
Harriot (Hariot), T. (*c.* 1600), 322, 430, 431, 471, 527
Harrison, J. (*c.* 1750), 674
Harun al-Rashid (*c.* 800), 672
Haskins, C. H., 27, 189
Ḥaṣṣâr, al- (*c.* 12th century), 118
Havet, J., 15
Hay, R., 541
Hayashi, T., 476
Heath, Sir T. L., 676, 680, 681; edition of Euclid, 14; *History*, 16
Hebrews, measures, 635; mysticism, 596; numerals, 53, 59
Hegesippus (*c.* 370), 542
Heiberg, J. L., 5, 80, 338, 524, 681
Heidel, W. A., 670
Heilbronner J. C. (*c.* 1740), 17
Heliodorus of Larissa, 340
Helix, 329
Hellins, J., 331
Helmreich, A., 390
Henderson, E. F., 188
Henry, C., 10, 14, 322
Hérigone, P. (*c.* 1634), 431, 618
Hermann, J. M. (*c.* 1814), 206
Hermite, C. (*c.* 1870), 470
Herodianic numerals, 49
Heromides, 274
Heron (Hero) (*c.* 50, or possibly *c.* 200), 605
Herschel, J. F. W. (*c.* 1840), 618
Herundes, 274
Herzog, D., 525
Heteromecic numbers, 18
Heuraet, H. van (*c.* 1659), 330
Hexagonal number, 24
Hiao-tze (*c.* 350 B.C.), 169
Hieratic writing, 47, 68
Hieroglyphics, 45, 68
Higher series, 504
Hilbert, D., 277
Hill, G. F., 64, 76
Hiller, E., 17
Hilprecht, H. V., 37
Hindasi, 64, 118
Hindu-Arabic numerals, 42
Hindus, algebra, 379; astronomy, 625; equations, 434; instruments, 365; measures, 637, 642; observatories, 365; quadratics, 444; trigonometry, 608, 615, 625, 629; values of π, 308
Hipparchus (*c.* 140 B.C.), 524, 604, 614, 659
Hippias of Elis (*c.* 425 B.C.), 300, 305
Hippocrates of Chios (*c.* 460 B.C.), 679; lunes of, 304
Hippolytus (3d century?), 152
Ḥisab al-Khataayn. *See* False Position
Hobson, E. W., 304
Hoccleve (1420), 188
Hochheim, A., 123
Hock, C. F., 7
Hoffman, S. V., 350
Homans, S. (*c.* 1860), 530
Homology, 334
Hoppe, E., 230, 514
Horace, 16
Horner, W. G. (1819), 381
Horsburgh, E. M., 203
Horseshoe problem, 551
Hoshino Sanenobu (1673), 592, 594
Hostus, M., 55
Ho-t'u, 196
Hoüel, S., 165
Hound-and-hare problem, 546
Hour, 669
Hourglass, 671
Howard, H. H., 68
Huber, D., 336
Hudalrich Regius (1536), 181
Hudde, J. (1659), 466, 689
Hübner, M., 157
Hulsius, L., 354
Hultsch, F., 209, 251, 636
Hunger, K. G., 18
Hunrath, K., 10, 145
Huswirt, J. (1501), 83
Huygens, C. (*c.* 1670), 673
Hylles, T. (1592), 491
Hyperbola, 317, 454, 689
Hyperbolic functions, 613

i for $\sqrt{-1}$, 613
Iamblichus (*c.* 325), 432
Ibn al-Zarqâla (*c.* 1050), 609, 616

Ibn Khallikân (1256), 549
Ibn Yûnis the Younger (*c.* 1200), 673
Ibrâhîm ibn Yaḥyâ. *See* Zarqâla
Ideler, L., 655
I Hang (*c.* 800 ?), 549
I-king, 524, 591
Imaginary numbers, 261; graphic representation, 263; in trigonometry, 612
Inch, 642
Incommensurable lines and numbers, 251
Incomposite numbers, 20
Indeterminate equations, 451–453, 584
India, 364. *See also* Algebra, Hindus, Numerals
Indivisibles, 677, 686
Infinite products, 420, 506
Infinite series, 506, 679
Infinitesimal. *See* Calculus
Inscribed quadrilateral, 286
Instruments in geometry, 344, 368
Integral sign, 696
Integration, 679, 684, 691
Interest, 555, 559; compound, 564; origin of term, 563
Inverse proportion, 490
Inverted fractional divisor, 227
Irrationals, 251
Irreducible case in cubics, 461, 464
Italian practice, 492
Iwasaki Toshihisa (*c.* 1775), 537

Jabir ibn Aflah (*c.* 1130), 390, 609, 632
Jackson, L. L., 479
Jacob, Simon (*c.* 1550), 346
Jacob ben Machir (*c.* 1250), 665
Jacobi, C. G. J. (*c.* 1830), 477
Jacob's staff, 346
Jacobs, F., 532
Jacobus. *See* Jacopo
Jacopo da Firenze (1307), 71
Jahnke, E., 268
Jaipur observatory, 366
Jai Singh (*c.* 1730), 365
Janet, P., 562
Janszoon (Jansen), Z. (*c.* 1610), 373
Japan, 421, 614, 701
Jastrow, M., 560
Jebb, S., 340
Jeber. *See* Jabir
Jechiel ben Josef (1302), 665
Jefferson, T. (*c.* 1790), 649
Jellen, M. (*c.* 1779), 118
Jemshîd. *See* al-Kashî
Jenkinson, C. H., 194
Jeremias, A., 351
Jerrard (*c.* 1834), 470
Jesuit missionaries, 364
Jetons (jettons), 192. *See also* Counters
Jevons, F. B., 561
Jews, 555, 565
Joannes Philoponus (*c.* 640 ?), 314
Job ben Salomon, 442
Johannes Hispalensis (*c.* 1140), 382
Johnson, G., 6
Jolly, Von (1881), 376
Jones, T. E., 359
Jones, W. (*c.* 1706), 312
Jordanus Nemorarius (of Namur, de Saxonia) (*c.* 1225), 384
Josephus problem, 541–544
Judah ben Barzilai, 684
Junge, G., 289
Just, R., 400
Jyā (jīva), 615, 616

Kabul, 72
Kästner, A. G. (*c.* 1770), 464, 613
Kalinga numerals, 67
Kant, I., 336
Karagiannides, A., 335
Karat, 639
Karkhî, al- (*c.* 1020), 382, 388, 504
Karoṣṭhī numerals, 65
Karpinski, L. C., 64, 93, 188, 212, 232, 236, 382
Kaye, G. R., 152, 158, 308, 365
Kelland, P., 268
Kenyon, F. G., 50
Kepler, J. (*c.* 1610), 342, 431, 685
Kerbenrechnung, 194
Keyser, C. J., 282, 335
Khallikân, Ibn (1256), 549
Khayyam. *See* Omar Khayyam
Khowârizmî (*c.* 825), 9, 72, 382, 388, 446
Kilderkin, 645
Kircher, A. (*c.* 1650), 392
Kittredge, G. L., 193, 647
K'iu-ch'ang Suan-shu (*Arithmetic in Nine Sections*), 257, 380, 433
Kliem, F., 679
Klimpert, R., 270
Klos, T. (1538), 176
Knight, Madam, 570
Knossos, 48
Knott, C. G., 157, 245
Knotted cords, 59
Köbel, J. (*c.* 1520), 549, 553
Koelle, S. W., 197
Konen, H., 452

Košťál, C., 437
Koutorga, M. S., 575
Kōwa. *See* Seki
Kress, G. von, 555
Kṣatrapa numerals, 67
Kubitschek, W., 162
Kuckuck, A., 101, 165
Kühn, H. (1756), 265
Kuṣana numerals, 67

Lacaille, N. L. de (*c.* 1750), 374
La Condamine, 649
Lacouperie, A. T. de, 157
Lacroix, S. F. (*c.* 1800), 328
Lagny, T. F. de (*c.* 1710), 612, 627
Lagrange, J. L. (*c.* 1780), 469, 470, 476, 688
Lahire, P. de (*c.* 1690), 324
Laisant, C. A., 327
Lambert, J. H. (*c.* 1770), 310, 336, 613, 629
Lambo, Ch., 414
Lanciani, R. A., 165
Landa, D. de, 43
Lange, G., 526
Langland, W., 80
Lao-tze, 195
Laplace, P. S. (*c.* 1800), 476
Lapland, survey in, 375
La Roche. *See* Roche
Latitude, 316
Latitudines, 319
Latus, 407, 409, 430
Lauder, W. (1568), 564
Lauremberg, J. W., 587
Lautenschlager, J. F. (*c.* 1598), 487
Law of signs (equation), 471
Leap year. *See* Calendar
Least squares, 530
Lecat, M., 476
Lecchi, G. A., 415
Lefebvre, B., 58
Legendre, F. (*c.* 1725), 192
Legge, J., 40
Legnazzi, E. N., 345
Leibniz, G. W. Freiherr von (*c.* 1682), 476, 691, 696; on complex numbers, 264; priority dispute, 698
Lemniscate, 329
Lemoine, É. M. H. (*c.* 1873), 290
Leonardo of Cremona (*c.* 1425), 610
Leonardo Fibonacci (of Pisa). *See* Fibonacci
Leonardo da Vinci (*c.* 1500), 327, 342
Lepsius, K. R., 641
Lessing, G. E., 584
Letter of credit, 576
Leucippus (*c.* 440 B.C.), 677
Leupold, J. (*c.* 1720), 204
Leurechon, J. (1624), 535
Levels, 357–360
Levi ben Gerson (*c.* 1330), 526, 630
Lex Falcidia, 544
Leybourn, W. (*c.* 1670), 202
Lichtenfeld, G. J., 402
Liebermann, F., 188
Lietzmann, W., 304
Lilius (Lilio), A. (*c.* 1560), 662
Limaçon, 326, 329
Limits, 13, 694
Lindemann, F., 26, 295
Line, 274
Lippersheim (Lippershey), J. (1608), 373
Liter (litre), 646
Little, A. G., 340
Lituus, 329
Liu Hui (*c.* 263), 380
Livingstone, R. W., 381
Livre, 646
Li Yeh (*c.* 1250), 381
Lobachevsky, N. I. (*c.* 1825), 335, 336
Locke, L. L., 195
Löffler, E., 230
Logarithmic spiral, 329
Logarithms, 513
Logistic, 7, 10, 392
Longitude, 316, 673
Longitudines, 319
Loria, G., 211, 212, 261, 270, 324, 331, 335, 470, 471, 684
Lo-shu, 106, 591
Love, A. E. H., 688
Lowell, P., 174
Lucas, E., 290, 536, 541
Luchu Islands, 171
Lucky numbers, 17
Lucretius (*c.* 100), 332
Ludlam, W. (1785), 283
Ludolf (Ludolph) van Ceulen (*c.* 1580), 310
Ludus duodecim scriptorum, 166
Ludus latrunculorum, 166
Lunes of Hippocrates, 304
Lutz, H. F., 210
Lyte, H. (1619), 247

McClintock, E. (*c.* 1890), 335, 530
Macedo, M. Æbutius, 357
Macfarlane, A. (*c.* 1900), 268
Mac Guckin de Slane, 549
Machin, J. (*c.* 1706), 312

Machina, 361
Machinula, 361
Mackay, J. S., 290, 313
Maclaurin's Theorem, 512
Macrobius, 22
Magic circles, 592, 594
Magic squares, 591
Magister Johannes, 457
Mahaffy, J. P., 50, 160
Mahāvīra. *See* Mahāvīrācārya
Mahāvīrācārya (*c.* 850), 108, 380, 387; on quadratics, 445
Mahmoud Bey, 34
Maḥmûd ibn Mohammed al-Rûmî (*c.* 1520), 626
Man, E. H., 18
Mannheim, A. (*c.* 1850), 206
Mannheim and Moutard, 333
Månsson, P. (*c.* 1515), 8, 486
Mantissa, 514
Margarita philosophica, 578
Marquardt, J., 165
Marre, A., 84, 128, 200
Martin, Th., 12
Martines, D., 157
Martini, G. H., 669
Masahiro. *See* Murai
Mascart, J., 297
Maskeleyne, N. (*c.* 1770), 376
Maspero, G., 193, 569
Maspero, H., 286
Mass of earth, 376
Massoretes, 53
Masterson, T. (*c.* 1590), 386, 556
Mas'ûdî (*c.* 950), 550
Mathieu, E., 144
Matthiessen, L., 424, 471, 585
Maupertuis, P. L. M. de (1746), 375
Maurolico (Maurolycus), F. (1558), 622
Maximus Planudes (*c.* 1340), 81
Maya numerals, 43
Meadows of Gold, 550
Mean proportionals, 454, 483
Measures, 634
Mechanical calculation, 156
Mediation, 33
Mei Wen-ting (*c.* 1675), 170
Mellis, J. (1594), 249
Menæchmus (*c.* 350 B.C.), 454
Menant, J., 560
Menelaus (*c.* 100), 603, 606, 615
Mensa geometricalis, 177
Mensa Pythagorica, 177
Mensor, 361
Mercatello (*c.* 1522), 235
Merchants' Rule (or Key), 488
Messahala (*c.* 800), 353
Messier, C. (1775), 649
Metius. *See* Adriaen
Meton (*c.* 432 B.C.), 658
Metric system, 242, 376, 648
Metrodorus (*c.* 500 ?), 532
Metropolitan Museum, 48, 49, 634
Miju Rakusai (1815), 628
Mikami, Y., 40, 124, 215
Mile, 641
Milham, W. I., 673
Milhaud, G. (*c.* 1900), 444
Miller, G. (1631), 518
Miller, G. A., 597
Miller, J. (1790), 649
Millet, J., 323
Milliard, 85
Million, 80
Milne, Joshua (*c.* 1830), 530
Mint problems, 589
Minuend, 96
Minus sign, 396, 397; in the Rule of False, 397, 441
Minute, 218, 232
Mîram Chelebî (*c.* 1520), 626
Mirror, 356, 358
Misrachi, Elia (*c.* 1500), 33
Mixed numbers, 14
Mixtures, 588
Miyake Kenryū (*c.* 1715), 173, 543
Mochinaga (1687), 702
Močnik, 101
Modern geometry, 331
Modulus, 267
Moirai (μοῖραι), 232, 615, 617
Moivre. *See* De Moivre
Mommsen, T. (*c.* 1850), 55
Money changers, 575. *See also* Bank, Check, Currency, Exchange
Monforte, A. di (*c.* 1700), 689
Monge, G. (*c.* 1800), 332
Monier-Williams, M., 387
Montauzan, C. G. de, 359
Month, 653
Months, names of, 659
Montucci (1846), 330
Moore, Jonas (1674), 622
Morland, S. (*c.* 1670), 204
Morley, F. V., 322
Morley, S. G., 43
Morse, H. B., 40
Mortality table, 530
Moulton, Lord, 514
Mouton, G. (*c.* 1670), 512, 649
Moya, J. P. de (*c.* 1562), 198

Müller, J. H. T., 274
Müller, T., 519
Muir, T., 476
Mule-and-ass problem, 552
Multiple angles, functions, 629
Multiplication, 101; by aliquot parts, 123; *per bericocoli*, 107; Bhāskara's plan, 107; cancellation, 118; *per casteluccio*, 111; *per colonna*, 124; column, 112; of common fractions, 224; complementary, 119–122; contracted, 123; *per coppa*, 119; cross, 112; *per gelosia*, 114; Greek and Roman, 106; left-to-right, 118; *per organetto*, 108; Pacioli's plans, 107; Polish, 120; process of, 106; quadrilateral, 114; quarter squares, 123; *per repiego*, 117; Russian, 106, 120; *per scacchero*, 108; *per scapezzo*, 117; of sexagesimals, 232; short methods, 119, 122; sign of, 114; Spanish, 107; symbols, 402; table, 123
Munro, D. C., 665
Murai Chūzen (*c.* 1765), 511
Murai Masahiro (*c.* 1732), 358, 359, 614
Muramatsu Kudayū Mosei (*c.* 1663), 542, 593
Muratori, L. A., 200
Myriad, 308

Naber, H. A., 290
Nāgarī numerals, 67
Nagl, A., 109, 158, 162
Name, R. van, 168
Nānā Ghāt inscriptions, 65
Napier, J. (*c.* 1614), 431, 514, 611, 632
Napier's rods, 202
Napier's Rules, 632
Narducci, E., 3, 8
Nasik numerals, 66, 67
Naṣîr ed-dîn (*c.* 1250), 609, 630, 632
Nasmith, J., 566
Neander, M. (*c.* 1570), 428
Negative exponents, 414
Negative numbers, 257, 396
Neile, W. (*c.* 1665), 330
Nepal numerals, 67
Nesselmann, G. H. F., 232
Neuberg, J., 327
Newton, Sir Isaac (*c.* 1680), 324, 344, 472, 511, 612, 692; priority dispute, 698
Newton, John (*c.* 1658), 612, 619
Nicholas Cusa (*c.* 1450), 327
Nicholson, W. (1787), 206
Nick sticks, 194
Nicolas Petri (1567), 459, 465, 468
Nicomachus (*c.* 100), 5
Nicomedes, conchoid of, 298, 327
Nider, J., 563
Nielsen, N., 611
Nine Sections, 215, 380, 432
Nines, casting out, 151
Ninni, A. P., 64
Nobutake. *See* Baba
Noel, E., 650
Noether, M., 335
Nokk, A., 604
Non-Euclidean geometry, 335
Norm, 267
Norton, R., 240
Norwood, R. (1631), 618
Notation, 33. *See also* Numerals
Number puzzles, 582
Number theory, 4. *See also* Arithmetica
Numbers, artificial, 208; complex, 261; composite, 12, 14; compound, 14; conjugate, 267; cube, 19; even, 16, 18; heteromecic, 18; imaginary, 261, 263; irrational, 251; large, 86; mixed, 14; negative, 257; oblong, 251; odd, 16, 18; perfect, 20; plane, 18; prime, 5, 20, 30; reading, 36, 86; square, 18, 24; writing, 36, 86
Numerals, Arabic, 68, 69, 70; Attic, 49; Babylonian, 36, 68; Chinese, 39, 67, 68; Cretan, 48; Cuneiform, 36, 68; Cypriote, 48; Egyptian, 45, 68; Etruscan, 58, 64; Gobâr, 73; Greek, 47, 49, 164; Hebrew, 53, 59; Herodianic, 49; Hindu-Arabic, 42–88; Hindu variants, 67, 70, 71; Roman, 54; Sanskrit, 42, 70; Spanish, 86; Sumerian, 67
Numeration, 33
Numerator, 220
Numerical higher equations, 471
Nunes, P. (*c.* 1530), 465
Núñez. *See* Nunes

Obenrauch, F. J., 331
Oblique coordinates, 324
Oblong numbers, 251
Odd numbers, 16, 18
Ôhashi (1687), 702
Oliva, A., 460
Omar Khayyam (*c.* 1100), 382, 426, 442, 447, 456, 508
Operations, fundamental, 32, 35, 416
Oppert, G., 64, 165

Optics, 338
Ordinal numbers, 26
Ordinate, 318, 324
Oresme (*c.* 1360), 319, 414, 526, 689
Otto (Otho), V. (*c.* 1573), 310, 627
Oughtred, W. (*c.* 1630), 205, 413, 430, 611
Ounce, 636
Oval of Cassini, 329
Ovio, G., 339
Ozanam, J. (1691), 326, 535

π, 270, 307–313, 702; the symbol, 312
Pacioli, L. (*c.* 1494), 384, 427, 443
Pan chu tsih, 168
Panton, A. W., 473
Paolo Dagomari, dell' Abaco (*c.* 1340), 123, 136, 216
Papias (*c.* 1050), 178
Pappus (*c.* 300), 689
Pappus-Guldin theorem, 296
Parabola, 317, 454, 679, 680
Paraboloid, 685
Parallelepiped, 291
Parallels, 279, 335, 336; postulate of, 282
Parent, A. (*c.* 1710), 325
Parentheses, 416
Partnership, 554
Pascal, B. (*c.* 1650), 203, 332, 508, 528, 529, 691
Pascal, E., 326
Pascal's Triangle, 508
Passions (operations), 36
Pasturage problems, 557
Paton, W. R., 532
Paul (Paolo) of Pisa, 81
Pauly-Wissowa, 209
Pearls (curve), 330
Peck, 645
Peckham, John (*c.* 1280), 341
Pecunia, 645
Peet, T. Eric, 34
Peetersen, N. (1567), 459
Peking, instruments at, 364
Peletier (Peletarius), J. (*c.* 1560), 439
Pell, J. (*c.* 1650), 406, 413
Pell Equation, 452
Pellos (Pellizzati) (*c.* 1492), 238
Pena (Péna, de la Pêne), J. (*c.* 1557), 338
Penny, 647
Pentagon, 290
Pentagonal number, 24
Per cent sign, 250
Percentage, 247
Perch, 644
Perez (modern Pérez). *See* Moya
Perfect numbers, 20
Periodic fractions, 30
Periods in notation, 86
Periphery, 278
Permillage, 250
Permutations, 524, 528
Persia, 364, 455, 608
Perspective, 338
Peru, mission to, 375
Peter, B., 654
Petri, Nicolas (1567), 459, 465, 468
Petrie, W. M. F., 293, 634, 642, 652
Petrie Papyrus, 432
Peurbach, G. von (*c.* 1460), 609
Picard, J. (*c.* 1670), 322, 374
Pinches, T. G., 560
Pint, 645
Pitiscus, B. (*c.* 1595), 611, 622
Pittarelli, G., 342
Place value, 43, 44
Plane numbers, 18
Plane surface, 276
Plane table, 356
Planets, 657
Planisphere, 351
Planudes, Maximus (*c.* 1340), 81
Platea, F. de (*c.* 1300), 563
Plato (*c.* 380 B.C.), 2, 5
Platonic bodies, 295
Playfair, J. (*c.* 1795), 283
Plimpton, G. A., 383, 391, 397, 404, 406, 427, 485, 551, 571, 666, 667
Plücker, J. (*c.* 1850), 325
Plunket, E. M., 672
Plural proportion, 492
Plus and minus signs, 397, 398, 402
Plus sign, variants, 402; in the Rule of False, 397, 441
Plutarch, 602
Poincaré, H. (*c.* 1900), 335
Point, 274
Points, harmonic, 332
Polar coordinates, 326
Poleni, G. (*c.* 1740), 204
Poll tax, 572
Polygonal numbers, 24, 27, 499
Polygons, area of, 606; regular, 301
Polyhedron theorem, 296
Polyhedrons, 295; regular, 296; stellar, 296
Pomodoro, G. (1624), 356, 358
Poncelet, J. V. (*c.* 1830), 333
Pondera, 361
Pons asinorum, 284, 289

Poole, R. L., 189
Porta, G. della (1558), 373
Portius, L., 637
Poseidonius (Posidonius) (*c.* 100 B.C.), 5, 371
Position. *See* False Position
Postulate of parallels, 282
Postulates, 280, 281
Pott, A. F., 200
Potts, R., 637
Poudra, N. G., 338
Pound, 636, 638, 646, 647
Powel, J., 567
Powers, 393
Poynting, J. H. (1891), 376
Practica (pratica, pratiche), 11
Practice, 492
Prayer sticks, 196
Prime meridian, 317
Prime number, 5, 20, 30
Printing, effect on numerals, 77
Priscian (6th century), 54
Prism, 291
Probability, 528
Problem, chessboard, 549; cistern, 536; hare-and-hound, 546; of Hiero's crown, 590; horseshoe, 551; Josephus, 542; testament, 544; Turks-and-Christians, 541. *See also* Problems
Problems, algebraic, 582; commercial, 552; economic, 552; elementary, 532; famous and fanciful, 297, 501, 532, 536; of Metrodorus, 532; of pursuit, 546; typical, 536
Product, 90
Profit and loss, 557
Progressions, 494, 496
Projectiles, 192
Projective geometry, 331, 332
Proportion, 413, 477, 479; arrangement of terms in, 483, 488, 489; compound, 491; divine, 291; inverse, 490; relation to series, 497; terms of a, 483; types of a, 482. *See also* Rule of Three
Proportional compasses, 347
Proportionality, 478
Propositions of geometry, typical, 284
Prosdocimo de' Beldamandi (*c.* 1410), 502
Psammites, 5
Ptolemy, Claude (*c.* 150), 371, 607, 615, 624, 629, 631, 632
Ptolemy's Theorem, 624
Puliśa, 308
Pulverizer, 387
Pure quadratic, 450
Pursuit, curve of, 327; problems of, 546
Puzzles, 582. *See also* Problems
Pyramid, 292; frustum of, 293
Pyramidal number, 25
Pythagoras (*c.* 540 B.C.), 4
Pythagorean numbers, 288, 451
Pythagorean table, 124
Pythagorean Theorem, 288

Quadrant, 352–357
Quadratic equation, 443–451; Hindu rules for, 444–446
Quadratrix, 300, 305
Quadratum geometricum, 345
Quadrature, 298, 302. *See also* Circle
Quadrivium (Quadruvium), 3
Quart, 644
Quarter squares, 123
Quaternions, 267
Quentos, 88
Quipu, 195
Quotient, 131

Rabuel, C., 323
Radical sign, 408, 409
Radius, with abacus, 138, 178; geometric, 278
Radulph of Liège (*c.* 1010), 177
Rahn, J. H. (*c.* 1660), 406, 411, 431, 474
Rainbow, 343
Ramsay, J. H., 189
Ramsay and Lanciani, 165
Ramus, P. (*c.* 1550), 342, 430
Range finder, 363
Raphson (Ralphson), J. (*c.* 1715), 692, 698
Rara Arithmetica, 34
Ratio, 477, 478, 678; anharmonic, 333, 334
Ratios compounded, 481
Rechenmeisters, 190
Rechenpfennig, 191
Reckoning on the lines, 183
Recorde, R. (*c.* 1542), 386, 411, 412, 439
Recreations, mathematical, 532. *See* Problem *and* Problems
Rectifications, 330
Refraction, 343
Regiomontanus (*c.* 1470), 427, 429, 609, 626, 630
Regius, H. (1536), 181

Regula, augmenti et decrementi, 441; bilancis, 440; coecis, 586; falsi, 437–442; infusa, 442; lancium, 440; positionis, 437–442; potatorum, 587; quatuor quantitatum, 607; sex quantitatum, 607; virginum, 587
Regular polygons, 301
Rehatsek, E., 597
Reiff, R., 506
Reimer, N. T., 298
Reinach, S., 50
Reinaud, J., 10
Relation, symbols of, 410
Remainder, 132
Requeno, V., 200
Res, 408, 427
Rhæticus, G. J. (*c.* 1550), 610, 621, 622, 623, 627, 629
Rhind, A. H. (papyrus), 34
Rhonius. *See* Rahn
Riccati, V. (*c.* 1750), 613
Richardson, L. J., 198
Richter (*c.* 1850), 311
Riemann, G. F. B. (*c.* 1850), 338
Right-angled triangle, 288
Risner, F. (*c.* 1570), 342
Robbins, F. E., 5, 552
Robert of Chester (*c.* 1140), 382, 426
Roberts, E. S., 50
Robertus Anglicus (*c.* 1231), 621
Roberval, G. P. de (*c.* 1640), 688
Robinson, L. G., 681
Roche, E. de la (*c.* 1520), 407
Rod, 642
Rod numerals, 40, 45
Rodet, L., 168, 278, 423
Rogimbold (*c.* 1010), 177
Rollandus (*c.* 1424), 77
Romans, calendar, 659; fractions, 208; measures, 636, 641; numerals, 54
Rood, 644
Roomen, A. van (*c.* 1593), 310
Roots, 144; abbreviated methods, 150; approximate, 253; cube, 144, 148; higher, 149; meaning of the term, 150; square, 144, 253
Rosaries, 196
Roscoe, translation, 342
Rosen, F., 388
Rosenhagen, G., 193
Rossi, G., 345
Roth, P. (*c.* 1610), 474
Roulette, 328
Round, J. H., 189
Rudio, F., 303, 312
Rudolff, C. (*c.* 1525), 384, 408, 428, 458, 520
Ruffini, P. (*c.* 1800), 469
Rule of False Position, 437–442. *See also* Regula
Rule of Five, Seven, etc., 491
Rule of Mixtures, 588
Rule of Three, 477, 483; compound, 491
Ruled surfaces, 326
Rûmî, al- (*c.* 1520), 626

Sa'adia ben Joseph (*c.* 930), 212
Saalfeld, G. A., 167
Saccheri, G. (1733), 335
Sachau, E. C., 650
Sagitta, 619
Saglio, E., 209
Saint-Vincent, G. de (*c.* 1650), 689
Saka forms, 67
Salamis abacus, 162
Salomon, J., 101
Salvemini. *See* Castillon
Salvianus Julianus (*c.* 125), 545
Salvino degli Armati (1317), 372
Sanchu, 171
Sand Reckoner, 5, 80
Sand table, 156
Sang Hung (*c.* 118 B.C.), 170
Sangi, 41, 171–173
Santa Maria de Ripoll, 75
Sāradā numerals, 71
Sargon (*c.* 2750 B.C.), 601
Satō Shigeharu (1698), 171
Sawaguchi Kazuyuki (*c.* 1665), 701
Sayce, A. H., 230, 554, 560
Scales, method of the, 440; of counting, 9, 41
Scaphe, 370
Scarburgh (Scarborough), C. (*c.* 1660), 618
Schack-Schackenburg, H., 432, 444
Schepp, A., 326
Scheubel (Scheybel), J. (*c.* 1550), 428
Schilling, F., 594
Schisare, 221
Schlegel, V., 268
Schmid, W. (1539), 292
Schmidt, J. J., 23
Schoner, A. (*c.* 1560), 430
Schooten, F. van, the Younger (*c.* 1656), 344, 469, 474
Schotten, H., 274
S'choty, 176
Schubert, H., 191
Schülke, A., 231

Schütte, F., 270
Schulenberg, W. von, 194, 195
Schumacher, H. C. (*c.* 1830), 337
Scipio Nasica (*c.* 159 B.C.), 672
Score, 192
Sebestyén, 194
Sebokht (*c.* 650), 64, 72
Secant, 622
Second, 232
Sector compasses, 347
Sedgwick, W. T., 337
Sédillot, L. P. E. Am., 626
Seebohm, F., 637
Sefer Jezira, 684
Seki Kōwa (*c.* 1680), 433, 476, 592, 701
Semidiameter, 279
Seqt (seqet, śḳd), 600, 619
Series, 494; antitrigonometric, 513; arithmetic, 498; convergency of, 507; extent of treatment of, 497; geometric, 500; Gregory's, 513; harmonic, 503; higher, 504; infinite, 506, 679; kinds of, 495; logarithmic, 513; names for, 496; relation to proportion, 497; sum of, 17, 497–505; trigonometric, 512
Serret, J. A. (*c.* 1865), 692
Seven, 2; check of, 154
Seven liberal arts, 3
Sexagesimal fractions, 228; symbols of, 234
Shadows, 17, 602, 620; tables of, 621
Shanks, W. (*c.* 1853), 311
Sheffers, G., 692
Shekel, 636
Shilling, 647
Shotwell, J. T., 651
Sié Fong-tsu (*c.* 1650), 523
Sieve of Eratosthenes, 5
Sign, 233
Significant figures, 15
Signs, law of, 396
Silberberg, M., 100, 152
Simon, M. (*c.* 1890), 432
Simson, R. (*c.* 1750), 469
Simultaneous equations, 431
Simultaneous quadratic equations, 450
Sine, 614; abbreviations for, 618; name for, 616
Sines, addition theorem of, 628; tables of, 626; theorem of, 630, 632
Single False, 440
Sinus totus, 627
Sissah ibn Dahir (*c.* 1250), 549
Slane, Mac Guckin de, 549
Slate, 179
Sloman, H., 692
Sluze, R. F. W., Baron de (*c.* 1660), 689; conchoid of, 327
Smethurst, 168
Smith, A. H., 168
Smith, D. E., 64, 124, 182, 210, 290, 312, 385, 536, 552, 676
Smith, W., 340
Smogolenski (*c.* 1650), 523
Smyly, J. G., 53, 164, 256, 294
Snell, W. (1627), 631
Snelling, T., 182
Solid analytic geometry, 325
Solid geometry, 291
Solid numbers, 19, 25
Solidus, 646
Sommerville, D. M. Y., 335
Soreau, R., 156
Spaces (spacia), 183
Species, 35
Speculum, 356, 358
Speidell, J. (1620), 517
Sphere, 294; astronomical, 603–608
Spherical numbers, 25
Spherical triangle, 631
Spinoza, B. (*c.* 1670), 528
Spirals, 329
Square, geometric, 345, 355
Square numbers, 18, 24
Square root, 144, 253
Squares, criteria for, 256; sums of, 17, 504
Squaring the circle, 298, 302
Śrīdhara (*c.* 1020), 446
Stadium, 340, 641
Stäckel, P., 335, 337
Stark, W. E., 345
Steele, R., 32, 78
Steiner, J. (*c.* 1840), 334
Steinschneider, M. (*c.* 1850), 34, 36, 127, 200, 437, 665
Stephano Mercatello (*c.* 1522), 235
Stereographic projection, 344, 351
Stevin, S. (*c.* 1590), 430
Stifel, M. (*c.* 1525), 384, 403, 502, 519, 520
Stitt, S. S., 595
Stock, 194, 576
Stone, E. (1740), 326
Stoy, H., 197
Straight line, 275
Studnička, F. J., 77, 310
Study, E., 261
Sturm, A., 298
Sturm, L. C. (*c.* 1710), 85
Suan-hio-ki-mŏng, 257

Suan-pan, 168, 203
Substractio, 95
Subtraction, 94, 184; of fractions, 223; methods of, 98; symbols of, 395
Subtrahend, 96
Suevus, 12
Sum, 89, 90; of a series, 17, 497–505
Sumario Compendioso, 385, 392, 590
Sundial, 370, 601, 620, 669–671
Sun-tzï (1st century ?), 380, 433
Superstitions, 17
Supputandi ars, 11
Surds, 251, 252, 257
Surface, 276
Surfaces, 325; curvature of, 325; ruled, 326
Surveying, 317, 344, 363
Su-shuh ki-i, 168
Suter, H. (*c.* 1890), 34, 74, 92, 118, 164, 437, 481, 550
Swan pan. *See* Suan-pan
Sylvester, J. J. (*c.* 1850), 477
Sylvester, Pope. *See* Gerbert
Symbolism, poor, 417. *See also* Symbols
Symbols, of addition, 395–398, 402; of aggregation, 416; of Diophantus, 422–424; of division, 406; of equality, 410; of equations, 421, 434; of imaginary numbers, 266; of inequality, 413; of multiplication, 402; of negative number, 259, 395–397; Oriental, 424; of per cent and per mill, 250; of proportion, 413; of ratio, 406; of relation, 410; of roots, 407; in the Rule of False, 397, 441; of subtraction, 395–397; of the unknown, 422, 428; of Vieta, 430, 449

Tâbit ibn Qorra (*c.* 870), 455, 685
Tables, 609 seq. *See also* Chords, Mortality, Multiplication, Powers, Roots, Shadows, Sines, Tangents, Trigonometric functions
Tabula geometricalis (abaci), 177
Tait, P. G. (*c.* 1880), 267, 268
Takebe Hikojirō Kenkō (*c.* 1722), 702
Takebe Kemmei (*c.* 1722), 702
Takeda Shingen (1824), 536
Talent, 635, 636
Tally sticks, 171, 192
Tangent, 620
Tangential coordinates, 326
Tangents, abbreviations for, 622; problem of, 689; tables of, 624; theorem of, 611, 631
Tangutans, 169
Tannery, P. (*c.* 1900), 3, 101, 124, 126, 254, 255, 313, 322, 453, 478, 604, 605
Tanstetter, G. (*c.* 1520), 341
Tanto, 393
Tare and tret, 567
Tariff, 571
Tartaglia, Nicolo (*c.* 1545), 26, 384, 428, 460, 494
Taurinus, 337
Tautochronous curve, 328
Taxes, 571
Taylor, I., 398
Taylor, J., 92
Taylor's Theorem, 512
Teixeira, F. G., 313, 326
Telescope, 372
Temple, R. C., 637
Tessera, 195
Testament problem, 544
Thales (*c.* 600 B.C.), 4, 602; measuring distances, 285
Theologumena, 27
Theon of Smyrna (*c.* 125), 5, 6, 453
Theophrastus (*c.* 350 B.C.), 677
Theory of numbers, 2, 29
Thibaut, G., 288
Thibetan "wheel of life," 595
Thierfelder, C. (1587), 439
Thirteen, fear of, 17
Three, 2
Thymaridas (*c.* 380 B.C.), 432
Thynne, F., 564
Tieffenthaler, J. (*c.* 1750), 366
Time, 651
Tithes, 572
Titulus, 61
Tod, M. N., 50, 162
Todhunter, I. (*c.* 1850), 528
Ton, 645
Topics for Discussion, 31, 155, 207, 269, 377, 531, 599, 633, 675, 703
Touraeff (Turajev), B. A., 293
Townsend, E. J., 277
Tractatus de uniformitate, etc., 319
Transcendental numbers, 268. *See also* Logarithms, π, Trigonometric functions
Transversals, theory of, 333
Treichel, A., 195
Tret, 567
Treutlein, P., 197, 255, 497
Triangle, 288, 290; arithmetic (Pascal's), 508; right-angled, 288; spherical, 604–608, 631
Triangular numbers, 24

Trigonometric functions, 623. *See also* Sine, Cosine, etc.
Trigonometric solutions of equations, 474
Trigonometry, 357, 600; analytic, 613
Trisection of an angle, 297, 298
Trivium, 3
Tropfke, J., 32
Troy weight, 638
Truck (troquer), 570
Truel, H. D. (1786), 266
Tschirnhausen, E. W., Graf von (*c.* 1690), 470
Tseu pan tsih, 168
Tun, 645
Turajev. *See* Touraeff
Turchillus (*c.* 1200), 177
Turetsky, M., 526
Turks-and-Christians problem, 541
Tycho Brahe, 310
Tyler, H. W., 337
Tylor, E. B., 16, 195

Ulugh Beg (*c.* 1435), 609
Umbra recta, 17, 602, 620
Umbra versa, 620
Uncial interest, 562
Unger, F., 84
Unit fraction, 210, 212
Unitary method, 494
United States, arithmetic in, 86
United States money, 647
Unity, 13, 26
University of Pennsylvania, 635
Unknown quantity, 393
Uranius, H., 637

Vacca, G., 302, 549
Vākāṭaka numerals, 67
Valerianus, J. P. (Bellunensis), 200
Valerio, Luca (1606), 685
Valhabī numerals, 67
Value, 645
Vander Hoecke (*c.* 1515), 399, 401, 427
Vandermonde, A. T. (*c.* 1775), 470, 476
Vanhée, L., 40, 42, 425, 448, 585
Varro (*c.* 60 B.C.), 3, 215
Vassilief, 335
Vaux, Carra de, 64, 229, 587
Vega, G. (*c.* 1775), 311
Veratti, B., 3
Verse, rules in, 439, 487, 488, 491, 500
Versed sine, 618
Verses in notation, 53
Vicesima libertatis, 247
Vicesimatio, 542
Viedebantt, O., 371
Vieta, F. (*c.* 1580), 310, 392, 430, 449, 465, 469, 472, 474, 503, 610, 623, 627, 629, 631
Vigarié, E., 290
Villicus, F., 194
Vincent, A. J. H., 162, 165, 339
Vinculum, 60
Visierer, 581
Vissière, 42
Vitale, G. (Vitalis, H.) (*c.* 1690), 393
Vitello. *See* Witelo
Vivanti, G., 692
Vlacq (Vlack), A. (*c.* 1650), 518
Vogt, H., 251, 289, 686
Volusius Maecianus (2d century), 215
Von Schulenberg, W., 194, 195
Vysierer. *See* Visierer

Waard, C. de, 688
Wade, H. T., 673
Wäschke, H., 81
Wallace, W. (*c.* 1810), 613
Wallis, J. (*c.* 1650), 7, 263, 311, 413, 415, 420, 431, 503, 612, 689, 695
Wallner, C. R., 686
Walters, R. C. S., 359
Wampum, 196
Wang Jung (3d century), 170
Wappler, H. E., 260
Ward, J., 29
Ward, S. (1654), 618
Warren, C., 640
Wattenbach, W., 194
Wax tablet, 178
Weber, C. F., 542
Week, 655
Wegener, A., 232
Weidner, E. F., 351
Weight, 634
Weissenborn, H., 27, 123, 164, 692
Welsh (Welsch) practice, 493
Wenceslaus (Wentsel), M. (1599), 493
Wertheim, G., 33, 435, 452
Wessel, C. (1797), 265
Weyr, E., 270
Whipple, F. J. W., 203
Whish, C. M., 309
Whitford, E. E., 452
Widman (Widmann), J. (1489), 573
Wiedemann, E., 340, 434, 489
Wieleitner, H., 322, 342, 415, 506, 526
Wiener, C., 331
Wilkens, M., 12

William of Malmesbury, 175
Williams, S. W., 40
Willichius (1540), 19
Willsford, T. (*c.* 1662), 236
Wilson, T., 578
Wilson's Theorem, 29
Wing, V. (*c.* 1648), 407
Wingate, E., 205
Witelo (*c.* 1270), 341
Witt, J. de (1658), 324
Witt, R., 247, 565
Witting, A., 427
Woepcke, F., 10, 34, 74, 115, 118, 157, 160, 175, 320, 437, 467, 508, 626
Wolf, R., 604
Woolhouse, W. S. B., 650
Wordsworth, C., 652
Wren, Sir C. (*c.* 1670), 649
Wright, E. (*c.* 1600), 404
Wright, S. (*c.* 1614), 404
Writing material, 36, 45
Wu-ts'ao Suan-king (*c.* 1st century), 499
Wylie, A., 42

Xenocrates (*c.* 350 B.C.), 524

Yard, 642
Year, 654, 661
Yenri (circle principle), 701
Young, J. W. A., 301

Zamberto (*c.* 1505), 338
Zangemeister, K., 56
Zarqâla, Ibn al- (*c.* 1050), 609, 616
Zeitschrift, 109
Zeno of Elea (*c.* 450 B.C.), 677
Zeno of Sidon (1st century B.C.), 281
Zenso. *See* Censo
Zero, 44, 69, 71, 74, 78; equating to, 431
Zeuthen, H. G. (*c.* 1900), 274, 296, 378
Ziegler, T., 665
Zuanne de Tonini da Coi (*c.* 1530), 460, 467
Zuñi Indians, 59
Zuzzeri, G. L., 670